



Cite this: *Phys. Chem. Chem. Phys.*,
2016, **18**, 12896

The salts of chloronium ions $R\text{-Cl}^+\text{-R}$ ($R = \text{CH}_3$ or CH_2Cl): formation, thermal stability, and interaction with chloromethanes†

Evgenii S. Stoyanov^{ab}

The interaction of $\text{CH}_3\text{Cl}/\text{CD}_3\text{Cl}$ or $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ with the carborane acid $\text{H}(\text{CHB}_{11}\text{Cl}_{11})$ (abbreviated as $\text{H}\{\text{Cl}_{11}\}$) generates the salts of $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ and $\text{CH}_2\text{Cl-}\{\text{Cl}_{11}\}$ and their deuterio analogs, respectively, which are analogs of the salts of asymmetric chloronium cations. Next, salts of chloronium cations $\text{CH}_3\text{-Cl}^+\text{-CH}_3$, $\text{ClCH}_2\text{-Cl}^+\text{-CH}_2\text{Cl}$, and $\text{ClCH}_2\text{-Cl}^+\text{-CH}_3$ and their deuterio analogs were obtained from the above compounds. The asymmetric $\text{ClCH}_2\text{-Cl}^+\text{-CH}_3$ cation was found to be unstable, and at ambient temperature, slowly disproportionated into symmetric cations $(\text{CH}_3)_2\text{Cl}^+$ and $(\text{CH}_2\text{Cl})_2\text{Cl}^+$. At a high temperature (150 °C), disproportionation was completed within 5 minutes, and the resulting cations further decomposed into $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ and $\text{CH}_2\text{Cl-}\{\text{Cl}_{11}\}$. The molecular fragment $\text{ClCH}_2\text{-}(X)$ of the compounds ($X = \{\text{Cl}_{11}\}$, $\text{-Cl}^+\text{-CH}_2\text{Cl}$, or $\text{-Cl}^+\text{-CH}_3$) is involved in exchange reactions with CH_2Cl_2 and CHCl_3 , converting into $\text{CH}_3\text{-}(X)$ with the formation of chloroform and CCl_4 , respectively.

Received 11th February 2016,
Accepted 4th April 2016

DOI: 10.1039/c6cp00946h

www.rsc.org/pccp

Halonium ions (R_2Hal^+) are well-recognized reactive intermediates in electrophilic chemistry.^{1,2} Their stability increases in the order $\text{Hal} = \text{F}, \text{Cl}, \text{Br}, \text{I}$. Recently, the evidence of the formation of the symmetrical fluoronium ions in solutions was obtained,^{3,4} and the nature of carbon-halogen bonds in the halonium ions was studied.⁵ Mostly stable dimethylbromonium and -iodonium salts are presently commercialized and widely used in chemical ionization mass spectroscopy (gas phase chemistry) as effective methylating⁶⁻¹² and protonating agents^{13,14} for a variety of compounds. Nevertheless, the chemistry of dialkylhalonium ions in condensed phases is virtually unknown. Recently, the salts of $(\text{CH}_3)_2\text{Cl}^+$ and $(\text{C}_2\text{H}_5)_2\text{Cl}^+$ with the exceptionally stable and inert toward reactive cations undecachlorocarborane ions, $\text{CHB}_{11}\text{Cl}_{11}^-$, were obtained and studied using X-ray and infrared (IR) spectroscopy.¹⁵ The solid salt $(\text{CH}_3)_2\text{Cl}^+(\text{CHB}_{11}\text{Cl}_{11}^-)$ is stable even at elevated temperatures and decomposes at 140 °C releasing $\text{CH}_3(\text{CHB}_{11}\text{Cl}_{11})$ and CH_3Cl . This is an important method for the isolation of pure $\text{CH}_3(\text{CHB}_{11}\text{Cl}_{11})$, which can be viewed as a neutral analogue of an asymmetric chloronium ion related to the dimethylchloronium ylide, CH_3ClCH_2 .^{16,17}

In the present work we obtained the salts of symmetric and asymmetric chloronium cations, $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ and

$\text{CH}_2\text{Cl-Cl}^+\text{-CH}_2\text{Cl}$, both protio and deuterio analogs, with the $\text{CHB}_{11}\text{Cl}_{11}^-$ counterion (hereafter abbreviated as $\{\text{Cl}_{11}^-\}$, Fig. S1, ESI†). We also explored their thermal stability and interaction with some simple chloromethanes. The carborane ion $\{\text{Cl}_{11}^-\}$ was chosen as a counterion for chloronium salts because of its exceptionally low basicity and high thermal stability, which ensure the stability of chloronium salts at room temperature and above.¹⁵

Experimental

Carborane acid $\text{H}\{\text{Cl}_{11}\}$ was prepared as previously described.¹⁸ The acid was sublimed at 150–160 °C under a pressure of 10^{-5} Torr on cold Si windows of a specially designed IR cell-reactor as a very thin translucent layer.¹⁹ The spectrum of the sublimed acid showed no traces of the H_3O^+ cation.²⁰ Dry gaseous chloromethanes (CH_3Cl , CH_2Cl_2 , and CHCl_3) were injected anaerobically into the IR cell. The values of their partial pressure were calculated as the ratio of absorption intensity to that of the standard spectrum recorded in the same cell filled with vapors at atmospheric pressure.

All procedures were performed in a Vacuum Atmospheres Corp. glovebox in the atmosphere of N_2 (O_2 and $\text{H}_2\text{O} < 0.5$ ppm). The IR spectra were recorded on a PerkinElmer Spectrum-100 spectrometer inside a dry box in the transmission mode ($400\text{--}4000\text{ cm}^{-1}$). The IR data were processed in the GRAMMS/A1 (7.00) software program from Thermo Fisher Scientific.

^a Vorozhtsov Institute of Organic Chemistry, Siberian Branch of Russian Academy of Sciences (SB RAS), Novosibirsk 630090, Russia. E-mail: evgenii@nioch.nsc.ru

^b Department of Natural Science, National Research University – Novosibirsk State University, Novosibirsk 630090, Russia

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c6cp00946h




Results and discussion

To explore the chemical processes involving chloronium ions by means of IR spectroscopy, their detailed IR spectra must be obtained and interpreted.

Chloronium salts and IR spectra

The $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ cation was obtained by introducing the CH_2Cl_2 vapors at a partial pressure ($P_{\text{CH}_2\text{Cl}_2}$) of 0.4 atm into an evacuated IR cell containing a film of $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ sublimed on their Si-windows. Reaction (1) of $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ formation proceeds very slowly, without a release of HCl.



After 40 minutes, the spectrum of the cation was isolated by subtracting the original spectrum of $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ multiplied by the scaling factor $f = 0.933$ (Fig. 1, red). This means that only 6.7% of the $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ salt (100–93.3%) has reacted.

If the partial pressure of the injected CH_2Cl_2 vapors was twofold higher (1 atm), then the reaction was accelerated significantly and completed within 6 minutes. The spectrum of the resultant $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ cation slightly differs from that of the previous sample (Fig. 1), indicating that the cation is sensitive to changes in the environment.

The $\text{CH}_3\text{-Cl}^+\text{-CD}_2\text{Cl}$ cation is formed when the vapors of CD_2Cl_2 are introduced into the IR cell with sublimed $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$.

The IR spectrum of the salt of this cation is shown in Fig. 2 (red; the gas phase was evacuated).

The $\text{CD}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ and $\text{CD}_3\text{-Cl}^+\text{-CD}_2\text{Cl}$ cations were formed when vapors of CH_2Cl_2 or CD_2Cl_2 respectively, were injected into the IR cell with a film of sublimed $\text{CD}_3\text{-}\{\text{Cl}_{11}\}$ salt on the Si windows. Their IR spectra are presented in Fig. S2 and S3 (ESI[†]) and in Table 1.

Interpretation of the IR spectra

IR spectra of the salts of the cations $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ (red), $\text{CD}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ (blue), and $\text{CH}_3\text{-Cl}^+\text{-CD}_2\text{Cl}$ (green) that are normalized to the intensity of the anion are shown in Fig. 3. The figure shows that in the frequency range of CH stretch vibrations, the sum of the spectra of the fragments $\text{ClH}_2\text{C-(Cl}^-\text{)}$ (blue) and $\text{CH}_3\text{-(Cl}^-\text{)}$ (green) matches the spectrum of the $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ cation. The same result was observed in the frequency range of the CH bend vibrations (Fig. 3, right). Interpretations of the IR spectra for all cations follow from the above data (Tables 1 and 2), taking into account that spectra of $\text{CH}_3\{\text{Cl}_{11}\}$ and $(\text{CH}_3\text{-Cl}^+\text{-CH}_3)\{\text{Cl}_{11}^-\}$ were interpreted earlier.¹⁵ It was important to determine the existence of specific absorption bands for each compound under study; we will use these bands (shown in Tables 1 and 2) as markers for the identification of these compounds in the mixtures. Intensity of the marked bands allowed us to estimate relative amounts of the compounds.




Fig. 1 Red: IR spectra of $(\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl})\{\text{Cl}_{11}^-\}$ formed at low (red) and high (blue) partial pressure of CH_2Cl_2 (0.4 and 1 atm respectively). Both spectra are normalized to intensity of the anion. The red spectrum was isolated after subtracting the spectrum of the unreacted $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$. Intense bands (marked by an asterisk) belong to the $\{\text{Cl}_{11}^-\}$ anion.





Fig. 2 IR spectra of the initial $\text{CH}_3\text{-}\{\text{Cl}_{11}\}$ salt (blue) and the salt of the $\text{CH}_3\text{-Cl}^+\text{-CD}_2\text{Cl}$ cation (red).



Table 1 IR frequencies of the methyl and bridged C–Cl⁺–C groups of the compounds analyzed, in comparison with the IR spectrum of CH₃Cl (the most characteristic or intense bands, used as markers to identify the cations in their mixtures, are underlined)

Compound	$\nu_{\text{as}}\text{CH}_3$	$\nu_{\text{s}}\text{CH}_3$	$\delta_{\text{as}}\text{CH}_3$	$\delta_{\text{s}}\text{CH}_3$	CH ₃ rock	$\nu_{\text{as}}(\text{CClC})$	$\nu_{\text{s}}(\text{CClC})$
CH ₃ Cl (gas) ²¹	3043	<u>3039</u>	2968	2879	1452	1355	732 (ν_{CCl})
CH ₃ {Cl ₁₁ } ¹⁵	3083	<u>3071</u>	2962	1409	1335	—	—
CH ₃ –Cl ⁺ –CH ₃ ¹⁵		<u>3068</u>	2957	1417	1324	<u>636</u>	
H ₃ C–Cl ⁺ –CD ₂ Cl	3064	<u>3054</u>	2947	1406	1330	934	632



**Fig. 3** IR spectra of salts of cations: CH₃–Cl⁺–CH₂Cl (red), CD₃–Cl⁺–CH₂Cl (blue) and CH₃–Cl⁺–CD₂Cl (green).**Table 2** IR frequencies of the methylene chloride and bridged C–Cl⁺–C groups of the compounds analyzed, in comparison with the IR spectrum of CH₂Cl₂ (the most characteristic or intense bands, used as markers to identify the cations in their mixtures, are underlined)

Compound	$\nu_{\text{as}}\text{CH}_2$	$\nu_{\text{s}}\text{CH}_2$	δCH_2 scissor	δCH_2 waggle	$\nu_{\text{as}}\text{CCl}_2/\nu_{\text{C-Cl}}$	$\nu_{\text{as}}(\text{CClC})$					
CH ₂ Cl ₂ (liquid) ²²	3045	2990	1424	1265vs	739	—					
ClH ₂ C–{Cl ₁₁ }	3079	<u>3063</u>	2978	1391	<u>1243vs</u>	793					
ClH ₂ C–Cl ⁺ –CH ₂ Cl IIa	3073	<u>3064</u>	3048	2980	1416	1324	<u>1284vs</u>	1248	770	635	
ClH ₂ C–Cl ⁺ –CH ₂ Cl IIb		<u>3044</u>	2964br	1406	1338	1283vs	1248	798	783	672	593
ClH ₂ C–Cl ⁺ –CH ₃	3044	2962	<u>1407</u>	1386	1261	<u>1248s</u>	1226	786	^a		
ClH ₂ C–Cl ⁺ –CD ₃	3044	2962	<u>1398</u>	1387	1263	<u>1251s</u>	1226	787	^a		


^a Not determined.

Interaction of mono- and dichloromethane with H{Cl₁₁} and chloronium cations

CH₃Cl interacts with the H{Cl₁₁} acid in two stages. At first, it is protonated with the release of HCl and the formation of CH₃{Cl₁₁} (eqn (2)):




Then, CH₃Cl binds to CH₃{Cl₁₁}, thus forming (CH₃)₂Cl⁺{Cl₁₁[–]} (eqn (3)). The formation of the resultant CH₃{Cl₁₁} can be observed by monitoring the intensity of the IR absorption of the released HCl (I_{HCl}). The amount of the resulting ion (CH₃)₂Cl⁺ can be traced from the intensity of its band $\nu_{\text{as}}(\text{CCl}^+\text{C})$ at 636 cm^{–1} (I_{636} ; Fig. S4, ESI[†]). Dependence of I_{636} on I_{HCl} was initially proportional (Fig. 4), which means that both reactions proceed at constant speeds. When the thickness of the layer of the formed surface products increases sufficiently to slow down the diffusion of CH₃Cl molecules to the acid, the rate of


**Fig. 4** Dependence of the intensity of band $\nu_{\text{as}}(\text{CClC})$ at 636 cm^{–1} of the (CH₃)₂Cl⁺ cation on the intensity of HCl absorption [reflects the dependence of the (CH₃)₂Cl⁺ formation on the CH₃{Cl₁₁} formation].

$\text{CH}_3^+\{\text{Cl}_{11}^-\}$ formation decreases and the slope increases (Fig. 4). The cation of the ultimate salt $(\text{CH}_3)_2\text{Cl}^+\{\text{Cl}_{11}^-\}$ does not show any further interaction with CH_3Cl .

Dichloromethane vapors (at a partial pressure of 0.65 atm) interacted with sublimed acid in the same way as chloromethane did. At the first stage, IR spectra registered the emergence of absorbance of the gaseous HCl and the surface $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ compound, which are formed in accordance with eqn (4):



The IR spectrum of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ showed the characteristic band of the C–Cl stretch at 793 cm^{-1} , and the absence of the bands of C–Cl⁺–C group vibrations in the frequency range $650\text{--}500\text{ cm}^{-1}$ (Fig. 5 and Table 2). Intensity of HCl absorption (I_{HCl}) was used to quantify the total amount of the formed $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$, whereas intensity of the band of the terminal C–Cl stretch (I_{CCl}) reflects the current amount of this compound. Dependence of I_{CCl} on I_{HCl} was linear for the first 5 hours of the reaction (Fig. 6); this result means that only $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ was formed. Then, the dependence started to drop (point 22 in Fig. 6), indicating the second stage of the reaction: involvement of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ in the formation of the $\text{ClH}_2\text{C}-\text{Cl}^+-\text{CH}_2\text{Cl}$ cation (eqn (5)).



Extrapolation of I_{CCl} from I_{HCl} to the moment of the reaction stoppage (at the 1380th minute, when CH_2Cl_2 was pumped out) yields $I_{\text{CCl}} = 0.084$ (arbitrary units; Fig. 6). This value corresponds to the amount of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ that was formed in accordance with the amount of HCl production (eqn (4)). Nevertheless, because $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ is further consumed (reaction (5)), I_{CCl} decreases to 0.040 arbitrary units (Fig. 6), meaning that 48% ($0.040/0.084$) of this compound survived, and 52% was converted to the chloronium ion.

An IR spectrum of $(\text{CH}_2\text{Cl})_2\text{Cl}^+$ was obtained by subtracting the spectrum of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ from the spectrum of the mixture of $(\text{CH}_2\text{Cl})_2\text{Cl}^+$ with the $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ up to complete compensation




Fig. 6 Dependence of I_{CCl} on I_{HCl} , pointing to $\text{ClH}_2\text{C}-\{\text{Cl}_{11}\}$ formation and its further consumption on the formation of the $\text{ClH}_2\text{C}-\text{Cl}^+-\text{CH}_2\text{Cl}$ cation.

of the bands $\nu\text{CCl} = 792\text{ cm}^{-1}$ and $\delta\text{CH}_2 = 1243\text{ cm}^{-1}$, which are specific to $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ (Fig. 5, blue; Table 2). The frequencies of the $(\text{CH}_2\text{Cl})_2\text{Cl}^+$ cation are very close to those of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ except for one intense band at 1284 cm^{-1} , which can be used as a marker of this cation (Table 2). Furthermore, we will denote $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ as compound **I** and the $(\text{CH}_2\text{Cl})_2\text{Cl}^+$ salt as compound **II**.

After stoppage of the reaction at minute 1380, an IR spectrum of the sample retained a strong absorption pattern of the unreacted acid, which constituted 70% of the original spectrum of the acid.

To continue reactions (4) and (5) at a higher speed, the CH_2Cl_2 vapors were reintroduced into our IR cell at higher partial pressure (1 atm). Under these conditions, the formation of compound **I** was completed after 8 minutes with full utilization of the acid and termination of the HCl release (point 30 in Fig. 7). An IR spectrum of this sample is shown in Fig. 8 (black).

Further interaction of compound **I** with CH_2Cl_2 and the formation of **II** is manifested in IR spectra as a decrease in the intensity of compound **I** and the upregulation of compound **II** with the appearance of isosbestic points (Fig. 8 and Fig. S5, ESI[†]).




Fig. 5 IR spectra of the $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ compound (red) and the salt of $\text{CH}_2\text{Cl}-\text{Cl}^+-\text{CH}_2\text{Cl}$ cation (blue). The spectra of unreacted $\text{H}\{\text{Cl}_{11}\}$ acid, gaseous HCl and CH_2Cl_2 were subtracted. The region of strong absorption of gaseous CH_2Cl_2 (C–Cl stretches), where compensation does not work, is shaded.






Fig. 7 Dependence of $\text{CH}_2\text{Cl}-(\text{Cl}_{11})$ formation (determined by means of intensity of its band $\nu_{\text{as}}\text{CH}_2$ at 3063 cm^{-1}) on the amount of released HCl (determined by means of intensity of the νHCl band).




Fig. 8 A change in intensity of the IR spectra of compounds $\text{CH}_2\text{Cl}-(\text{Cl}_{11})$ and $(\text{CH}_2\text{Cl}-\text{Cl}^+-\text{CH}_2\text{Cl})(\text{Cl}_{11}^-)$ as reaction (5) proceeds. The arrows indicate isosbestic points. The spectrum of gaseous CH_2Cl_2 was subtracted.

The spectrum of the resultant cation **II** differed from that of the same cation formed during a slow reaction of $\text{CH}_2\text{Cl}-(\text{Cl}_{11})$ with CH_2Cl_2 (Fig. 9 and Table 2). Thus, two isomers of $\text{CH}_2\text{Cl}-\text{Cl}^+-\text{CH}_2\text{Cl}$




Fig. 9 IR spectra of $(\text{CH}_2\text{Cl})_2\text{Cl}^+(\text{Cl}_{11})$, formed by slow (**IIa**, some hours, blue) and quick interaction of $\text{CH}_2\text{Cl}-(\text{Cl}_{11})$ with CH_2Cl_2 (**IIb**, some minutes, red). The region of strong absorption of gaseous CH_2Cl_2 (C–Cl stretches), where compensation does not work, is shaded.

are formed: compound **IIa** from the slowly proceeding reaction (5) and compound **IIb** from the rapid reaction (5). They differ in frequencies of stretches CH and CCl. Nonetheless, their bend CH vibrations are very similar and have one specific band δCH_2 (1284 cm^{-1}), which does not overlap with the bands of other types of cations. Therefore, the intensity of this band (I_{1284}) can be used for the estimation of the amount of the **IIa** + **IIb** mixture.

Fig. 10 shows the time dependence of the intensity of absorption bands of compounds **I** (I_{3063}) and **II** (I_{1284}) that reflects the formation of the corresponding cations. One can see that the formation of compound **II** passes through a maximum at point 34, when compound **I** disappears, and then decreases.

Simultaneously, the bands indicative of the cation $\text{CH}_3-\text{Cl}^+-\text{CH}_2\text{Cl}$ at 1261 and 1248 cm^{-1} made an appearance (Table 2) and increased in intensity. We will denote this cation as compound **III**. Its narrow band $\delta\text{CH}_2 = 1248\text{ cm}^{-1}$ (I_{1248}) does not overlap with the bands of other cations and can be used for the assessment of the relative amount of compound **III** (Fig. 10, green). Along with the spectrum of compound **III**, the absorption bands of the cation $(\text{CH}_3)_2\text{Cl}^+$ also develop (hereafter denoted as compound **IV**). Intensity of its single band at 1324 cm^{-1} (I_{1324}) was used to assess the formation of compound **IV**. Fig. 10 summarizes the sequence of the formation of compounds **I**, **II**, **III**, and **IV** and the relative amounts of each compound, but does not describe the quantitative relations among them.

Cations **III** and **IV** can be formed only if the reaction of **I** or **II** with CH_2Cl_2 is accompanied by the formation of gaseous products. The spectra of the gas phase revealed the band of H–C–Cl bend vibration of chloroform at 1219 cm^{-1} . The time dependence of its intensity shows that the formation of chloroform is associated with the formation of compounds **I** and **II** (Fig. 10).




Fig. 10 Kinetic curves of the formation of compounds **I–IV** and chloroform; these curves were constructed by means of the intensity values of their bands at 3068 (compound **I**), 1284 (**II**), 1248 (**III**), 1322 (**IV**), and 1220 cm^{-1} (chloroform). Curves **I–IV** do not quantitatively indicate the proportion of a cation in the mixture of compounds **I–IV**.








Fig. 11 Dependence of CHCl_3 formation (shown as intensity of the band at 1219 cm^{-1}) on $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ formation (indicated by intensity of its band at 1242 cm^{-1}) under the conditions when only compound **I** is formed. Cation **II** starts to form after point 18.

In the situation when only **I** is formed ($P_{\text{CH}_2\text{Cl}_2} = 0.65\text{ atm}$), the dependence of the chloroform formation (I_{1219}) on the formation of **I** (I_{1242}) increases both with an increase in the amount of compound **I** and with an increase in the contact time of **I** with dichloromethane (Fig. 11). Subtraction of the spectrum of compound **I** from the spectrum of the products (at point 18 in Fig. 11) leads to the manifestation of the spectrum of compound $\text{CH}_3\{\text{Cl}_{11}\}$ (Fig. S6, ESI[†]). Hence, CH_2Cl_2 reacts with $\text{CH}_2\text{Cl}\{\text{Cl}_{11}\}$ according to eqn (6):



With the rapid formation of $\text{CH}_2\text{Cl}\{\text{Cl}_{11}\}$ in reaction (4) at $P_{\text{CH}_2\text{Cl}_2} = 1\text{ atm}$, chloroform is formed symbatically (Fig. 12),






Fig. 12 The link between formation of $\text{CH}_2\text{Cl}-\{\text{Cl}_{11}\}$ (indicated by intensity of its band at 3063 cm^{-1}) and CHCl_3 (indicated by intensity of its band at 1219 cm^{-1}).

confirming reaction (6). In contrast, starting from point 32, chloroform formation stopped, whereas from point 34, it begins to decrease, when the amount of **II** passes through a maximum, and compound **I** is exhausted (Fig. 10). The subsequent expenditure of chloroform occurs simultaneously with the consumption of compound **II** and the increasing amount of **III**. This finding points to reaction (7):



(anions are omitted), which is suggestive of CCl_4 formation.

In the same way, compound **IV** can form:



The detection of CCl_4 was carried out as follows. After the completion of the reaction, the gaseous phase and all surface-adsorbed molecules were removed by pumping. The difference in IR spectra before and after the evacuation represented the spectrum of removed molecules. It consists of a strong absorption pattern of the original dichloromethane, the characteristic band at 1219 cm^{-1} of chloroform, and a weak band at 790 cm^{-1} , which may belong to the C-Cl stretch of CCl_4 . The latter frequency is lower than that of gaseous CCl_4 (795 cm^{-1}), but equals that of CCl_4 solvated with dichloromethane in its solutions (789 cm^{-1}). Therefore, the traces of the formed CCl_4 are adsorbed by superficial chloronium salts.

To sum up, we can conclude that CH_2Cl_2 and CHCl_3 interact with the molecular fragment $\text{CH}_2\text{Cl}-$, whose reactivity is highest in $\text{CH}_2\text{Cl}\{\text{Cl}_{11}\}$ and is consistently reduced in cations **II** and **III**. In general, these interactions can be expressed as:


Decreasing the reactivity of the molecular fragment $\text{CH}_2\text{Cl}-$ in compounds **I**, **II**, and **III** is correlated with its decreasing polarizability, which in turn is determined by the change in ionicity of the bonds in chloronium bridge $\text{C}-\text{Cl}^+-\text{C}$.

Ionicity/covalency of the bonds in the $\text{C}-\text{Cl}^+-\text{C}$ bridge of chloronium ions


The stretch frequencies of the methyl group, especially $\nu_{\text{as}}\text{CH}_3$, are sensitive to CH_3 polarization: the higher these frequencies, the stronger the CH_3 group is polarized and the higher is ionicity of the $\text{CH}_3-\text{Cl}(\text{X})$ bond. In compounds $\text{CH}_3-\{\text{Cl}_{11}\}$ and $\text{CH}_3-\text{Cl}^+-\text{CH}_3$, the CH stretches differ insignificantly (Table 1), pointing to almost the same ionicity of CH_3^+ bonding to $\{\text{Cl}^-\}$ and $\text{Cl}-\text{CH}_3$, respectively. Nonetheless, one would expect weaker ionicity (stronger covalency) of the bonds in $\text{CH}_3-\text{Cl}^+-\text{CH}_3$ than in $\text{CH}_3-\{\text{Cl}_{11}\}$. The crystal structure of the $(\text{CH}_3-\text{Cl}^+-\text{CH}_3)\{\text{Cl}_{11}^-\}$ salt shows (Fig. S7 in ESI[†]) that the chloronium Cl-atom forms six ionic bonds with Cl-atoms of the four $\{\text{Cl}_{11}^-\}$ anions of its



environment. This situation favors an increase in the ionicity of C–Cl⁺–C bonds and contributes to the convergence of polarizability of CH₃ groups in CH₃–{Cl₁₁[–]} and (CH₃–Cl⁺–CH₃){Cl₁₁[–]} salts. If we now examine the salt (CH₃–Cl⁺–CD₂Cl){Cl₁₁[–]}, one can see that CH stretches significantly decrease (Table 1). It is evident that this salt is amorphous and has a disordered structure that may reduce the number of linkages between the chloronium Cl atom and neighboring {Cl₁₁[–]} anions, thus increasing the covalency of the C–Cl⁺–C group. In any case, the ionicity of the CH₃–Cl(–X) bond decreases in the following order:



Just as the methyl group in salts CH₃–{Cl₁₁[–]} and (CH₃–Cl⁺–CH₃){Cl₁₁[–]}, the CH₂Cl group in salts ClH₂C–{Cl₁₁[–]} and **IIa** shows identical CH stretch frequencies (Table 2). Probably, in the case of the slow formation (hours) of the (ClH₂C–Cl⁺–CH₂Cl){Cl₁₁[–]} salt (**IIa**), its structure is the densest, close to that of the single crystal with a maximal number of interactions between the chloronium Cl atom and atoms of the nearest {Cl₁₁[–]} anions. In the case of quick formation (minutes), the (ClH₂C–Cl⁺–CH₂Cl){Cl₁₁[–]} salt is amorphous with a loose structure and fewer contacts between the chloronium Cl atom and Cl atoms of the neighboring anions. This situation increases the covalency of the bonds in the C–Cl⁺–C bridge and decreases the polarizability of CH₂Cl groups and their CH stretches. We named this salt “isomer **IIb**”. Polarizability and CH frequencies of the CH₂Cl group of **IIb** coincide with those of the salt (ClH₂C–Cl⁺–CH₃){Cl₁₁[–]}. Thus, ionicity of the ClH₂C–Cl(–X) bond decreases in the following order:



in accordance with decreasing frequencies of CH stretches, which indicate the polarization of the ClH₂C group. Isomers **IIa** and **IIb** differ in frequencies of stretch vibrations of C–Cl⁺–C



bridges (Table 2); this observation confirms that this is the starting point of their differences.

The proposed model seems to be a reasonable explanation for the existence of **IIa** and **IIb** isomers when they are formed slowly (within hours) or quickly (in minutes), respectively.

Thermal stability of chloronium cations

Ambient temperature. The salt of the symmetric cation (CH₃)₂Cl⁺ can be stored indefinitely (weeks or months) without any changes in the IR spectrum. The spectra of the salts of asymmetric cations are time dependent at ambient temperature: exposure of freshly prepared salt (CD₃–Cl⁺–CH₂Cl){Cl₁₁[–]} to vacuum for one day results in disappearance of the bands of its cation (Fig. 13, blue). They are replaced with the bands of cation **IIa**: CH₂Cl–Cl⁺–CH₂Cl [labeled (2) in Fig. 13, red]. Weak bands of cations CD₂Cl–Cl⁺–CD₂Cl and (CH₃)₂Cl⁺ also appeared (labeled 2 and 3 respectively), as did the traces of HCl absorption.

In the range of CD stretch vibrations, the band $\nu_{\text{as}}\text{CD}_3 = 2297 \text{ cm}^{-1}$ of the cation CD₃–Cl⁺–CH₂Cl was replaced with the intense $\nu_{\text{as}}\text{CD}_3$ band at 2307 cm^{-1} of the cation CD₃–Cl⁺–CD₃. Thus, the CD₃–Cl⁺–CH₂Cl cation in the solid phase at ambient temperature is unstable, and during the day, disproportionates into symmetrical cations, predominantly according to eqn (9) and to a slight extent, according to eqn (10)



The disproportionation looks as if at the chlorine atom, there is an intermolecular exchange of groups CD₃– and –CH₂Cl that is difficult to imagine for the solid phase. Intermediates with the molecular fragments CHDCl– and CH₂D– were not detected by IR spectroscopy.

Increased temperature. Heating of the salt (CH₃–Cl⁺–CH₂Cl){Cl₁₁[–]} to –100 °C for 5 min in a sealed cell led to disappearance of its spectrum (Fig. 14 and Fig. S8, ESI,† blue) and to the emergence of overlapping spectra of the symmetric cation (CH₃)₂Cl⁺ and isomer (CH₂Cl)₂Cl⁺, **IIa** (with the marked




Fig. 13 IR spectra of the freshly prepared salt (CD₃–Cl⁺–CH₂Cl){Cl₁₁[–]} (blue) and after one or three days of storage (red). The bands of the cations (CH₂Cl)₂Cl⁺, CD₂Cl–Cl⁺–CD₂Cl and (CH₃)₂Cl⁺ are marked respectively as (1), (2) and (3).









Fig. 14 IR spectra in the frequency region of CH bend vibrations of the salt $(\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl})\{\text{Cl}_{11}^-\}$ before (blue) and after heating for 5 minutes at 100 °C (red) and 150 °C (green). The most characteristic bands of the formed compounds are marked with (a) for $(\text{CH}_3)_2\text{Cl}^+$, (b) for $(\text{CH}_2\text{Cl})_2\text{Cl}^+$ (IIa) and (c) for $\text{CH}_3\text{-Cl}_{11}$.

bands a and b, respectively, Fig. 14 and Fig. S8, ESI,† red). The absorption bands of gaseous HCl did not appear.


Therefore, at 100 °C, the asymmetric cation $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ quickly disproportionates into the more stable symmetric cations:



Further heating of the sample for 5 minutes at 150 °C led to an equal (threefold) reduction in intensity of the bands at 1324 and 1284 cm^{-1} of the cations $(\text{CH}_3)_2\text{Cl}^+$ and $(\text{CH}_2\text{Cl})_2\text{Cl}^+$, respectively, and appearance and an increase in intensity of the bands of $\text{CH}_3\text{-Cl}_{11}$ (1335 cm^{-1} ; Fig. 14, green) and $\text{ClH}_2\text{C-Cl}_{11}$ ($\nu_{\text{CCl}_{\text{term}}} = 793 \text{ cm}^{-1}$). Additionally, bands with a rotational structure at 1270 and 761 cm^{-1} of gaseous CH_2Cl_2 were observed. Consequently, an increase in temperature facilitates the decomposition of chloronium ions according to eqn (12) and (13); this change should lead to increasing intensity of the IR spectrum of released dichloromethane.



Longer heating of the sample at 150 °C did not enhance the spectrum of CH_2Cl_2 . It remained virtually unchanged. In contrast, the bands of CHCl_3 appeared (1219 and 772 cm^{-1}) and were enhanced with heating and a very weak absorption pattern of gaseous HCl developed. The solid-phase spectrum shows increased intensity of the $\text{CH}_3\text{-Cl}_{11}$ band at 1335 cm^{-1} and disappearance of the adsorption of $\text{ClH}_2\text{C-Cl}_{11}$. Taken together, our data indicate that the CHCl_3 formation is caused by the interaction of $\text{ClH}_2\text{C-Cl}_{11}$ with dichloromethane:




To determine the reason for the formation of trace amounts of HCl, additional studies are needed.

Conclusions

The salts of symmetric methylchloronium cations, $(\text{CH}_3)_2\text{Cl}^+\{\text{Cl}_{11}^-\}$ and $(\text{CH}_2\text{Cl})_2\text{Cl}^+\{\text{Cl}_{11}^-\}$, and their neutral analogs $\text{CH}_3\text{-Cl}_{11}$ and $\text{CH}_2\text{Cl-Cl}_{11}$, are stable at ambient and increased temperatures. Nevertheless, the asymmetric cation $\text{ClCH}_2\text{-Cl}^+\text{-CH}_3$ even at ambient temperature disproportionates into symmetrical $(\text{CH}_3)_2\text{Cl}^+$ and $(\text{CH}_2\text{Cl})_2\text{Cl}^+$. The molecular fragment $\text{CH}_2\text{Cl-}$ of the chloronium ions enters exchange reactions with CH_2Cl_2 and CHCl_3 with increasing reactivity in the order $\text{ClCH}_2\text{-Cl}^+\text{-CH}_3$, $\text{ClCH}_2\text{-Cl}^+\text{-CH}_2\text{Cl}$, and $\text{CH}_2\text{Cl-Cl}_{11}$, yielding more stable and less reactive $(\text{CH}_3)_2\text{Cl}^+\{\text{Cl}_{11}^-\}$ and $\text{CH}_3\{\text{Cl}_{11}\}$ compounds.

If we take into account the reactivity of $\text{H}\{\text{Cl}_{11}\}$ acid or its chloronium salts with CH_2Cl_2 , and the simultaneous disproportionation of the $\text{CH}_3\text{-Cl}^+\text{-CH}_2\text{Cl}$ cation (eqn (11); initiated by the elevated temperature), then the end products of interactions at ambient, or slightly elevated temperature, are the chemically inert $\text{CH}_3\text{-Cl}^+\text{-CH}_3$ and CCl_4 :



Reactions (9)–(13) (as determined in this study) proceed with the formation of a trace amount of HCl, thus indicating the negligible presence of parallel reactions. Under the other conditions, they may appear to be significant and interesting for future research.

Acknowledgements

This work was supported by grant # 16-13-10151 from the Russian Science Foundation and by the Ministry of Education and Science of the Russian Federation within the Project of the joint Laboratories of the Siberian Branch of the Russian Academy of Sciences and National Research Universities. The author thanks Irina S. Stoyanova for providing the carborane acids and technical support.

References

- 1 G. A. Olah, *Halonium Ions*, Wiley, New York, 1975.
- 2 G. A. Olah, K. K. Laali, Q. Wang and G. K. S. Prakash, *Onium Ions*, Wiley, New York, 1998, ch. 6.
- 3 M. D. Struble, M. T. Scerba, M. A. Siegler and T. Lectka, *Science*, 2013, **340**, 57.
- 4 M. D. Struble, M. G. Holl, M. T. Scerba, M. A. Siegler and T. Lectka, *J. Am. Chem. Soc.*, 2015, **137**, 11476.
- 5 R. Kalescky, W. Zou, E. Kraka and D. Cremer, *J. Phys. Chem. A*, 2014, **118**, 1948.
- 6 G. A. Olah and J. R. DeMember, *J. Am. Chem. Soc.*, 1970, **92**, 718.
- 7 G. A. Olah and J. R. DeMember, *J. Am. Chem. Soc.*, 1970, **92**, 2562.



- 8 G. A. Olah and Y. K. Mo, *J. Am. Chem. Soc.*, 1974, **96**, 3560.
- 9 H. W. Zappey, T. Drewello, S. Ingemann and N. M. M. Nibbering, *Int. J. Mass Spectrom. Ion Processes*, 1992, **115**, 193.
- 10 T. Partanen and P. Vainiotalo, *Rapid Commun. Mass Spectrom.*, 1997, **11**, 881.
- 11 M. A. Freitas, R. A. J. O'Hair and T. D. Williams, *J. Org. Chem.*, 1997, **62**, 6112.
- 12 D. K. Sen Sharma and P. Kebarle, *J. Am. Chem. Soc.*, 1982, **104**, 19.
- 13 J. L. Beachamp, D. Holtz, S. D. Woodgate and S. L. Patt, *J. Am. Chem. Soc.*, 1972, **74**, 2798.
- 14 G. Bouchoux, F. Caunan, D. Leblanc, M. T. Nguyen and J. Y. Salpin, *ChemPhysChem*, 2001, **10**, 604.
- 15 E. S. Stoyanov, I. V. Stoyanova, F. S. Tham and C. A. Reed, *J. Am. Chem. Soc.*, 2010, **132**, 4062.
- 16 L. A. Noronha, T. J. L. Judson, J. F. Dias, L. S. Santos, M. N. Eberlin and C. J. A. Mota, *J. Org. Chem.*, 2006, **71**, 2625.
- 17 A. Jubert, N. Okulik, M. C. Michelini and C. J. A. Mota, *J. Phys. Chem. A*, 2008, **112**, 11468.
- 18 M. Juhasz, S. Hoffmann, E. S. Stoyanov, K. Kim and C. A. Reed, *Angew. Chem., Int. Ed.*, 2004, **43**, 5352.
- 19 E. S. Stoyanov, I. V. Stoyanova and C. A. Reed, *J. Am. Chem. Soc.*, 2011, **133**, 8452.
- 20 E. S. Stoyanov, K.-C. Kim and C. A. Reed, *J. Am. Chem. Soc.*, 2006, **128**, 1948.
- 21 L. M. Sverdlov, M. A. Kovner and E. P. Krainov, *Vibrational Spectra of Polyatomic Molecules*, Nauka, Moscow, 1970, p. 340.
- 22 T. Shimanouchi and I. Suzuki, *J. Mol. Spectrosc.*, 1962, **8**, 222.

