

CrossMark
click for updatesCite this: *Anal. Methods*, 2016, 8, 4354

The soybean lipoxygenase-fluorescein reaction may be used to assess antioxidant capacity of phytochemicals and serum

M. Soccio, M. N. Laus, M. Alfarano and D. Pastore*

Evaluation of the putative beneficial effects of food antioxidants by *in vitro* assays of the antioxidant capacity (AC) of food extracts sometimes appears questionable. A more realistic evaluation of antioxidant effectiveness may derive from the integration of *in vitro* assays with *ex vivo* assays of blood/serum/plasma AC after food intake. The aim of this study was to develop the novel lipoxygenase-fluorescein (LOX-FL) method to assay the AC of both food extracts and serum/plasma. This method was applied on both extracts from the antioxidant-rich dietary cereal supplement, Lisosan G, and serum (seven subjects) for 240 min after the intake of 20 g of supplement. The widely used ORAC and TEAC methods were used for comparison. The new LOX-FL method is based on the reaction between the soybean LOX-1 isoenzyme and FL in the presence of linoleic acid, which undergoes hydroperoxidation, thus generating physiological reactive species, including LOO[•], LO[•], HO[•] and ¹O₂, which are able to quench FL. The quenching rate is slowed by antioxidants; this inhibition may be calibrated in terms of Trolox equivalents to assess the AC. Interestingly, the LOX-FL method discriminated the *in vitro* AC of four different Lisosan G extracts similarly to the ORAC and TEAC methods. In contrast, only the LOX-FL method was able to highlight a general increase in the serum AC (up to 40% after 30 min) after Lisosan G intake, thus confirming its physiological effectiveness by *ex vivo* serum assay. This quality of the LOX-FL method probably comes from the ability to highlight simultaneously different antioxidant mechanisms and to effectively show synergy among food phenols and endogenous serum antioxidants.

Received 6th April 2016

Accepted 3rd May 2016

DOI: 10.1039/c6ay01002d

www.rsc.org/methods

Introduction

A direct measurement of the antioxidant capacity (AC) of antioxidant-rich foodstuffs, in particular, fruits, vegetables, whole cereal grains, wine, tea and chocolate, has been the subject of a very large number of articles in the last few years. The AC has been evaluated by a lot of different *in vitro* systems (ORAC, TEAC, TRAP, FRAP, and DPPH).^{1–3} The interest in these measurements derives from the idea that food AC is related to prevention of diseases. Consistently, recent population studies found significant associations between the AC of the food consumed (obtained from food-frequency questionnaires) and healthy effects, including positive effects on chronic diseases and several tumours as well as a lower risk of heart failure and strokes.^{4–7} On the other hand, methods for AC measurement show many technical and conceptual limitations. The technical problems are mostly related to the chemistry behind the different AC assays making it difficult to compare AC values obtained using different methods and different experimental conditions.² From a conceptual point of view, an important limitation is that the values indicating *in vitro* AC have not been

demonstrated to be relevant to the biological effects of specific bioactive compounds.⁸ Moreover, these assays do not measure bioavailability, *in vivo* stability, retention of antioxidants by tissues, and reactivity *in situ*. Finally, there is growing evidence that the metabolic pathways associated with the prevention or amelioration of chronic diseases by bioactive compounds are often dependent on enzyme/protein and/or gene expression regulation rather than a true antioxidant effect.^{9,10} In the light of these considerations, suggesting that the AC values of foods have no relevance to the effects of specific bioactive compounds on human health, the U.S. Department of Agriculture (USDA) removed its ORAC database for selected foods from its nutrient data laboratory website in 2012.¹¹ However, this choice was not fully shared and remains to a certain extent questionable.¹²

A possible different, more physiological approach may be to move the focus from the analysis of the AC of foods to the analysis of the AC of serum/plasma after food intake. This measure may take into account bioavailability and metabolism, so giving integrated information on a true effect on blood antioxidant status, which is beyond the original AC of the ingested food. In addition, the assessment of the antioxidant status of blood may be *per se* an important target. In fact, blood plays a central role in the homeostasis of the cellular redox status by carrying and releasing antioxidants in the body;

Dipartimento di Scienze Agrarie, degli Alimenti e dell'Ambiente, Università degli Studi di Foggia, Via Napoli, 25, 71122, Foggia, Italy. E-mail: donato.pastore@unifg.it



moreover, the maintenance of the blood physiological antioxidant status may preserve the endothelial function, which is thought to be an essential determinant of healthy aging.¹³ Given the simplicity of AC assays and the easy access to human blood, AC measurements of serum/plasma flourished in the last years, generating a large amount of data.¹⁴ Unfortunately, in short-term studies (minutes or hours after ingestion), foods very rich in antioxidants often induced a limited AC increase.¹⁵ These results may be, at least in part, dependent on some weakness in the analytical methods. For example, TEAC performs less well than other methods.¹⁴ Therefore, in this paper, we describe the use of the novel well-performing lipoxygenase (LOX)-fluorescein (FL) method. It essentially derives from the LOX/4-nitroso-*N,N*-dimethylaniline (LOX/RNO) method, which is advisable for use with food extracts,^{16–18} but it shows low sensitivity towards serum (unpublished data). Similarly to the LOX/RNO method, the LOX-FL assay utilizes the soybean LOX-1 reaction with linoleate to generate linoleate hydroperoxide (LOOH) and several physiologically reactive species, mainly the linoleate alkoxy (LO[•]) and peroxy (LOO[•]) derivatives, HO[•] and ¹O₂,¹⁹ which are able to cause the quenching of FL.

Here, we compared the LOX-FL method with the widely used ORAC and TEAC methods to measure the AC of food extracts from the antioxidant-rich dietary wheat grain supplement Lisosan G and of serum after Lisosan G intake. Lisosan G was chosen in the light of its well-documented bioactivity, including its protective role in carbon tetrachloride- and cisplatin-induced toxicity in rat tissues,²⁰ the induction of antioxidant and detoxifying systems in rat hepatocytes,²¹ the antimutagenic and antioxidant activity in *Saccharomyces cerevisiae*²² and the improvement of human endothelial progenitor cell (EPC) function.²³

Interestingly, the LOX-FL method, similarly to the ORAC and TEAC methods, was found to be very suitable for assessing the AC of hydrophilic and phenolic extracts of Lisosan G and was able to discriminate among the different extracts. On the other hand, only the new method was able to evaluate strong synergistic effects among phenols and serum and to effectively highlight an increase in serum AC after Lisosan G intake. So, the LOX-FL method may be successfully applied to both *in vitro* analysis of food extracts and *ex vivo* analysis of serum.

Experimental section

Chemicals

All reagents used were of the highest commercially available purity and were purchased from SIGMA Chemical Co. (St. Louis, MO, USA).

The chemicals, 3',6'-dihydroxyspiro[isobenzofuran-1[3*H*],9'[9*H*]-xanthen]-3-one (fluorescein, FL), 2,2'-azobis(2-methylpropionamide) dihydrochloride (AAPH) and (±)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid (Trolox) were dissolved in different media, depending on the assay used to determine the AC. The chemicals, 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), ascorbate, bilirubin, urate, and albumin were dissolved in deionized water. An ammonium sulfate suspension of soybean LOX type V (LOX-1

isoenzyme, E.C. 1.13.11.12) was used, properly diluted with 100 mM Na-borate buffer, pH 9.0. The linoleate solution was prepared and assayed as reported in Pastore *et al.*²⁴

Antioxidant-enriched food

Lisosan G, an antioxidant-enriched wheat grain preparation produced by Agrisan Company (Larciano, PT, Italy), was kindly provided by Dr Vincenzo Longo from "Istituto di Biologia e Biotecnologia Agraria" (CNR, Pisa, Italy). It is registered by the Italian Ministry of Health as a dietary supplement. In the production process, the bran and germ are separated, collected, mixed with water and inoculated with selected microbial starter cultures, typically consisting of a mix of lactobacillus and natural yeast strains. Finally, once the product is fermented, it is dried and packaged.²¹

Oxygen uptake catalyzed by soybean LOX-1

Oxygen uptake was monitored at 37 °C by means of a 5300A YSI (Yellow Spring, OH, USA) oxygraph equipped with a 5331 YSI Clark-type electrode. The reaction medium (2 mL) consisted of 100 mM Na-borate buffer, pH 9.0, 400 μM Na-linoleate, and 1 μL Tween 20 per μmol linoleate; the reaction was started by adding 0.5 enzymatic units (EU) of soybean LOX-1.

FL quenching catalyzed by soybean LOX-1

FL quenching in the course of linoleate hydroperoxidation by soybean LOX-1 was fluorometrically monitored by measuring the FL fluorescence decrease (λ_{ex} 485 nm; λ_{em} 515 nm) at 37 °C using a Perkin Elmer LS 55 fluorescence spectrometer. The reaction mixture (2 mL) contained 100 mM Na-borate buffer, pH 9.0, 400 μM Na-linoleate, 1 μL Tween 20 per μmol linoleate and 6.3 nM FL; the reaction was started by adding 0.5 EU of soybean LOX-1. The rate of the reaction, expressed as Δ Arbitrary Units of Fluorescence (AUF)/s, was calculated as the highest slope to the experimental curve.

As shown in Fig. 1C, FL quenching was monitored spectrophotometrically at 485 nm and 37 °C using the reaction medium reported above, with the exception of the FL concentration that, in this case, ranged from 1 to 12.5 μM. A Perkin Elmer LAMBDA 45 spectrophotometer was used. Rates of reaction were expressed as $\Delta A_{485}/\text{min}$.

Extraction and assay of antioxidant compounds

Extraction of hydrophilic, lipophilic and phenolic compounds from Lisosan G. Hydrophilic extracts were prepared as described in Laus *et al.*,¹⁷ by extracting Lisosan G with deionized water at a (w/v) ratio equal to 1 g/5 mL in an ice-water bath for 1 h. Lipophilic compounds were extracted as in Laus *et al.*¹⁷ and reconstituted in ethanol. Insoluble bound (IB) phenolic and free soluble (FS) phenolic compounds from Lisosan G were extracted as reported in Laus *et al.*¹⁷ and reconstituted in water. All extracts were used daily.

Determination of total phenols and flavonoids. Total phenols were determined using the colorimetric Folin-Ciocalteu method and quantified by means of a proper calibration






Fig. 1 Kinetics of the FL quenching catalyzed by soybean LOX-1. The LOX-1-dependent FL quenching (A and B, trace a) was fluorometrically monitored (λ_{ex} 485 nm; λ_{em} 515 nm) at 37 °C in 2 mL 100 mM Na-borate buffer, pH 9.0, containing 6.3 nM FL, 400 μM linoleate and 1 μL Tween 20 per μmol linoleate (see Experimental section). In (A) the reaction was started by adding 0.05 enzymatic units (EU) soybean LOX-1 and monitored by recording spectra every 90 s. In (B) the reaction was monitored as a time course and simultaneously oxygen uptake (dotted line, curve b) was oxygraphically measured (see Experimental section). At the time indicated by the arrow, 0.5 EU of LOX was added. The rate of the FL quenching (v) and the lag phase are also indicated. In (C), measurements were carried out spectrophotometrically at 485 nm by using the same reaction medium described in (B), but in the presence of the reported FL concentrations. The rates of FL quenching, expressed as $\Delta A_{485\text{ nm}}/\text{min}$, are reported as Michaelis-Menten and Lineweaver-Burk (inset) plots.

on a gallic acid basis, as reported in Laus *et al.*¹⁷ Flavonoids were determined as reported in Laus *et al.*¹⁷ and quantified by means of a proper calibration curve on a catechin basis. The protein content in the hydrophilic extract from Lisosan G was evaluated as reported in Laus *et al.*,¹⁷ using bovine serum albumin as a standard.

Sera collection

Sera were obtained from seven volunteers (4 men and 3 women with ages ranging from 24 to 33 years). Each volunteer provided informed written consent to participate in this study in accordance with the Declaration of Helsinki. The research was performed in compliance with relevant Italian laws and institutional ethical policies. In particular, the experiment was approved by both the Board of the SAFE Department of the University of Foggia, Italy, and the Scientific Committee of the Regional Technological District DARE-Puglia, Italy.

For 2 days before analysis, the volunteers followed a low antioxidant diet, avoiding phenolic-rich foods, namely all fresh fruits and vegetables and derived products (tea, coffee, fruit juices, wine, and chocolate). The working window was free from interference by different variables (*e.g.* physical exercise and energy expenditure).

After an overnight fast, the volunteers ingested 20 g of Lisosan G suspended in 500 mL of water. This is an amount able to induce a serum AC increase, as verified in preliminary experiments. Venous samples at baseline (T_0) and at 30, 60, 90, 120 and 240 min after Lisosan G consumption were collected in vacutainers for serum collection. Sera were obtained by centrifugation (3000 \times g, 5 min), distributed in portions and frozen at -80 °C until analysis.

AC determination by means of the LOX-FL, ORAC and TEAC methods

LOX-FL method. The LOX-FL reaction was carried out at 37 °C as already described for the LOX-1-dependent FL quenching, in both the absence (control) and the presence of the sample (extract or serum or pure/standard antioxidant). The inhibition of the LOX-FL reaction was determined by calculating the (%) decrease of the rate of the FL quenching measured in the presence of the sample (or standard antioxidant) (v_a) with respect to the control (v_c), according to the equation: inhibition (% of the control) = $[1 - (v_a/v_c)] \times 100$. The AC was calculated by means of a dose-response curve obtained with Trolox by plotting the (%) decrease of the rate of FL quenching as a function of the standard antioxidant concentration.

Since the rate of the LOX-FL reaction is affected by ethanol, when the lipophilic extract of Lisosan G, reconstituted in ethanol, was evaluated, a constant volume of ethanol was maintained in the assay mixture.

ORAC method. The ORAC protocol described in Ou *et al.*²⁵ was applied with some modifications by using a CLARIOstar – BMG Labtech microplate reader. Every working well of a 96-well plate contained the assay mixture (final volume 0.2 mL), consisting of 75 mM Na-phosphate buffer, pH 7.4, 10 nM FL and an



appropriate volume of sample (extract or serum or pure/standard antioxidant). The reaction was started by adding 40 mM AAPH. The FL fluorescence intensity decay was monitored at 37 °C at excitation and emission wavelengths of 483 nm (bandwidth 14 nm) and 530 nm (bandwidth 30 nm), respectively. Since lipophilic antioxidants extracted from Lisosan G were reconstituted in ethanol, in this case a constant volume of ethanol was maintained in the assay mixture.

TEAC method. The TEAC assay described in Re *et al.*²⁶ was applied with slight modifications as reported in Laus *et al.*²⁷ The ABTS^{•+} solution was diluted with 5 mM Na-phosphate buffer, pH 7.4 (or ethanol for lipophilic extract of Lisosan G).

For all three AC methods, measurements were carried out in triplicate for three different amounts of sample and the AC was quantified by means of a dose–response curve obtained with Trolox.¹⁸

Results and discussion

Characterization and setting of the LOX-FL reaction as new tool to evaluate AC

A first characterization of the LOX-1-dependent FL quenching is reported in Fig. 1. The addition of the enzyme to the reaction mixture, containing an excess of linoleate (400 μM) and a limited amount of oxygen (200 μM at 37 °C, temperature of measurement), caused the disappearance of both excitation and emission FL fluorescence spectra in the vis region (Fig. 1A). The time course of this reaction can be easily monitored by continuously measuring the FL fluorescence decrease (Fig. 1B). In this experiment, the LOX-1-mediated FL quenching (curve a) and oxygen uptake (dotted curve, b) were simultaneously monitored. It should be noted that LOX-1 addition to the reaction mixture caused rapid oxygen consumption due to linoleate dioxygenation, whereas no simultaneous FL fluorescence change was observed. The FL quenching started about 20–25 s after enzyme addition, that is, only when a very low oxygen concentration, approaching anaerobiosis, was reached in the test sample. Therefore, the observed lag phase represents the time necessary to consume oxygen during the primary aerobic LOX-1 reaction of linoleate peroxidation.^{18,19} On the contrary, FL quenching is essentially a secondary anaerobic reaction.

A study of the kinetics of the LOX-1-dependent FL quenching was also carried out; to do this, since the high FL concentration necessary in this experiment is incompatible with fluorescence measurements, the reaction was monitored spectrophotometrically at 485 nm. By plotting the rate of FL quenching vs the FL concentration, a hyperbolic relationship was obtained (Fig. 1C). Data were also plotted as Lineweaver–Burk (inset), Hanes, Eadie–Hofstee and Eadie–Scatchard plots (not shown). All plots were linear, thus confirming the hyperbolic kinetics, with K_m and V_{max} equal to 1.53 ± 0.08 (SD) μM and 0.28 ± 0.01 (SD) $\Delta A_{485 \text{ nm}}/\text{min}$, respectively, calculated by means of GRAFIT 5.0 software.

The FL sensitivity to some relevant physiological reactive species, some of which were generated during the course of LOX-1-mediated anaerobic reactions,^{18,19} was investigated (not

shown). In particular, FL was insensitive to H₂O₂ as well as to O₂^{•−} and LOOH, generated *via* xanthine/xanthine oxidase²⁸ and LOX/linoleate systems, respectively. On the contrary, FL was sensitive to LO[•] and HO[•] generated *via* the Fenton reaction from LOOH or H₂O₂, respectively, in the presence of CoCl₂.²⁹ Similarly, FL quenching was also observed in the presence of ¹O₂ generated *via* the Na₂MoO₄/H₂O₂ system.³⁰

In another set of experiments, the sensitivity of the FL quenching reaction to antioxidants was evaluated. Firstly, the sensitivity of FL quenching to the standard antioxidant, Trolox, an α -tocopherol analogue with enhanced water solubility, was tested. In Fig. 2A, the experimental curves of the FL quenching reactions carried out in both the absence (control, trace a) and presence of Trolox at different concentrations (10, 15 and 20 μM; traces b, c and d, respectively) are shown. Trolox was found to inhibit FL quenching by causing a decrease in the reaction rate consistent with its antiradical activity. In the inset of Fig. 2A, Trolox-dependent inhibition, expressed as (%) decrease with respect to the control trace, is reported as a function of the standard antioxidant concentration: a linear dependence of the inhibition from about 15% to 45% on the Trolox concentration, ranging between 5 and 20 μM, was obtained (see the equation in the figure caption).

Interestingly, kinetic analysis highlights the competitive inhibition mechanism of Trolox on the LOX-1-mediated FL quenching with a K_i value equal to about 5 μM (not shown). This result implies that the antioxidant/oxidant interaction occurs at the level of the enzyme macromolecule rather than in the bulk phase of the solution.

Since the LOX-1-mediated FL quenching reaction is sensitive to reactive species and also shows inhibition by Trolox, which may be easily quantified, we adopted this reaction as a new tool to measure the AC. Interestingly, this new LOX-FL method shows several advantages. The hyperbolic dependence of the reaction rate on the FL concentration indicates that the reaction cannot occur in the bulk phase of the reaction mixture, but at the LOX surface, in particular, at the level of a definite number of FL binding sites, generating a ternary enzyme-radical-FL complex. Therefore, the effect of an antioxidant should be considered to be more physiological, as it refers to a reaction taking place within a biological macromolecule. In addition, the LOX-FL assay evaluates the oxidant–antioxidant competition at low oxygen concentration, which often occurs in human cells. Moreover, the LOX-FL method may evaluate antioxidants acting in a different manner: as scavengers of different physiological radicals, including LO[•] and LOO[•] radical derivatives of linoleate, HO[•] as well as ¹O₂, but also as reducing agents or chelators of the non-heme Fe³⁺ necessary for LOX catalysis or as direct inhibitors of the peroxidation reaction catalysed by LOX, thus providing a comprehensive AC evaluation.¹⁸

A first application of the new LOX-FL method was carried out on the dietary supplement, Lisosan G. The capability of the LOX-FL assay to evaluate the AC in FS and IB phenols, hydrophilic and lipophilic extracts of Lisosan G, was tested. In particular, in Fig. 2B, typical experimental traces of FL quenching in the presence of FS phenols of Lisosan G are shown. Small amounts of extract (obtained from 2.5 to 10 mg






Fig. 2 Inhibition of the LOX-1-dependent FL quenching by Trolox (A), FS phenols from Lisosan G (B) and human serum (C). The LOX-FL reaction was monitored as the FL fluorescence decreased at 37 °C as shown in Fig. 1B and in the Experimental section. Measurements were carried out in both the absence (control, traces a, e and i) and in the presence of (A) Trolox (10, 15 and 20 μM ; traces b, c and d, respectively), (B) FS phenols from Lisosan G (obtained from 5, 7.5 and 10 mg of Lisosan G, d.w.; traces f, g and h, respectively) and (C) human serum (10, 20 and 35 μL ; traces l, m and n, respectively). The rates of the FL quenching, expressed as percentage decrease with respect to the control, are reported as a function of the Trolox concentration (inset of (A)), FS phenols from Lisosan G (inset of (B)) and volume of serum (inset of (C)). The resulting equations are: inhibition (%) = $1.920[\text{Trolox}] + 3.23$ ($r = 0.9996$, $p < 0.001$) (inset of (A)); inhibition (%) = $4.154[\text{Lisosan G}] + 8.12$ ($r = 0.9985$, $p < 0.001$) (inset of (B)); and inhibition (%) = $1.165[\text{serum}] + 7.10$ ($r = 0.9989$, $p < 0.001$) (inset of (C)), where Trolox, Lisosan G and serum are expressed in μM , mg d.w. and μL , respectively.

dry weight, d.w., of Lisosan G) were found to inhibit the LOX-FL reaction. In this experiment, the FS phenols inhibited the rate of the FL quenching from about 20% to 50% with a linear relationship between the inhibition and the amount of sample (inset of Fig. 2B). Comparison with the linear regression of Trolox, carried out as reported in Pastore *et al.*,¹⁸ gave an AC value of 4.3 ± 0.4 (SD) μmol Trolox eq. per g (d.w.) (see also Table 2).

The LOX-FL assay was also tested to evaluate the AC in a human serum sample. Analogously, as shown in Fig. 2C, typical experimental traces in the presence of serum are reported. Also in this case, small volumes of serum (ranging from 10 μL to 40 μL) were found to linearly inhibit (from 20% to 50%) the LOX-FL reaction (inset of Fig. 2C). In this experiment, the AC value, in terms of Trolox, was 1.21 ± 0.03 (SD) μmol Trolox eq. per mL serum.

Interestingly, the LOX-FL method may be used on both serum and plasma (not shown). However, caution has to be paid when plasma is obtained using EDTA or citrate as anticoagulants, since they may chelate iron that is necessary to LOX activity. On the contrary, the LOX/RNO assay was found to require too high a volume of both biological fluids (unpublished data).

In order to check the capability of the LOX-FL assay to evaluate the AC of the main antioxidant compounds contained in serum samples, *i.e.* albumin, bilirubin, urate and ascorbate,^{2,11} they were individually investigated (Table 1). Trolox was used as an analogue of α -tocopherol. A comparison with the ORAC assay is also reported. All tested compounds were found to show the AC, as measured by both the LOX-FL and ORAC methods. Interestingly, the IC_{50} values are 4–160 fold higher for the LOX-FL than for the ORAC assay, thus indicating a general lower sensitivity of LOX-FL.

Table 1 Effect of different serum antioxidants on the LOX-FL and ORAC reactions

LOX-FL		
Antioxidant	Range (μM)	IC_{50}^a (μM)
Trolox	5–20	26 ± 2
Albumin	7.5–30	25 ± 2
Bilirubin	1.5–6.0	7.6 ± 0.6
Urate	10–25	40 ± 2
Ascorbate	150–1000	1360 ± 10
ORAC		
Antioxidant	Range (μM)	IC_{50}^b (μM)
Trolox	2.5–10	2.5 ± 0.2
Albumin	0.075–0.225	0.16 ± 0.02
Bilirubin	0.15–0.90	2.0 ± 0.2
Urate	1–4	1.6 ± 0.1
Ascorbate	1.5–18	11.1 ± 0.1

^a For LOX-FL, the IC_{50} values represent the antioxidant concentration able to halve the rates of the LOX-FL reaction. ^b As for ORAC, the IC_{50} values represent the antioxidant concentration able to double the area of the blank ORAC reaction.



Consistently, AC values of sera measured by means of the LOX-FL method show a reference interval from about 1 to 1.6 $\mu\text{mol Trolox eq. per mL}$ (see also caption of Fig. 4), about tenfold lower than that of ORAC as well as about twofold lower than TEAC. Interestingly, this ability to measure the lower basal AC in serum may help to better highlight AC changes after food antioxidant ingestion. In another set of experiments, the effect of the addition of an FS phenolic extract to serum (Fig. 3A and C) and the possible synergism among the antioxidant compounds reported in Table 1 (Fig. 3B and D) were also evaluated. Once again, LOX-FL and ORAC were compared. In each experiment, the AC of each sample was measured and the sum of the AC values of all tested compounds was calculated (AC_{sum}). Then, the same samples were mixed and the AC was determined again (AC_{mix}).

In the experiment shown in Fig. 3A, two equal amounts (0.5 mL) of serum and FS phenols were mixed. For the resultant mixture (1 mL), an AC_{mix} value of $4.41 \pm 0.01 \mu\text{mol Trolox eq.}$,

124% higher than the AC_{sum} ($1.97 \pm 0.08 \mu\text{mol Trolox eq.}$), was measured. In Fig. 3B, to mimic serum composition,³¹ the mix was a solution (1 mL) containing 250 μM urate, 50 μM ascorbate, 10 μM bilirubin, 600 μM albumin and 30 μM Trolox. Trolox was used instead of α -tocopherol due to its solubility in aqueous solution. Interestingly, an AC_{mix} value equal to 1.48 ± 0.05 (SD) $\mu\text{mol Trolox eq.}$ was obtained, 74% higher than the AC_{sum} ($0.85 \pm 0.05 \mu\text{mol Trolox eq.}$).

By using the same approach, low synergism (16% increase, Fig. 3C) or no significant synergistic effect (Fig. 3D) was observed using the ORAC method.

Therefore, the LOX-FL assay may highlight the synergistic effect among food and serum antioxidants as well as among serum antioxidants better than ORAC.

This is consistent with the ability of the LOX-FL reaction to generate, as reported above, more than one oxidant species having relevant physiological significance and to evaluate




Fig. 3 Synergism among human serum and FS phenols (A and C) and among ascorbate, Trolox, bilirubin, urate and albumin (B and D), evaluated by means of the LOX-FL (A and B) and ORAC (C and D) methods. In each plot, the AC values of each sample, as well as the sum (AC_{sum}) and the AC of the Mix (AC_{mix}) are shown as $\mu\text{mol Trolox eq.}$ and as percentages of the AC_{sum} . Synergism is reported as a percentage increase in the AC_{mix} value with respect to the AC_{sum} value according to the equation: synergistic increase (%) = $[(AC_{\text{mix}}/AC_{\text{sum}}) - 1] \times 100$. Data are reported as mean value \pm SD. The probability level (**, $p \leq 0.01$; ***, $p \leq 0.001$; n.s. = not significant) according to the Student's t -test is also reported.



simultaneously different antioxidant mechanisms. These findings are in accordance with the ability of the LOX/RNO method to effectively highlight antioxidant synergisms.^{17,18}

An interesting result evident from Table 1 and Fig. 3 is the major role exerted by albumin as serum antioxidant, whatever the AC method used for the assay; this is in good agreement with previous literature findings.³¹

Ability of the new LOX-FL method to evaluate both the AC of food extracts (*in vitro* measurements) and the effect of food intake on the AC of serum (*ex vivo* measurements)

To better investigate the ability of the new LOX-FL method to measure the *in vitro* AC of food extracts, three different antioxidant extracts from Lisosan G were investigated in some detail. The AC was also evaluated by means of the ORAC and TEAC methods (Table 2). As expected, depending on the different chemistry of the assays, the absolute AC values measured by the different methods are very different from each other, but all three methods measured the highest AC values in the hydrophilic extract of Lisosan G, probably mainly attributable to proteins, as already demonstrated for wheat grain.¹⁷ With regard to the phenolic extracts of Lisosan G, all three methods measured much higher AC values in the IB phenolic extract with respect to the FS phenolic one; these data are in good agreement with numerous literature data reporting IB phenolic compounds as the major phenolic component in cereal whole grain.³² Consistently, the total phenolic and flavonoid contents were higher in the IB phenolic extract than in the FS phenolic one. The lipophilic extract was also evaluated, but only negligible activity was measured (not shown), with AC values equal to about 0.6%, 0.6% and 4% of the total activity as evaluated by the LOX-FL, ORAC and TEAC methods, respectively. On the whole, these results indicate a very high total AC of Lisosan G, which was about 2- to 10-fold higher than that of whole grain from different wheat species.^{16,17}

This result is in agreement with the well-documented bioactivity of Lisosan G. In fact, it has been already shown that Lisosan G protects rats against toxicity induced in the liver by carbon tetrachloride and in the liver, kidney and testis by cisplatin, possibly by attenuating oxidative stress and by preserving antioxidant enzymes.²⁰ Moreover, it induces a decrease in the intracellular ROS concentration and H₂O₂-dependent mutagenesis in *Saccharomyces cerevisiae*,²² activates the NAD(P)H:quinone oxidoreductase and heme oxygenase-1 in

rat hepatocytes,²¹ glutathione peroxidase-1 and superoxide dismutase-2 expression in EPCs exposed to oxidative stress²³ and decreases H₂O₂-induced toxicity in rat hepatocytes.²¹ Furthermore, activation of the nuclear Nrf-2/ARE in rat hepatocytes²¹ and EPCs exposed to oxidative stress²³ has been reported. So, the bioactivity of phytochemicals from Lisosan G may be exerted in several different modes, possibly including AC properties, but also *via* direct modulation of metabolic pathways and/or gene expression.

In order to evaluate the ability of the LOX-FL method to assess changes in the AC of blood, in comparison with ORAC and TEAC, *ex vivo* measurements of serum were carried out before and 30, 60, 90, 120 and 240 min after the intake of 20 g of Lisosan G (18 g in terms of d.w.). On the basis of data reported in Table 2, this implies the intake of about 1700, 3700 and 1600 μ mol Trolox eq. as evaluated by means of the LOX-FL, ORAC and TEAC methods, respectively. The effect on the serum AC is reported as the percentage change in the initial (T_0) AC (Fig. 4). As shown, the ingestion of Lisosan G caused an increase of about 40% in the AC, evaluated by means of the LOX-FL assay, after 30 min. Then, the AC decreased after 90 min, until it reached an AC value similar to T_0 , and increased again at 120 min. This behaviour might be compatible with a prompt release and absorption of FS phenols (see Table 2) within 30–60 min in the small intestine, followed by either a delayed release of IB phenols at 120 min, probably due to the activity of gut microbiota, or to metabolism-increasing antioxidant activity. Finally, the AC value gradually decreased and reached about the same value as that at T_0 after 240 min. On the contrary, the ORAC method did not evaluate a significant increase in the AC after Lisosan G intake, even showing an incoherent decrease at 120 min (about 20%). The TEAC method also did not show relevant changes in the serum AC. Therefore, under the same experimental conditions, different methods may give very different results, thus suggesting that the use of different assays represents a main source of incoherence in a lot of literature data.

Overall, it appears that the ORAC and TEAC methods perform less well with respect to LOX-FL; once again, this is probably due to the ability of this method to simultaneously evaluate different antioxidant mechanisms.

On the contrary, the ORAC assay only measures the chain-breaking AC against a peroxy radical according to a hydrogen atom transfer (HAT) reaction, whereas the TEAC method evaluates the capability of antioxidants to induce the quenching of the non-physiological ABTS^{•+} radical cation, mainly according

Table 2 AC of Lisosan G extracts evaluated by means of the LOX-FL, ORAC and TEAC methods, and content in some antioxidant compounds

Extract	AC (μ mol Trolox eq. per g d.w.)			Antioxidant compounds		
	LOX-FL	ORAC	TEAC	Protein content ^a	Phenolic content ^b	Flavonoid content ^c
Hydrophilic	81.7 \pm 4.7 ^d	123 \pm 6	48 \pm 3	131.7 \pm 2.6	n.d.	n.d.
FS phenolic	4.3 \pm 0.4	25.6 \pm 0.7	7.1 \pm 1.2		34.5 \pm 1.1	2.76 \pm 0.15
IB phenolic	7.1 \pm 0.1	56.6 \pm 4.8	29.5 \pm 2.1		81.1 \pm 0.8	26.6 \pm 1.1

^a Data are expressed as mg per g d.w. ^b Data are expressed as mg of gallic acid eq. per 100 g d.w. ^c Data are expressed as mg of catechin eq. per 100 g d.w. ^d Data are reported as mean value \pm SD. n.d., not determined.






Fig. 4 Percent changes of serum AC, evaluated by means of the LOX-FL, ORAC and TEAC methods, during 240 min after consumption of 20 g of Lisosan G in seven subjects. A different curve was obtained for each subject, having a T_0 value ranging from 1.04 to 1.65, from 9.32 to 26.18 and from 1.78 to 3.71 μmol Trolox eq. per mL of serum, as calculated by LOX-FL, ORAC and TEAC methods, respectively. Then, for each AC method, the seven curves were mediated. Data are reported as mean value \pm SE.

to a single electron transfer (SET) redox reaction, this latter having a minor role *in vivo*.¹ The different rationales for the different methods easily explain why the results of LOX-FL, ORAC and TEAC are unrelated to each other.

Conclusions

The LOX-FL method is able to give a physiological and comprehensive evaluation of the AC, therefore better highlighting the synergistic interactions among the different classes of antioxidants. This novel tool may be used to obtain both *in vitro* measurements of food extracts and *ex vivo* measurements of serum after food ingestion. This ability allows an integrated evaluation of food, using the same assay method, by testing whether foods showing a strong AC may give realistic physiological effects after ingestion. Interestingly, by using this approach on the antioxidant-rich dietary wheat grain supplement, Lisosan G, it was shown that its consumption may significantly improve the antioxidant status of serum.

This information cannot be obtained when the ORAC and TEAC methods are used. In fact, although all methods are able to give coherent information about the AC of *in vitro* extracts of Lisosan G, LOX-FL was found to be able to highlight a remarkable AC increase in serum after Lisosan G ingestion, while under the same experimental conditions, ORAC and TEAC failed to do this.

Acknowledgements

This work was supported by the following research projects: the Apulian Region Project "FutureInResearch – Study of antioxidant

action of plant foods using innovative *in vitro* and *ex vivo* methods for valorisation of health properties of apulian products (code 2GE9US7)", the MiUR projects PON01_01145 "ISCOCEM" and PON02_00186_2937475 "Pro.Ali.Fun." and the Apulian Region Project "LAIFF-Laboratories Network for the Innovation on Functional Foods". We thank Dr Gaetano Pio Spera who participated as a PhD student in the present work. Publication supported by funds "5 \times 1000 dell'IRPEF a favore dell'Università degli Studi di Foggia" in memory of Gianluca Montel.

References

- 1 D. Huang, O. U. Boxin and R. L. Prior, *J. Agric. Food Chem.*, 2005, **53**, 1841–1856.
- 2 E. Niki, *Free Radical Biol. Med.*, 2010, **49**, 503–515.
- 3 F. Shahidi and Y. Zhong, *J. Funct. Foods*, 2015, **18**, 757–781.
- 4 D. Del Rio, A. Rodriguez-Mateo, J. P. E. Spencer, M. Tognolini, G. Borges and A. Crozier, *Antioxid. Redox Signaling*, 2013, **18**, 1818–1892.
- 5 S. Rautiainen, S. Larsson, J. Virtamo and A. Wolk, *Stroke*, 2012, **43**, 335–340.
- 6 S. Rautiainen, E. B. Levitan, M. A. Mittleman and A. Wolk, *Am. J. Med.*, 2013, **126**, 494–500.
- 7 S. Rautiainen, E. B. Levitan, M. A. Mittleman and A. Wolk, *Eur. J. Heart Failure*, 2015, **17**, 20–26.
- 8 C. G. Fraga, P. I. Oteiza and M. Galleano, *Biochim. Biophys. Acta, Gen. Subj.*, 2014, **1840**, 931–934.
- 9 C. Bumke-Vogt, M. A. Osterhoff, A. Borchert, V. Guzman-Perez, Z. Sarem, A. L. Birkenfeld, V. Bähr and A. F. H. Pfeiffer, *PLoS One*, 2014, **9**, e104321.
- 10 M. Marimoutou, F. Le Sage, J. Smadja, C. L. D'Hellencourt, M.-P. Gonthier and C. R. Da Silva, *J. Inflammation*, 2015, **12**, 10.
- 11 A. Pompella, H. Sies, R. Wacker, F. Brouns, T. Grune, H. K. Biesalski and J. Frank, *Nutrition*, 2014, **30**, 791–793.
- 12 R. L. Prior, *J. Funct. Foods*, 2015, **18**, 797–810.
- 13 M. El Assar, J. Angulo and L. Rodríguez-Mañas, *Free Radical Biol. Med.*, 2013, **65**, 380–401.
- 14 D. Lettieri-Barbato, F. Tomei, A. Sancini, G. Morabito and M. Serafini, *Br. J. Nutr.*, 2013, **109**, 1544–1556.
- 15 M. S. Fernández-Panchón, D. Villano, A. M. Troncoso and M. C. Garcia-Parrilla, *Crit. Rev. Food Sci. Nutr.*, 2008, **48**, 649–671.
- 16 M. N. Laus, A. Gagliardi, M. Soccio, Z. Flagella and D. Pastore, *J. Food Sci.*, 2012, **77**, C1150–C1155.
- 17 M. N. Laus, D. Tozzi, M. Soccio, A. Fratianni, G. Panfili and D. Pastore, *J. Cereal Sci.*, 2012, **56**, 214–222.
- 18 D. Pastore, M. N. Laus, D. Tozzi, V. Fogliano, M. Soccio and Z. Flagella, *J. Agric. Food Chem.*, 2009, **57**, 9682–9692.
- 19 D. Pastore, D. Trono, L. Padalino, N. Di Fonzo and S. Passarella, *Plant Physiol. Biochem.*, 2000, **38**, 845–852.
- 20 V. Longo, P. G. Gervasi and V. Lubrano, *Food Chem. Toxicol.*, 2011, **49**, 233–237.
- 21 M. La Marca, P. Beffy, A. Pugliese and V. Longo, *PLoS One*, 2013, **8**, e83538.
- 22 S. Frassinetti, C. M. Della Croce, L. Caltavuturo and V. Longo, *Food Chem.*, 2012, **135**, 2029–2034.



- 23 D. Lucchesi, R. Russo, M. Gabriele, V. Longo, S. Del Prato, G. Penno and L. Pucci, *PLoS One*, 2014, **9**, e109298.
- 24 D. Pastore, D. Trono, L. Padalino, S. Simone, D. Valenti, N. Di Fonzo and S. Passarella, *J. Cereal Sci.*, 2000, **31**, 41–54.
- 25 B. Ou, M. Hampsch-Woodill and R. L. Prior, *J. Agric. Food Chem.*, 2001, **49**, 4619–4626.
- 26 R. Re, N. Pellegrini, A. Proteggente, A. Pannala, M. Yang and C. Rice-Evans, *Free Radical Biol. Med.*, 1999, **26**, 1231–1237.
- 27 M. N. Laus, N. A. Di Benedetto, R. Caporizzi, D. Tozzi, M. Soccio, L. Giuzio, P. De Vita, Z. Flagella and D. Pastore, *Plant Foods Hum. Nutr.*, 2015, **70**, 207–214.
- 28 D. Trono, M. Soccio, M. N. Laus and D. Pastore, *Plant Sci.*, 2013, **199–200**, 91–102.
- 29 B. Ou, M. Hampsch-Woodill, J. Flanagan, E. K. Deemer, R. L. Prior and D. Huang, *J. Agric. Food Chem.*, 2002, **50**, 2772–2777.
- 30 J. M. Aubry and S. Bouttemy, *J. Am. Chem. Soc.*, 1997, **119**, 5286–5294.
- 31 G. Cao and R. L. Prior, *Clin. Chem.*, 1998, **44**, 1309–1315.
- 32 N. Okarter and R. H. Liu, *Crit. Rev. Food Sci. Nutr.*, 2010, **50**, 193–208.

