



Cite this: *Dalton Trans.*, 2016, **45**, 10118

Received 14th April 2016,
Accepted 28th April 2016

DOI: 10.1039/c6dt01439a

www.rsc.org/dalton

Stabilization of molecular lanthanide polysulfides by bulky scorpionate ligands†‡

Marcel Kühling,^a Robert McDonald,^b Phil Liebing,^a Liane Hilfert,^a Michael J. Ferguson,^b Josef Takats*^b and Frank T. Edelmann*^a

Well-defined lanthanide polysulfide complexes containing S_4^{2-} and S_5^{2-} ligands, the samarium(III) pentasulfide complex $Sm(Tp^{iPr_2})(\kappa^1-3,5\text{-}iPr_2Hpz)(S_5)$ and the tetrasulfide-bridged binuclear ytterbium(III) complex $(\mu-S_4)[Yb(Tp^{iPr_2})(\kappa^1-3,5\text{-}iPr_2Hpz)(\kappa^2-3,5\text{-}iPr_2pz)]_2$ (Tp^{iPr_2} = hydro-tris(3,5-diisopropylpyrazolyl)borate), have been synthesized and structurally characterized by single-crystal X-ray diffraction.

Introduction

Polysulfides of the rare-earth metals (Ln) have been known for over 100 years.^{1,2} The most common types of these materials are LnS_2 ,^{3,4} and LnS_{2-x} ,^{5,6} but well-defined, more complicated species such as Ln_8S_{14} ,⁹ (Ln = Dy, Ho) have also been described.⁷ The synthetic routes to such lanthanide polysulfides normally require high-temperature and/or high-pressure techniques.^{8,9} In sharp contrast, molecular lanthanide complexes containing polysulfide ligands remain little explored, although a number of interesting inorganic^{10–17} and organometallic^{18–20} sulfido clusters have been reported in recent years. Most of these contain the S_2^{2-} ligand, while $(Cp^*_2Sm)_2(S_3)(THF)$ (Cp^* = pentamethylcyclopentadienyl) appears to be the only lanthanide complex with an S_3^{2-} ligand.¹⁰ We describe here the formation and structural characterization of the first lanthanide polysulfide complexes containing the S_4^{2-} and S_5^{2-} ligands, namely $Sm(Tp^{iPr_2})(\kappa^1-3,5\text{-}iPr_2Hpz)(S_5)$ (**3**) and $(\mu-S_4)[Yb(Tp^{iPr_2})(\kappa^1-3,5\text{-}iPr_2Hpz)(\kappa^2-3,5\text{-}iPr_2pz)]_2$ (**5**)

(Tp^{iPr_2} = hydro-tris(3,5-diisopropylpyrazolyl)borate; $3,5\text{-}iPr_2pz$ = 3,5-diisopropylpyrazolate; $3,5\text{-}iPr_2Hpz$ = 3,5-diisopropylpyrazole), which are stabilized by bulky scorpionate ligands.

It is well established from the chemistry of the bent metallocenes Cp^*_2Ln (Ln = Sm, Eu, Yb) that these strongly reducing lanthanide(II) precursors are easily oxidized by elemental sulfur or diorganodisulfides (RSSR), resulting in the formation of trivalent metallocenes containing S^{2-} or RS^- ligands.^{10,21,22} Thus, it was of interest to investigate the reactivity of the recently reported “bent sandwich-like” divalent lanthanide scorpionates, $Ln(Tp^{iPr_2})_2$ (Ln = Sm, Eu, Tm, Yb)^{23,24} with S_8 with a view to establish whether the presence of bulky scorpionate ligands might lead to the formation of polysulfide-ligated species. Here we report the results of the reactions of $Ln(Tp^{iPr_2})_2$ (Ln = Sm (**1**), Yb (**2**)) with sulfur.

Synthesis and characterization

The reaction of $Sm(Tp^{iPr_2})_2$ (**1**) with sulfur was first carried out in toluene. Addition of one equiv. of S_8 to a toluene solution of **1** at r.t., resulted in a rapid color change from dark green to orange, with some unreacted sulfur remaining at the bottom of the flask. Recovery of the sulfur showed that only half an equivalent of sulfur (S_8) reacted with **1**. Solvent removal produced a somewhat sticky yellow solid, the usual form of crude products due to the greasy, lipophilic nature of the Tp^{iPr_2} ligand. Curiously, crystallization attempts from various solvent mixtures (Et_2O/THF ; Et_2O /hexane) first produced small amounts of yellow crystals, which proved to be elemental sulfur, orthorhombic and monoclinic forms. From further crystallization attempts only a few suitable crystals for X-ray analysis could be isolated, which were shown to be $Sm(Tp^{iPr_2})(\kappa^1-3,5\text{-}iPr_2Hpz)(S_5)$ (**3**), Scheme 1. The observation that the first crystalline materials were elemental sulfur indicated either that the amount of sulfur in solution was more than needed for the reaction or that some intermediate Sm-polysul-


^aChemisches Institut der Otto-von-Guericke-Universität, 39106 Magdeburg, Germany. E-mail: frank.edelmann@ovgu.de

^bDepartment of Chemistry, University of Alberta, Edmonton, Alberta, AB, Canada T6G 2G2. E-mail: joe.takats@ualberta.ca

†Dedicated to Professor R. Dieter Fischer on the occasion of his 80th birthday.

‡Electronic supplementary information (ESI) available: Full crystallographic data for **3**, **4**, and **5**. CCDC 1472924, 1472923 and 1472811. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6dt01439a





Scheme 1 Reactions of $\text{Ln}(\text{Tp}^{\text{iPr}_2})_2$ ($\text{Ln} = \text{Sm}, \text{Yb}$) with elemental sulfur.

fide species underwent sulfur extrusion. To answer the question as to the stoichiometric amount of sulfur needed to oxidize the precursor $\text{Sm}(\text{Tp}^{\text{iPr}_2})_2$ compound, a second reaction was carried out which showed that bleaching of the dark green THF solution of **1** at RT occurred after the addition of only 1 equiv. of sulfur (S) per Sm(II). Crystallization of the crude, yellow, sticky solid produced a small amount of white crystals which proved to be $\text{Sm}(\text{Tp}^{\text{iPr}_2})_2(\kappa^2\text{-}3,5\text{-}^i\text{Pr}_2\text{pz})$ (**4**, Scheme 1). Unfortunately, numerous other crystallization attempts from various mixed solvent system did not yield crystals suitable for X-ray analysis, hence the nature of the sulfur containing species from this second reaction remain unknown.

The analogous reaction of $\text{Yb}(\text{Tp}^{\text{iPr}_2})_2$ (**2**) with sulfur was found to be more straightforward. Stirring of equimolar amounts (Yb : S) of **2** and sulfur in toluene for 24 h caused a color change from bright red to amber.

Recrystallization of the reaction product from *n*-pentane afforded orange, cube-like single crystals of $5 \cdot 2\text{C}_5\text{H}_{12}$ suitable for X-ray diffraction.

The constitution and molecular structure of compounds **3**, **4** and **5** were revealed by single crystal X-ray diffraction studies. The structures of **3** and **5** are shown in Fig. 1 and 2, respectively. The structure of compound **4** can be found in the ESI.† The structure of **3** consists of a normal $\kappa^3\text{-Tp}^{\text{iPr}_2}$ ligand, a coordinated, neutral $\kappa^1\text{-}3,5\text{-}^i\text{Pr}_2\text{Hpz}$ ligand, and the electronic demand of the Sm(III) ion is satisfied by the dianionic S_5^{2-} polysulfide ligand. The most remarkable feature of the compound is the conformation of the SmS_5 six-membered ring. As opposed to the classical chair conformation found in transition-metal complexes containing the MS_5 moiety, such as Cp_2MS_5 ($\text{M} = \text{Ti}, \text{Zr}, \text{Hf}$), $[\text{FeS}_5(\mu\text{-S})]_2^{2-}$ and $[\text{Pt}(\text{S}_5)_3]^{2-}$,^{25–30} and that in cyclo-S_6 ,³¹ the SmS_5 six-membered ring adopts the twist-boat conformation. It is noteworthy that a similar ring conformation was also observed in the related actinide f-element compound, $(\text{C}_5\text{Me}_5)_2\text{ThS}_5$,³² and may signal that the twist-boat conformation is a common feature of An/Ln S_5 containing compounds. The similarities between the two structures extend to the M–S distances as well. Interestingly not only the terminal S(alpha) atoms are bonded to Sm (Sm–S1/S5




Fig. 1 Molecular structure of **3** in the crystal showing the atom-labeling scheme. Selected bond lengths [Å] and angles [°]: Sm–N12 2.533(2), Sm–N22 2.531(2), Sm–N32 2.509(2), Sm–N41 2.606(2), Sm–S1 2.755(1), Sm–S5 2.792(1), Sm–S2 2.952(1), Sm–S4 3.090(1); Sm–S1–S2 74.66(3), Sm–S5–S4 77.88(3), S1–Sm–S5 113.10(2), S1–S2–S3 106.63(5), S5–S4–S3 108.54(5), S2–S3–S4 101.42(4).




Fig. 2 Molecular structure of **5** in the crystal showing the atom-labeling scheme. Symmetry operator to generate equivalent atoms: ' 2 – x, y, 1 – z. Selected bond lengths [Å] and angles [°]: Yb–N12 2.423(7), Yb–N22 2.421(7), Yb–N32 2.406(6), Yb–N41 2.304(7), Yb–N42 2.244(7), Yb–N52 2.443(7), Yb–S1 2.703(2), S1–S2 2.072(3), S2–S2' 2.044(5), Yb–S1–S2 106.0(1), S1–S2–S2' 106.8(1), S1–S2–S2'–S1' dihedral angle 69.5(6).

2.755(1)/2.792(1) Å) but also the two S(beta) atoms (Sm–S2/S4 2.952(1)/3.090(1) Å); the corresponding distances in the C_2 symmetric $(\text{C}_5\text{Me}_5)_2\text{ThS}_5$ compound are Th–S1/S1' and Th–S2/S2', 2.768(4) and 3.036(4), respectively. In the transition-metal MS_5 compounds with the chair conformation only the terminal sulfurs of the chelating S_5^{2-} ligand are coordinated to the metal.

Compound **5** is a dinuclear ytterbium(III) polysulfide complex in which two $[\text{Yb}(\text{Tp}^{\text{iPr}_2})(\kappa^1\text{-}3,5\text{-}^i\text{Pr}_2\text{Hpz})(\kappa^2\text{-}3,5\text{-}^i\text{Pr}_2\text{pz})]$ moieties are bridged by a $\mu\text{-S}_4^{2-}$ ligand. Each formally 7-coordinated ytterbium ion is bonded to an original $\kappa^3\text{-Tp}^{\text{iPr}_2}$ ligand as well as a neutral $\kappa^1\text{-}3,5\text{-}^i\text{Pr}_2\text{Hpz}$ ligand and an anionic $\kappa^2\text{-}3,5\text{-}^i\text{Pr}_2\text{pz}$ ligand which were formed as a result of ligand fragmentation.³³ Certainly this accounts for the fairly low isolated yield of compound **5** (32%). In accordance with the overall C_2 -symmetry of the molecule, the ^{11}B NMR spectrum of **5** shows only a single resonance ($\delta = -4.4$ ppm). The conformation of the bridging $\mu\text{-S}_4^{2-}$ ligand closely resembles that of the tetrasulfide dianion in crystalline Na_2S_4 ,^{34,35} with a dihedral angle of 69.5(6)° (97.81° in Na_2S_4). The Yb–S bond length is 2.703(2) Å.



Conclusions

In summarizing the results reported here, although formation of compounds **3**, **4** and **5** was accompanied by Tp^{iPr_2} ligand fragmentation, their isolation underscores the following points: 1. The fact that $\text{Sm}(\text{Tp}^{\text{iPr}_2})_2(\kappa^2\text{-}3,5\text{-iPr}_2\text{pz})$ (**4**) contains a κ^2 -pyrazolide wedged between two $(\text{Tp}^{\text{iPr}_2})_2$ ligand, holds out the hope that under suitable conditions, highly reducing $\text{Ln}(\text{Tp}^{\text{iPr}_2})_2$ compounds might be able to bind and activate interesting small molecules like CO, CO_2 , etc. 2. The very rapid reaction of compound **1** with sulfur and the rather non-selective formation of compound **3** underlines the stronger reducing power of the $\text{Sm}(\text{II})$ precursor compared with $\text{Yb}(\text{II})$. Thus, the reactivity of $\text{Yb}(\text{Tp}^{\text{iPr}_2})_2$ is significantly reduced and allows for the more straightforward preparation of the tetrasulfide derivative **5**. 3. The successful isolation and structural characterization of the first lanthanide(III) polysulfide complexes, $\text{Sm}(\text{Tp}^{\text{iPr}_2})(\kappa^1\text{-}3,5\text{-iPr}_2\text{Hpz})(\text{S}_5)$ (**3**) and $(\mu\text{-S}_4)[\text{Yb}(\text{Tp}^{\text{iPr}_2})(\kappa^1\text{-}3,5\text{-iPr}_2\text{Hpz})(\kappa^2\text{-}3,5\text{-iPr}_2\text{pz})]_2$ (**5**) encourages the quest for more rational synthesis of similar lanthanide polysulfides and the study of their bonding characteristics. Such studies are underway in our laboratories.

This work was financially supported by the Otto-von-Guericke-Universität Magdeburg and the University of Alberta. JT thanks Jackie Kiplinger (Los Alamos National Laboratory) for useful discussions and we thank the reviewers for helpful comments.

Experimental section

Synthesis of complex **3**

(i) Sulfur flakes (56.7 mg, 0.22 mmol S_8) were added to a stirred toluene solution of **1** (239 mg, 0.22 mmol) at r.t. After only about a minute the color changed from dark green to orange, with substantial amount of unreacted sulfur (ca. 28 mg) at the bottom of the round-bottom flask. Decantation from the unreacted sulfur, followed by solvent removal gave a slightly sticky, yellow solid. Extraction with *n*-pentane left behind a small amount of pale yellow solid, shown to be sulfur (orthorhombic) by X-ray diffraction on crystals obtained from $\text{Et}_2\text{O}/\text{THF}$ solvent mixture. Crystallization attempts from *n*-pentane only produced powdery solids. Crystallization from $\text{Et}_2\text{O}/n$ -hexane solvent mixtures, two attempts, produced small amounts of yellow crystals which were again shown to be sulfur (orthorhombic and monoclinic forms). Ultimately from $\text{THF}/\text{hexamethylidisiloxane}$ (HMDSO), a small number of pale yellow crystals were harvested, which proved to be compound **3**.

(ii) Small amounts of sulfur flakes were added to a stirred THF solution of **1** (523 mg, 0.484 mmol) at r.t. After the addition of only 16 mg (0.5 mmol of S) the original dark green color changed to orange. Solvent removal gave a slightly sticky, yellow solid. Crystallizations from $\text{Et}_2\text{O}/n$ -hexane solvent mixtures, twice produced crystals of compound **3**. Numerous other crystallization attempts failed to produce crystals suitable for X-ray analysis and failed to give sulfur-containing material.

Synthesis of complex **5**

Powdered sulfur (14 mg, 0.44 mmol) was added to a solution of compound **2** (484 mg, 0.44 mmol) in toluene (20 ml). After 24 h stirring at r.t. the color had changed from bright red to amber. Removal of the solvent and recrystallization from *n*-pentane (15 ml, -5°C) afforded 152 mg (32%) of **5** as orange, cube-like crystals. Mp. 130°C . ^1H NMR (toluene- d_6 , 400 MHz): δ 10.53 (s br, 6H, pz), 5.40 (s br, 2H, C-H ^iPr), 3.04 (s, 2H, pz), 2.0–2.2 (s br, 2H, B-H), 2.38 (s, 12H, CH_3 ^iPr), 1.23 (m, 12H, CH_2 of *n*-pentane), 0.87 (t, 12H, CH_3 of *n*-pentane), -0.20 (s br, 6H, C-H ^iPr), -0.65 (s, 2H, pz), -0.99 (s, 2H, C-H ^iPr), -2.33 (s br, 12H, CH_3 ^iPr), -6.22 (s br, 48H, CH_3 ^iPr), -12.00 (s br, 48H, CH_3 ^iPr). ^{13}C NMR (toluene- d_6 , 100 MHz): δ = 173.8 (q-C pz), 150.9 (q-C pz), 102.3 (CH pz) 87.3 (CH pz), 34.4 (CH_2 of *n*-pentane), 27.3 (CH ^iPr), 23.5 (CH_3 ^iPr), 22.7 (CH_2 of *n*-pentane), 21.5 (CH_3 ^iPr), 19.5 (C-H ^iPr), 15.6 (CH_3 ^iPr), 14.1 (CH_3 of *n*-pentane). ^{11}B NMR (toluene- d_6 , 128 MHz): δ = -4.4 ppm. Elemental analysis calcd for $\text{C}_{100}\text{H}_{178}\text{B}_2\text{N}_{20}\text{S}_4\text{Yb}_2$ ($5\text{-}2\text{C}_5\text{H}_{12}$): C 55.69, H 8.32, N 12.99; found C 56.04, H 8.64, N 12.50.

References

- 1 W. Biltz, *Z. Elektrochem.*, 1911, **17**, 668.
- 2 J. Flahaut, M. Guitard and M. Patrie, *Bull. Soc. Chim. Fr.*, 1959, 1917.
- 3 Y. Yanagisawa and S. Kume, *Mater. Res. Bull.*, 1986, **21**, 379.
- 4 L. C. Otero-Diaz, M. J. Torralvo and R. M. Rojas, *Solid State Ionics*, 1993, **63–65**, 318.
- 5 R. Tamazyan, S. van Smaalen, I. G. Vasilyeva and H. Arnold, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2003, **59**, 709.
- 6 T. Doert, C. Graf, P. Laux and T. Schleid, *Z. Anorg. Allg. Chem.*, 2007, **633**, 2719.
- 7 T. Doert, C. Graf, I. G. Vasilyeva and W. Schnelle, *Inorg. Chem.*, 2012, **51**, 282.
- 8 A. W. Webb and H. T. Hall, *Inorg. Chem.*, 1970, **9**, 1084.
- 9 C. J. Müller, U. Schwarz and T. Doert, *Z. Anorg. Allg. Chem.*, 2012, **638**, 2477, and references cited therein.
- 10 W. J. Evans, G. W. Rabe, J. W. Ziller and R. J. Doedens, *Inorg. Chem.*, 1994, **33**, 2719.
- 11 J. H. Melman, T. J. Emge and J. G. Brennan, *Chem. Commun.*, 1997, 2269.
- 12 D. Freedman, T. J. Emge and J. G. Brennan, *Inorg. Chem.*, 1999, **38**, 4400.
- 13 J. H. Melman, T. J. Emge and J. G. Brennan, *Inorg. Chem.*, 1999, **38**, 2117.
- 14 M. Fitzgerald, T. J. Emge and J. G. Brennan, *Inorg. Chem.*, 2002, **41**, 3528.
- 15 Z. Zhang, L. Zhang, Y. Li, L. Hong, Z. Chen and X. Zhou, *Inorg. Chem.*, 2010, **49**, 5715.
- 16 K. Norton, S. Banerjee, S. Das, L. Hübner, T. J. Emge and J. G. Brennan, *Dalton Trans.*, 2010, **39**, 6794.
- 17 J. F. Corbey, M. Fang, J. W. Ziller and W. J. Evans, *Inorg. Chem.*, 2015, **54**, 801.



- 18 Y. Li, C. Pi, J. Zhang, X. Zhou, Z. Chen and L. Weng, *Organometallics*, 2005, **24**, 1982.
- 19 N. Arleth, S. Bestgen, M. T. Gamer and P. W. Roesky, *J. Am. Chem. Soc.*, 2014, **136**, 14023.
- 20 J. Li, J. Hao and C. Cui, *Dalton Trans.*, 2015, **44**, 767.
- 21 D. J. Berg, C. J. Burns, R. A. Andersen and A. Zalkin, *Organometallics*, 1989, **8**, 1865.
- 22 A. Recknagel, M. Noltemeyer, D. Stalke, U. Pieper, H.-G. Schmidt and F. T. Edelmann, *J. Organomet. Chem.*, 1991, **411**, 347.
- 23 A. Momin, L. Carter, Y. Yang, R. McDonald, S. Essafi, F. Nief, I. Del Rosal, A. Sella, L. Maron and J. Takats, *Inorg. Chem.*, 2014, **53**, 12066.
- 24 M. Kühling, C. Wickleder, M. J. Ferguson, C. G. Hrib, R. McDonald, M. Suta, L. Hilfert, J. Takats and F. T. Edelmann, *New J. Chem.*, 2015, **39**, 7617.
- 25 H. Köpf, B. Block and M. Schmidt, *Chem. Ber.*, 1968, **101**, 272.
- 26 E. F. Epstein, I. Bernal and H. Köpf, *J. Organomet. Chem.*, 1971, **26**, 229.
- 27 E. G. Muller, J. L. Petersen and L. F. Dahl, *J. Organomet. Chem.*, 1976, **111**, 91.
- 28 M. Draganjac and T. B. Rauchfuss, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 742, and references therein.
- 29 A. Müller and E. Diemann, *Adv. Inorg. Chem.*, 1987, **31**, 89.
- 30 N. Takeda, N. Tokito and R. Okazaki, *Top. Curr. Chem.*, 2003, **231**, 153.
- 31 M. Schmidt, B. Block, H. D. Block, H. Köpf and E. Wilhelm, *Angew. Chem., Int. Ed. Engl.*, 1968, **7**, 632.
- 32 D. A. Wroblewski, D. T. Cromer, J. V. Ortiz, T. B. Rauchfuss, R. R. Ryan and A. P. Sattelberger, *J. Am. Chem. Soc.*, 1986, **108**, 174.
- 33 Fragmentation of substituted Tp ligands is well documented (A. Domingos, M. R. J. Elsegood, A. C. Hillier, G. Lin, S. Y. Liu, I. Lopes, N. Marques, G. H. Maunder, R. McDonald, A. Sella, J. W. Steed and J. Takats, *Inorg. Chem.*, 2002, **41**, 6761, and references therein), but it seems to be even more prevalent with the Tp^{iPr}₂ ligand as seen in reactivity studies of Ln(Tp^{iPr}₂)₂ (Ln = Sm, Tm) (J. Takats, unpublished work.).
- 34 R. Tegman, *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.*, 1973, **29**, 1463.
- 35 S. J. Chadwell, D. Rickard and G. W. Luther III, *Electroanalysis*, 2001, **13**, 21.

