

# Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: W. Tang, J. Liu, M. Nie, Q. Zhou, S. Gao, W. Jiang, L. W. Chung and K. Ding, *Chem. Sci.*, 2017, DOI: 10.1039/C7SC01254C.




This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



- chiral tertiary boronic ester
- perfect regioselectivity
- Pd loading as low as 0.2 mol %
- excellent yields and enantioselectivities
- first enantioselective synthesis of brassinazole
- concerted oxidative addition and allene insertion

Enantioselective palladium-catalyzed diboration of 1,1-disubstituted allenes is developed for the first time by employing a P-chiral monophosphorus ligand BI-DIME.





## Chemical Science

## EDGE ARTICLE

## Enantioselective Palladium-Catalyzed Diboration of 1,1-Disubstituted Allenes

Jiawang Liu,<sup>a,§</sup> Ming Nie,<sup>a,§</sup> Qinghai Zhou,<sup>b,d</sup> Shen Gao,<sup>a</sup> Wenhao Jiang,<sup>a</sup> Lung Wa Chung,<sup>b,\*</sup> Wenjun Tang,<sup>a,\*</sup> Kuiling Ding<sup>a,c,d,\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A practical and enantioselective palladium-catalyzed diboration of 1,1-disubstituted allenes is developed by employing a P-chiral monophosphorus ligand BI-DIME to form a series of diboronic esters containing a chiral tertiary boronic ester moiety in excellent yields and ee's with the palladium loading as low as 0.2 mol %. DFT calculations revealed a concerted mechanism of oxidative addition of bis(pinacolato)diboron and allene insertion, as well as a critical dispersion effect on the origins of the enantioselectivity. The method is successfully applied to a concise and enantioselective synthesis of brassinazole.

Chiral boronic esters have become versatile building blocks in synthetic organic chemistry.<sup>1</sup> Synthesis of chiral tertiary boronic esters have attracted considerable interests recently and some notable methods are developed (Figure 1) including Aggarwal-Matteson lithiation-borylation methodology from chiral secondary alcohols,<sup>2</sup> asymmetric hydroboration of 1,1-disubstituted alkenes,<sup>3</sup> and asymmetric borylation of allylic carbonates,<sup>4</sup> Michael acceptors,<sup>5</sup> as well as tertiary halides.<sup>6</sup> The asymmetric diboration of 1,1-disubstituted alkenes or allenes<sup>7</sup> would not only provide a chiral tertiary boronic ester moiety, but also form an additional boronic ester component for further transformations. Work by Morken on asymmetric diboration<sup>8,9</sup> with various transition metal catalysts (Rh, Pt, or Pd) has provided significant progress in forming secondary boronic ester products with excellent ee's. However, the asymmetric diboration of 1,1-disubstituted alkenes or allenes to form chiral tertiary boronic esters remains challenging with either low ee's or low yields.<sup>9b</sup> Herein we communicate our results on palladium-catalyzed asymmetric diboration of 1,1-disubstituted allenes that have led to a series of diboronic esters containing a chiral tertiary boronic ester moiety in excellent enantioselectivities and yields with the employment of a P-chiral monophosphorus ligand BI-DIME. The chiral

diboronic ester products are useful chiral building blocks which have led to the first enantioselective synthesis of a specific brassinosteroid biosynthetic inhibitor—brassinazole.<sup>19</sup>

<sup>a</sup> J. Liu, M. Nie, Shen Gao, Wenhao Jiang, Prof. Dr. W. Tang, and Prof. Dr. K. Ding State Key Laboratory of Organometallic Chemistry and State Key Laboratory of Bio-Organic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Ling Ling Rd, Shanghai 200032. E-mail: tangwenjun@sioac.ac.cn; kding@sioac.ac.cn.

<sup>b</sup> Dr. Q. Zhao, Prof. Dr. L. W. Chung Department of Chemistry, South University of Science and Technology of China, Shenzhen 518055. E-mail: oscarchung@sustc.edu.cn.

<sup>c</sup> University of Chinese Academy of Sciences, Beijing 100049 (China)

<sup>d</sup> Collaborative Innovation Center of Chemical Science and Engineering, Tianjin 300071 (China)

<sup>§</sup> J. Liu and M. Nie contributed equally

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

**Figure 1.** Formation of chiral tertiary boronic esters by asymmetric diboration.

We chose to use buta-2,3-dien-2-ylbenzene (**1a**) as the substrate to investigate the palladium-catalyzed asymmetric diboration of 1,1-disubstituted allenes (Table 1). The reactions were carried out at rt in cyclohexane for 24 h with bis(pinacolato)diboron as the reagent in the presence of Pd<sub>2</sub>(dba)<sub>3</sub> (1 mol %) and a chiral phosphorus ligand (2.5 mol %)





## ARTICLE


## Journal Name


(Table 1). It was found that chelating phosphorus ligands such as BINAP and SDP did not provide any reactivities (entries 1-2).


**Table 1.** Enantioselective Pd-catalyzed diboration of buta-2,3-dien-2-ylbenzene (**1a**) with bis(pinacolato)diboron (**2**)





  
**1a**


  
**2** (1.2 equiv)


  
**3a**


  
(*R*)-BINAP (**L1**)


  
(*R*)-SDP (**L2**)


  
(*S,S*)-Me-DuPhos (**L3**)

  
(*R,R,R*)-SKP (**L4**)

  
**L5**

  
(*S*)-MonoPhos (**L6**)

  
(*S*)-AntPhos (**L7**)

  
(*S*)-BI-DIME (**L8**)

Entry <sup>a</sup>	L*	solvent	yield (%) <sup>b</sup>	ee (%) <sup>c</sup>
1	<b>L1</b>	CyH	0	--
2	<b>L2</b>	CyH	0	--
3	<b>L3</b>	CyH	15	52
4	<b>L4</b>	CyH	95	59
5	<b>L5</b>	CyH	76	79
6	<b>L6</b>	CyH	45	12
7	<b>L7</b>	CyH	60	91
8	<b>L8</b>	CyH	98	94
9	<b>L8</b>	toluene	82	92
10	<b>L8</b>	THF	51	91
11	<b>L8</b>	Dioxane	27	92
12	<b>L8</b>	DCM	0	--
13 <sup>d</sup>	<b>L8</b>	CyH	33	92
14 <sup>e</sup>	<b>L8</b>	CyH	97	94

<sup>a</sup>Unless otherwise specified, the reactions were performed under nitrogen at rt for 24 h with **1a** (0.20 mmol), **2** (0.24 mmol), L\* (2.5 mol %) and Pd<sub>2</sub>(dba)<sub>3</sub> (1.0 mol %) in the specified solvent. Product **3a** was the only detectable product. The *R* absolute configuration of **3a** was determined by comparing its optical rotation with reported data.<sup>14</sup> <sup>b</sup>Isolated yields. <sup>c</sup>Determined by HPLC on a chiral IC-3 column. <sup>d</sup>Pd(OAc)<sub>2</sub> instead of Pd<sub>2</sub>(dba)<sub>3</sub> was employed as the precursor. <sup>e</sup>**1a** (36.0 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (0.1 mmol %), **L8** (0.25 mmol %), 72 h.

A low yield (15%) and ee (52% ee) were observed when Me-DuPhos was employed as the ligand (entry 3). It should be noted that the diboration occurred exclusively on the substituted double bond of the allene, provided the product **3a** containing both a tertiary boronic ester and an alkenyl boronic ester moieties. Interestingly, the SKP ligand (**L4**) with a large bite angle led to an excellent yield (95%) and a moderate ee (59% ee) (entry 4). We thus predicted that the reaction could be better promoted with a monophosphorus ligand, as observed by Morken in diboration of monosubstituted allenes.<sup>9</sup> Thus, a TADDOL-derived monophosphoramidite ligand **L5** led to the formation of **3a** in 76% yield and in 79% ee (entry 5). Another monophosphoramite ligand **L6** derived from a chiral BINOL backbone proved to be less effective, indicating the importance of the ligand scaffold on both reactivity and enantioselectivity of the reaction. Encouragingly, the P-chiral

monophosphorus ligand AntPhos<sup>11</sup> (**L7**) provided a moderate yield (60%) and an excellent ee (91% ee) (entry 7). Further study of the P-chiral phosphorus ligands developed in our laboratory showed that BI-DIME<sup>12</sup> (**L8**) provided an almost quantitative yield and a highest ee (94%) (entry 8). Screening of the solvent showed the reaction was facilitated with a nonpolar and non-coordinating solvent, as a diminished yield was observed in either toluene, THF or dioxane. No reaction was observed when dichloromethane was employed (entries 8-12). A Pd(0) precursor appeared to be advantageous for the reaction since a diminished yield was observed when Pd(OAc)<sub>2</sub> was applied (entry 13). Finally, the diboration was studied at a low catalytic loading (0.2 mol % Pd, 0.25 mol % **L8**) and at a gram scale (36 mmol **1a**, 4.8 g). The product **3a** (13.6 g) was obtained in 97% yield and in 94% ee (entry 14), demonstrating the practicality of this asymmetric transformation.

The substrate scope of this asymmetric diboration was then investigated. As depicted in Table 2, a series of diboronic esters with various electronic properties and substitution patterns on the benzene ring (**3b-p**) were smoothly formed at rt in excellent ee's and yields with Pd-**L8** as the catalyst. The enantioselectivities obtained were slightly higher with substrates having electron-donating substituents, but substituents such as fluoro (**3d**) and trifluoromethyl group (**3g**) were well applicable. Substrates with an *ortho* substituent (**3n-p**) were also tolerable. The reactions of both 1- and 2-naphthyl substrates provided excellent yields and ee's (**3q-r**). A chiral furyl product **3s** was also synthesized successfully. Substrates with multiple substituents on the benzene ring were equally effective for the transformation (**3t-u**). In order to test the chemoselectivity between an allene and an olefin, a substrate containing both moieties was subjected for diboration and only the allene moiety was reactive under the current conditions to form product **3v** in 99% yield and 96% ee. Both cyclic and heterocyclic substrates were applicable to smoothly afford **3w** and **3x**, respectively, in excellent yields and ee's. Switch of the methyl substituent with an ethyl group on allene resulted in an inferior ee (72% ee, **3y**). Introduction of a cyclopropyl group instead of the methyl substituent was less effective (**3z**). 1,1-Dialkylallenes were also applicable. While a moderate ee (67%) was obtained on product **3aa** bearing two primary alkyl substituents on the quaternary stereocenter, high ee's (87% and 91%) were obtained on products **3ab** and **3ac** containing both a methyl and a secondary alkyl groups at the chiral center. A good ee was also achieved on **3ad** bearing a tertiary alkyl group. Finally, the diboration of 1,1-diaryllallenes were studied. While a low ee was obtained on product **3ae** indicating little difference between a phenyl substituent and a *para*-tolyl group, a moderate ee (75%) was achieved on product **3af** bearing both a phenyl and an *ortho*-tolyl substituents at the quaternary stereocenter.

To understand the mechanism of this catalytic asymmetric reaction, the diboration of **1a** was investigated with a scalemic mixture of ligand **L8**. A perfect linear relationship between ee of **L8** and ee of product **3a** was observed, indicating the reaction catalyzed by a palladium catalyst composed of a



single monophosphorus ligand **L8**. Variation of the Pd:**L8** ratio from 1:1, 1:2, and 2:1 did not lead to a significant change of both the yield and the enantioselectivity, further demonstrating the


**Table 2.** Enantioselective Pd-catalyzed diboration of 1,1'-disubstituted allenes<sup>a</sup>

<b>3b</b> 91% yield, 93% ee	<b>3c</b> 90% yield, 94% ee	<b>3d</b> 98% yield, 90% ee	<b>3e</b> 93% yield, 92% ee
<b>3f</b> 85% yield, 90% ee	<b>3g</b> 94% yield, 87% ee	<b>3h</b> 93% yield, 90% ee	<b>3i</b> 93% yield, 91% ee
<b>3j</b> 91% yield, 92% ee	<b>3k</b> 97% yield, 92% ee	<b>3l</b> 80% yield, 86% ee	<b>3m</b> 95% yield, 93% ee
<b>3n</b> 91% yield, 92% ee	<b>3o</b> 95% yield, 94% ee	<b>3p</b> 90% yield, 94% ee	<b>3q</b> 98% yield, 94% ee
<b>3r</b> 86% yield, 91% ee	<b>3s</b> 93% yield, 88% ee	<b>3t</b> 86% yield, 90% ee	<b>3u</b> 98% yield, 94% ee
<b>3v</b> 99% yield, 96% ee	<b>3w<sup>c</sup></b> 98% yield, 91% ee	<b>3x<sup>c</sup></b> 96% yield, 90% ee	<b>3y<sup>d</sup></b> 76% yield, 72% ee
<b>3z<sup>c,d</sup></b> 28% yield, 34% ee	<b>3aa<sup>c</sup></b> 92% yield, 67% ee	<b>3ab<sup>c</sup></b> 95% yield, 87% ee	<b>3ac<sup>c</sup></b> 92% yield, 91% ee
<b>3ad<sup>c</sup></b> 81% yield, 89% ee	<b>3ae<sup>c</sup></b> 97% yield, 8% ee	<b>3af<sup>c</sup></b> 93% yield, 75% ee	<b>3f</b>

<sup>a</sup>Unless otherwise specified, the reactions were performed under nitrogen at rt for 24 h with **1** (0.20 mmol), **2** (0.24 mmol), **L8** (2.5 mol %) and Pd<sub>2</sub>(dba)<sub>3</sub> (1.0 mol %) in cyclohexane (2.0 mL). The yields of isolated products were shown here. The ee values were determined by HPLC on a chiral column. The *R* absolute configuration of **3f** was determined by X-ray crystallographic analysis,<sup>13</sup> the others were assigned by analogy. <sup>b</sup>Pd<sub>2</sub>(dba)<sub>3</sub> (2.0 mol %) and **L8** (5.0

mol %) were employed. <sup>c</sup>The absolute configurations were not determined. <sup>d</sup>Incomplete conversions.

composition of a single **L8** in the active palladium catalyst. To understand the perfect regioselectivity and the stereochemical model of this asymmetric diboration, energetics of the catalytic transformation was calculated at B3LYP-D3/6-31G(d)+SDD level. As shown in Figure 2a, the reaction initiates from the palladium species **RC**, which undergoes oxidative addition of bis(pinacolato)diboron concerted with allene insertion where the boryl group migrates to the middle carbon of the allene to give an η<sup>3</sup> Pd(II)-allyl intermediate **INT<sub>Sr</sub>** or **INT<sub>Re</sub>**. Notably, this initial oxidative boryl migration<sup>16</sup> is in well agreement with the perfect regioselectivity of diboration on the internal double bond of allene, and such process is an irreversible and stereo-determining step of the transformation.<sup>15</sup> **TS1<sub>Re</sub>** is computed to be lower in free energy than **TS1<sub>Sr</sub>** by 1.4 kcal/mol, in qualitative agreement with the observed enantioselectivity (ΔG<sub>exp</sub>: 2.1 kcal/mol).



**Figure 2.** (a) Free-energy profile (in kcal/mol) for the Pd-catalyzed asymmetric diboration of buta-2,3-dien-2-ylbenzene (**1a**) with (*S*)-BI-DME (**L8**) as the ligand at B3LYP-D3/6-31G(d)+SDD level.<sup>15</sup> (b) VDW representation of the optimized



## ARTICLE

## Journal Name


**TS1<sub>Re</sub>** and **TS1<sub>Si</sub>** with relative free and electronic energies in kcal/mol

Distortion/interaction analysis<sup>17</sup> reveals that there is a greater dispersion interaction between the phenyl group<sup>17b</sup> on the allene and one boryl group in **TS1<sub>Re</sub>** than **TS1<sub>Si</sub>**, which is considered as the key factor for the enantioselectivity (Figure 2b).<sup>15</sup> When excluding the dispersion contribution, the enantioselectivity is computed to be reversed. Finally, the reductive elimination via **TS2<sub>Re</sub>** proceeds from **INT<sub>Re</sub>** to produce **PC<sub>Re</sub>** with a barrier of about 14.9 kcal/mol. Apart from Morcken's proposal where the oxidative addition of diboron to Pd proceeded prior to migratory insertion and the oxidative addition of diboron to Pd was computed as the rate-determining step,<sup>9i</sup> our calculations on diboration of **1a** revealed a concerted mechanism of oxidative addition of bis(pinacolato)diboron and allene insertion, provided the first computational insight on the origins of the enantioselectivity (molecular details and a critical dispersion effect), and disclosed the final reductive elimination step as the rate-determining step.

The chiral diboronic ester products are versatile building blocks in organic synthesis. For example, Aggarwal reported a stereospecific allylation between the diboronic ester (**S**)-**3a** and benzaldehyde to form a tetrasubstituted alkene after a Suzuki-Miyaura cross-coupling.<sup>14,18,9k</sup> To further explore the synthetic applications of such chiral diboronic esters, an enantioselective synthesis of brassinazole,<sup>19</sup> a specific inhibitor of brassinosteroid biosynthesis, was studied with (**R**)-**3a** as the starting material (Scheme 1). Surprisingly, despite its significant biological properties, its enantioselective synthesis was not reported to our knowledge. We envisioned that the diboration product (**R**)-**3a** would provide a rapid and efficient access to brassinazole through simple transformations. Thus, (**R**)-**3a** was subjected to a one-pot, three step sequence: a) hydroboration by treatment with 9-BBN;<sup>20</sup> b) Suzuki-Miyaura coupling with 1-chloro-4-iodobenzene; and c) oxidation under conditions of H<sub>2</sub>O<sub>2</sub>/NaOH. The chiral diol **4** was formed smoothly at a *cis/trans* ratio of 4:1 in 79% overall yield. Under Ley's conditions,<sup>21</sup> the diol **4** was readily oxidized to form the corresponding sulfate **5**, which was isolated as a pure *cis* product in 78% yield. Finally, treatment of sulfate **5** with 1,2,4-triazole under conditions of NaH/DMF yielded (**2R,3R**)-brassinazole (**6**) in 80% yield, whose absolute configurations were confirmed by X-ray crystallography.<sup>13</sup> It should be noted that the sulfate **5** proceeded first through an intramolecular S<sub>N</sub>2 reaction to form an epoxide, which was subsequently attacked by 1,2,4-triazole to undergo a 2<sup>nd</sup> S<sub>N</sub>2 reaction, yielding product **6** with the net retention of stereochemistry. Compound **4** was also transformed to its diastereomer **7** through an oxidation-reduction procedure, which ultimately led to the formation of (**2R,3S**)-brassinazole (**8**) via sulfate formation and nucleophilic substitution. Thus, we accomplished a concise and first enantioselective synthesis of brassinazole.

## Conclusions

In summary, we have developed a practical and enantioselective palladium-catalyzed diboration of 1,1-disubstituted allenes which have led to the synthesis of a series of diboronic esters containing a chiral tertiary boronic ester in excellent yields and enantioselectivities with the palladium loading as low as 0.2 mol %. The reaction enjoys a broad substrate scope and good functional group compatibility. The chiral ligand BI-DIME has proven to be crucial for the success of the reaction. DFT calculations have identified a concerted mechanism of oxidative addition of bis(pinacolato)diboron and allene insertion, and revealed a critical dispersion effect on the origins of the enantioselectivity. Finally, the application of the chiral diboronic ester to a concise and first enantioselective synthesis of brassinazole has been successfully demonstrated.



**Scheme 1.** Asymmetric synthesis of brassinazole

## Acknowledgments

This work is supported by the Strategic Priority Research Program of the Chinese Academy of Sciences XDB20000000, NSFC-21432007, 21272254, 21572246, 21232009, 20421091, CAS (QZDY-SSW-SLH012, SLH029).

## Notes and references

- For reviews, see: (a) A. H. Cherney, N. T. Kadunce, S. E. Reisman, *Chem. Rev.* 2015, **115**, 9587. (b) D. Leonori, V. K. Aggarwal, *Angew. Chem. Int. Ed.* 2015, **54**, 1082. (c) H. K. Scott, V. K. Aggarwal, *Chem. Eur. J.* 2011, **17**, 13124.
- (a) D. Leonori, V. K. Aggarwal, *Acc. Chem. Res.* 2014, **47**, 3174. (b) J. L. Stymiest, V. Bagutski, R. M. French, V. K. Aggarwal, *Nature* 2008, **456**, 778. (c) V. Bagutski, R. M. French, V. K. Aggarwal, *Angew. Chem. Int. Ed.* 2010, **49**, 5142. (d) A. P. Pulis, D. J. Blair, E. Torres, V. K. Aggarwal, *J. Am. Chem. Soc.* 2013, **135**, 16054.
- (a) N. Hu, G. Zhao, Y. Zhang, X. Liu, G. Li, W. Tang, *J. Am. Chem. Soc.* 2015, **137**, 6746. (b) V. M. Shoba, N. C. Thacker,



- A. J. Bochat, J. M. Takacs, *Angew. Chem. Int. Ed.* 2016, **55**, 1465.
- 4 A. Guzman-Martinez, A. H. Hoveyda, *J. Am. Chem. Soc.* 2010, **132**, 10634.
- 5 (a) I-H. Chen, L. Yin, W. Itano, M. Kanai, M. Shibasaki, *J. Am. Chem. Soc.* 2009, **131**, 11664. (b) J. M. O'Brien, K.-S. Lee, A. H. Hoveyda, *J. Am. Chem. Soc.* 2010, **132**, 10630. (c) X. Feng, J. Yun, *Chem. Eur. J.* 2010, **16**, 13609. (d) S. Rodomkit, A. H. Hoveyda, *Angew. Chem. Int. Ed.* 2014, **53**, 3387.
- 6 A. S. Dudnik, G. C. Fu, *J. Am. Chem. Soc.* 2012, **134**, 10693.
- 7 For reviews of allenes, see: (a) J. Ye, S. Ma, *Acc. Chem. Res.* 2014, **47**, 989. (b) P. Koschker, B. Breit, *Acc. Chem. Res.* 2016, **49**, 1524. (c) E. Soriano, I. Fernández, *Chem. Soc. Rev.* 2014, **43**, 3041. (d) S. Yu, S. Ma, *Angew. Chem. Int. Ed.* 2012, **51**, 3074. (e) S. Ma, *Acc. Chem. Res.* 2009, **42**, 1679. (f) M. Jeganmohan, C.-H. Cheng, *Chem. Commun.* 2008, 3101. (g) S. Ma, *Chem. Rev.* 2005, **105**, 2829. (h) S. Ma, *Acc. Chem. Res.* 2003, **36**, 701. (i) R. Zimmer, C. U. Dinesh, E. Nandan, F. A. Khan, *Chem. Rev.* 2000, **100**, 3067. (j) Y. Yamamoto, U. Radhakrishnan, *Chem. Soc. Rev.* 1999, **28**, 199.
- 8 For reviews of diboration, see: (a) J. Takaya, N. Iwasawa, *ACS Catal.* 2012, **9**, 1993. (b) L. Dang, Z. Lin, T. B. Marder, *Chem. Commun.* 2009, 3987. (c) J. Ramírez, C. Lillo, A. M. Segarra, E. C. R. Fernández, *Chim.* 2007, **10**, 138. (d) H. E. Burks, J. P. Morken, *Chem. Commun.* 2007, 4717. (e) I. B. Beletskaya, C. Moberg, *Chem. Rev.* 2006, **106**, 2320. (f) T. Ishiyama, N. Miyaoura, *Chem. Rev.* 2004, **3**, 271. (g) T. B. Marder, N. C. Norman, *Top. Catal.* 1998, **5**, 63. (h) E. C. Neeve, S. J. Geier, I. A. I. Mkhalid, S. A. Westcott, T. B. Marder, *Chem. Rev.* 2016, **116**, 9091.
- 9 For asymmetric rhodium-catalyzed diborations: (a) J. B. Morgan, S. P. Miller, J. P. Morken, *J. Am. Chem. Soc.* 2003, **125**, 8702. (b) S. Trudeau, J. B. Morgan, M. Shrestha, J. P. Morken, *J. Org. Chem.* 2005, **70**, 9538. (c) S. P. Miller, J. B. Morgan, F. J. Nepveux, J. P. Morken, *Org. Lett.* 2004, **6**, 131. (d) K. Toribatake, H. Nishiyama, *Angew. Chem. Int. Ed.* 2013, **52**, 11011. For platinum-catalyzed diboration: (e) L. T. Kliman, S. N. Mlynarski, J. P. Morken, *J. Am. Chem. Soc.* 2009, **131**, 13210. (f) L. T. Kliman, S. N. Mlynarski, G. E. Ferris, J. P. Morken, *Angew. Chem. Int. Ed.* 2012, **51**, 521. (g) J. R. Coombs, D. Haefner, L. T. Kliman, J. P. Morken, *J. Am. Chem. Soc.* 2013, **135**, 11222. (h) S. N. Mlynarski, C. H. Schuster, J. P. Morken, *Nature* 2014, **505**, 386. For asymmetric palladium-catalyzed diboration: (i) X. Guo, A. K. Nelson, C. Slebodnick, W. L. Santos, *ACS Catal.* 2015, **5**, 2172. (j) N. F. Pelz, A. R. Woodward, H. E. Burks, J. D. Sieber, J. P. Morken, *J. Am. Chem. Soc.* 2004, **126**, 16328. (k) J. D. Sieber, J. P. Morken, *J. Am. Chem. Soc.* 2006, **128**, 74. (l) H. E. Burks, S. Liu, J. P. Morken, *J. Am. Chem. Soc.* 2007, **129**, 8766.
- 10 (a) Z.-Y. Cao, X. M. Wang, C. Tan, X.-L. Zhao, J. Zhou, K. Ding, *J. Am. Chem. Soc.* 2013, **135**, 8197. (b) X. M. Wang, P. Guo, Z. Han, X. B. Wang, Z. Wang, K. Ding, *J. Am. Chem. Soc.* 2014, **136**, 405. (c) X. M. Wang, F. Meng, Y. Wang, Z. Han, Y. Chen, L. Liu, Z. Wang, K. Ding, *Angew. Chem. Int. Ed.* 2012, **51**, 9276. (d) X. M. Wang, Z. Han, Z. Wang, K. Ding, *Angew. Chem. Int. Ed.* 2012, **51**, 936. (e) J. Liu, Z. Han, X. Wang, Z. Wang, K. Ding, *J. Am. Chem. Soc.* 2015, **137**, 15346.
- 11 (a) W. Fu, M. Nie, A. Wang, Z. Cao, W. Tang, *Angew. Chem. Int. Ed.* 2015, **54**, 2520. (b) N. Hu, K. Li, Z. Wang, W. Tang, *Angew. Chem. Int. Ed.* 2016, **55**, 5044. (c) G. Xu, Q. Zhao, W. Tang, *Chin. J. Org. Chem.* 2014, **34**, 1919. (d) C. Li, D. Chen, W. Tang, *Synlett* 2016, **27**, 2183.
- 12 (a) W. Tang, N. D. Patel, G. Xu, X. Xu, J. Savoie, S. Ma, M. -H. Hao, S. Keshipeddy, A. G. Capacci, X. Wei, Y. Zhang, J. J. Gao, W. Li, S. Rodriguez, B. Z. Lu, N. K. Yee, C. H. Senanayake, *Org. Lett.* 2012, **14**, 2258. (b) G. Xu, W. Fu, G. Liu, C. H. Senanayake, W. Tang, *J. Am. Chem. Soc.* 2014, **136**, 570. (c) K. Du, P. Guo, Y. Chen, Z. Cao, Z. Wang, W. Tang, *Angew. Chem. Int. Ed.* 2015, **54**, 3033. (d) M. Nie, W. Fu, Z. Cao, W. Tang, *Org. Chem. Front.* 2015, **2**, 1322.
- 13 CCDC 1517472 and 1517578 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html) (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB21EZ, UK; fax: (+44)1223-336-033; or [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).
- 14 M. J. Hesse, S. Essafi, C. G. Watson, J. N. Harvey, D. Hirst, C. L. Willis, V. K. Aggarwal, *Angew. Chem. Int. Ed.* 2014, **53**, 6145.
- 15 See Supporting Information for computational details, results and discussion.
- 16 X. Zhang, L. W. Chung, Y.-D. Wu, *Acc. Chem. Res.* 2016, **49**, 1302.
- 17 (a) C. Y. Legault, Y. Garcia, C. A. Merlic, K. N. Houk, *J. Am. Chem. Soc.* 2007, **129**, 12664. (b) A dispersion effect could also be applied to 1,1-dialkyl substrates **3ab**, **3ac**, and **3ad** for the observed high ees.
- 18 A. R. Woodward, H. E. Burks, L. M. Chan, J. P. Morken, *Org. Lett.* 2005, **7**, 5505.
- 19 (a) T. Asami, Y. K. Min, N. Nagata, K. Yamagishi, S. Takatsuto, S. Fujioka, N. Murofushi, I. Yamaguchi, S. Yoshida, *Plant Physiol.* 2000, **123**, 93. (b) S. Yoshida, A. Fujioka, T. Asami, *Jpn. Kokai Tokkyo Koho.* 2000, JP2000053657 (A). (c) Y. K. Min, T. Asami, S. Fujioka, N. Murofushi, E. Yamaguchi, S. Yoshida, *Bioorg. Med. Chem. Lett.* 1999, **9**, 425.
- 20 N. F. Pelz, J. P. Morken, *Org. Lett.* 2006, **8**, 4557.
- 21 (a) Y. Gao, Y. K. B. Sharpless, *J. Am. Chem. Soc.* 1988, **110**, 7538. (b) J. E. Baldwin, R. M. Adlington, D. W. Gollins, *Tetrahedron.* 1995, **51**, 5169.

