


 Cite this: *New J. Chem.*, 2018, 42, 1521

## Correction: Revisiting syntheses of Ti(IV)/H<sub>2</sub>PO<sub>4</sub>–HPO<sub>4</sub> functional ion-exchangers, properties and features

 Mylène Trublet,<sup>\*a</sup> Daniela Rusanova<sup>a</sup> and Oleg N. Antzutkin<sup>ab</sup>

DOI: 10.1039/c7nj90090b

 Correction for 'Revisiting syntheses of Ti(IV)/H<sub>2</sub>PO<sub>4</sub>–HPO<sub>4</sub> functional ion-exchangers, properties and features' by Mylène Trublet *et al.*, *New J. Chem.*, 2017, DOI: 10.1039/c7nj03065g.

rsc.li/njc

The authors would like to correct Table 1, as the initial TiO<sub>2</sub> content for samples SA3 and SA4 is incorrect in the published article. The correct Table 1 is shown below.

**Table 1** Synthesis conditions for TiP1 sorbents synthesized using different titanium sources

Ti source name	Source	Initial TiO <sub>2</sub> content (%)	State	[TiO <sub>2</sub> ] <sub>f</sub> (g L <sup>-1</sup> )	[H <sub>2</sub> SO <sub>4</sub> ] <sub>f</sub> (g L <sup>-1</sup> )	Yield <sup>a</sup> (%)
A	Apatity, Russia	~ 7	Liquid	62	405	89
SA1	Sigma Aldrich	~ 29	Powder	60	441	64
SA2	Sigma Aldrich	~ 29	Powder	66	397–410	65
SA3	Sigma Aldrich	~ 7	Liquid	76	394	71
SA4	Sigma Aldrich	~ 8	Liquid	110	356–409	95

[ ]<sub>f</sub>: concentration before adding H<sub>3</sub>PO<sub>4</sub> during the synthesis process.<sup>a</sup> Calculated based on the initial titanium concentration.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup> *Chemistry of Interfaces, Luleå University of Technology, 97187 Luleå, Sweden. E-mail: Mylene.Trublet@ltu.se; Tel: +46-(0)920-493448*
<sup>b</sup> *Department of Physics, Warwick University, CV47AL, Coventry, UK*
