

Cite this: *Chem. Sci.*, 2018, 9, 238Received 2nd October 2017
Accepted 23rd October 2017

DOI: 10.1039/c7sc04283c

rsc.li/chemical-science

Z-Selective iridium-catalyzed cross-coupling of allylic carbonates and α -diazo esters†

Bryce N. Thomas, Patrick J. Moon, Shengkang Yin, Alex Brown^{ID}
and Rylan J. Lundgren^{ID}*


A well-defined Ir-allyl complex catalyzes the Z-selective cross-coupling of allyl carbonates with α -aryl diazo esters. The process overrides the large thermodynamic preference for E-products typically observed in metal-mediated coupling reactions to enable the synthesis of Z,E-dieneoates in good yield with selectivities consistently approaching or greater than 90 : 10. This transformation represents the first productive merger of Ir-carbene and Ir-allyl species, which are commonly encountered intermediates in allylation and cyclopropanation/E-H insertion catalysis. Potentially reactive functional groups (aryl halides, ketones, nitriles, olefins, amines) are tolerated owing to the mildness of reaction conditions. Kinetic analysis of the reaction suggests oxidative addition of the allyl carbonate to an Ir-species is rate-determining. Mechanistic studies uncovered a pathway for catalyst activation mediated by NEt_3 .

Introduction


Metal-catalyzed cross-couplings involving carbene intermediates are a valuable set of transformations to generate new carbon-carbon bonds and molecular complexity from simple building blocks.^{1–3} A commonly exploited pathway to generate substituted olefins from carbene-precursors, such as α -diazo esters or N-tosylhydrazones, involves β -hydride elimination from a metal-alkyl species generated *via* a migratory insertion process. These reactions are initiated by the formation of a metal carbene intermediate, with the β -hydrogen containing substrate being either the cross-coupling partner (as depicted in Fig. 1a) or the carbene precursor.^{2b} This general concept has been used in a broad series of Pd-catalyzed coupling reactions of *in situ* generated carbenes with various reaction partners to generate polysubstituted E-alkenes and E,E-dienes. In these processes, the stereoselectivity of the olefination is controlled by the relative energetics of species leading to *syn* β -hydride elimination, as they are in classical Heck-type processes.⁴ Wang and co-workers have demonstrated the ability of allylic halides,⁵ allenes,⁶ vinyl boronic acids,⁷ and vinyl cyclopropanes⁸ to undergo interception with various metal carbene intermediates to generate E-dienes.⁹ While these are powerful synthetic transformations, control over the stereochemistry of the newly formed carbon-carbon double bond that overrides the inherent preference for E-alkene products remains an unaddressed challenge in cross-coupling methodology involving metal carbenes.

Inspired by the rapid development of Pd-, Cu-, and Rh-catalyzed carbene cross-coupling type reactions, we questioned whether alternative transition metals could afford unique reactivity and selectivity in alkene-generating processes. Given the widespread use of both Ir-allyl and Ir-carbene intermediates in allylic substitution and carbene insertion chemistry respectively (Fig. 1b), we sought to merge these species to enable a new reactivity mode in Ir-catalysis.¹⁰ Herein


a. Carbene Cross-Coupling Olefination



b. Common Intermediates in Ir-Catalysis



c. This Work: Ir-Catalyzed Carbene-Allyl Electrophile Cross-Coupling



- new reactivity mode for Ir/carbene intermediates
- Z-selectivity **increases** with R-size (ester)
- **reverses** large thermodynamic preference for E,E diene

Department of Chemistry, University of Alberta, Edmonton, Alberta, T6G 2G2, Canada. E-mail: rylan.lundgren@ualberta.ca

† Electronic supplementary information (ESI) available: Full procedures, computational details and characterization data. See DOI: 10.1039/c7sc04283c

Fig. 1 Pathway for metal-catalyzed olefination reactions of carbene precursors, common Ir-intermediates in catalysis and overview of the process reported herein to generate Z,E-dienes.

we report that a well-defined Ir-allyl complex catalyzes the *Z*-selective olefination of allyl carbonates with α -aryl diazo esters to provide *Z,E*-dienoates (Fig. 1c). Increasing the steric demand of the ester substituent of the diazo compound was found to enhance the *Z*-selectivity of the reaction, overcoming the large thermodynamic preference for the formation of the *E,E*-diene products typically generated by conventional catalytic methods.^{11–14} This transformation represents a rare example of a formal migratory insertion process of an Ir-allyl intermediate^{15,16} and proceeds by a catalyst activation step promoted by NEt₃ involving the generation of an allylic ammonium species. The method helps to address a methodological gap in classical olefination chemistry, as modified Horner–Wadsworth–Emmons processes fail to deliver bulky dienyl esters with appreciable selectivity.

Results and discussion

Reaction development


In the attempt to intercept an Ir-allyl intermediate with a carbene precursor, a range of metal catalysts and ligands were explored to promote coupling between an α -aryl diazo ester¹⁷ and an allylic electrophile. Table 1A provides a snapshot of the performance of various Ir- (and Rh-) complexes previously established as active allylation catalysts. Only the use of [Ir(COD)Cl]₂ and the phosphoramidite ligand **L1** as the catalyst mixture formed an appreciable amount of product (12%). Given reports by Hartwig and Helmchen demonstrating the benefits of allyl-ligated, ligand-cyclometalated cationic Ir species featuring **L1** in allylic substitution reactions,^{18–20} complex **1** was prepared and used as the catalyst in the carbene coupling reaction. Use of **1** resulted in a dramatic improvement in yield of the desired product (85%) in ~2 : 1 selectivity for the *Z,E*-diene isomer. Other related catalyst structures, including **L1**Ir(COD)Cl, cyclometalated Ir(*i*)**L1** complexes without an allyl ligand, and ligand-modified versions of the standard catalyst **1** proved considerably less reactive, as was a variant of Krische's Ir-SEGPHOS allylation catalyst.²¹ *In situ* generation of **1** with [Ir(COD)Cl]₂, **L1**, cinnamyl carbonate and AgClO₄ did not provide significant product. Diene products arising from insertion into a branched Ir-allyl species were not observed in any case.

Reaction parameters important to observe high yields, including solvent, catalyst loading, and concentration are shown in Table 1B (entries 1–5). NEt₃ is vital for product formation (entry 6); other inorganic and amine bases provided effectively no product (entry 7). Alternative leaving groups were inferior to *tert*-butyl carbonate, however a phosphonate ester substrate yielded product with a reversal in selectivity (~2 : 1 *E*-selectivity entry 8). Branched allylic carbonates could be used as substrates, but in general provided reduced yields (entry 10). The formation of the linear carbonate over the course of the reaction was observed with branched allylic starting materials. The use of corresponding methyl carbonate substrate led to methyl ether formation instead of diene.

Increasing the steric demand of the R-group on the α -aryl diazo ester led to a significant improvement in the *Z*-selectivity of the newly generated olefin (entries 11 and 12, up to 90 : 10

Table 1 A. Catalyst identification for the *Z*-selective cross-coupling of α -diazo esters and allylic carbonates. B. Effect of key reaction parameters and ester group on reactivity. See the ESI for additional optimization details^{a,b}

Reaction development A: catalyst identification



Reaction development B: effect of reaction parameters

Reaction scheme showing the cross-coupling of an α -aryl diazo ester (Ph-CO₂R) with an allylic carbonate (BocO-CH₂-CH=CH-Ph) to form a *Z,E*-dienoate (Ph-CO₂R-CH=CH-CH=CH-Ph). The reaction conditions are NEt₃ in DMA at room temperature. The catalyst is **1** (10%).

Entry	Deviation from above	Conv. (%)	Yield (%)	[<i>Z,E</i> : <i>E,E</i>]
1	None (R = Me)	100	85	[65:35]
2	THF instead of DMA	33	21	[57:43]
3	DCE instead of DMA	32	24	[54:46]
4	5% 1 instead of 10%	35	30	[62:38]
5	0.05 M instead of 0.2 M	58	48	[65:36]
6	No NEt ₃	<10	<2	[nd]
7	DBU or K ₂ CO ₃ instead of NEt ₃	<10	<2	[nd]
8	OP(O)(OEt) ₂ instead of OBoc	150	35	[30:70]
9	Cl instead of OBoc	150	7	[77:23]
10	Branched OBoc instead of linear	150	72	[64:36]
11	CO ₂ <i>t</i> Bu diazo instead of CO ₂ Me	98	85	[87:13]
12	CO ₂ CH(<i>i</i> Pr) ₂ diazo instead of CO ₂ Me	100	71	[90:10]


^a 1.5 equiv. cinnamylOBoc, 3.0 equiv. NEt₃, 10% [catalyst], 18 h. Conv. and yields determined by calibrated ¹H NMR, *Z,E*:*E,E* = ~65 : 35, see ESI for full details. ^b 1.5 equiv. cinnamylOBoc, 3.0 equiv. NEt₃, 18–24 h. conv. (based on 150 mol% electrophile) and yields determined by ¹H NMR calibrated with an internal standard, see ESI for full details, bolded entries are isolated yields.

selectivity). The observation of *Z,E*-product formation is particularly remarkable in consideration of the large thermodynamic preference for the *E,E* isomer over the *Z,E* product, with the calculated relative $\Delta G = 6.5$ kJ mol^{–1} at 25 °C; *Z,E* : *E,E* ~ 5 : 95 for product **2** where R = *t*-Bu (see ESI† for details, the thermodynamic ratio is similar for the corresponding methyl ester). These results are in direct contrast to the well-established preference for related Pd-catalyzed reactions to generate *E*-products *via* β -hydride elimination processes.^{4,5,9}



In light of the scarcity of reports concerning the *Z*-selective olefination of carbonyls with α -aryl acetate-derived phosphates,^{13,22} the venerable Still–Gennari modification²³ of the Horner–Wadsworth–Emmons reaction was explored in an attempt to generate similar *Z,E*-diene products. Olefination of cinnamaldehyde with a bulky bis(trifluoroethyl)phosphonate reagent resulted in formation of diene **2** with poor selectivity (*Z,E* : *E,E* = 42 : 58), further highlighting the difficulty associated with the stereo-controlled preparation of this class of diene (eqn (1)).²⁴

Comparison to *Z*-Selective Olefination Method [Still–Gennari]



Reaction scope and limitations

The Ir-catalyzed carbene cross-coupling reaction provides synthetically useful yields and good *Z*-selectivities across a range of aryl-substituted allylic carbonates with bulky α -diazo esters (Table 2). The reaction tolerates halogen substitution at the 2, 3, and 4-positions of cinnamyl derivatives, including potentially reactive aryl bromide groups, and proceeds with similar effectiveness for electron-rich and electron-poor aryl allylic carbonates. A dienyl electrophile substrate delivered the desired *Z,E,E* triene with excellent *Z*-selectivity (**12**, 95 : 5). Halogen groups were also tolerated on the α -aryl diazo ester partner, as were electron-withdrawing groups (keto and cyano). A notable difference in selectivity was observed when electron-withdrawing groups (keto and cyano) are placed at either the 4-position (**15**, 95 : 5 *Z,E* : *E,E*) versus the 3-position (**22**, 63 : 37 *Z,E* : *E,E*). Less successful substrates include alkyl-substituted allylic carbonates (**13**) and thienyl α -diazo esters (**23**, see the ESI† for additional examples).

Variation of the ester group clearly demonstrated that an increase in the size of the alkyl group on the ester increased *Z*-selectivity, with isopropyl (**24**), diphenylmethyl (**25**), 4-*N*-Boc piperidinyll (**26**), or 1-adamantyl (**27**) groups giving better selectivity compared to methyl (Table 1B, entry 1), but less than that of the *tert*-butyl or diisopropylmethyl ester derivatives.

The selective generation of *Z,E*-dienyl ester products affords a useful synthetic building block, primed for further functionalization. For example, ester reduction with DIBAL-H, Sharpless' asymmetric dihydroxylation followed by lactonization, and regioselective Rh-catalyzed olefin reduction can each be used to obtain an array of new products in which the *Z*-olefin unit remains intact or directs reactivity (Fig. 2).

Stoichiometric and kinetic studies

While we do not have a complete mechanistic understanding of the Ir-catalyzed olefination process at this time, stoichiometric and kinetic experiments have provided valuable mechanistic insights. In contrast to Pd-catalyzed reactions of allyl

Table 2 Scope of the Ir-catalyzed cross-coupling of α -aryl diazo esters and allylic carbonates^a

Variation of electrophile aryl [R = <i>t</i> Bu or CH(<i>i</i> -Pr) ₂ , Ar = Ph]			
2 <i>t</i> -Bu 85% [87:13] 3 CH(<i>i</i> -Pr) ₂ 71% [90:10] [1 mmol scale: 74%] ^[b]	4 <i>t</i> -Bu 54% [92:8] ^[c] 5 CH(<i>i</i> -Pr) ₂ 48% [94:6]	6 <i>t</i> -Bu 63% [85:15] 7 CH(<i>i</i> -Pr) ₂ 70% [90:10]	
8 <i>t</i> -Bu 78% [82:18] 11 <i>t</i> -Bu 69% [85:15] ^[d]	9 <i>t</i> -Bu 50% [84:16] ^[e] 12 <i>t</i> -Bu 54% [>95:5] ^[e]	10 <i>t</i> -Bu 74% [81:19] 13 <i>t</i> -Bu <20%	
Variation of diazo aryl [R = <i>t</i> Bu, Ar' = Ph]			
14 95% [87:13] 18 60% [89:11] 19 80% [90:10] ^[b,f]	15 55% [95:5] 20 77% [86:14] 21 75% [84:16] ^[f]	16 89% [91:9] ^[g] 22 73% [63:37] ^[c,e]	17 80% [80:20] ^[f] 23 <10%
Variation of diazo ester [Ar' = Ph, Ar' = Ph]			
24 84% [79:21] 25 75% [84:16]	26 74% [76:24] 27 63% [84:16]		

^a 1.5 equiv. allylic carbonate, 0.3–0.4 M 16–28 h, yields are of combined diene isomers with *Z,E/E,E* indicated in brackets. ^b Pure *Z,E*-product, 7.5 mol% **1**. ^c Crude ratio, isolated >95.5. ^d At 35 °C. ^e Determined by ¹H NMR. ^f 15 mol% **1**. ^g R = CH(*i*Pr)₂.

electrophiles and α -diazo esters,⁵ direct reaction between an Ir-allyl species and the carbene precursor is not observed (Fig. 3a). When **1** is added to 1 or 10 equivalents of α -diazo ester,




Fig. 2 Use of *Z,E*-dienoate **3** as a precursor for selective transformations. (a) DIBAL-H (2.5 equiv.) CH₂Cl₂, rt (b) AD-mix, MeSO₂NH₂, *t*-BuOH/H₂O, rt (c) 2.5 mol% [Rh(COD)Cl]₂, 15 mol% PPh₃, HCO₂H/NEt₃ (5 : 2), MeCN, 35 °C.


unreacted starting materials are recovered. The same reaction conducted in the presence of NEt_3 leads to decomposition of **1** to a complex mixture and <5% product formation. In the absence of α -diazo ester, NEt_3 promotes allyl exchange between complex **1** and allylic carbonate **4s**, concurrently generating allylic ammonium species (Fig. 3b).

We have observed *N*-allyl ammonium species by ^1H NMR to form rapidly under the standard reaction conditions. Control experiments demonstrated that the allylic ammonium species is not converted into diene product under the standard conditions.

Taken together, these experiments suggest that the role of NEt_3 is to activate Ir-allyl species **1** toward Ir-carbene formation by generating an open coordination site, likely the first mechanistic step in the product forming cycle (Fig. 3c). Notably, bulkier amine bases (DIPEA, $\text{N}(i\text{-Bu})_3$) fail to generate allylic ammonium species under the standard conditions and product is not observed when using these bases.²⁵

In a reaction using the aryl fluoride allyl carbonate **4s** at high catalyst loading (20 mol% **1**), the catalyst cinnamyl cross-over product is observed in 5% yield along with 85% product from **4s** (Fig. 4a), supporting the series of catalyst activation steps and allylic exchange processes in Fig. 2c.²⁶ Stoichiometric reactions using 1 equivalent of **1** mirror these observations, as significant crossover product is observed between the catalyst bound allyl group (20% yield) in comparison to the allyl unit originating from the carbonate substrate (35% yield). The two diene products are generated concurrently (Fig. 4b). Collectively, these results suggest allyl exchange processes at Ir occur on the time

a. Product Distribution at High Catalyst Loadings



b. Product Distribution at 100 mol% Catalyst Loading






Fig. 4 Ir-allyl/allyl carbonate cross-over experiments. At high catalyst loadings (a), small amounts of Ir-allyl crossover diene product is observed, which increases at 100 mol% catalyst (b).

a. NEt_3 and External Electrophile Required for Reaction



b. NEt_3 Mediates Allyl Exchange on Ir



c. Catalyst Activation and Allyl Carbonate Exchange




Fig. 3 Mechanistic aspects of the Ir-catalyzed cross-coupling of α -aryl diazo esters and allylic carbonates based on stoichiometric reactions (steps drawn as irreversible for clarity).

scale with product forming steps involving a putative Ir-carbene intermediate.

Reaction progress kinetic analysis (RPKA) was used to gain additional insights into the nature of the mechanism.²⁷ A representative RPKA plot is shown in Fig. 5. The rapid formation of allylic ammonium **D** is observed with concurrent depletion of complex **1** (inset) in line with the observations that NEt_3 acts to activate Ir-precatalysts with bound allyl fragments. The amount of allylic ammonium **D** increases slowly after the initial burst, indicative of regeneration of Ir-allyl species in relatively low concentration. Attempts to characterize Ir-based species in solution during catalysis by ^{31}P NMR were not successful.

Using variable time normalization plots,²⁸ the rate law of the reaction was determined to be zero order in α -diazo ester and




Fig. 5 Representative kinetic profile of the Ir-catalyzed coupling of α -aryl diazo esters and allylic carbonates [$\text{R} = \text{CH}(i\text{-Pr})_2$].




Fig. 6 Rate law determination by variable time normalization plots for the Ir-catalyzed coupling of α -diazo esters and allylic carbonates.

NEt_3 , while partial positive order (~ 0.5) in allyl carbonate (Fig. 6, see Fig S2–S4 in ESI† for additional plots and discussion). The order in allylic carbonate could be rationalized by the electrophile being involved in a rate-determining product forming step, but also causing the regeneration of off-cycle Ir-allyl species similar to **1** (as in Fig. 3c); the slow growth of allylic ammonium salt (Fig. 5) also suggests this.

The reaction was observed to be first-order in catalyst **1** at early time points, but deviated over the course of the reaction (see Fig. S5 in the ESI†). Catalyst stability was examined by interrogating temporal concentration profiles for the process at varied initial concentrations of α -diazo ester (**A**) and diene product (**3**, Fig. 7). For reactions conducted with half the typical concentration of α -diazo ester, a lack of overlay in the time-shifted profile with the standard reaction conditions is observed (the reaction proceeds faster), indicating catalyst deactivation or substrate inhibition. Reactions conducted with an initial product concentration of 0.09 M (corresponding to $\sim 35\%$ diene formation) to mimic the time-adjusted experiment




Fig. 7 Kinetic profile for the Ir-catalyzed coupling of α -diazo esters and allylic carbonates employing varying initial concentrations of α -diazo ester (**A**), and product (**3**).




Fig. 8 Potential mechanism for the Ir-catalyzed cross-coupling of α -aryl diazo esters and allylic carbonates.

resulted in direct overlay with the same experiment without added product, ruling out catalyst inhibition by diene product.

Given these observations, we propose a potential mechanism for the Ir-catalyzed cross-coupling of allylic carbonates and α -aryl diazo esters in which NEt_3 acts as a nucleophile to activate the pre-catalyst **1** (Fig. 8). Activation generates a coordinatively unsaturated Ir species that can engage the diazo compound to generate an Ir-carbene or react with allylic carbonate to form off-cycle species (**1'** in Fig. 2). The carbene intermediate could undergo rate-determining oxidative addition of the allylic carbonate to generate an η^1 -allyl/carbene intermediate. Migratory insertion followed by β -hydride elimination generates the diene product and liberates free catalyst. *Z*-selectivity could arise from the large coordination sphere of the catalyst to influence elimination from an otherwise unfavorable conformer, or alternatively base-induced anti-elimination could occur. The allylic leaving group impacts *Z*-selectivity (Table 1B), indicative that the resulting *tert*-butoxide or diethyl phosphate anion may play a role in selectivity determining elimination in a potential E2-type pathway.

Conclusions

In summary, we have reported the development and mechanistic features of an Ir-catalyzed cross-coupling reaction between allylic carbonates and α -aryl diazo esters to form substituted *Z,E*-dienoates. *Z*- α -aryl acrylates with sterically demanding ester groups can be generated with good selectivity, suggesting this new reactivity mode may serve as a stereo-selective complement to other metal-catalyzed carbene cross-coupling reactions. The *Z,E*-diene products are not readily accessible *via* traditional modifications of carbonyl olefination methods. Use of an Ir-allyl (pre)catalyst containing a cyclo-metallated ligand is essential for product formation which undergoes activation with amine base. Allylic carbonate oxidative addition appears to be the rate determining step, while catalyst deactivation processes help to explain the necessity for relatively high catalyst loadings. While the exact origin of *Z*-



selectivity is not well understood at this stage, the steric demand of the ester group and the nature of the allylic leaving group, which likely plays a role as the terminal base, significantly influence the selectivity of the bond-forming process. We believe these studies set the stage for future development in the area of Ir-catalyzed carbene coupling reactions and may have broader implications in designing Z-selective Heck-type reactions.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We acknowledge the University of Alberta and NSERC Canada for support (Discovery Grants to R. J. L. and A. B., CGS-D fellowship for P. J. M.). Computational work enabled through the support of Westgrid (<http://www.westgrid.ca>) and Compute Canada (<http://www.computeCanada.ca>). Shoshana Bachman (Caltech) is acknowledged for helpful discussions.

Notes and references

- Pioneering studies: (a) K. L. Greenman, D. S. Carter and D. L. Van Vranken, *Tetrahedron*, 2001, **57**, 5219–5225; (b) K. L. Greenman and D. L. Van Vranken, *Tetrahedron*, 2005, **61**, 6438–6441; (c) J. Barluenga, P. Moriel, C. Valdes and F. Aznar, *Angew. Chem., Int. Ed.*, 2007, **46**, 5587–5590; (d) C. Peng, Y. Wang and J. Wang, *J. Am. Chem. Soc.*, 2008, **130**, 1566–1567.
- Reviews: (a) J. Barluenga and C. Valdes, *Angew. Chem., Int. Ed.*, 2011, **50**, 7486–7500; (b) Q. Xiao, Y. Zhang and J. Wang, *Acc. Chem. Res.*, 2013, **46**, 236–247.
- Selected recent examples: (a) G. P. Qin, L. X. Li, J. W. Li and H. M. Huang, *J. Am. Chem. Soc.*, 2015, **137**, 12490–12493; (b) W. D. Chu, L. Zhang, Z. K. Zhang, Q. Zhou, F. Y. Mo, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, 2016, **138**, 14558–14561; (c) J. Feng, B. Li, Y. He and Z. H. Gu, *Angew. Chem., Int. Ed.*, 2016, **55**, 2186–2190; (d) A. Gutierrez-Bonet, F. Julia-Hernandez, B. de Luis and R. Martin, *J. Am. Chem. Soc.*, 2016, **138**, 6384–6387; (e) Y. Y. Li, Z. S. Qi, H. Wang, X. F. Yang and X. W. Li, *Angew. Chem., Int. Ed.*, 2016, **55**, 11877–11881; (f) A. Das, D. Wang, M.-C. Belhomme and K. Szabo, *Org. Lett.*, 2015, **17**, 4754–4757; (g) D. Wang and K. J. Szabo, *Org. Lett.*, 2017, **19**, 1622–1625; (h) B. H. Ye and N. Cramer, *Angew. Chem., Int. Ed.*, 2014, **53**, 7896–7899.
- I. P. Beletskaya and A. V. Cheprakov, *Chem. Rev.*, 2000, **100**, 3009–3066.
- (a) S. F. Chen and J. Wang, *Chem. Commun.*, 2008, 4198–4200; (b) K. Wang, S. F. Chen, H. Zhang, S. Xu, F. Ye, Y. Zhang and J. Wang, *Org. Biomol. Chem.*, 2016, **14**, 3809–3820.
- Q. Xiao, B. L. Wang, L. M. Tian, Y. Yang, J. Ma, Y. Zhang, S. F. Chen and J. Wang, *Angew. Chem., Int. Ed.*, 2013, **52**, 9305–9308.
- Y. M. Xia, Y. Xia, Z. Liu, Y. Zhang and J. Wang, *J. Org. Chem.*, 2014, **79**, 7711–7717.
- S. Feng, F. Mo, Y. Xia, Z. Liu, Z. Liu, Y. Zhang and J. Wang, *Angew. Chem., Int. Ed.*, 2016, **55**, 15401–15405.
- For the Pd-catalyzed, E-selective allylic C–H olefination with α -diazo esters see: P. S. Wang, H. C. Lin, X. L. Zhou and L. Z. Gong, *Org. Lett.*, 2014, **16**, 3332–3335.
- (a) Ir-catalyzed allylic substitution: J. F. Hartwig and L. M. Stanley, *Acc. Chem. Res.*, 2010, **43**, 1461–1475; (b) G. Helmchen, in *Iridium Complexes in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2009, pp. 211–250; (c) W. B. Liu, J. D. Xia and S. L. You, *Top. Organomet. Chem.*, 2012, **38**, 155–208; (d) Ir-catalyzed cyclopropanation: H. Suematsu, S. Kanchiku, T. Uchida and T. Katsuki, *J. Am. Chem. Soc.*, 2008, **130**, 10327–10337; (e) Ir-catalyzed C–H insertion: H. Suematsu and T. Katsuki, *J. Am. Chem. Soc.*, 2009, **131**, 14218–14219. For the Ir(III) catalyzed insertion of carbenes into aryl C–H bonds mediated by ortho-directing groups see: (f) W. Ai, X. Y. Yang, Y. X. Wu, X. Wang, Y. C. Li, Y. X. Yang and B. Zhou, *Chem.-Eur. J.*, 2014, **20**, 17653–17657; (g) Y. Xia, Z. Liu, S. Feng, Y. Zhang and J. Wang, *J. Org. Chem.*, 2015, **80**, 223–236.
- Z-selective Au-catalyzed cross-diazo ester coupling: D. M. Zhang, G. Y. Xu, D. Ding, C. H. Zhu, J. Li and J. T. Sun, *Angew. Chem., Int. Ed.*, 2014, **53**, 11070–11074.
- Z-selective Au-catalyzed diazo coupling with enolates: F.-M. Liao, Z.-Y. Cao, J.-S. Yu and J. Zhou, *Angew. Chem., Int. Ed.*, 2017, **56**, 2459–2463.
- For a general overview of Z-olefin synthesis see: W. Y. Siau, Y. Zhang, Y. Zhao in *Stereoselective Alkene Synthesis*, ed. J. Wang, Springer-Verlag Berlin, Berlin, vol. 327, 2012, pp. 33–58.
- Recent methods that enable Z,E-diene synthesis: (a) X. H. Hu, J. Zhang, X. F. Yang, Y. H. Xu and T. P. Loh, *J. Am. Chem. Soc.*, 2015, **137**, 3169–3172; (b) E. C. Yu, B. M. Johnson, E. M. Townsend, R. R. Schrock and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2016, **55**, 13210–13214; (c) S. X. Luo, J. S. Cannon, B. H. L. Taylor, K. M. Engle, K. N. Houk and R. H. Grubbs, *J. Am. Chem. Soc.*, 2016, **138**, 14039–14046; (d) C. Souris, A. Misale, Y. Chen, M. Luparia and N. Maulide, *Org. Lett.*, 2015, **17**, 4486–4489; (e) R. K. Feng, W. L. Yu, K. Wang, Z. X. Liu and Y. H. Zhang, *Adv. Synth. Catal.*, 2014, **356**, 1501–1508; (f) F. F. Li, C. B. Yu, J. Zhang and G. F. Zhong, *Org. Lett.*, 2016, **18**, 4582–4585.
- A rare case of migratory insertion of a Ir-allyl species with CO gives linear product: R. Takeuchi and Y. Akiyama, *J. Organomet. Chem.*, 2002, **651**, 137–145.
- For the Ir-catalyzed allylation of 2-aminostyrenes to form 1,4-dienes see: (a) H. He, W. Liu, L. Dai and S. L. You, *J. Am. Chem. Soc.*, 2009, **131**, 8346–8347; (b) K. Ye, H. He, W. Liu, L. Dai, G. Helmchen and S. L. You, *J. Am. Chem. Soc.*, 2011, **133**, 19006–19014.
- H. M. L. Davies and J. R. Manning, *Nature*, 2008, **451**, 417–424.



- 18 C. A. Kiener, C. T. Shu, C. Incarvito and J. F. Hartwig, *J. Am. Chem. Soc.*, 2003, **125**, 14272–14273.
- 19 G. Lipowsky, N. Miller and G. Helmchen, *Angew. Chem., Int. Ed.*, 2004, **43**, 4595–4597.
- 20 Notes: (a) the catalyst can be prepared in one-step and is readily purified by filtration in air; no counter ion effects were observed; ; (b) No product is formed in the absence of catalyst or base, see the ESI† for more details; ; (c) see ESI† for additional details concerning solvent, concentration, and other non-productive electrophiles and diazo substrates.
- 21 Y. J. Zhang, J. H. Yang, S. H. Kim and M. J. Krische, *J. Am. Chem. Soc.*, 2010, **132**, 4562–4563.
- 22 *Modern Carbonyl Olefination*, ed. T. Takeda, Wiley-VCH, Weinheim, Germany, 2004.
- 23 (a) W. C. Still and C. Gennari, *Tetrahedron Lett.*, 1983, **24**, 4405–4408. The Still-Gennari modification provides exceptional selectivity for Z-olefinations with alkyl-substituted α -phosphonate esters, aryl substituted variants have not been reported in detail. For an example of aryl aldehyde olefination using aryl-substituted α -phosphonate esters with modest Z-selectivity (3 : 1–4 : 1) see: (b) C. M. Mapes, N. S. Mani, X. Deng, C. R. Pandit, K. J. McClure, M. C. W. Pippel, C. A. Schon, L. Gomez, S. Shinde, J. G. Breitenbucher and R. K. Jones, *J. Org. Chem.*, 2010, **75**, 7950–7953.
- 24 Reaction under alternative conditions (silylamide base at -78°C in THF) provided similar selectivities [55 : 45].
- 25 Reactions with catalytic NEt_3 (30 mol%) provide nearly identical results to reactions conducted with 3 equivalents. High yields of allylic ammonium were observed with smaller *N*-methylpyrrolidine (>90% after 1 h). No amination of the allylic carbonate occurs in the absence of catalyst. It remains unclear as to why only linear ammonium species are observed, the branched cinnamyl ammonium could be generated first then undergo rapid isomerization. To the best of our knowledge, the observation of Ir-catalyzed allylation of tertiary amines has not been reported. The regiochemical outcome of the reaction suggests an η -1 coordination mode, however we do not have firm evidence against η -cinnamyl intermediates. For a rare example of Pd-catalyzed intermolecular allylation of tertiary amines followed by [2,3] Stevens rearrangement see: A. Soheili and U. K. Tambar, *J. Am. Chem. Soc.*, 2011, **133**, 12956–12959.
- 26 Carbonate cross-over in Ir-catalyzed allylation: O. V. Singh and H. Han, *J. Am. Chem. Soc.*, 2007, **129**, 774–775.
- 27 D. G. Blackmond, *J. Am. Chem. Soc.*, 2015, **137**, 10852–10866.
- 28 J. Burés, *Angew. Chem., Int. Ed.*, 2016, **55**, 2028–2031.

