

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: C. Pan, W. Guo and Z. Gu, *Chem. Sci.*, 2018, DOI: 10.1039/C8SC01657G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Unusual Biaryl Torsional Strain Promotes Reactivity in Cu-Catalyzed Sommelet-Hauser Rearrangement

Chongqing Pan, Wenjing Guo and Zhenhua Gu *

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


A Cu-catalyzed Sommelet-Hauser dearomatization of dihydrophenanthridine and diazo compounds is reported for the synthesis of spiro-indolines. A spiro-structure with adjacent quaternary and tertiary carbon centers was constructed in one step as a single isomer. Increasing steric hindrance by introducing *ortho*-substituents dramatically improved substrate reactivity in this transformation.

Introduction


Dearomatization is a robust method for construction of polyfunctionalized cyclic compounds; it can generate quaternary carbon centers and multi-substituted C=C double bonds in one or two steps. Due to high stabilization of aromaticity, the dearomatization reactions are traditionally performed under harsh conditions, such as oxidation in the presence of hypervalent iodine,¹ Birch reduction,² enzymatic conditions,³ or other conditions.⁴ Dearomatization has recently been achieved using photochemistry and transition metal catalysis. In particular, catalytic asymmetric dearomatization (CADA) has attracted substantial attention, such as from the You group, for constructing optically pure molecules from arenes.⁵

The Sommelet-Hauser reaction is an intramolecular rearrangement of benzyl ammonium salts to give *ortho*-substituted benzylamines (Scheme 1a, **path A**).⁶⁻⁸ Sommelet-Hauser dearomatization can efficiently generate valuable three-dimensional compounds bearing quaternary carbon centers if the two *ortho* positions have non-hydrogen substitutions (Scheme 1b). In a variation known as thio-Sommelet rearrangement, dearomatized compounds bearing thioether functionalities can be prepared by replacing the ammonium ylides with sulfur ylides.⁹⁻¹⁰


a) Sommelet-Hauser Rearrangement and Stevens Rearrangement




b) Sommelet-Hauser Rearrangement for Construction of Quaternary Carbon Center




c) Sommelet-Hauser Rearrangement for Construction of Spiro-Compounds



- # Limited Substrate Scope
- # Pre-synthesis of ammonium salts
- # Poor yields, poor selectivity



d) This Work: Cu-Catalyzed Sommelet-Hauser Rearrangement



- § Broad Substrate Scope
- § Highly Efficient Spiro-compound Synthesis; Excellent Selectivity
- § Molecular Torsional Strain Increasing the Reactivity

Scheme 1. The Sommelet-Hauser Rearrangement

A challenge to exploiting the Sommelet-Hauser reaction is preventing competitive Stevens rearrangement,¹¹⁻¹² in which formal intramolecular nucleophilic attack at the benzylic

Department of Chemistry, Center for Excellence in Molecular Synthesis, and Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, 96 Jinzhai Road, Hefei, Anhui 230026 (China)
E-mail: zhgu@ustc.edu.cn

*Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



position affords 2-phenylethanamine analogues (Scheme 1a, path B). Singleton has argued that these two types of rearrangement compete dynamically because they can occur via the same transition state.¹³

Sommelet-Hauser rearrangement can be used to produce spiro-compounds from benzocyclic ammonium salts, but the reported reactions usually give low yields and wide product distribution.¹⁴ For example, the reaction usually produces two regioisomers together with ring-enlarged products from Stevens rearrangement (Scheme 1c). Ring strain in seven- and eight-membered structures lowers the rearrangement rate, significantly increasing the yields of desilylation by-products C. In addition, the requirement for pre-synthesized ammonium salts considerably limits the substrate scope of the reaction. Third, the Sommelet-Hauser rearrangement appears incompatible with ammonium salts with fewer than seven members. As a result, the reaction cannot generate rings smaller than six-membered *N*-heterocycles (Scheme 1c).

Herein we report the copper-catalyzed generation of nitrogen ylide intermediates capable of Sommelet-Hauser rearrangement, in which *N*-containing six-membered rings efficiently participate. We further report that molecular torsional strain in biaryl substrates increases their reactivity in this rearrangement.

We wondered whether we could adapt dearomatization via Sommelet-Hauser rearrangement to generate the important spirocyclohexaneindoline structural motif, found in several natural products including gelsemine,¹⁵ 21-oxogelsemine¹⁶ and koumine.¹⁷ It also comprises the core structural unit of various biologically active molecules.¹⁸

Results and discussion

Optimization

We were unable to dearomatize biaryl compound **1a** using diazo compound **2a** in the presence of silver, rhodium or gold salts as catalysts, which are frequent catalysts in reactions involving carbene (Table 1, entries 1-3). We were pleased to find that **1a** reacted with **2a** in the presence of Cu(OTf) \cdot 0.5PhH at 90 °C to give 27% of spiro product **3a** as a single isomer, along with a small amount of **4a** (entry 4). The reaction generated a spiro-structure with adjacent quaternary and tertiary carbon centers in one




Figure 1. Natural Products Containing a Spirocyclohexaneindoline Skeleton

Table 1. Optimization of Reaction Conditions [a]

entry	catalyst	solvent	T/°C	Yield of 3a /%	Yield of 4a /%
1	Rh ₂ (OAc) ₄ (5 mol%)	toluene	90	< 5	-
2	(Ph ₃ P)AuCl	toluene	90	- ^[b]	-
3	AgOTf	toluene	90	NR	-
4	Cu(OTf) \cdot 0.5PhH	toluene	90	27	< 5
5	Cu(OTf) ₂	toluene	90	NR	-
6	Cu(MeCN) ₄ PF ₆	toluene	90	32	< 5
7	CuTC	toluene	90	47	< 5
8	Cu(acac) ₂	toluene	90	63	< 5
9	Cu(hfacac) ₂	toluene	90	< 5	-
10	Cu(acac) ₂	toluene	50	72	-
11	Cu(acac) ₂	MTBE	50	83	-
12	Cu(acac) ₂	DCE	50	53	-
13	Cu(acac) ₂	THF	50	72	-
14	Cu(acac) ₂ (5 mol %)	MTBE	50	79	-


[a] Unless stated otherwise, the reaction was conducted with **1a** (0.10 mmol), **2a** (0.20 mmol), and metal catalyst (10 mol%). [b] An unidentified product was formed.

step. Screening of copper salts showed Cu(acac)₂ to perform the best in terms of yield and reaction time (entries 5-9). Lowering the reaction temperature increased selectivity for **3a** over the Stevens rearrangement product **4a**, improving the isolated yield of **3a** to 72% (entry 10). Solvent screening identified methyl *tert*-butyl ether (MTBE) as the best (entries 11-13). The reaction proceeded smoothly with only 5 mol% of Cu(acac)₂, albeit with slightly lower yield (entry 14).

Substrate scope

With the optimal conditions in hand, we tested the generality of this Cu(acac)₂-catalyzed Sommelet-Hauser rearrangement. Diazo compounds with different ester groups, such as methyl, *tert*-butyl and substituted benzyl esters, reacted smoothly (**3b-f**). A diazo substrate with an electron-withdrawing group (CF₃) led to slightly lower yield (**3e**). The reaction tolerated additional methyl groups, a fluorine atom and methoxy groups in the biaryl aniline skeleton (**3g-3j**, **3t**). Naphthalene-based substrates and a binaphthalene derivative reacted uneventfully to give the spiro compounds in decent to excellent yields (**3k-3r**, **3t-3z** and **3bb**). Stereoinduction did not occur with the (*R*)-1-phenylethanol-derived diazo compound, such that a pair of diastereoisomers were isolated in a 1:1 ratio (**3t**). This rearrangement was compatible with ketones and amides, which afforded the desired products in decent to good yields (**3u-3x**). Introducing a methyl group adjacent to the C-N bond completely inhibited this transformation (**3cc**). The reaction tolerated an aryl-Br bond. Scale-up also proved possible, with



Table 2. Substrate Scope^[a]


[a] The reaction was conducted with **1** (0.10 mmol), diazo compound **2** (0.20 mmol), and Cu(acac)₂ (0.005 mmol, 5 mol%) in MTBE at 50 °C. [b] The reactions were performed in toluene.

compound **3x** isolated in 69% yield on the 1.0-mmol scale.


The two *ortho*-substituents (R¹ and R² in Table 2) strongly influenced the efficiency of the Sommelet-Hauser reaction and selectivity for this type of rearrangement over Stevens rearrangement (Scheme 2). For example, under the standard conditions at 50–90 °C, the conversions of **5** were less than 55%, while isolated yields of **6** were no more than 35%. Product **6** was much less stable during purification than products **3a–3bb**. Performing the reaction at higher temperature in toluene gave Stevens rearrangement product **7** as the major product, with **5** recovered at 55% yield.

Synthetic applications

Preliminary synthetic studies were performed to demonstrate



Scheme 2. Substituent Effects



Scheme 3. Control Experiment and Synthetic Applications

the utility of this rearrangement. Under 1 atm H₂, the diene compound **3k** underwent chemoselective 1,4-hydrogenation in the presence of Pd/C to give **8k** in 83% yield, whose structure was unambiguously confirmed by single-crystal x-ray diffraction analysis. The *N*-methyl group could be further modified via rhodium-catalyzed C–H functionalization to afford **9k**. Finally, the ester functionality could be reduced to the corresponding alcohol by DIBAL–H at –78 °C without eliminating the diene moiety. The diene was further cyclized in the presence of MeNH·OH·HCl catalyst to yield polycyclic compound **10k** in 57% yield (Scheme 3).

Mechanistic studies


In a plausible catalytic cycle (Scheme 4), the reaction begins when copper salt interacts with the diazo compound to give copper-carbene complex **D**, releasing one molecule of nitrogen gas. Subsequent reaction of copper-carbene complex with amine **1** affords the ylide **E**. Intramolecular nucleophilic attack in **E** delivers the spiro product **3** either before or after dissociation of the copper catalyst. To obtain structural information about ammonium salt **E** and explain the different reactivity between **1** and **5**, two ammonium compounds **F** and **G** were synthesized and analyzed by single-crystal x-ray diffraction. The torsional angle (C5–C6/C7–C13) was around 41 ° in **F** and 29 ° in **G**.¹⁹ Steric repulsion between the two *ortho*-substituents increased the distortion in the biaryl structure. We reasoned that the two *ortho*-substituents in **E** should distort its structure to a similar extent as in **F**. As a result, the



ARTICLE

Journal Name

relatively high-energy intermediate **E** undergoes rapid intramolecular



Scheme 4. Plausible Mechanism

nucleophilic reaction to deliver spiro compounds. Alternatively, the increased torsion reduces delocalization of the aniline lone-pair, which increases basicity and accelerates ammonium formation. Substituents on the substrates strongly influence the ratio of [2,3]-rearrangement product to Stevens rearrangement product;¹³ the latter rearrangement usually proceeds via a bi-radical pathway under thermal conditions.²⁰ These considerations imply that less distorted substrates such as **5** generate lower-energy ylide intermediates, which therefore must overcome a higher activation barrier to give spiro products. Consistent with this, compound **5** was associated with lower conversion rates at the similar conditions.

Conclusions

We have reported a copper-catalyzed Sommelet-Hauser rearrangement to construct spiroindoline compounds bearing adjacent quaternary and tertiary carbon centers. Increasing the torsional strain in the biaryl substrate increases its reactivity in this transformation. This is, to our knowledge, the first report in which Sommelet-Hauser rearrangement was used to generate spiro compounds bearing five-membered *N*-heterocycles.

Conflicts of interest


There are no conflicts to declare.

Acknowledgements

This work was supported by the NSFC (21622206, 21472179), '973' project from the MOST of China (2015CB856600), Strategic Priority Research Program of the Chinese Academy of Sciences (XDB20000000), and Fundamental Research Funds for the Central Universities (WK2060190086).


Notes and references

- (a) A. Yoshimuar, V. V. Zhdankin, *Chem. Rev.* 2016, **116**, 3328. (b) R. D. Richardson, T. Wirth, *Angew. Chem. Int. Ed.* 2006, **45**, 4402. (c) M. Uyanik, K. Ishihara, *ChemCatChem* 2012, **4**, 177. (d) T. Dohi, Y. Kita, *Chem. Commun.* 2009, 2073.
- P. W. Rabideau, Z. Marcinow, *Org. React.* 1992, **42**, 1–334.
- (a) M. Sono, M. P. Roach, E. D. Coulter, J. H. Dawson, *Chem. Rev.* 1996, **96**, 2841. (b) R. G. Harvey, *Acc. Chem. Res.* 1981, **14**, 218. (c) T. Hudlicky, D. Gonzalez, D. T. Gibson, *Aldrichimica Acta*, 1999, **32**, 35. (d) F. L. Ortiz, M. J. Iglesias, I. Fernández, C. M. A. Sánchez, G. R. Gómez, *Chem. Rev.* 2007, **107**, 1580.
- For some typical examples: (a) J. Barluenga, A. A. Trabanco, J. Flórez, S. García-Granda, E. Martín, *J. Am. Chem. Soc.* 1996, **118**, 13099. (b) J. García-Fortanet, F. Kessler, S. L. Buchwald, *J. Am. Chem. Soc.* 2009, **131**, 6676. (c) C. Shen, R.-R. Liu, R.-J. Fan, Y.-L. Li, T.-F. Xu, J.-R. Gao, Y.-X. Jia, *J. Am. Chem. Soc.* 2015, **137**, 4936. (d) J. Nan, Z. Zuo, L. Luo, L. Bai, H. Zheng, Y. Yuan, J. Liu, X. Luan, Y. Wang, *J. Am. Chem. Soc.* 2013, **135**, 17306. (e) A. Seoane, N. Casanova, N. Quiñones, J. L. Mascareñas, M. Gulías, *J. Am. Chem. Soc.* 2014, **136**, 7607. (f) F. C. Pigge, J. J. Coniglio, R. Dalvi, *J. Am. Chem. Soc.* 2006, **128**, 3498. (g) L. Bai, Y. Yuan, J. Liu, J. Wu, L. Han, H. Wang, Y. Wang, X. Luan, *Angew. Chem. Int. Ed.* 2016, **55**, 6946. (h) Z. Zuo, H. Wang, L. Fan, J. Liu, Y. Wang, X. Luan, *Angew. Chem. Int. Ed.* 2017, **56**, 2767.
- (a) S. You, S.-L. You, *Asymmetric Dearomatization Reactions*, Wiley-VCH, 2016. (b) C. Zheng, S.-L. You, *Chem* 2016, **1**, 830. (c) C.-X. Zhuo, C. Zheng, S.-L. You, *Acc. Chem. Res.* 2014, **47**, 2558.
- (a) M. Sommelet, *Compt. Rend.* 1937, **205**, 56. (b) S. W. Kantor, C. R. Hauser, *J. Am. Chem. Soc.* 1951, **73**, 4122–4131.
- E. Tayama, *Chem. Rec.* 2015, 789–800.
- (a) E. Tayama, R. Sato, K. Takedachi, H. Iwamoto, E. Hasegawa, *Tetrahedron* 2012, **68**, 4710. (b) E. Tayama, H. Kimura, *Angew. Chem. Int. Ed.* 2007, **46**, 8869. (c) E. Tayama, K. Orihara, H. Kimura, *Org. Biomol. Chem.* 2008, **6**, 3673. (d) E. Tayama, K. Watanabe, Y. Matano, *Eur. J. Org. Chem.* 2016, 3631. (e) E. Tayama, K. Watanabe, S. Sotome, *Org. Biomol. Chem.* 2017, **15**, 6668–6678.
- (a) T.-J. Lee, W. J. Holtz, *Tetrahedron Lett.* 1983, **24**, 2071. (b) R. Berger, J. W. Ziller, D. L. Van Vranken, *J. Am. Chem. Soc.* 1998, **120**, 841. (c) Y. Li, Y. Shi, Z. Huang, X. Wu, P. Xu, J. Wang, Y. Zhang, *Org. Lett.* 2011, **13**, 1210.
- For a reaction involving Sommelet-Hauser rearrangement: Gassman, P. G.; van Bergen, T. J.; Gilbert, D. P.; Cue, B. W., Jr. *J. Am. Chem. Soc.* 1974, **96**, 5495–5517.
- (a) A. Jończyk, D. Lipiak, *J. Org. Chem.* 1991, **56**, 6933–6937. (b) A. Jończyk, D. Lipiak, K. Sienkiewicz, *Synlett* 1991, 493–496. (c) T. Zdrojewski, A. Jonczyk, *J. Org. Chem.* 1998, **63**, 452. (d) S. Harthong, R. Bach, C. Besnard, L. Guénée, J. Lacour, *Synthesis*, 2013, **45**, 2070.



- 12 For some related reactions: (a) Z. Zhang, Z. Sheng, W. Yu, G. Wu, R. Zhang, W.-D. Chu, Y. Zhang, J. Wang, *Nat. Chem.* 2017, **9**, 970-976. (b) A. Soheili, U. K. Tambar, *J. Am. Chem. Soc.* 2011, **133**, 12956. (c) C.-Y. Zhou, W.-Y. Yu, P. W. H. Chan, C.-M. Che, *J. Org. Chem.* 2004, **69**, 7072. (d) M. S. Holzwarth, I. Alt, B. Plietker, *Angew. Chem. Int. Ed.* 2012, **51**, 5351-5354.
- 13 B. Biswas, S. C. Collins, D. A. Singleton, *J. Am. Chem. Soc.* 2014, **136**, 3740-3643.
- 14 K. Narita, N. Shirai, Y. Sato, *J. Org. Chem.* 1997, **62**, 2544.
- 15 (a) Sonnenschein, F. L. *Ber. Dtsch. Chem. Ges.* 1876, **9**, 1182 - 1186. (b) Conroy, H. & Chakrabarti, J. K. *Tetrahedron Lett.* 1959, **1**, 6 - 13. (c) Y. Schun, G. A. Cordell, *J. Nat. Prod.* 1985, **48**, 969 - 971. (d) F. M. Lovell, R. Pepinsky, A. J. C. Wilson, *Tetrahedron Lett.* 1959, **1**, 1 - 5.
- 16 Y. Schun, G. A. Cordell, M. Garland, *J. Nat. Prod.* 1986, **49**, 483 - 487.
- 17 Y.-K. Xu, L. Yang, S.-G. Liao, P. Cao, B. Wu, H.-B. Hu, J. Guo, P. Zhang, *J. Nat. Prod.* 2015, **78**, 1511.
- 18 (a) A. Fensome, M. Koko, J. Wrobel, P. Zhang, Z. Zhang, J. Cohen, S. Lundeen, K. Rudnick, Y. Zhu, R. Winneker, *Bioorg. Med. Chem. Lett.* 2003, **13**, 1317. (b) H. Venkatesan, M. C. Davis, Y. Altas, J. P. Snyder, D. C. Liotta, *J. Org. Chem.* 2001, **66**, 3653.
- 19 K. Zhao, L. Duan, S. Xu, J. Jiang, Y. Fu, Z. Gu, *Chem* 2018, **4**, 599-612.
- 20 (a) W. D. Ollis, M. Rey, I. O. Sutherland, *J. Chem. Soc. Chem. Commun.* 1975, 543. (b) Y. Maeda, Y. Sato, *J. Chem. Soc., Perkin Trans. 1*, 1997, 1491. (c) G. Ghigo, S. Cagnina, A. Maranzana, G. Tonachini, *J. Org. Chem.* 2010, **75**, 3608.





- § Highly Efficient Spiro-compound Synthesis; Excellent Selectivity
- § Molecular Torsional Strain Increasing the Reactivity

A Cu-catalyzed Sommelet-Hauser for the synthesis of spiro-indolines is reported, where the reactivity was significantly improved by molecular torsional strain.

