



Cite this: *Chem. Commun.*, 2020, 56, 1605

Received 9th November 2019,
Accepted 6th January 2020

DOI: 10.1039/c9cc08756g

rsc.li/chemcomm

A palladium-catalyst stabilized in the chiral environment of a monoclonal antibody in water†

Yuichiro Kobayashi,^{ib} a Keisuke Murata,^a Akira Harada^{ib} *^b and
Hiroyasu Yamaguchi^{ib} *^a

We report the first preparation of a monoclonal antibody (mAb) that can immobilize a palladium (Pd)-complex. The allylic amination reaction using a supramolecular catalyst consisting of the Pd-complex and mAb selectively gives the (R)-enantiomer product with an enantiomeric excess (ee) of 98 ± 2%. This is in sharp contrast to the reaction catalyzed by a conventional Pd-catalyst (ee < 2%).

Palladium (Pd)-complexes have attracted much attention because they promote various cross-coupling reactions such as Negishi coupling,^{1–3} Suzuki–Miyaura cross-coupling,^{4–6} and the Mizoroki–Heck reaction.^{7–10} Recent efforts have focused on controlling the synthesis products and increasing the catalytic activity due to the importance of chirality for living systems. For structural control of a product, both the first coordination sphere, which is formed by metal ions and ligands, and the second coordination sphere must be appropriately constructed. Synthesizing a ligand with the desired structure can realize the desired first coordination sphere. However, the second coordination sphere is composed by multiple non-covalent interactions such as hydrogen bonding, hydrophobic effects, and electro-static interactions. Hence, it is difficult to produce the desired second coordination sphere with a low molecular weight system.

To this end, hybrid catalysts with various combinations of Pd-complexes and biomolecules,¹¹ such as ferritin,¹² lipase,¹³ and biotin-streptavidin,^{14,15} have been developed to obtain structure-controlled products. A Pd-complex anchored in a biomolecule provides asymmetric environmental fields in the second coordination sphere. In this case, the Pd-complex is incorporated into biomacromolecules by employing a non-direct method. One example is the utilization of avidin–biotin

interactions during the complex formation of avidin with a biotinylated Pd-complex.^{14,15} Because the space originally possessed by the biomacromolecules is used as the second coordination sphere of the Pd-complex, it is not suitable for the Pd-complex. If biomacromolecules can directly recognize Pd-complexes, complexation of the Pd-complex and biomacromolecules can construct an appropriate second coordination sphere, realizing a novel platform for a hybrid catalyst of the Pd-complex and biomacromolecules.

Our work has focused on monoclonal antibodies (mAbs) because the binding site of mAbs can be tailored to the antigen specifications.^{16–29} This intriguing feature allows asymmetric environmental fields to be constructed as the second coordination sphere for transition metal complexes, enabling a stereo-specific reaction.^{30,31} However, to the best of our knowledge, a hybrid catalyst consisting of a Pd-complex and a mAb is yet to be reported because typical Pd-complexes are unstable in water and degrade during immunization. Currently, there is only one report on the complexation of a Pd-complex of a porphyrin.³² This may be due to the difficulty in preparing mAbs capable of recognizing Pd-complexes.

In this study, we describe our efforts to create a supramolecular catalyst using a mAb to immobilize an unstable Pd-complex (**1**) (Fig. 1 and Fig. S1–S3, Scheme S1, ESI†) for the allylic amination reaction. Because **1** degrades during immunization, a mAb capable of recognizing **1** is prepared by the cross-reactivity of mAb binding to the rhodium (Rh)-complex (**2**) (Fig. 1 and Fig. S4, S5, Scheme S2, ESI†) with the same ligand as **1**. A supramolecular catalyst consisting of **1** and mAb catalyzes the allylic amination reaction, which selectively affords an (R)-enantiomer product.

We chose a η^3 -allyl palladium(II) complex as a late transition metal complex because it promotes various coupling reactions such as allylic amination reactions.³³ To obtain mAbs capable of recognizing **1**, we prepared antigens in which **1** was modified to keyhole limpet hemocyanin (KLH) (**1**-KLH, Scheme S3 and Table S1, ESI†) for immunization and immunized **1**-KLH in saline emulsified 1 : 1 in Freund's complete adjuvant for Balb/c mice four times at two-week intervals. To investigate antibody

^a Department of Macromolecular Science, Graduate School of Science, Osaka University, 1-1 Machikaneyama-cho, Toyonaka, Osaka, 560-0043, Japan. E-mail: hiroyasu@chem.sci.osaka-u.ac.jp

^b The Institute of Scientific and Industrial Research, Osaka University, 8-1 Mihogaoka, Ibaraki 567-0047, Japan. E-mail: harada@chem.sci.osaka-u.ac.jp

† Electronic supplementary information (ESI) available: Experimental details, ¹H and ¹³C NMR and affinities between the mAb and the catalyst. See DOI: 10.1039/c9cc08756g






Fig. 1 Chemical structures of 1–5.

production, **1** was introduced into bovine serum albumin (BSA) to obtain antigens for assays (1-BSA), and then enzyme-linked immunosorbent assay (ELISA) measurements of the blood from immunized and non-immunized mice using a 1-BSA coated plate were performed. However, there were no differences in the absorbance ascribable to the enzymatic reaction product in ELISA for the immunized and non-immunized mice, showing that no antibodies for **1** have been produced (Fig. 2a). This is most probably due to the low stability of the Pd-complex in water. In fact, the proton nuclear magnetic resonance (^1H NMR) spectrum of **1** was changed by immersing it in *N,N*-dimethylformamide- d_7 /deuterium oxide for 7 days, showing decomposition of **1**. (Fig. S6, ESI †). The **1** was decomposed during immunization, which could not allow for creation of antibodies.

To overcome this problem, we employed cross-reactivity of antibodies. Because Rh-complexes are stable in water, we immunized **2** with the same ligand as **1**. The **2** was prepared from the chloro(1,5-cyclooctadiene)rhodium(i) dimer. The **2** was introduced to the proteins, such as KLH and BSA, in 0.1 M PBS buffer (pH 7.0) to obtain the antigens for immunization and

assays, respectively, (2-KLH and 2-BSA) (Scheme S3, ESI †). The modification ratio was estimated by using the 2,4,6-trinitrobenzenesulfonic acid (TNBS) method,^{34–36} and over six hundred or eight of **2** were introduced into one KLH or BSA, respectively (Table S1, ESI †). Balb/c mice were immunized with 2-KLH in saline emulsified 1 : 1 in Freund's complete adjuvant four times at two-week intervals. To confirm antibody production, we performed ELISA measurement of the blood from immunized and non-immunized mice using a 2-BSA coated plate. In contrast to **1**, the absorbance ascribable to the enzymatic reaction product in ELISA for an immunized mouse was higher than that of a non-immunized mouse (Fig. 2b). Moreover, in the case of an immunized mouse, the absorbance of 2-BSA was higher than that of BSA (Fig. S7, ESI †). These results clearly show the preparation of antibodies for **2**. The hybridomas secreting antibodies specific to **2** were cloned twice by the limiting dilution method and then the obtained mAbs were purified from the ascites fluid by affinity chromatography using a HiTrap IgM Purification HP column (GE healthcare). The class of mAbs was determined using the Iso Strip Mouse Monoclonal Antibody Isotyping Kit. As a result, the obtained mAb was immunoglobulin M (IgM), and the light chain was the kappa chain.

To quantitatively investigate the affinity of the mAb to **1**, we determined the dissociation constant (K_d) of the complex between the mAb and **1** or **2**. The K_d values of the complexes between the mAb and **1** or **2** were found to be 2.0 or 7.6×10^{-5} M, respectively (Fig. S8a and b, ESI †), showing that the affinities of mAb to **1** and that of mAb to **2** were almost the same. To understand this reason, the binding form of mAb to **2** was estimated using K_d of the complex between the mAb and the ligand of **2** (**3**), benzoic acid (**4**), or the Rh-complex (**5**) whose structure is different from **2** (Fig. 1). The mAb showed no affinity to **4** and **5** ($>1.5 \times 10^{-2}$ and $>1.0 \times 10^{-3}$ M, respectively), whereas the mAb showed affinity to **3** (3.4×10^{-4} M) (Fig. S8c and S9, ESI †). This result indicates that the mAb recognizes the molecular elements of **3** in complex **2**. The ligand structure of **2** and that of **1** were the same, and thus the mAb can form complexes with **1**. We prepared for the first time the mAb for a Pd-complex using the cross-reactivity of the mAb.

A reaction in a nanoconfined geometry often gives products whose symmetries differ from those obtained in a solution reaction. We carried out an allylic amination reaction using **1** in the absence and presence of the mAb (Table 1). Firstly, the reaction concentrations of **1** and mAb were optimized. 3-Buten-2-yl acetate (**6**) (10 mM) and benzyl amine (**7**) (20 mM) were reacted with **1** (1 μM) and the mAb (0.1 μM , binding site: 1 μM) in a mixed solution of 0.1 M phosphate borate buffer (PBB; pH 9.0) and DMSO (9 : 1) at r.t. for 24 h (Fig. S10 and Scheme S4, ESI †). Under these conditions, the binding site of the mAb and **1** is the same. The yield and ee of the product (**8**) were determined by comparing the area ratio of **8** and allylbenzene as an internal standard in HPLC analysis (Fig. S11, ESI †) and calculated from the peak area ratio of (*R*)- and (*S*)-enantiomers of **8** of HPLC spectra (Fig. S12, ESI †). The yield and ee of **8** decreased with increasing concentration of **1** and mAb. This result suggests that the aggregation of IgM causes **1** to be





Fig. 2 Results of ELISA for the blood obtained from the mouse immunized with transition-metal complex modified KLH (red) and the non-immunized mouse (black) [(a) **1**-KLH and (b) **2**-KLH].

Table 1 Results of the 1-catalyzed allylic amination reaction



Entry	Catalyst	Conc. of Pd (μM)/mAb (μM)	Yield of 8 ^a (%)	ee ^b (%)
1	1	1/0	38	< 2
2	1 + mAb	1/0.1 ^c	6	98 $\pm$ 2
3	1 + BSA	1/0.1	11	< 2

^a Calculated from the peak area ratio of HPLC spectra using allylbenzene as an internal standard. ^b Calculated from the peak area ratio of (*R*)- and (*S*)-enantiomers of **8** of HPLC spectra. ^c Since the mAb is IgM with 10 binding sites, the concentrations of **1** and the binding site of mAb are equivalent.

adsorbed at locations other than where the product structure is controlled. The optimum reaction concentration of **1** and mAb was found to be 1 and 0.1 μM , respectively. The yield of **8** in the presence of the mAb (6%) was lower than that in the absence of the mAbs (38%) (Table 1, entries 1 and 2). Owing to the complex formation between **1** and mAb, it is difficult for the substrates such as **6** and **7** to access **1**. The decline in the yield of **8** was also observed in the presence of BSA (11%) (Table 1, entry 3), suggesting that the presence of protein reduces the catalytic activity of **1**. In the case of the absence of mAb, **1** gave racemic **8** (ee < 2%) (Table 1, entry 1 and Fig. S12a, ESI[†]). In contrast, interestingly, the reaction catalysed by **1** incorporated into the binding site of the mAb was found to proceed with excellent (*R*)-enantioselectivity with a 98 $\pm$ 2% ee (Table 1, entry 2 and Fig. S12b, ESI[†]). These results indicate that no catalytic reaction occurred outside the binding sites. Since the catalytic activity of free **1** is reduced by the presence of protein, the catalytic activity of free **1** would be negligible in the mAb system. This asymmetric allylic amination was not observed in the presence of BSA (ee < 2%) (Table 1, entry 3 and Fig. S12c, ESI[†]). These results indicate that the binding site of mAb functions as a reaction field for the asymmetric reaction.

In summary, we have demonstrated that Pd-complex catalysed allyl amination in the presence of mAb is a useful synthetic strategy to control the resultant product asymmetry. The substrate is converted to a racemic product by a Pd-catalyst without a mAb. In contrast, the (*R*)-enantiomer (98 $\pm$ 2% ee) is obtained in the presence of the complex between the mAb and the Pd-catalyst.

This work was supported by the JSPS KAKENHI grant number JP15H05807 in Precisely Designed Catalysts with Customized Scaffolding.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 E. Negishi, Q. Hu, Z. H. Huang, M. X. Qian and G. W. Wang, *Aldrichimica Acta*, 2005, **38**, 71.
- 2 E. Negishi, G. W. Wang, H. H. Rao and Z. Q. Xu, *J. Org. Chem.*, 2010, **75**, 3151.
- 3 D. Haas, J. M. Hammann, R. Greiner and P. Knochel, *ACS Catal.*, 2016, **6**, 1540.
- 4 N. Miyaara and A. Suzuki, *Chem. Commun.*, 1979, 866.
- 5 N. Miyaara, K. Yamada and A. Suzuki, *Tetrahedron Lett.*, 1979, **20**, 3437.
- 6 N. Miyaara and A. Suzuki, *Chem. Rev.*, 1995, **95**, 2457.
- 7 T. Mizoroki, K. Mori and A. Ozaki, *Bull. Chem. Soc. Jpn.*, 1971, **44**, 581.
- 8 R. F. Heck and J. P. Nolley, *J. Org. Chem.*, 1972, **37**, 2320.
- 9 I. P. Beletskaya and A. V. Chepurkov, *Chem. Rev.*, 2000, **100**, 3009.
- 10 A. B. Dounay and L. E. Overman, *Chem. Rev.*, 2003, **103**, 2945.
- 11 F. Schwizer, Y. Okamoto, T. Heinisch, Y. F. Gu, M. M. Pellizzoni, V. Lebrun, R. Reuter, V. Kohler, J. C. Lewis and T. R. Ward, *Chem. Rev.*, 2018, **118**, 142.
- 12 S. Abe, J. Niemeyer, M. Abe, Y. Takezawa, T. Ueno, T. Hikage, G. Erker and Y. Watanabe, *J. Am. Chem. Soc.*, 2008, **130**, 10512.
- 13 M. Filice, O. Romero, A. Aires, J. M. Guisan, A. Rumbero and J. M. Palomo, *Adv. Synth. Catal.*, 2015, **357**, 2687.
- 14 J. Pierron, C. Malan, M. Creus, J. Gradinaru, I. Hafner, A. Ivanova, A. Sardo and T. R. Ward, *Angew. Chem., Int. Ed.*, 2008, **47**, 701.
- 15 A. Chatterjee, H. Mallin, J. Klehr, J. Vallapurackal, A. D. Finke, L. Vera, M. Marsh and T. R. Ward, *Chem. Sci.*, 2016, **7**, 673.
- 16 B. L. Iverson and R. A. Lerner, *Science*, 1989, **243**, 1184.
- 17 A. Harada, K. Okamoto, M. Kamachi, T. Honda and T. Miwatani, *Chem. Lett.*, 1990, 917.
- 18 A. G. Cochran and P. G. Schultz, *Science*, 1990, **249**, 781.
- 19 V. A. Roberts, B. L. Iverson, S. A. Iverson, S. J. Benkovic, R. A. Lerner, E. D. Getzoff and J. A. Tainer, *Proc. Natl. Acad. Sci. U. S. A.*, 1990, **87**, 6654.
- 20 R. A. Lerner, S. J. Benkovic and P. G. Schultz, *Science*, 1991, **252**, 659.
- 21 P. Ghosh, D. Shabat, S. Kumar, S. C. Sinha, F. Grynspan, J. Li, L. Noodleman and E. Keinan, *Nature*, 1996, **382**, 339.
- 22 D. A. Blake, P. Chakrabarti, M. Khosraviani, F. M. Hatcher, C. M. Westhoff, P. Goebel, D. E. Wylie and R. C. Blake, *J. Biol. Chem.*, 1996, **271**, 27677.
- 23 A. Harada, H. Fukushima, K. Shiotsuki, H. Yamaguchi, F. Oka and M. Kamachi, *Inorg. Chem.*, 1997, **36**, 6099.
- 24 M. Khosraviani and R. C. Blake, *Bioconjugate Chem.*, 2000, **11**, 267.
- 25 K. M. Nicholas, P. Wentworth, C. W. Harwig, A. D. Wentworth, A. Shafton and K. D. Janda, *PNAS*, 2002, **99**, 2648.
- 26 R. Ricoux, H. Sauriat-Dorizon, E. Girgenti, D. Blanchard and J. P. Mahy, *J. Immunol. Methods*, 2002, **269**, 39.
- 27 H. Yamaguchi, K. Tsubouchi, K. Kawaguchi, E. Horita and A. Harada, *Chem. – Eur. J.*, 2004, **10**, 6179.
- 28 R. C. Blake, A. R. Pavlov, M. Khosraviani, H. E. Ensley, G. E. Kiefer, H. Yu, X. Li and D. A. Blake, *Bioconjugate Chem.*, 2004, **15**, 1125.
- 29 J. P. Mahy, J. D. Marechal and R. Ricoux, *Chem. Commun.*, 2015, **51**, 2476.
- 30 H. Yamaguchi, T. Hirano, H. Kiminami, D. Taura and A. Harada, *Org. Biomol. Chem.*, 2006, **4**, 3571.
- 31 T. Adachi, A. Harada and H. Yamaguchi, *Sci. Rep.*, 2019, **9**, 13551.
- 32 A. P. Savitsky, M. V. Demcheva, E. Y. Mantrova and G. V. Ponomarev, *FEBS Lett.*, 1994, **355**, 314.
- 33 M. Johannsen and K. A. Jorgensen, *Chem. Rev.*, 1998, **98**, 1689.
- 34 K. Satake, T. Okuyama, M. Ohashi and T. Shinoda, *J. Biochem.*, 1960, **47**, 654.
- 35 L. C. Mokrasch, *Anal. Biochem.*, 1967, **18**, 64.
- 36 S. L. Snyder and P. Z. Sobocinski, *Anal. Biochem.*, 1975, **64**, 284.

