


Environmental Science Nano

Volume 8
Number 5
May 2021
Pages 1123–1484

rsc.li/es-nano



ISSN 2051-8153

PAPER

Craig J. Dedman, Joseph A. Christie-Oleza,
Gemma-Louise Davies *et al.*
Environmentally relevant concentrations of titanium dioxide
nanoparticles pose negligible risk to marine microbes

PAPER

View Article Online
View Journal | View Issue



Cite this: *Environ. Sci.: Nano*, 2021, 8, 1236

Environmentally relevant concentrations of titanium dioxide nanoparticles pose negligible risk to marine microbes†

Craig J. Dedman,^a Aaron M. King,^c
Joseph A. Christie-Oleza^a and Gemma-Louise Davies^a

Nano-sized titanium dioxide (nTiO₂) represents the highest produced nanomaterial by mass worldwide and, due to its prevalent industrial and commercial use, it inevitably reaches the natural environment. Previous work has revealed a negative impact of nTiO₂ upon marine phytoplankton growth, however, studies are typically carried out at concentrations far exceeding those measured and predicted to occur in the environment currently. Here, a series of experiments were carried out to assess the effects of both research-grade nTiO₂ and nTiO₂ extracted from consumer products upon the marine dominant cyanobacterium, *Prochlorococcus*, and natural marine communities at environmentally relevant and supra-environmental concentrations (i.e., 1 µg L⁻¹ to 100 mg L⁻¹). Cell declines observed in *Prochlorococcus* cultures were associated with the extensive aggregation behaviour of nTiO₂ in saline media and the subsequent entrapment of microbial cells. Hence, higher concentrations of nTiO₂ particles exerted a stronger decline of cyanobacterial populations. However, within natural oligotrophic seawater, cultures were able to recover over time as the nanoparticles aggregated out of solution after 72 h. Subsequent shotgun proteomic analysis of *Prochlorococcus* cultures exposed to environmentally relevant concentrations confirmed minimal molecular features of toxicity, suggesting that direct physical effects are responsible for short-term microbial population decline. In an additional experiment, the diversity and structure of natural marine microbial communities showed negligible variations when exposed to environmentally relevant nTiO₂ concentrations (i.e., 25 µg L⁻¹). As such, the environmental risk of nTiO₂ towards marine microbial species appears low, however the potential for adverse effects in hotspots of contamination exists. In future, research must be extended to consider any effect of other components of nano-enabled product formulations upon nanomaterial fate and impact within the natural environment.

Received 22nd August 2020,
Accepted 6th April 2021

DOI: 10.1039/d0en00883d

rs.c.li/es-nano

Environmental significance

Titanium dioxide (nTiO₂) represents the highest produced engineered nanomaterial worldwide, yet its impact in the marine environment is poorly understood. Despite 70–80% use in the cosmetics industry, typically only pristine research-grade materials are examined. Herein, the effects of research-grade and nTiO₂ extracted from common consumer goods upon marine microbial species were investigated. We show cell decline of the cyanobacterium *Prochlorococcus* during short-term exposure (72 h), but ultimate recovery under environmentally relevant conditions after 10 days. Cell decline following nTiO₂ exposure appear to result from nanoparticle aggregation and entrapment of cyanobacteria, while little molecular features of toxicity are identified. Exposure of natural communities reveals negligible effect of nTiO₂ on marine microbial community structure. Therefore, the risk of nTiO₂ exposure in the marine environment appears low.

^a School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry, CV4 7AL, UK. E-mail: C.Dedman@warwick.ac.uk

^b Department of Chemistry, University of Warwick, Gibbet Hill, Coventry, CV4 7EQ, UK

^c UCL Department of Chemistry, University College London, 20 Gordon Street, London, WC1H 0AJ, UK. E-mail: gemma-louise.davies@ucl.ac.uk

^d Department of Biology, University of the Balearic Islands, Ctra. Valldemossa, km 7.5, CP: 07122, Palma, Spain

^e IMEDEA (CSIC-UIB), CP: 07190, Esporles, Spain. E-mail: joseph.christie@uib.eu

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0en00883d

1. Introduction

The fate and effects of engineered nanomaterials (NMs) within the natural environment has become an ecological concern over the past decade.^{1–3} Nano-sized titanium dioxide (nTiO₂) represents the highest produced NM worldwide, with annual production predicted to reach 2.5 million tonnes by 2025.^{4,5} Due to its prevalent use across a wide range of industries, including plastic production, paints, foods and



cosmetics,^{5,6} nTiO₂ is highly likely to enter the natural environment.⁷ The cosmetics industry dominates the use of nTiO₂, accounting for approximately 70–80% of nTiO₂ use.⁶ Typically product formulations are made up of 2–14 wt% nTiO₂,^{8,9} however, the exact physicochemical properties of NMs used vary greatly. In fact, it is estimated that the chemical composition of NMs utilised for over 50% of commercial products is not widely publicised.⁴

Domestic use of consumer products such as sunscreen and toothpaste is predicted to release considerable volumes of TiO₂ into the aquatic environment.^{6,10} Extraction of metal oxide NMs from wastewater treatment works has been recorded at an efficiency of approximately 95%,¹¹ resulting in the entry of approximately 5% of metal oxide NMs into the aquatic environment *via* wastewater effluent. Given that 4100 tonnes of TiO₂ is predicted to enter US wastewaters as a result of toothpaste use every year,⁶ we can estimate that up to 205 tonnes will enter wastewater effluent and be transported to natural systems, where their ultimate fate remains unknown.

In recent years, increasing efforts have been made to uncover the environmental concentrations of engineered NMs. Difficulties in sampling techniques and varying effectiveness of methods for characterising nano-pollutants, particularly at the low concentrations predicted in the environment, has led to a lack of environmental data.^{12,13} ‘Environmental Fate Models’ represent a powerful tool for this purpose,¹⁴ and surface water concentrations of nTiO₂ are estimated in the range of 0.021–10.000 µg L⁻¹.¹⁵ However, more recently, environmental sampling has revealed concentrations of up to 40 µg L⁻¹ close to major transport infrastructure and locations heavily impacted by tourism.^{16,17} For example, during the peak tourist season, TiO₂ derived from sunscreen use has been measured in the range of 7–40 µg L⁻¹ in surface waters off the Mallorcan coast.¹⁷ As such, it appears that the environmental risk from nano-sized pollutants is likely focused within localised areas of contamination which may vary temporally, where particle-specific properties govern their subsequent fate and transport.¹⁸

The microbial community plays a fundamental role in the functioning of the marine ecosystem, contributing approximately 50% of global primary productivity and influencing major climatic and biogeochemical cycles.^{19,20} Hence, understanding the potential effects of contaminants upon marine microbes is key to evaluating their likely ecosystem-wide impact. Previous research has revealed a toxic effect of nTiO₂ exposure upon marine microbial species,^{6,21–30} although studies where little or no adverse effect is recorded also exist.^{31,32} The primary effect of exposure appears to be growth inhibition, however EC₅₀ values are typically recorded in the mg L⁻¹ range, far greater than those measured in the environment (up to 40 µg L⁻¹).^{23–25} In addition to growth inhibition, negative effects such as the induction of oxidative stress pathways, physical damage to cells, and entrapment of cells within aggregates of

NMs have been recorded following exposure, however results vary greatly.^{23–25,32,33} Indeed, physical interaction between phytoplankton and aggregates of nTiO₂ appears widespread in laboratory exposure^{34–37} and may represent a possible pathway for phototrophic removal from epipelagic layers³⁵ or transport of NMs to higher trophic levels.

Much of the toxicological work of nTiO₂ has so far been carried out using standardised research-grade NMs, which are not usually surface modified. More recently, studies have begun to emerge focussing upon the toxicity of surface functionalised nTiO₂ (*e.g.* sunscreens which often possess surfactant/polymer surface modifications) upon marine phytoplankton.^{6,27} In these studies, adverse effects were observed in phytoplankton exposed to sunscreens containing nTiO₂, associated with increased production of reactive oxygen species (ROS), membrane damage and possible genotoxicity.²⁷ However, biostimulating effects of nTiO₂-containing sunscreens have also been recorded, attributed to other organic components of the sunscreen formulation.⁸ Interestingly, nTiO₂ derived from commercial products have been reported to result in greater growth inhibition than research-grade nanoparticles.⁶ It should be noted, though, that much of the work investigating metal oxide NMs are carried out using exposure concentrations in the mg L⁻¹ range (*i.e.* 1–30 mg L⁻¹),^{6,21–30} far exceeding those predicted and measured in the environment (0.021–40 µg L⁻¹).^{6,8,15,17} Increased research is required to examine the end-products of consumer goods which may enter the aquatic environment.⁸⁹ Given the great variation in physicochemical properties between specific NMs, and hence varied fate in aquatic media, it is important that representative NMs derived from consumer goods are utilised alongside research-grade materials during experimentation.^{9,38} To provide a greater understanding of the environmental impact of nTiO₂, research must also be directed to simulate environmental conditions as effectively as possible.

Since much of the previous research in this field has focussed upon phytoplankton species such as diatoms and green algae, comparatively little evidence for the effects of NM exposure upon photosynthetic cyanobacteria exist. Marine cyanobacteria, mainly *Prochlorococcus* and *Synechococcus* species, are the most abundant photosynthetic organisms on earth and major contributors to global primary productivity.^{39,40} Furthermore, among phytoplankton taxa, cyanobacteria appear particularly sensitive to nano-pollutants *e.g.* silver nanoparticles.⁴¹ Herein, we aimed to provide a broad-spectrum analysis of both commercially available research-grade (non-surface modified) nTiO₂, as well as nTiO₂ extracted from common consumer products, and examined their impact upon marine phytoplankton at both the organism- and community-level. Short-term (72 h) and medium-term (10 d) toxicity of nTiO₂ was examined using the ecologically significant cyanobacterium *Prochlorococcus* sp. MED4 under environmentally relevant conditions (*i.e.*, at ambient cell densities (10⁴–10⁵ cells per mL),^{42,43} in oligotrophic natural seawater). The behaviour of nTiO₂ in



seawater and its interaction with cyanobacteria was investigated through the use of dynamic light scattering (DLS), flow cytometry and fluorescent microscopy. Molecular features of toxicity were assessed by shotgun proteomic analysis and appeared negligible. Finally, an additional experiment was conducted using natural coastal seawater to characterise the whole community response towards consumer nTiO₂ exposure. Amplicon sequencing of the 16S rRNA and 18S rRNA genes revealed little impact of extracted nTiO₂ derived from sunscreen upon natural marine microbial community structure at environmental concentrations. This multi-OMIC study provides a comprehensive assessment of the differing effects that result from exposure to various types of nTiO₂, representative of materials likely to enter the marine environment, thus facilitating effective evaluation of their likely interaction with marine microbial species and overall environmental risk.

2. Methods

2.1 Materials

Research-grade nTiO₂ used during experimentation was purchased from Sigma Aldrich (21 nm (19.9 ± 6.6 nm, TEM)). Three consumer products; Skincenticals™ sunscreen (S1), Boots Soltan™ sunscreen (S2), and The Body Shop™ liquid foundation (P1) were selected based on nTiO₂ being listed as an ingredient and purchased from a high street retailer. Natural seawater (NSW) obtained from Station L4 (Plymouth, UK, 50°15.0'N; 4°13.0'W) was routinely used during experimental work. Prior to use, NSW was autoclaved and filtered (0.22 µm, polyethersulfone membrane Corning®). 'Neat' sunscreen stocks were prepared by diluting the cream directly in NSW. For all experimental work, nTiO₂ stocks were prepared in NSW and sonicated for 15–30 min (Branson 1210 Sonicator, 40 kHz) prior to addition to experimental media to avoid extensive aggregation. Glassware was acid-washed before use. Axenic *Prochlorococcus* sp. strain MED4 was routinely grown using Pro99 media⁴⁴ and maintained at 23 °C before use in experimentation.

2.2 Extraction of nTiO₂ from consumer products

Methods adapted from those described by Galletti *et al.* (2016) were utilised to extract nTiO₂ from consumer goods.⁶ Briefly, 1.5–2.5 g of product was soaked in 20 mL hexane for 2 h. Suspensions were then shaken manually and centrifuged at 4400 rpm for 5 min. The supernatant was subsequently discarded, and 20 mL ethanol was added and shaken manually. The solution was then centrifuged at 4400 rpm for 5 min and the resultant supernatant was discarded. Following this, the remaining product was washed in Milli-Q ultrapure H₂O (0.22 µm filter operated at 18.2 MΩ at 298 K) three times using centrifugation; first centrifuging at 11 000 rpm for 5 min, extending this to 10 min for the final two washes. Extracted pellets were subsequently dried in a 60 °C oven to form a powder. Samples were stored in darkness

prior to characterisation and use in experiments with marine phytoplankton.

2.3 Characterisation of materials

Transmission electron microscopy (TEM) was utilised to determine primary nanoparticle size. A JEOL 2100 TEM, 200 kV, LaB₆ instrument operated with a beam current of ~115 mA was used to obtain images using a Gatan Orius 11-megapixel camera. Samples were prepared by deposition and drying of nanoparticle samples (10 µL aqueous suspension) onto formvar-coated 300 mesh copper TEM grids (EM Resolutions). Diameters were measured using ImageJ version 3.2; average values were calculated by measuring the diameter of >100 particles with errors represented as standard deviations. Energy-dispersive X-ray spectroscopy (EDS) was collected using an Oxford Instruments X-Max 80T detector. Additionally, samples were ground to fine powders before powder X-ray diffraction (P-XRD) was performed using a STOE Stadi-P diffractometer with a molybdenum X-ray source (operated at 50 kV and 30 mA), λ = 0.7093 Å. The 2θ scan range was 2–40.115° at a step size of 0.495° and 5 seconds per step. Samples were prepared using STOE zero scattering foils before being inserted into the transmission sample holder.

2.4 Short-term (72 h) consumer nTiO₂ exposure

Prochlorococcus MED4 was inoculated into oligotrophic NSW at ambient cell densities (~10⁴ cells per mL) and incubated 72 h for pre-adaptation to the oligotrophic conditions prior to experimentation (*i.e.*, 23 °C at constant 10 µmol photons m⁻² s⁻¹ light intensity, using a Lifelite™ full spectrum bulb with UV, and with shaking at 100 rpm). The light intensity used is optimal for cultured *Prochlorococcus* sp. MED4 and ensured cyanobacteria were not affected by light stress during experiments. 30 mL of pre-adapted *Prochlorococcus* culture was aliquoted into 50 mL tissue culture flasks and spiked with nTiO₂ stocks to make up test concentrations of 0, 5, 50 and 500 µg L⁻¹, all in triplicate. Four nTiO₂ treatments were investigated: nTiO₂ nanopowder (Sigma Aldrich), nTiO₂ derived from sunscreen S1 and S2, and nTiO₂ derived from liquid foundation P1. Additionally, a 'neat' sunscreen treatment was tested, where 0.1 g of sunscreen S2 was immersed in NSW and mixed *via* manual shaking and 15 min sonication. Cultures were subsequently spiked with a defined volume of the sunscreen suspension to make up equivalent test concentrations based on nTiO₂ making up ~10 wt% of the sunscreen formulation.^{8,9} After the addition of each treatment, cultures were further incubated under the conditions described above and monitored by flow cytometry using a Becton Dickinson Fortessa flow cytometer at time points 0, 24, 48 and 72 h. Cell densities were calculated in respect to reference beads (2.2 µm high intensity fluorescent Nile Red particles (Spherotech FH-2056-2)), added to samples at a defined concentration. For additional details of flow



cytometric analyses used throughout experimentation see Section S2, ESI.†

2.5 Medium-term (10 d) nTiO₂ exposure

To investigate the medium-term effects of nTiO₂ exposure upon *Prochlorococcus* MED4, research-grade nTiO₂ (Sigma Aldrich, 19.9 ± 6.6 nm) was utilised for experimentation, based on results of earlier experiments described in section 2.4. Two culture conditions were tested: i) cell-dense cultures (~10^{6–7} cells per mL) in nutrient-rich Pro99 media, and ii) cultures grown to ambient cell densities (~10^{4–5} cells per mL) in oligotrophic NSW, representing environmental conditions. Cultures were set up as described in section 2.4. Culture flasks were subsequently spiked with a nTiO₂ stock achieving final concentrations in the µg L^{−1} (1, 10 and 100 µg L^{−1}) and mg L^{−1} range (1, 10 and 100 mg L^{−1}), all in triplicate; representing environmental and supra-environmental concentrations respectively. Cell counts were monitored at 0, 24, 48, 72, 192 and 240 h by flow cytometry as previous and compared to that of an untreated control. In addition to monitoring of cell density, flow cytometric analysis was utilised to infer the behaviour of nTiO₂ within test media and their interaction with cyanobacterial cells (see Section S2.1,† for further information).

2.6 Imaging of nTiO₂-cyanobacterial aggregates by fluorescent microscopy

To investigate the hetero-aggregation between nTiO₂ and *Prochlorococcus*, a 200 µL sample was collected from the bottom of culture flasks from one replicate of each treated group during medium-term experiments (section 2.5). This sub-sample was stained with 1X SYBR Gold nuclear stain (ThermoFisher) and imaging was carried out at 40× magnification using a Nikon widefield fluorescence microscope under brightfield and GFP fluorescence. Images were captured from both channels and subsequently merged to assess the presence and extent of aggregation between cyanobacterial cells and nTiO₂. Controls containing nTiO₂ only (100 mg L^{−1}) were included to prove the nanoparticles did not get stained by the dye. In addition, NSW samples and untreated *Prochlorococcus* culture in the absence of nTiO₂ were imaged to confirm aggregates were indeed nTiO₂ rather than other particulate material.

2.7 Examining the behaviour of nTiO₂ within natural seawater by dynamic light scattering

The aggregation behaviour of nTiO₂ (Sigma Aldrich, 19.9 ± 6.6 nm) within NSW was assessed by z-average size (d.nm) over a period of 336 h (14 d). Here, hydrodynamic particle size measurements were determined by dynamic light scattering (DLS) using a Malvern Zetasizer Nano ZS instrument, equipped with a 4 mW He-Ne 633 nm laser module. A stock of nTiO₂ was sonicated for 15–30 min prior to addition to NSW. Concentrations of 1 mg L^{−1} and 100 mg L^{−1} were utilised due to limitations of the DLS at lower

concentrations, hence hindering the ability to assess nTiO₂ aggregation at the environmentally relevant concentrations (*i.e.*, in the µg L^{−1} range) used in this study during toxicity testing. nTiO₂ suspensions were made up in 20 mL autoclaved and filtered (0.22 µm) NSW in 50 mL tissue culture flasks and placed on an orbital shaker (100 rpm) to simulate natural movement of water. DLS measurements were carried out upon a 200 µL sub-sample collected from the mid-point of flasks at set timepoints (0, 1, 2, 4, 24, 48, 72, 168, 240, 336 h). For each sample the average was taken from 3 measurements made up of 11 sampling runs lasting 10 s each.

2.8 Shotgun proteomic analysis

To ensure sufficient biological material was obtained for proteomic analyses, cell-dense *Prochlorococcus* MED4 cultures were grown in Pro99 media. Following 72 h preadaptation to experimental conditions, described above (section 2.4), triplicate cultures were spiked with research-grade nTiO₂ stock (Sigma Aldrich, 19.9 ± 6.6 nm) to achieve a test concentration of 100 µg L^{−1}. Untreated cultures were also prepared as controls. Following the addition of nTiO₂, cultures were incubated 24 h, after which samples were immediately centrifuged for 10 min at 4 °C at 4000 × *g* and pellets were immediately frozen in dry ice until further processing. The supernatants containing the extracellular proteome were filtered (0.22 µm) and stored at −20 °C. Supernatants were thawed at room temperature and underwent trichloroacetic acid (TCA) protein precipitation as previously described.⁴⁵ Subsequently, cell and extracellular proteome pellets were resuspended in 1x LDS buffer (ThermoFisher) containing 1% beta-mercaptoethanol and run on a NuPage 4–12% Bis-Tris precast polyacrylamide gels as previously done.⁴⁶ In-gel trypsin digestion and peptide recovery was performed⁴⁷ followed by a nanoLC-ESI-MS/MS analysis using an UltiMate 3000 RSLCnano System coupled to an Orbitrap Fusion (Thermo Scientific) using conditions and settings as previously described.⁴⁸ RAW mass spectral files were processed using MaxQuant version 1.5.5.1 (ref. 49) for peptide identification and protein label-free quantification using the *Prochlorococcus* sp. MED4 UniProt coding domain sequences (downloaded on 16/01/2018). For further details of the laboratory and bioinformatics protocols used for proteomics analysis see Section S4, ESI.†

2.9 16S rRNA/18S rRNA amplicon sequencing

A site for experimental work was selected in the Balearic Islands, Spain (39.493868, 2.739820). Field experiments were carried out during April 2018. Here, coastal seawater (NSW) containing its natural microbial community was collected at a depth of approximately 1 m, representing those microorganisms most likely to interact with nTiO₂ derived from sunscreen in the coastal system. Subsequently, 500 mL was transferred to pre-washed 1 L Nalgene plastic bottles, leaving sufficient volume empty for air exchange. Microbial



communities were exposed to one of three treatments: 1) untreated control, where no nTiO₂ was added; 2) nTiO₂ extracted from sunscreen S2; 3) 'Neat' sunscreen S2 dispersed in NSW. Extracted nTiO₂ from S2 were obtained as described above (section 2.2) and added at a final concentration of 25 µg L⁻¹. 'Neat' sunscreen stock was made up in NSW and a defined volume was added to achieve ~25 µg L⁻¹ of nTiO₂, assuming that nTiO₂ typically makes up ~10 wt% of such products.^{8,9} Bottles were mixed by inversion three times and incubated over two days in an outdoor water container to provide temperature stability while being exposed to natural sunlight to best replicate natural conditions. Bottle caps were loosened to ensure sufficient gas exchange and bottles were shaken manually (15–30 s) at regular intervals throughout the experiment to ensure water was well mixed. Following exposure, microbial cells were collected by filtering the 500 mL through a 0.22 µm filter (Millipore). Filters were transferred to a 2 mL Eppendorf containing lysis buffer (Qiagen) and stored at -20 °C. DNA extraction was carried out using the DNeasy Power Biofilm extraction kit (Qiagen) according to the manufacturer's instructions, including a bead beating step as previously described.⁵⁰ DNA quantification was obtained by Qubit® HS DNA kit (Life Technologies Corporation). Extracted DNA samples were stored at -20 °C. Prokaryotic and eukaryotic community analysis was performed by amplicon sequencing using the 515F-Y and 926R primers to amplify the 16S rRNA v4–5 regions, and V8F and 1510R primers to amplify the 18S rRNA v8–9 regions, respectively.^{51,52} PCR products were purified, indexed, normalized and analysed by 2 × 300 bp paired-end sequencing using the MiSeq system with v3 reagent kit (Illumina) as described in Wright *et al.* 2019.⁵⁰ Raw sequencing data was analysed using the DADA2 bioinformatic pipeline based on its enhanced taxonomic resolution compared to alternative methods.^{50,53–55} For additional details of the amplicon sequencing methods and data analysis used, see Section S5.1, ESI.†

2.10 Statistical analysis

To identify significant alterations in cell density recorded during toxicity tests with *Prochlorococcus* MED4 (sections 2.4 and 2.5) two-way *t*-tests were carried out between untreated controls and cultures exposed to various nTiO₂ treatments at each timepoint. Downstream statistical analysis of shotgun proteomics data (section 2.8) was carried out using Perseus version 1.5.5.3,⁵⁶ following the pipeline described previously.⁵⁷ The mass spectrometry data have been deposited to the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org>) via the PRIDE partner repository⁵⁸ with the dataset identifier PXD024726. For amplicon sequencing data collected as described in section 2.9, taxonomically assigned data in the form of amplicon sequencing variants (ASVs) were analysed using MicrobiomeAnalyst software.^{59,60} Briefly, following normalisation by total sum scaling, principle coordinates

analysis (PCA) based on Bray–Curtis dissimilarity, in conjunction with permutational multivariate analysis of variance (PERMANOVA) was utilised to assess significant alterations in community composition between treatments at the individual ASV level. Subsequently, two-way *t*-tests were utilised to identify significant variations in relative abundance of various taxonomic groups between control and treated samples. Sequence files have been deposited in the NCBI Short Read Archive (SRA) database under Bioproject: PRJNA690209.

3. Results and discussion

3.1 Characterisation of research-grade and consumer nTiO₂

nTiO₂ was initially extracted from commercial sunscreen and cosmetics products, S1, S2, and P1, as detailed in the Methods section, and characterised. The extraction method utilised to extract particles from the three product formulations was largely effective, yielding a visible powder that could be dried and stored for experimental use. Characterisation of research-grade and extracted materials was carried out using a combination of TEM and EDS mapping, revealing primary particle size, morphology and elemental composition (Table 1 and Fig. 1).

Research-grade nTiO₂ (Sigma Aldrich) possessed an average particle size of 19.9 ± 6.6 nm, close to the manufacturer's advertised size (21 nm), whilst materials extracted from consumer products ranged in average primary particle size 50.0 to 158.1 nm. Interestingly, all primary particles extracted from commercially available products had large standard deviations in sizes, indicating a wide size range in particle populations (Table 1). Typically, samples appeared as small aggregates, a common feature often observed as a result of drying of the samples onto TEM grids (Fig. 1). EDS mapping confirmed that all primary particles were entirely composed of Ti and O (Table 1 and Fig. S1†). P-XRD was carried out on all samples (Fig. S2†). This showed research-grade nTiO₂ to have a mixture of anatase and rutile phases, with peaks corresponding with the JCPDS patterns 21-1272 (ref. 61) and 21-1276 (ref. 62) for the tetragonal structure of anatase and rutile TiO₂, respectively. The sunscreens S1 and S2 samples appeared to be present as the rutile phase only, whilst the product P1 sample presented only anatase phase. All samples showed some peak broadening, indicative of the presence of nano-sized crystallites, as previously observed by electron microscopy. Previous research has indicated that nTiO₂ phase has impacted upon toxicity towards biota,^{22,63} hence, such information is important to fully evaluate outcomes of toxicity testing.

In accordance with previous studies, variation in physical properties of extracted particles is observed.^{9,38} For example, primary nTiO₂ particles extracted from consumer products S1 and S2 showed significant differences in morphology (Fig. 1B and C), despite both being utilised for UV protection in sunscreen formulations. These results emphasize the



Table 1 Summary of material characteristics as determined by TEM, EDS mapping and P-XRD

| nTiO ₂ source | Primary particle characteristics | | | | Secondary particle characteristics | | |
|--------------------------|----------------------------------|-----------------------|--------------------|----------------------------|------------------------------------|-----------------------|------------------|
| | Size (nm) (TEM) | Elemental composition | Phase | Morphology (TEM) | Size (nm) (TEM) | Elemental composition | Morphology (TEM) |
| Sigma Aldrich | 19.9 ± 6.6 | TiO ₂ | Anatase and rutile | Mixed cuboid and spherical | n/a | n/a | n/a |
| Sunscreen S1 | 50.0 ± 32.9 | TiO ₂ | Rutile | Needle shaped | 294.3 ± 37.5 | Carbon-based | Spherical |
| Sunscreen S2 | 64.6 ± 26.4 | TiO ₂ | Rutile | Cuboid | n/a | n/a | n/a |
| Product P1 | 158.1 ± 68.7 | TiO ₂ | Anatase | Cuboid | 2705.8 ± 1333.8 | SiO ₂ | Spherical |

difficulty researchers face in the field of nano-ecotoxicology in selecting appropriate NMs for investigation, where NMs belonging to the same class of material vary extensively in physicochemical properties which will inevitably alter their fate and behaviour in the environment. As a result, it is difficult, or impossible, to effectively compare between studies utilising such materials. This issue is exacerbated by the fact that for >50% of NMs used commercially, the chemical structure is unknown.⁴ Additionally, other components of the product formulation may be difficult to separate from nanoparticles during the extraction process.⁶⁴ In studies carried out by Philippe *et al.* (2018), all nTiO₂ extracted from sunscreen was believed to be coated, most commonly with Al or Si.⁹ However, herein, based on EDS mapping, primary particles appeared to be present as pure TiO₂. Product P1 additionally showed the presence of secondary particles – large micron-sized spheres (Fig. 1D) composed of SiO₂ (from EDS measurements, Fig. S1Dii†), a common filler used in cosmetics. On the other hand, S1 showed the additional presence of hollow carbon-based

organic spheres (Fig. 1B; EDS, Fig. S1B†), believed to be micellar structures from additional organic components present in the sunscreen which were not completely removed during extraction. No alumina (Al₂O₃), another common filler in cosmetics, was found during EDS measurements of all samples. Efforts must be directed at producing materials for ecotoxicological research that are representative of those that are likely to enter the natural environment, such as those displayed here from common user products.

3.2 Investigating the toxicity of research-grade and consumer nTiO₂ on the marine cyanobacterium *Prochlorococcus*

The numerically most abundant phototroph on Earth, the marine cyanobacterium *Prochlorococcus*, was grown in the presence of research-grade nTiO₂ (Sigma Aldrich) and nanoparticles extracted from common consumer products, as well as ‘neat’ sunscreen under environmentally relevant conditions, *i.e.*, natural oligotrophic seawater with relevant cell density ($\sim 10^4$ cells mL⁻¹) and nTiO₂ concentrations (1, 50 and 500 $\mu\text{g L}^{-1}$). Following 72 h exposure under simulated natural conditions (Fig. 2), significant declines in population size were experienced by *Prochlorococcus* strain MED4 in response to three out of four nTiO₂ treatments when compared to the untreated control (*i.e.*, research-grade nTiO₂, and nTiO₂ extracted from S1 and P1, two-way *t*-test; $p \leq 0.05$). Despite previous research indicating a key role of TiO₂ phase in determining toxicity,^{22,63} no clear influence of varying nTiO₂ phase was observed. Toxic effects were recorded in both rutile and anatase treatments, as well as mixed phase nTiO₂. It must be acknowledged that due to the methods used to extract nanoparticles, materials may not accurately reflect those released into the environment within product matrixes. Here, behaviour of nanoparticles may be altered by the presence of other components of the product formulation, which require consideration. The process of nanoparticle release from consumer products upon their entry into the environment requires further investigation to comprehensively evaluate their impact.

Upon comparing the response displayed by cultures exposed to research-grade and extracted materials, recorded cell decline of *Prochlorococcus* was greatest during exposure to research-grade nTiO₂ at higher concentrations (*i.e.*, 50 and 500 $\mu\text{g L}^{-1}$). Here, exposure resulted in a significant reduction in cell density of up to 70% compared to the control after 72 h (two-way *t*-test, $p \leq 0.05$), recorded in the 500 $\mu\text{g L}^{-1}$




Fig. 1 TEM images of nTiO₂ utilised in experimental work; A – research-grade nTiO₂ purchased from Sigma Aldrich; B – nTiO₂ extracted from SkinCeuticals™ sunscreen (S1); C – nTiO₂ extracted from Boots Soltan™ sunscreen (S2); D – nTiO₂ extracted from The Body Shop™ liquid foundation (P1).






Fig. 2 Cell density of *Prochlorococcus* MED4 when exposed to nTiO₂ (0–500 µg L⁻¹) for a period of 72 h under simulated natural conditions as measured by flow cytometry; A – research-grade nTiO₂ nanopowder purchased from Sigma Aldrich; B – nTiO₂ extracted from Skinceuticals™ sunscreen (S1); C – nTiO₂ extracted from The Body Shop™ liquid foundation (P1); D – nTiO₂ extracted from Boots Soltan™ sunscreen (S2); E – ‘Neat’ Boots Soltan™ sunscreen immersed in NSW. Data points are presented as the mean ± standard error (*n* = 3). Markers indicate where two-way *t*-tests revealed cell density of treated groups to be significantly lower (crosses) or higher (circles) than the untreated control at each timepoint (*p* ≤ 0.05).

treatment (Fig. 2A). Significant declines were also recorded in the extracted S1 and P1 treatments; however, these were less severe, resulting in decreases in cell density of up to 46%, and 56% compared to the untreated control respectively (two-way *t*-test, *p* ≤ 0.05), at the highest concentration (500 µg L⁻¹) (Fig. 2B and C). This result differs from trends described in previous works in literature, where materials extracted from sunscreen and toothpaste were recorded to exert stronger adverse effects than pristine research-grade nTiO₂.⁶ In that work, it is likely that residual components of product formulations, not fully removed during the extraction process, are toxic at the relatively high concentrations which were tested, are toxic at the relatively high concentrations which were tested (1–5 mg L⁻¹).⁶ Whilst, the highest concentration (500 µg L⁻¹) tested here is greater than that predicted in the environment,¹⁵ some evidence of

Prochlorococcus decline was also observed at 50 µg L⁻¹. This concentration does not far exceed the concentration of TiO₂ derived from sunscreen recorded in areas of high tourism within the natural environment (7–40 µg L⁻¹).¹⁷ Therefore, potential exists for such materials to exert an adverse effect upon marine cyanobacteria such as *Prochlorococcus* in those areas more susceptible to localised pollution.

No adverse effect of exposure was recorded in response to S2 extracted nanoparticles. In fact, after 72 h average cell density of cultures exposed to S2 nanoparticles reached slightly higher values than control cultures (Fig. 2D). Interestingly, herein, cultures exposed nTiO₂ extracted from S1 and P1 also displayed evidence of enhanced growth during early stages (24 h) of experimentation (Fig. 2B and C). Here, an approximate 10% increase in cell density of both the S1 and P1 treatment was observed relative to the untreated



control following 24 h exposure (two-way t -test, $p \leq 0.05$). A beneficial effect of some sunscreen formulations upon phytoplankton growth has been observed in previously published research.⁸ Such increases in cell density may arise from possible biostimulating effects of residual product components such as antioxidants, preservatives and moisturisers, believed able to enhance or stimulate growth of phytoplankton;⁸ or essential nutrients such as N, P and Si, reportedly released by sunscreen upon entry into seawater.¹⁷

Whilst evidence of significant cell decline was observed in the presence of both research-grade and consumer nTiO₂ (after 72 h), these declines were not as severe as those experienced by cultures exposed to 'neat' sunscreen S2 immersed in seawater (Fig. 2E). Here, decline of the cyanobacterial population was the most rapid, and resulted in an 81% decrease of the population by the end of the 72 h incubation at the highest concentration (500 $\mu\text{g L}^{-1}$) relative to the untreated control (two-way t -test, $p \leq 0.05$). However, given 'neat' sunscreen treatments were established based on estimated nTiO₂ content, exact nTiO₂ concentrations may be higher or lower than intended. Nevertheless, sunscreens containing nTiO₂ have previously been recorded to exert toxicity upon phytoplankton *via* generation of ROS,²⁷ which are particularly toxic to *Prochlorococcus* due to its lack of catalase.⁶⁵ Given that no adverse effect was observed when exposing *Prochlorococcus* to nTiO₂ extracted from sunscreen S2, it is likely that toxic effects arise from the other components of the sunscreen formulation which *Prochlorococcus* is highly sensitive to, such as organic compounds or metals.^{66,67} For example, the organic UV-filter octocrylene, found in many sunscreen products, is toxic to a range of marine species.⁶⁸ These findings support the belief that whilst examining the effects of NMs utilised in consumer goods, understanding the exact chemical characteristics of materials is vital. It is important to consider any unknown components of product formulations which may too exert adverse (or stimulating) effects upon biota,⁶⁹ or alter outcomes of toxicity testing. Without such information, we are unable to attribute potential toxicity solely to NMs.

3.3 Medium-term exposure of *Prochlorococcus* to research-grade nTiO₂

Typically, ecotoxicological studies are carried out for a number of days (3–4 d),^{6,21–23,26–28,31,37} thus missing the opportunity to assess chronic effects (>5 d) of a particular substance.⁷⁰ However, examples of longer-term studies do exist.^{71–73} To address this and assess the ability of *Prochlorococcus* MED4 populations to recover from short-term (72 h) stress, incubations with research-grade nTiO₂ observed to exert strongest effects during short-term exposure, were extended to 10 d (Fig. 3). These nanoparticles were identified as mixed rutile and anatase phase by P-XRD (see section 3.1) and hence represent all TiO₂ phase types investigated. In this work, the nTiO₂ concentrations examined were altered to span the predicted environmental ($\mu\text{g L}^{-1}$) and supra-




Fig. 3 Medium-term exposure of *Prochlorococcus* MED4 to research-grade nTiO₂ (Sigma Aldrich, 19.9 ± 6.6 nm (TEM)) in both NSW (A) and nutrient rich Pro99 media (B) at concentrations representing the environmental (0, 1, 10, 100 $\mu\text{g L}^{-1}$) to supra-environmental range (1, 10, 100 mg L^{-1}). Data points are presented as the mean \pm standard error ($n = 3$). Markers indicate where two-way t -tests revealed cell density of treated groups to be significantly lower (crosses) or higher (circles) than the untreated control at each timepoint ($p \leq 0.05$).

environmental range (mg L^{-1}); hence, results are not directly comparable to short-term (72 h) experiments. Additionally, we tested cultures grown in nutrient rich Pro99 media to assess the effect of altered experimental conditions. When monitoring cultures exposed to research-grade nTiO₂ in the mg L^{-1} range, deposited material was visible in the culture flasks, which we suggest to be hetero-aggregations of nTiO₂ and cyanobacterial cells (NP-cell). Due to precipitation arising from increased density, any cells entrapped in NP-cell aggregates would likely be removed from the water column, thus reducing the planktonic cyanobacterial population. Given this, only freely suspended *Prochlorococcus* cells were used for the calculation of cell density, although NP-cell aggregates were also monitored.

Significant declines in the free-living *Prochlorococcus* population were observed at a range of concentrations of added nTiO₂ $\geq 10 \mu\text{g L}^{-1}$ at specific timepoints throughout the 10 d incubation in both NSW and Pro99 media (see Fig. 3). In NSW (Fig. 3A), at the 24 h timepoint only the 10 and 100 mg L^{-1} treatments caused a significant decline in cell density compared to the untreated control (two-way t -test, $p \leq 0.05$). After 48 h the 100 mg L^{-1} continued to exert



significant declines in the cyanobacterial population, additionally the cell density of the $10\ \mu\text{g L}^{-1}$ treatment was also significantly lower than the untreated control (two-way t -test, $p \leq 0.05$). Following 72 h exposure, significant declines were recorded in the $10\ \mu\text{g L}^{-1}$, $100\ \mu\text{g L}^{-1}$, $10\ \text{mg L}^{-1}$ and $100\ \text{mg L}^{-1}$ treatments (two-way t -test, $p \leq 0.05$). However, in later stages of exposure (*i.e.*, 192 and 240 h) only $100\ \text{mg L}^{-1}$ nTiO₂ caused a significant decline in the *Prochlorococcus* population in NSW compared to control cultures (two-way t -test, $p \leq 0.05$). Cultures grown in nutrient rich Pro99 media (Fig. 3B) experienced a decrease in cell density compared to the control in response to $100\ \text{mg L}^{-1}$ nTiO₂ at the each timepoint 24–72 h (two-way t -test, $p \leq 0.05$). Additionally, at the 24 h timepoint, the $10\ \mu\text{g L}^{-1}$ treatment also caused a significant decline in the cyanobacterial population (two-way t -test, $p \leq 0.05$). In later stages of exposure (*i.e.*, 192–240 h) cultures grown in Pro99 media experienced significant declines in cell density in response to 10 and $100\ \text{mg L}^{-1}$ nTiO₂ compared to the untreated control, where at the 192 h timepoint $10\ \mu\text{g L}^{-1}$ was also observed to significantly reduce cell number (two-way t -test, $p \leq 0.05$).

Adverse effects in terms of cell decline were augmented when concentrations were increased to those in the supra-environmental range (*i.e.*, $100\ \text{mg L}^{-1}$, see Fig. 3A and B), likely due to increased rate of encounter between nanoparticles and cyanobacteria. Following 10 d, exposure to $100\ \text{mg L}^{-1}$ nTiO₂ drove cell declines of 79% and 84% in NSW and Pro99 media when compared to the untreated control respectively (two-way t -test, $p \leq 0.05$). Cultures grown to higher cell densities in nutrient-rich media were observed to suffer greater adverse effects of exposure than those grown in NSW (see Fig. 3B). This likely arises due to an increased rate of encounter between nTiO₂ and cyanobacteria in cell-dense cultures. Unexpectedly, no negative effect was observed at the concentration of $1\ \text{mg L}^{-1}$ in NSW or Pro99 media, possibly representing an anomalous result. This being said, evidence of increased growth following incubations with metal oxide NMs has previously been recorded in phytoplankton exposed to nCeO₂ ($<5\ \text{mg L}^{-1}$).^{26,74}

Despite evidence of early declines ($<72\ \text{h}$) in cell number displayed by *Prochlorococcus* following exposure to research-grade nTiO₂ (*i.e.*, $\geq 10\ \mu\text{g L}^{-1}$ and up to 72 h), the ability of the cyanobacterial culture to overcome initial stress in NSW was revealed in extended incubations when exposed to both environmentally relevant and supra-environmental concentrations $\leq 10\ \text{mg L}^{-1}$ (Fig. 3A). The recovery of microbial populations following metal oxide NM exposure has recently been reported.^{70,72,73} For example, decline of *Picochlorum* sp. during early exposure in response to nTiO₂ and ZnO nanoparticles ($10\ \text{mg L}^{-1}$) was reversed in later stages, believed due to aggregation and sedimentation of particles, thus reducing direct exposure of phytoplankton to stable non-sedimenting particles.⁷³ Herein, growth of *Prochlorococcus* was positive in the majority of treatments investigated, despite being significantly lower than the untreated control at specific concentrations, particularly in

early stages. This suggests that although metal oxide NMs such as nTiO₂ may remove a fraction of the microbial population through processes of aggregation and sedimentation, the remaining planktonic population continues to grow and is able to recover to ambient cell densities in extended exposure.

3.4 Physical toxicity and entrapment of cyanobacteria by nTiO₂

Previous research has highlighted that aggregation between metal oxide NMs and phytoplankton is widespread during laboratory exposure.^{23–25,32,75} The physical interaction between NMs and cells, and their entrapment by aggregates is often considered the primary driver of adverse effects experienced by marine microbial organisms.^{23,25,36,75} Adherence of metal oxide NMs to the cell surface is believed to enhance oxidative stress by increasing ROS production^{33,76} and may cause physical damage to the cell wall or membrane.⁷⁵ Aggregation is also proposed to reduce light availability in water, although such shading effects have not




Fig. 4 Estimated number of NP-cell aggregates mL⁻¹ comprised of nTiO₂ and *Prochlorococcus* MED4 observed during flow cytometric analysis of *Prochlorococcus* MED4 exposed to nTiO₂ (Sigma Aldrich) in A) NSW, and B) Pro99 media for a period of 240 h ($n = 3$).






Fig. 5 Hetero-aggregation of *Prochlorococcus* MED4 (stained with SYBR Gold (green)) and research-grade nTiO₂ (Sigma Aldrich; non-stained particles) as observed by fluorescent microscopy; A1–3: untreated *Prochlorococcus* culture; B1–3 *Prochlorococcus* culture exposed to 10 mg L^{−1} nTiO₂; C1–3 *Prochlorococcus* culture exposed to 100 mg L^{−1} nTiO₂; D1–3 natural seawater control (no nTiO₂); E1–3 natural seawater nTiO₂ suspension (100 mg L^{−1}). Images represent those merged from GFP and Brightfield channels.



been conclusively proven.^{34,37,77} Through the subsequent process of sedimentation, hetero-aggregation of NMs and cells effectively removes phytoplankton from the water column, and this is believed to be the primary cause of cell decline in the work presented herein. During flow cytometric analysis of cyanobacterial cultures, it was possible to estimate the number of NP-cell aggregates present within samples in mg L^{-1} treatments (Fig. 4).

The estimated number of NP-cell aggregates was recorded to decline rapidly within the early stages of incubation, indicating their rapid settling and hence removal of cyanobacteria from the water column. In NSW (Fig. 4A) the number of aggregates in the 100 mg L^{-1} treatment declined by 64% between 24 and 48 h. This rapid precipitation and hence decrease in estimated NP-cell aggregate number was also observed in Pro99 supplemented cultures (Fig. 4B), and as expected the total aggregate count far exceeded those recorded in NSW, likely due to increased rate of encounter between nTiO_2 and cyanobacteria at higher cell densities. In previous work, this trend has also been observed, with highest rates of aggregation associated with highest cell density and maximised detrimental effect to cells.³⁴ Such increased rates of aggregation and interaction between nTiO_2 and cyanobacteria may have driven the enhanced decline in cell number observed in cell-dense cultures observed in this study. However, findings vary, and it has also previously been recorded that aggregation and subsequent sedimentation of metal oxide NMs can reduce the negative effects associated with exposure.³⁷ This process may explain the patterns of *Prochlorococcus* recovery observed when exposures were extended in NSW, where following aggregation and subsequent sedimentation of the majority of nTiO_2 , free-living cyanobacterial populations are able to recover.

The hetero-aggregation and subsequent entrapment of *Prochlorococcus* by nTiO_2 was confirmed using fluorescent microscopy (Fig. 5). Here, aggregates of nTiO_2 and cells readily formed in both NSW and nutrient-rich media, observed by microscopy at concentrations of 10 and 100 mg L^{-1} . Control images obtained from NSW and *Prochlorococcus* samples in the absence of nanoparticles, supported the belief that precipitated material observed was indeed nTiO_2 , rather than other particulate matter. The entrapment of phytoplankton by nTiO_2 has previously been observed using microscopic techniques in work carried out with *P. tricornutum* in response to concentrations similar to those used in our study (50 mg L^{-1}).²⁵ Hetero-aggregates were recorded to reach sizes in the micron-range when measured along their longest axis ($2.4\text{--}133.5 \mu\text{m}$, ImageJ analysis), and displayed an average size of $28.0 \mu\text{m}$ ($n = 100$). Such aggregations of research-grade nTiO_2 (25 nm) have been previously recorded to form within 30 minutes of addition into saline f/2 media within this concentration range.³²

In response to NM exposure, marine phytoplankton have been observed to enhance the production of exopolymeric substances (EPS).^{78–82} The secretion of EPS is associated with microbial stress and believed a defence mechanism against

NMs.⁷⁸ For example, EPS was shown to protect the freshwater cyanobacterium *Synechocystis* PCC6803 from nTiO_2 -mediated stress by quenching toxic ROS species and reducing direct contact of nanoparticles with the cell surface.⁸¹ Whilst cyanobacteria are able to produce EPS, *Prochlorococcus* lacks most EPS related proteins, believed lost through the evolutionary streamlining of the *Prochlorococcus* genome due to energy costs of EPS production likely outweighing any benefits.⁸³ As such, any enhancement to EPS production displayed by *Prochlorococcus* during exposures is likely to be limited, and is not believed to be an influential factor in the hetero-aggregation process proposed. However, as the presence of EPS is believed likely to enhance aggregation of nTiO_2 and hence reduce its bioavailability over time,⁸² any increase in EPS production may act to promote the processes of entrapment and co-precipitation described. Indeed, while EPS was recorded to mitigate toxicity in *Synechocystis*, hetero-aggregation between the EPS-producing cyanobacteria and nTiO_2 was still observed.⁸¹

In the natural environment, processes of entrapment and sedimentation may reduce the availability of phytoplankton prey,³⁵ potentially causing an energy deficit in higher trophic levels. Additionally, through ingestion of entrapped phytoplankton, or indirect ingestion during feeding, trophic transport and possible bioaccumulation of metal oxide NMs such as nTiO_2 may occur. Evidence of trophic transfer of nTiO_2 from marine invertebrates to fish has been observed under laboratory conditions.⁸⁴ Once ingested, metal oxide NMs may exert adverse effects upon higher organisms, such as disruption to immune processes and antioxidant activities, as has been recorded in marine invertebrates following exposure to nCeO_2 .^{85,86} Given that nTiO_2 has been recorded to interact with other contaminants within the water column including tributyltin, phenanthrene, polycyclic aromatic hydrocarbons; enhancing their uptake and toxicity to biota,^{87,88} it is important to also consider the potential of metal oxide NM aggregates to adsorb such pollutants which may also be passed to higher trophic levels of the marine food web.

3.5 Behaviour of nTiO_2 in seawater

The fate and behaviour of NMs upon entry into the environment largely determines their bioavailability and mechanism of toxicity towards biota, and is therefore a key consideration during nano-ecotoxicological research.^{73,75,82} Upon entry into saline media with high ionic strength ($\sim 0.7 \text{ eq L}^{-1}$),⁸⁹ metal oxide NMs have generally been observed to aggregate freely and undergo sedimentation, at a faster rate than bulk material.^{36,89} Herein, we utilised dynamic light scattering (DLS) methods to confirm that research-grade nTiO_2 (1 and 100 mg L^{-1}) displays rapid aggregation upon entry into saline media. Hence, supporting our belief that the processes of nTiO_2 aggregation and subsequent co-precipitation with cells is the key driver of cyanobacterial



Table 2 Summary of results obtained from DLS analysis of research-grade nTiO₂ (Sigma Aldrich, 19.9 ± 6.6 nm (TEM)) suspensions in NSW

| Time (h) | 1 mg L ⁻¹ | | 100 mg L ⁻¹ | |
|----------|-----------------------|----------------------------|------------------------|----------------------------|
| | Z-Average size (d.nm) | Polydispersity index (PDI) | Z-Average size (d.nm) | Polydispersity index (PDI) |
| 0 | 1580 ± 124 | 0.804 ± 0.170 | 1624 ± 113 | 0.580 ± 0.156 |
| 1 | 5626 ± 586 | 1.000 | 6560 ± 464 | 1.000 |
| 2 | 2262 ± 129 | 0.890 ± 0.191 | 4994 ± 198 | 1.000 |
| 4 | 1393 ± 284 | 0.907 ± 0.081 | 1569 ± 129 | 0.737 ± 0.108 |
| 24 | 2367 ± 665 | 1.000 | 5636 ± 670 | 0.980 ± 0.035 |
| 48 | 2253 ± 519 | 1.000 | 4133 ± 956 | 0.957 ± 0.055 |
| 72 | 2741 ± 670 | 1.000 | 1966 ± 936 | 0.978 ± 0.025 |
| 168 | Undetectable | | 1994 ± 141 | 1.000 |
| 240 | Undetectable | | 1564 ± 168 | 0.953 ± 0.060 |
| 336 | Undetectable | | 3133 ± 215 | 1.000 |

decline recorded in toxicity experiments described above (sections 3.3 and 3.4).

DLS revealed that nTiO₂ reached sizes far exceeding those of primary particles analysed by TEM (19.9 ± 6.6 nm), displaying an average size of 1547–6560 nm throughout the 14 d experiment (Table 2). These sizes are expected to be somewhat larger than those recorded by TEM (19.9 ± 6.6 nm), since dynamic light scattering measures the hydrodynamic size of colloidal nanoparticles, taking into account van der Waals and interparticle interactions. However, the sizes observed for particles in NSW are far larger than would be expected due to their colloidal behaviour, indicating their rapid aggregation. It should be noted that supra-environmental concentrations have been investigated here due to the poor sensitivity of the equipment to the lower, environmentally relevant concentrations. Nanoparticles were recorded to aggregate immediately upon entry into NSW and DLS measurements displayed that variation in z-average size between replicates was large, suggesting a high variability in aggregate size and low uniformity of this process. This is reflected in the polydispersity index (PDI) values, which remained in the range of 0.58–1.00 throughout the 14 d experiment observed, indicating the presence of a largely polydisperse system. It must be noted, though, that due to biases towards larger particles through use of DLS analysis, smaller particles are likely under-represented.⁹⁰ Precipitation of nTiO₂ from the water column as a result of increased aggregation was recorded, and deposits of nTiO₂ were visible at the bottom of flasks after 24 h incubation with media and throughout the remaining timepoints of the experiment (see Fig. S4†).

The mean z-average size (d.nm) of nTiO₂ at T₀ was 1580 and 1624 nm for 1 and 100 mg L⁻¹ samples respectively, indicating the rapid aggregation of nTiO₂ upon entry into NSW. The PDI at the start of the experiment also indicated a high level of aggregation between nTiO₂ particles, 0.804 and 0.580 for 1 and 100 mg L⁻¹ samples, respectively. Over the first hour, aggregation of nTiO₂ was most rapid, reaching respective peaks in z-average size of 5626 and 6560 nm in the 1 and 100 mg L⁻¹ treatments, whilst PDI reached a value of

1.000. Following this, z-average size in the 1 mg L⁻¹ group appeared to show a slight decrease for subsequent timepoints, although variation remained high, likely indicating that particles are aggregating and sinking, with larger particles no longer within the measurement window of the DLS. PDI values remained high during this period, with a value of 1.000 observed at 24, 48 and 72 h. After 72 h, samples at the 1 mg L⁻¹ concentration were undetectable and therefore unsuitable for data acquisition by DLS. The removal of nTiO₂ from the water column *via* precipitation, observable in flasks after 24 h (see Fig. S3†), and hence reduction in suspended nTiO₂ particles is believed to cause the lack of data acquisition at these timepoints following 72 h during DLS measurement at the 1 mg L⁻¹ concentration.

Mean z-average size of nTiO₂ added to NSW at the 100 mg L⁻¹ concentration varied throughout the 14 d experiment but displayed clear evidence of extensive aggregation of nTiO₂, also indicated by the high PDI values recorded. Overall, z-average size varied 1547–6560 nm during the experiment. Hence, the aggregation of nTiO₂ within NSW appears a rapid, non-uniform and highly variable process. As with lower concentration samples, a large extent of deposition of nTiO₂ aggregates were visible after 24 h and continued throughout the 14 d experiment (see Fig. S3†). In later stages (>72 h), mean z-average size appeared to show a slight decrease. Such data suggests that the size of aggregates remaining in suspension during later stages of the experiment were smaller in size as larger aggregates had likely undergone precipitation and were deposited at the bottom of flasks. Despite the clear evidence of extensive aggregation and sedimentation behaviour of nanoparticles recorded, it should be noted that these observations were made at concentrations considerably higher than those predicted in the environment (due to the low sensitivity of equipment to µg L⁻¹ concentrations). As such, at lower environmental concentrations the rate of homo-aggregation is likely to be reduced, due to effects of dilution and a decreased rate of encounter between individual particles.

Additional information regarding nTiO₂ sedimentation was obtained during flow cytometric analysis by recording the event number of nTiO₂ of varying concentration in NSW in the absence of cyanobacteria (see Section S2.1 and Fig. S3†). Here, particularly in the mg L⁻¹ range, nTiO₂ was observed to undergo rapid sedimentation following entry into seawater, correlating with previous research^{32,36,37,89} and the results acquired by DLS. Up to 58% loss of nTiO₂ *via* precipitation within 6 h has previously been observed using UV-vis spectrophotometry.³² Herein, nTiO₂ counts were recorded to decrease 57%, 81% and 89% in 1, 10 and 100 mg L⁻¹ samples respectively during the initial 24 h of entry into NSW. Following 48–72 h, the number of nTiO₂ aggregates observed in suspension during our work was comparable between all concentrations.

It is clear from DLS and flow cytometry experiments that rapid aggregation and sedimentation of nTiO₂ occurs in marine water. As a result, the bioavailability of nTiO₂ towards



planktonic organisms displays a continual decrease, as has been recorded previously.⁸² As such, the risk of nTiO₂ towards marine microbial species appears lowered during extended exposure. However, at high concentrations, negative effects of exposure may be experienced for longer periods as populations take time to recover or nTiO₂ takes longer durations to enter lower fractions of the water column. In the natural environment the process of reduced bioavailability will likely be concentration-specific, with a lowered rate of encounter between individual nTiO₂ particles at lower concentrations and hence likely lowered homo-aggregation. Extent of sedimentation observed will also be influenced by other environmental variables not investigated in detail here, including pH, and the presence of natural organic and particulate matter.⁹¹ However, despite the high levels of sedimentation recorded in this study, previous environmental assessment by Gondikas *et al.* (2018) has revealed nTiO₂ derived from seasonal use of sunscreen remain in suspension for a period of weeks before settling, where particles may interact with planktonic organisms.¹³

3.6 Identifying molecular features of nTiO₂ toxicity: shotgun proteomic analysis

Alongside observations of cell decline, and direct physical entrapment of cyanobacteria, we performed a shotgun proteomic analysis of *Prochlorococcus* to inform on any other potential toxic effects on this phototroph other than physical entrapment in NM aggregates. Here, exposure to research-grade nTiO₂ did not produce metabolic alterations in *Prochlorococcus*, such as oxidative and nutrient stress or reduction in photosynthetic machinery.

A number of toxic modes of action have been proposed for metal oxide NMs against microbial species, although largely these have been performed at much higher concentrations. Oxidative stress due to the photocatalytic generation of reactive oxygen species (ROS) by nTiO₂ is believed a key driver of stress.^{21,32,92,93} Presence of UV light is therefore a key consideration in experimental design,²⁷ hence in our study full spectrum bulbs with environmentally-relevant levels of UV were utilised. Environmental levels of UV have been previously recorded to induce phototoxic effects of nTiO₂ against phytoplankton with ROS concentration increasing with increased nTiO₂ concentration.²¹ Therefore, it can be proposed that oxidative stress due to ROS production may be a feature of toxicity observed herein during incubations with nTiO₂. Photosynthetic processes have also been observed to be disrupted in response to nTiO₂, such as decreased oxygen generation³⁷ and damage to photosystems.⁹⁴ Metal oxide NMs such as ZnO have been recorded to cause toxicity due to the release of toxic Zn ions,^{31,95,96} however the release of dissolved Ti from nTiO₂ is believed to be negligible under experimental conditions.^{23,97,98} Additionally, TiO₂ has been reported to absorb Zn and P from experimental media,⁹⁹ as

well as to effectively reduce total N and P during microcosm experiments carried out in freshwater.³⁴

Cultures were exposed to an environmentally relevant concentration (100 µg L⁻¹) of research-grade nTiO₂ previously recorded to induce a decline in average cell density in both cell dense and ambient cell densities, for a period of 24 h (7–8% decline (24 h), see Fig. 3). Here, experiments were carried out using cell-dense cultures only grown in Pro99 media to ensure sufficient biological material was collected. Following analysis of both cellular and extracellular proteomes, little difference was observed between treated and control samples (Fig. S5†). Indeed, no individual proteins were significantly altered in abundance following the addition of nTiO₂. Protein function was classified using the Uniprot database and KEGG assignment. Here, little difference was also apparent upon classifying proteins based upon their biological function (see ESI† data Tables S1 and S2, and Fig. S6). The cellular proteome was largely comprised of proteins associated with basic cellular processes, making up ~33% of proteins identified. Proteins involved with central metabolic processes were also relatively abundant in cellular proteome samples and made up ~20% of proteins identified in control and nTiO₂-treated cultures. Despite previous research suggesting disruption to photosynthesis and induction of oxidative stress pathways is likely associated with nTiO₂ exposure, evidence of such effects was not recorded. Herein, proteins associated with photosynthetic processes and oxidative stress represented approximately 6% and 3% of the cellular proteome respectively. Similarly, any evidence of nutrient stress was not apparent. In the extracellular proteome, nTiO₂ exposure was associated with a slight decrease in membrane transport proteins (40% (control) to 36% (nTiO₂)), and slight increase in proteins associated with basic cellular processes (20% (control) to 22% (nTiO₂)) and metabolism (17% (control) to 21% (nTiO₂)). The increase in cytoplasmic-associated proteins may be indicative of a slight increase in cell lysis in treated samples, which may result from hetero-aggregation and entrapment by nTiO₂. Based on the proteomic data reported here, it can be suggested that molecular features of nTiO₂ toxicity may be negligible at the population scale when exposed to environmentally relevant concentrations. Rather, it appears cell decline is primarily driven by physical effects of exposure.

3.7 Effects of consumer nTiO₂ upon natural marine communities

Given the complex nature of the marine microbial community, it is important to consider the community-wide response towards environmental contaminants. Exposure of a natural marine microbial community to nTiO₂ extracted from sunscreen (S2) and 'neat' nTiO₂-containing sunscreen (S2) immersed in seawater at an environmentally relevant concentration of 25 µg L⁻¹,¹⁷ revealed negligible effects of nTiO₂ exposure upon community composition under environmental conditions. Rather, any observed alterations



in community structure were primarily associated with the presence of 'neat' sunscreen, and likely attributed to the other components of the product formulation.

Amplicon sequencing data for bacterial and eukaryotic communities exposed to either treatment was compared to that obtained from an untreated control where no nTiO₂ or sunscreen was added. Principle coordinates analysis based on Bray–Curtis dissimilarity, in conjunction with PERMANOVA analysis (Fig. 6A and B), revealed no significant differences in the 16S rRNA bacterial or 18S rRNA eukaryotic community composition at the ASV level between individual treatments. Similarly, measures of species richness and evenness for both 16S and 18S rRNA datasets, displayed no statistical variation between treatments (see Tables S2 and S3†).

To gain additional insight into the composition of bacterial and eukaryotic communities, the relative abundance of major bacterial and eukaryotic phyla present in control and treated samples (25 µg L⁻¹) was calculated (see Fig. S7 and S8†). Here, bacterial communities were dominated by the Bacteroidetes and Proteobacteria phyla, making up approximately 90% of the total community (Fig. S7†). In the 'neat' sunscreen treatment the proportion of Bacteroidetes to Proteobacteria appeared to differ from both the untreated control and nTiO₂ treatments. In control and extracted nTiO₂-exposed samples average relative abundance of Proteobacteria and Bacteroidetes was approximately 41% and 52–53% respectively, compared to 48% and 44% in 'neat' sunscreen samples (Fig. 6C and D). However, due to large variation between individual samples belonging to the 'neat'




Fig. 6 PCoA plot of 16S rRNA (A) and 18S rRNA (B) data based on Bray–Curtis dissimilarity. Coloured circles indicate samples belonging to each treatment; red circles represent control samples, where no nTiO₂ was added; green circles represent samples treated with nTiO₂ extracted from sunscreen (S2); blue circles represent samples treated with 'neat' sunscreen. PERMANOVA analysis was performed on both datasets respectively; 16S rRNA: F -value = 1.4814, R -squared = 0.27025, p -value < 0.06; 18S rRNA: F -value = 1.2109, R -squared = 0.21203, p -value < 0.144. Panel A and B were adapted from output generated by MicrobiomeAnalyst software. Bar charts presented in panels C–E display the average relative abundance (%) of the Bacteroidia, Gammaproteobacteria and Oxyphotobacteria classes in terms of individual taxonomic orders in each of the treatments respectively.



sunscreen treatment this difference was not significant. Here, the increase in relative abundance of Proteobacteria in the sunscreen treatment was largely due to an increase in relative abundance of Gammaproteobacteria from 25–34%. Specifically, an increase in the Cellvibrionales order made up much of this change (Fig. 6D). The relative decrease in the Bacteroidetes phylum was largely attributed to a decline in relative abundance of the Flavobacteria in response to neat sunscreen, reducing on average from 51% in the untreated control to 41% (Fig. 6C). The reduction in the ratio of Bacteroidetes to Proteobacteria has previously been recorded in marine biofilm communities exposed to nTiO₂-treated surfaces.¹⁰⁰ Similarly, in riverine environments, Bacteroidetes members such as Flavobacteria have been recorded to display increased susceptibility to nTiO₂ during community exposure.¹⁰¹ In this example, Actinobacteria also displayed a reduced relative abundance in the presence of nTiO₂, whereas comparatively Betaproteobacteria were observed to increase by approximately 40% under the same conditions in comparison to the untreated control.¹⁰¹ These results were obtained at a test concentration of 100 mg L⁻¹, far exceeding that measured in the environment.¹⁷ It is possible that should nTiO₂ concentrations have been increased in this experiment, any decline in abundance of Flavobacteria in response to nTiO₂-containing sunscreen would have been exacerbated and perhaps observed in the extracted nTiO₂ treatment. However, in our study it appears these alterations arise as a result of other components of the sunscreen formulation rather than the action of nTiO₂, as no such changes were observed in the extracted nTiO₂ treatment. Despite *Prochlorococcus* displaying declines in cell density in earlier testing (sections 3.2 and 3.3), no significant effect on relative abundance of cyanobacteria was recorded in either treatment during community exposure. In fact, on average a slight increase in cyanobacteria was observed in both extracted nTiO₂ and neat sunscreen treatments (Fig. 6E). Overall, only two bacterial phyla were observed to alter significantly in relative abundance between control and extracted nTiO₂ or sunscreen-treated samples as a result of two-way *t*-tests. Both were observed in the extracted nTiO₂ (S2) treatment and belonged to low abundant phyla; here, Fusobacteria displayed a significant increase in relative abundance from 0.007% in the control to 0.035% in the extracted S2 nTiO₂ treatment (*p* = 0.043); whilst Lentisphaerae, reduced in relative abundance from 0.0015% in the control to 0.0011% in S2 nTiO₂ samples (*p* = 0.024).

Eukaryotic communities were principally made up of phototrophic organisms belonging to the Ochrophyta, Prymnesiophyceae, Chlorophyta and Protalveolata phyla, with Arthropoda also relatively abundant in control and 'neat' sunscreen treatments (Fig. S8†). Species belonging to the Ochrophyta phylum were most abundant in all treatments, representing 28–30% of ASVs identified. The Ochrophyta largely comprises phototrophic organisms, and here diatoms made up the majority of taxa in this phylum and represented 24–25% of eukaryotic species identified across all treatments.

Once again, differences in relative abundance of eukaryotic phyla between control and treated samples were tested by means of two-way *t*-tests. Here, no significant differences were identified despite evidence being available upon the negative effect of nTiO₂ exposure towards eukaryotic phytoplankton in the literature, albeit largely recorded at greater concentrations (1–13 mg L⁻¹).^{6,21–30} Previous research has revealed that at such concentrations, varying eukaryotic taxa display differential sensitivity to nTiO₂ exposure.^{21,27} For example, Sendra *et al.* (2017) present that following exposure to nTiO₂, and nTiO₂-containing sunscreens the marine microalgae *Nannochloropsis gaditana* displays highest sensitivity to all treatments compared to the diatom, *Chaetoceros gracilis*, the dinoflagellate, *Amphidinium carterae*, and coccolithophore, *Pleurochrysis roscoffensis*.²⁷ Whilst such evidence exists, our results suggests that marine eukaryotic communities are little affected by exposure to environmentally-relevant concentrations of nTiO₂ extracted from, or in the presence of nTiO₂-containing sunscreen.

The findings presented suggest that current concentrations nTiO₂ are unlikely to drive alterations to the structure and biodiversity of natural marine communities and is therefore unlikely to impact upon community functioning. Based on this, we can predict that the likely environmental risk of nTiO₂ derived from consumer products such as sunscreen towards marine microbial communities is low. However, it remains that should contamination of the marine environment by NMs continue to increase, evidence suggests that phototrophic organisms may be negatively affected within hotspots of pollution comprising higher concentrations, mainly due to hetero-aggregation and removal from the water column *via* sinking.

4. Conclusions

Overall, we have shown that the interaction between marine microbial species and nTiO₂ appears a highly dynamic process influenced largely by the behaviour of nanoparticles in saline media. Toxic endpoints appear dependent upon the length of exposure, where risks associated with nTiO₂ exposure to marine microbial species appear low at currently predicted environmental concentrations. We have shown for the first time that the ecologically significant cyanobacterium, *Prochlorococcus*, suffers short-term (72 h) adverse effects following exposure to nTiO₂, mainly due to hetero-aggregation with agglomerated nTiO₂, and subsequent sinking out of the water column. Nevertheless, populations could recover when incubations were extended in natural oligotrophic seawater. No other sign of metabolic stress was observed by high throughput proteomics at a concentration relevant to those predicted in the environment, suggesting that physical interactions between nanoparticles and cyanobacteria are responsible for cell declines recorded. Monitoring of natural marine bacterial and eukaryotic communities exposed to nTiO₂ and nTiO₂-containing sunscreen, revealed no effect of either treatment at current



environmental concentrations, suggesting that neither is likely to alter marine community structure or diversity in the natural environment.

The works presented here support the belief that the current environmental risk of nTiO₂ is low, however uncertainties exist in the environmental concentrations of NMs currently predicted,¹⁰² and hotspots of contamination may serve to produce localised areas of cell decline given that severity of negative effects is enhanced with increasing nTiO₂ concentration. The interaction and synergistic effect of NMs and other contaminants also requires attention given the high affinity of materials such as nTiO₂ with other contaminants,^{88,103,104} which may act to enhance toxicity at lower concentrations. It is important that we continue to develop analytical techniques to reveal NM concentrations within the natural environment and enhance experimental work by studying materials that represent those likely to be entered into aquatic systems. As such, the use of extracted materials as we have examined in the above study provides great scope for future research in the field. However, there is a need to consider the whole product formulation in research on nano-enabled products, where specific components may play key roles in toxicity which are not identified when using uncoated nanoparticles. In particular, processes of NM release from product matrixes and their resultant state requires attention and is important to fully assess the environmental impact of engineered nanomaterials.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

C. J. D. was supported by the NERC CENTA DTP studentship NE/L002493/1. J. A. C.-O. was funded by a NERC Independent Research Fellowship NE/K009044/1, Ramón y Cajal contract RYC-2017-22452 (co-funded by the Ministerio de Ciencia, Innovación y Universidades, the Agencia Estatal de Investigación (10.13039/100014440), and the European Social Fund (10.13039/501100004895)) and project PID2019-109509RB-I00 / AEI / 10.13039/501100011033. In addition, we thank the BBSRC/EPSC Synthetic Biology Research Centre WISB (grant ref.: BB/M017982/1) for access to equipment. Particular thanks are given to Dr Steve Firth at University College London for assistance with EDS measurements, as well as Dr Robyn Wright and Dr Gabriel Erni Cassola for assistance with 16S and 18S rRNA amplicon sequencing and analysis.

References

- 1 A. Nel, T. Xia, L. Madler and N. Li, Toxic potential of materials at the nanolevel, *Science*, 2006, **311**(5761), 622–627.
- 2 B. Nowack and T. D. Bucheli, Occurrence, behavior and effects of nanoparticles in the environment, *Environ. Pollut.*, 2007, **150**(1), 5–22.
- 3 Y. Zhu, X. Liu, Y. Hu, R. Wang, M. Chen, J. Wu, Y. Wang, S. Kang, Y. Sun and M. Zhu, Behavior, remediation effect and toxicity of nanomaterials in water environments, *Environ. Res.*, 2019, **174**, 54–60.
- 4 M. E. Vance, T. Kuiken, E. P. Vejerano, S. P. McGinnis, M. F. Hochella Jr., D. Rejeski and M. S. Hull, Nanotechnology in the real world: Redeveloping the nanomaterial consumer products inventory, *Beilstein J. Nanotechnol.*, 2015, **6**, 1769–1780.
- 5 F. Gottschalk, T. Sonderer, R. W. Scholz and B. Nowack, Modeled environmental concentrations of engineered nanomaterials (TiO₂, ZnO, Ag, CNT, Fullerenes) for different regions, *Environ. Sci. Technol.*, 2009, **43**(24), 9216–9222.
- 6 A. Galletti, S. Seo, S. H. Joo, C. Su and P. Blackwelder, Effects of titanium dioxide nanoparticles derived from consumer products on the marine diatom *Thalassiosira pseudonana*, *Environ. Sci. Pollut. Res.*, 2016, **23**(20), 21113–21122.
- 7 A. Menard, D. Drobne and A. Jemec, Ecotoxicity of nanosized TiO₂. Review of in vivo data, *Environ. Pollut.*, 2011, **159**(3), 677–684.
- 8 S. Schiavo, M. Oliviero, A. Philippe and S. Manzo, Nanoparticles based sunscreens provoke adverse effects on marine microalgae *Dunaliella tertiolecta*, *Environ. Sci.: Nano*, 2018, **5**, 3011–3022.
- 9 A. Philippe, J. Kosik, A. Welle, J.-M. Guigner, O. Clemens and G. E. Schaumann, Extraction and characterization methods for titanium dioxide nanoparticles from commercialized sunscreens, *Environ. Sci.: Nano*, 2018, **5**, 191–202.
- 10 R. Danovaro, L. Bongiorni, C. Corinaldesi, D. Giovannelli, E. Damiani, P. Astolfi, L. Greci and A. Pusceddu, Sunscreens cause coral bleaching by promoting viral infections, *Environ. Health Perspect.*, 2008, **116**(4), 441–447.
- 11 L. K. Limbach, R. Bereiter, E. Muller, R. Krebs, R. Galli and W. J. Stark, Removal of oxide nanoparticles in a model wastewater treatment plant: influence of agglomeration and surfactants on clearing efficiency, *Environ. Sci. Technol.*, 2008, **42**(15), 5828–5833.
- 12 R. F. Domingos, M. A. Baalousha, Y. Ju-Nam, M. M. Reid, N. Tufenkji, J. R. Lead, G. G. Leppard and K. J. Wilkinson, Characterizing manufactured nanoparticles in the environment: multimethod determination of particle sizes, *Environ. Sci. Technol.*, 2009, **43**(19), 7277–7284.
- 13 A. Gondikas, F. von der Kammer, R. Kaegi, O. Borovinskaya, E. Neubauer, J. Navratilova, A. Praetorius, G. Cornelis and T. Hofmann, Where is the nano? Analytical approaches for the detection and quantification of TiO₂ engineered nanoparticles in surface waters, *Environ. Sci.: Nano*, 2018, **5**, 313–326.
- 14 A. Praetorius, M. Scheringer and K. Hungerbühler, Development of environmental fate models for engineered nanoparticles—a case study of TiO₂ nanoparticles in the Rhine River, *Environ. Sci. Technol.*, 2012, **46**(12), 6705–6713.



- 15 M. A. Maurer-Jones, I. L. Gunsolus, C. J. Murphy and C. L. Haynes, Toxicity of engineered nanoparticles in the environment, *Anal. Chem.*, 2013, **85**(6), 3036–3049.
- 16 P. S. Bauerlein, E. Emke, P. Tromp, J. Hofman, A. Carboni, F. Schooneman, P. de Voogt and A. P. van Wezel, Is there evidence for man-made nanoparticles in the Dutch environment?, *Sci. Total Environ.*, 2017, **576**, 273–283.
- 17 A. Tovar-Sanchez, D. Sanchez-Quiles, G. Basterretxea, J. L. Benede, A. Chisvert, A. Salvador, I. Moreno-Garrido and J. Blasco, Sunscreen products as emerging pollutants to coastal waters, *PLoS One*, 2013, **8**(6), e65451.
- 18 A. C. Johnson and B. Park, Predicting contamination by the fuel additive cerium oxide engineered nanoparticles within the United Kingdom and the associated risks, *Environ. Toxicol. Chem.*, 2012, **31**(11), 2582–2587.
- 19 C. B. Field, M. J. Behrenfeld, J. T. Randerson and P. Falkowski, Primary production of the biosphere: integrating terrestrial and oceanic components, *Science*, 1998, **281**(5374), 237–240.
- 20 P. Flombaum, J. L. Gallegos, R. A. Gordillo, J. Rincon, L. L. Zabala, N. Jiao, D. M. Karl, W. K. Li, M. W. Lomas, D. Veneziano, C. S. Vera, J. A. Vrugt and A. C. Martiny, Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**(24), 9824–9829.
- 21 R. J. Miller, S. Bennett, A. A. Keller, S. Pease and H. S. Lenihan, TiO₂ nanoparticles are phototoxic to marine phytoplankton, *PLoS One*, 2012, **7**(1), e30321.
- 22 L. Clement, C. Hurel and N. Marmier, Toxicity of TiO₂ nanoparticles to cladocerans, algae, rotifers and plants - effects of size and crystalline structure, *Chemosphere*, 2013, **90**(3), 1083–1090.
- 23 B. Xia, B. Chen, X. Sun, K. Qu, F. Ma and M. Du, Interaction of TiO₂ nanoparticles with the marine microalga *Nitzschia closterium*: growth inhibition, oxidative stress and internalization, *Sci. Total Environ.*, 2015, **508**, 525–533.
- 24 S. Manzo, S. Buono, G. Rametta, M. Miglietta, S. Schiavo and G. Di Francia, The diverse toxic effect of SiO₂ and TiO₂ nanoparticles toward the marine microalgae *Dunaliella tertiolecta*, *Environ. Sci. Pollut. Res.*, 2015, **22**(20), 15941–15951.
- 25 Y. Wang, X. Zhu, Y. Lao, X. Lv, Y. Tao, B. Huang, J. Wang, J. Zhou and Z. Cai, TiO₂ nanoparticles in the marine environment: Physical effects responsible for the toxicity on algae *Phaeodactylum tricornutum*, *Sci. Total Environ.*, 2016, **565**, 818–826.
- 26 X. Y. Deng, J. Cheng, X. L. Hu, L. Wang, D. Li and K. Gao, Biological effects of TiO₂ and CeO₂ nanoparticles on the growth, photosynthetic activity, and cellular components of a marine diatom *Phaeodactylum tricornutum*, *Sci. Total Environ.*, 2017, **575**, 87–96.
- 27 M. Sendra, D. Sanchez-Quiles, J. Blasco, I. Moreno-Garrido, L. M. Lubian, S. Perez-Garcia and A. Tovar-Sanchez, Effects of TiO₂ nanoparticles and sunscreens on coastal marine microalgae: Ultraviolet radiation is key variable for toxicity assessment, *Environ. Int.*, 2017, **98**, 62–68.
- 28 B. Xia, Q. Sui, X. Sun, Q. Han, B. Chen, L. Zhu and K. Qu, Ocean acidification increases the toxic effects of TiO₂ nanoparticles on the marine microalga *Chlorella vulgaris*, *J. Hazard. Mater.*, 2018, **346**, 1–9.
- 29 M. Li, D. Chen, Y. Liu, C. Y. Chuang, F. Kong, P. J. Harrison, X. Zhu and Y. Jiang, Exposure of engineered nanoparticles to *Alexandrium tamarense* (Dinophyceae): Healthy impacts of nanoparticles via toxin-producing dinoflagellate, *Sci. Total Environ.*, 2018, **610–611**, 356–366.
- 30 V. Thiagarajan, S. Ramasubbu, C. Natarajan and A. Mukherjee, Differential sensitivity of marine algae *Dunaliella salina* and *Chlorella* sp. to P25 TiO₂ NPs, *Environ. Sci. Pollut. Res.*, 2019, **26**(21), 21394–21403.
- 31 R. J. Miller, H. S. Lenihan, E. B. Muller, N. Tseng, S. K. Hanna and A. A. Keller, Impacts of metal oxide nanoparticles on marine phytoplankton, *Environ. Sci. Technol.*, 2010, **44**(19), 7329–7334.
- 32 E. Morelli, E. Gabellieri, A. Bonomini, D. Tognotti, G. Grassi and I. Corsi, TiO₂ nanoparticles in seawater: Aggregation and interactions with the green alga *Dunaliella tertiolecta*, *Ecotoxicol. Environ. Saf.*, 2018, **148**, 184–193.
- 33 L. Fu, M. Hamzeh, S. Dodard, Y. H. Zhao and G. I. Sunahara, Effects of TiO₂ nanoparticles on ROS production and growth inhibition using freshwater green algae pre-exposed to UV irradiation, *Environ. Toxicol. Pharmacol.*, 2015, **39**(3), 1074–1080.
- 34 M. Bessa da Silva, N. Abrantes, V. Nogueira, F. Goncalves and R. Pereira, TiO₂ nanoparticles for the remediation of eutrophic shallow freshwater systems: Efficiency and impacts on aquatic biota under a microcosm experiment, *Aquat. Toxicol.*, 2016, **178**, 58–71.
- 35 B. Campos, C. Rivetti, P. Rosenkranz, J. M. Navas and C. Barata, Effects of nanoparticles of TiO₂ on food depletion and life-history responses of *Daphnia magna*, *Aquat. Toxicol.*, 2013, 174–183.
- 36 M. Sendra, M. P. Yeste, J. M. Gatica, I. Moreno-Garrido and J. Blasco, Homoagglomeration and heteroagglomeration of TiO₂, in nanoparticle and bulk form, onto freshwater and marine microalgae, *Sci. Total Environ.*, 2017, **592**, 403–411.
- 37 J. Hu, J. Wang, S. Liu, Z. Zhang, H. Zhang, X. Cai, J. Pan and J. Liu, Effect of TiO₂ nanoparticle aggregation on marine microalgae *Isochrysis galbana*, *J. Environ. Sci.*, 2018, **66**, 208–215.
- 38 J. M. Farner, R. S. Cheong, E. Mahe, H. Anand and N. Tufenkji, Comparing TiO₂ nanoparticle formulations: stability and photoreactivity are key factors in acute toxicity to *Daphnia magna*, *Environ. Sci.: Nano*, 2019, **6**, 2532–2543.
- 39 D. J. Scanlan, M. Ostrowski, S. Mazard, A. Dufresne, L. Garczarek, W. R. Hess, A. F. Post, M. Hagemann, I. Paulsen and F. Partensky, Ecological genomics of marine picocyanobacteria, *Microbiol. Mol. Biol. Rev.*, 2009, **73**(2), 249–299.
- 40 S. C. Bagby and S. W. Chisholm, Response of *Prochlorococcus* to varying CO₂:O₂ ratios, *ISME J.*, 2015, **9**(10), 2232–2245.



- 41 C. J. Dedman, G. C. Newson, G.-L. Davies and J. A. Christie-Oleza, Mechanisms of silver nanoparticle toxicity on the marine cyanobacterium *Prochlorococcus* under environmentally-relevant conditions, *Sci. Total Environ.*, 2020, 141229.
- 42 L. Ma, B. C. Calfee, J. J. Morris, Z. I. Johnson and E. R. Zinser, Degradation of hydrogen peroxide at the ocean's surface: the influence of the microbial community on the realized thermal niche of *Prochlorococcus*, *ISME J.*, 2018, **12**, 473–484.
- 43 D. Mella-Flores, S. Mazard, F. Humily, F. Partensky, F. Mahe, L. Bariat, C. Courties, D. Marie, J. Ras, R. Mauriac, C. Jeanthon, E. Mahdi Bendif, M. Ostrowski, D. J. Scanlan and L. Garczarek, Is the distribution of *Prochlorococcus* and *Synechococcus* ecotypes in the Mediterranean Sea affected by global warming?, *Biogeosciences*, 2011, **8**, 2785–2804.
- 44 L. R. Moore, A. F. Post, G. Rocap and S. W. Chisholm, Utilization of different nitrogen sources by the marine cyanobacteria *Prochlorococcus* and *Synechococcus*, *Limnol. Oceanogr.*, 2002, **47**(4), 989–996.
- 45 J. A. Christie-Oleza and J. Armengaud, In-depth analysis of exoproteomes from marine bacteria by shotgun liquid chromatography-tandem mass spectrometry: the *Ruegeria pomeroyi* DSS-3 case-study, *Mar. Drugs*, 2010, **8**(8), 2223–2239.
- 46 V. Zadjelovic, A. Chhun, M. Quareshy, E. Silvano, J. R. Hernandez-Fernaund, M. M. Aguilo-Ferretjans, R. Bosch, C. Dorador, M. I. Gibson and J. A. Christie-Oleza, Beyond oil degradation: enzymatic potential of *Alcanivorax* to degrade natural and synthetic polyesters, *Environ. Microbiol.*, 2020, **22**(4), 1356–1369.
- 47 A. Shevchenko, H. Tomas, J. Havlis, J. V. Olsen and M. Mann, In-gel digestion for mass spectrometric characterization of proteins and proteomes, *Nat. Protoc.*, 2006, **1**(6), 2856–2860.
- 48 J. A. Christie-Oleza, D. J. Scanlan and J. Armengaud, “You produce while I clean up”, a strategy revealed by exoproteomics during *Synechococcus*-*Roseobacter* interactions, *Proteomics*, 2015, **15**(20), 3454–3462.
- 49 J. Cox and M. Mann, MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification, *Nat. Biotechnol.*, 2008, **26**(12), 1367–1372.
- 50 R. J. Wright, M. I. Gibson and J. A. Christie-Oleza, Understanding microbial community dynamics to improve optimal microbiome selection, *Microbiome*, 2019, **7**(1), 85.
- 51 A. E. Parada, D. M. Needham and J. A. Fuhrman, Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples, *Environ. Microbiol.*, 2016, **18**(5), 1403–1414.
- 52 I. M. Bradley, A. J. Pinto and J. S. Guest, Design and Evaluation of Illumina MiSeq-Compatible, 18S rRNA Gene-Specific Primers for Improved Characterization of Mixed Phototrophic Communities, *Appl. Environ. Microbiol.*, 2016, **82**(19), 5878–5891.
- 53 B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. Johnson and S. P. Holmes, DADA2: High-resolution sample inference from Illumina amplicon data, *Nat. Methods*, 2016, **13**(7), 581–583.
- 54 B. J. Callahan, K. Sankaran, J. A. Fukuyama, P. J. McMurdie and S. P. Holmes, Bioconductor Workflow for Microbiome Data Analysis: from raw reads to community analyses, *F1000Research*, 2016, **5**, 1492.
- 55 L. W. Hugerth and A. F. Andersson, Analysing Microbial Community Composition through Amplicon Sequencing: From Sampling to Hypothesis Testing, *Front. Microbiol.*, 2017, **8**, 1561.
- 56 S. Tyanova, T. Temu, P. Sinitcyn, A. Carlson, M. Y. Hein, T. Geiger, M. Mann and J. Cox, The Perseus computational platform for comprehensive analysis of (prote)omics data, *Nat. Methods*, 2016, **13**(9), 731–740.
- 57 A. Kaur, J. R. Hernandez-Fernaund, M. D. M. Aguilo-Ferretjans, E. M. Wellington and J. A. Christie-Oleza, 100 Days of marine *Synechococcus*-*Ruegeria* *pomeroyi* interaction: A detailed analysis of the exoproteome, *Environ. Microbiol.*, 2018, **20**(2), 785–799.
- 58 Y. Perez-Riverol, A. Csordas, J. Bai, M. Bernal-Llinares, S. Hewapathirana, D. J. Kundu, A. Inuganti, J. Griss, G. Mayer, M. Eisenacher, E. Perez, J. Uszkoreit, J. Pfeuffer, T. Sachsenberg, S. Yilmaz, S. Tiwary, J. Cox, E. Audain, M. Walzer, A. F. Jarnuczak, T. Ternent, A. Brazma and J. A. Vizcaino, The PRIDE database and related tools and resources in 2019: improving support for quantification data, *Nucleic Acids Res.*, 2019, **47**(D1), D442–D450.
- 59 J. Chong, P. Liu, G. Zhou and J. Xia, Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data, *Nat. Protoc.*, 2020, **15**(3), 799–821.
- 60 A. Dhariwal, J. Chong, S. Habib, I. L. King, L. B. Agellon and J. Xia, MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data, *Nucleic Acids Res.*, 2017, **45**(W1), W180–W188.
- 61 Y. Liao, W. Que, Q. Jia, Y. He, J. Zhang and P. Zhong, Controllable synthesis of brookite/anatase/rutile TiO₂ nanocomposites and single-crystalline rutilenanorods array, *J. Mater. Chem.*, 2012, **22**, 7937–7944.
- 62 J.-G. Li, T. Ishigaki and X. Sun, Anatase, Brookite, and Rutile Nanocrystals via Redox Reactions under Mild Hydrothermal Conditions: Phase-Selective Synthesis and Physicochemical Properties, *J. Phys. Chem. C*, 2007, **111**(13), 4969–4976.
- 63 X. Chen, Y. Zhu, K. Yang, L. Zhu and D. Lin, Nanoparticle TiO₂ size and rutile content impact bioconcentration and biomagnification from algae to daphnia, *Environ. Pollut.*, 2019, **247**, 421–430.
- 64 A. Al-Kattan, A. Wichser, S. Zuin, Y. Arroyo, L. Golanski, A. Ulrich and B. Nowack, Behavior of TiO₂ released from Nano-TiO₂-containing paint and comparison to pristine Nano-TiO₂, *Environ. Sci. Technol.*, 2014, **48**(12), 6710–6718.



- 65 J. J. Morris, R. E. Lenski and E. R. Zinser, The Black Queen Hypothesis: evolution of dependencies through adaptive gene loss, *MBio*, 2012, **3**(2), e00036-12.
- 66 L. R. Moore, A. Coe, M. A. Saito, M. B. Sullivan, D. Lindell, K. Frois-Moniz, J. Waterbury and S. W. Chisholm, Culturing the marine cyanobacterium *Prochlorococcus*, *Limnol. Oceanogr.: Methods*, 2007, **5**(10), 353–362.
- 67 S. G. Tetu, I. Sarker, V. Schrameyer, R. Pickford, L. D. H. Elbourne, L. R. Moore and I. T. Paulsen, Plastic leachates impair growth and oxygen production in *Prochlorococcus*, the ocean's most abundant photosynthetic bacteria, *Commun. Biol.*, 2019, **2**, 184.
- 68 A. Giraldo, R. Montes, R. Rodil, J. B. Quintana, L. Vidal-Linan and R. Beiras, Ecotoxicological Evaluation of the UV Filters Ethylhexyl Dimethyl p-Aminobenzoic Acid and Octocrylene Using Marine Organisms *Isochrysis galbana*, *Mytilus galloprovincialis* and *Paracentrotus lividus*, *Arch. Environ. Contam. Toxicol.*, 2017, **72**(4), 606–611.
- 69 A. Tovar-Sanchez, D. Sanchez-Quiles and A. Rodriguez-Romero, Massive coastal tourism influx to the Mediterranean Sea: The environmental risk of sunscreens, *Sci. Total Environ.*, 2019, **656**, 316–321.
- 70 S. L. Mitchell, N. V. Hudson-Smith, M. S. Cahill, B. N. Reynolds, S. D. Frand, C. M. Green, C. Wang, Z. V. Feng, C. L. Haynes and E. E. Carlson, Chronic exposure to complex metal oxide nanoparticles elicits rapid resistance in *Shewanella oneidensis* MR-1, *Chem. Sci.*, 2019, **10**, 9768–9781.
- 71 G. K. Bielmyer-Fraser, T. A. Jarvis, H. S. Lenihan and R. J. Miller, Cellular partitioning of nanoparticulate versus dissolved metals in marine phytoplankton, *Environ. Sci. Technol.*, 2014, **48**(22), 13443–13450.
- 72 M. Li, Y. Jiang, C. Y. Chuang, J. Zhou, X. Zhu and D. Chen, Recovery of *Alexandrium tamarense* under chronic exposure of TiO₂ nanoparticles and possible mechanisms, *Aquat. Toxicol.*, 2019, **208**, 98–108.
- 73 L. J. Hazeem, M. Bououdina, S. Rashdan, L. Brunet, C. Slomianny and R. Boukherroub, Cumulative effect of zinc oxide and titanium oxide nanoparticles on growth and chlorophyll a content of *Picochlorum* sp, *Environ. Sci. Pollut. Res.*, 2016, **23**(3), 2821–2830.
- 74 M. Sendra, J. Blasco and C. V. M. Araujo, Is the cell wall of marine phytoplankton a protective barrier or a nanoparticle interaction site? Toxicological responses of *Chlorella autotrophica* and *Dunaliella salina* to Ag and CeO₂ nanoparticles, *Ecol. Indic.*, 2018, **95**, 1053–1067.
- 75 I. Rodea-Palomares, K. Boltes, F. Fernández-Piñas, F. Leganés, E. García-Calvo, J. Santiago and R. Rosal, Physicochemical characterization and ecotoxicological assessment of CeO₂ nanoparticles using two aquatic microorganisms, *Toxicol. Sci.*, 2010, **119**(1), 135–145.
- 76 D. M. Metzler, M. Li, A. Erdem and C. P. Huang, Responses of algae to photocatalytic nano-TiO₂ particles with an emphasis on the effect of particle size, *Chem. Eng. J.*, 2011, **170**, 538–546.
- 77 V. Aruoja, H. C. Dubourguier, K. Kasemets and A. Kahru, Toxicity of nanoparticles of CuO, ZnO and TiO₂ to microalgae *Pseudokirchneriella subcapitata*, *Sci. Total Environ.*, 2009, **407**(4), 1461–1468.
- 78 M. H. Chiu, Z. A. Khan, S. G. Garcia, A. D. Le, A. Kagiri, J. Ramos, S. M. Tsai, H. W. Drobenaire, P. H. Santschi, A. Quigg and W. C. Chin, Effect of Engineered Nanoparticles on Exopolymeric Substances Release from Marine Phytoplankton, *Nanoscale Res. Lett.*, 2017, **12**(1), 620.
- 79 S. Zheng, Q. Zhou, C. Chen, F. Yang, Z. Cai, D. Li, Q. Geng, Y. Feng and H. Wang, Role of extracellular polymeric substances on the behavior and toxicity of silver nanoparticles and ions to green algae *Chlorella vulgaris*, *Sci. Total Environ.*, 2019, **660**, 1182–1190.
- 80 K. Zhou, Y. Hu, L. Zhang, K. Yang and D. Lin, The role of exopolymeric substances in the bioaccumulation and toxicity of Ag nanoparticles to algae, *Sci. Rep.*, 2016, **6**, 32998.
- 81 M. Planchon, T. Jittawuttipoka, C. Cassier-Chauvat, F. Guyot, A. Gelabert, M. F. Benedetti, F. Chauvat and O. Spalla, Exopolysaccharides protect *Synechocystis* against the deleterious effects of titanium dioxide nanoparticles in natural and artificial waters, *J. Colloid Interface Sci.*, 2013, **405**, 35–43.
- 82 V. Thiagarajan, M. Pavani, S. Archanaa, R. Seenivasan, N. Chandrasekaran, G. K. Suraishkumar and A. Mukherjee, Diminishing bioavailability and toxicity of P25 TiO₂ NPs during continuous exposure to marine algae *Chlorella* sp, *Chemosphere*, 2019, **233**, 363–372.
- 83 S. B. Pereira, R. Mota, C. P. Vieira, J. Vieira and P. Tamagnini, Phylum-wide analysis of genes/proteins related to the last steps of assembly and export of extracellular polymeric substances (EPS) in cyanobacteria, *Sci. Rep.*, 2015, **5**, 14835.
- 84 Z. Wang, L. Yin, J. Zhao and B. Xing, Trophic transfer and accumulation of TiO₂ nanoparticles from clamworm (*Perinereis aibuhitensis*) to juvenile turbot (*Scophthalmus maximus*) along a marine benthic food chain, *Water Res.*, 2016, **95**, 250–259.
- 85 C. Falugi, M. G. Aluigi, M. C. Chiantore, D. Privitera, P. Ramoino, M. A. Gatti, A. Fabrizi, A. Pinsino and V. Matranga, Toxicity of metal oxide nanoparticles in immune cells of the sea urchin, *Mar. Environ. Res.*, 2012, **76**, 114–121.
- 86 M. Garaud, J. Trapp, S. Devin, C. Cossu-Leguille, S. Pain-Devin, V. Felten and L. Giamberini, Multibiomarker assessment of cerium dioxide nanoparticle (nCeO₂) sublethal effects on two freshwater invertebrates, *Dreissena polymorpha* and *Gammarus roeseli*, *Aquat. Toxicol.*, 2015, **158**, 63–74.
- 87 X. Zhu, J. Zhou and Z. Cai, TiO₂ nanoparticles in the marine environment: impact on the toxicity of tributyltin to abalone (*Haliotis diversicolor supertexta*) embryos, *Environ. Sci. Technol.*, 2011, **45**(8), 3753–3758.
- 88 J. Lu, S. Tian, X. Lv, Z. Chen, B. Chen, X. Zhu and Z. Cai, TiO₂ nanoparticles in the marine environment: Impact on the toxicity of phenanthrene and Cd²⁺ to marine zooplankton *Artemia salina*, *Sci. Total Environ.*, 2018, **615**, 375–380.



- 89 A. A. Keller, H. Wang, D. Zhou, H. S. Lenihan, G. Cherr, B. J. Cardinale, R. Miller and Z. Ji, Stability and aggregation of metal oxide nanoparticles in natural aqueous matrices, *Environ. Sci. Technol.*, 2010, **44**(6), 1962–1967.
- 90 R. Tantra, S. Jing, S. K. Pichaimuthu, N. Walker, J. Noble and V. A. Hackley, Dispersion stability of nanoparticles in ecotoxicological investigations: the need for adequate measurement tools, *J. Nanopart. Res.*, 2011, **13**, 3765–3780.
- 91 B. Collin, M. Auffan, A. C. Johnson, I. Kaur, A. A. Keller, A. Lazareva, J. R. Lead, X. Ma, R. C. Merrifield and C. Svendsen, Environmental release, fate and ecotoxicological effects of manufactured ceria nanomaterials, *Environ. Sci.: Nano*, 2014, **1**(6), 533–548.
- 92 N. J. Rogers, N. M. Franklin, S. C. Apte, G. E. Batley, B. Angel, J. R. Lead and M. Baalousha, Physico-chemical behaviour and algal toxicity of nanoparticulate CeO₂ in freshwater, *Environ. Chem.*, 2010, **7**, 50–60.
- 93 T. Xia, M. Kovichich, M. Liong, L. Madler, B. Gilbert, H. Shi, J. I. Yeh, J. I. Zink and A. E. Nel, Comparison of the mechanism of toxicity of zinc oxide and cerium oxide nanoparticles based on dissolution and oxidative stress properties, *ACS Nano*, 2008, **2**(10), 2121–2134.
- 94 D. Wu, S. Yang, W. Du, Y. Yin, J. Zhang and H. Guo, Effects of titanium dioxide nanoparticles on *Microcystis aeruginosa* and microcystins production and release, *J. Hazard. Mater.*, 2019, **377**, 1–7.
- 95 A. Castro-Bugallo, A. Gonzalez-Fernandez, C. Guisande and A. Barreiro, Comparative responses to metal oxide nanoparticles in marine phytoplankton, *Arch. Environ. Contam. Toxicol.*, 2014, **67**(4), 483–493.
- 96 A. J. Miao, X. Y. Zhang, Z. Luo, C. S. Chen, W. C. Chin, P. H. Santschi and A. Quigg, Zinc oxide-engineered nanoparticles: dissolution and toxicity to marine phytoplankton, *Environ. Toxicol. Chem.*, 2010, **29**(12), 2814–2822.
- 97 T. P. Dasari, K. Pathakoti and H. M. Hwang, Determination of the mechanism of photoinduced toxicity of selected metal oxide nanoparticles (ZnO, CuO, Co₃O₄ and TiO₂) to *E. coli* bacteria, *J. Environ. Sci.*, 2013, **25**(5), 882–888.
- 98 J. Hou, T. Li, L. Miao, G. You, Y. Xu and S. Liu, Effects of titanium dioxide nanoparticles on algal and bacterial communities in periphytic biofilms, *Environ. Pollut.*, 2019, **251**, 407–414.
- 99 J. S. Kuwabara, J. A. Davis and C. C. Y. Chang, Algal growth response to particle-bound orthophosphate and zinc, *Limnol. Oceanogr.*, 1986, **31**, 503–511.
- 100 J. L. Yang, Y. F. Li, X. P. Guo, X. Liang, Y. F. Xu, D. W. Ding, W. Y. Bao and S. Dobretsov, The effect of carbon nanotubes and titanium dioxide incorporated in PDMS on biofilm community composition and subsequent mussel plantigrade settlement, *Biofouling*, 2016, **32**(7), 763–777.
- 101 C. T. Binh, T. Tong, J. F. Gaillard, K. A. Gray and J. J. Kelly, Acute effects of TiO₂ nanomaterials on the viability and taxonomic composition of aquatic bacterial communities assessed via high-throughput screening and next generation sequencing, *PLoS One*, 2014, **9**(8), e106280.
- 102 P. A. Holden, F. Klaessig, R. F. Turco, J. H. Priester, C. M. Rico, H. Avila-Arias, M. Mortimer, K. Pacpaco and J. L. Gardea-Torresdey, Evaluation of exposure concentrations used in assessing manufactured nanomaterial environmental hazards: are they relevant?, *Environ. Sci. Technol.*, 2014, **48**(18), 10541–10551.
- 103 W. Fan, M. Cui, H. Liu, C. Wang, Z. Shi, C. Tan and X. Yang, Nano-TiO₂ enhances the toxicity of copper in natural water to *Daphnia magna*, *Environ. Pollut.*, 2011, **159**(3), 729–734.
- 104 X. Liu, J. Wang, Y.-W. Huang and T. Kong, Algae (*Raphidocelis*) reduce combined toxicity of nano-TiO₂ and lead on *C. dubia*, *Sci. Total Environ.*, 2019, **686**, 246–253.

