



Cite this: *Phys. Chem. Chem. Phys.*, 2022, 24, 6856

A reduced electrophilicity for simple Lewis acids A involved in non-covalent interactions with Lewis bases B†

Ibon Alkorta ^a and Anthony Legon ^b

Dissociation energies D_e for $B \cdots A = B + A$ can be written $D_e = c' N_B E_A$, where N_B and E_A are the nucleophilicities and electrophilicities of the Lewis base B and the Lewis acid A, respectively. A reduced electrophilicity is defined $\bar{E}_A = E_A/\sigma_{\max}$, where σ_{\max} is the maximum electrostatic surface potential on iso-surface of A, the atom directly involved in the non-covalent interaction. This definition is tested for halogen-bonded complexes $B \cdots YX$, with Lewis bases $B = N_2, CO, C_2H_2, C_2H_4, H_2S, HCN, H_2O$ or NH_3 . D_e plotted against N_B for several series $B \cdots YX$ yields straight lines of gradient E_A . When D_e/σ_{\max} is the ordinate, the straight lines conflate to a single line, gradient $\bar{E}_{IX} = E_{IX}/\sigma_{\max}$. Hydrogen-bonded complexes $B \cdots HX$ ($X = F, Cl, Br, I$), coinage-metal complexes $B \cdots MX$ ($M = Cu, Ag, Au; X = F, Cl, Br, I$), and alkali-metal bonded complexes $B \cdots MX$ ($M = Li, Na; X = F, H, CH_3$) behave similarly. \bar{E}_A is an intrinsic property of the atom immediately involved in the non-covalent bond.

Received 15th February 2022,
 Accepted 27th February 2022

DOI: 10.1039/d2cp00779g

rsc.li/pccp

1. Introduction

The use of reduced quantities in chemistry/physics to identify or focus attention on relationships among properties of otherwise disparate substances/topics is not uncommon. For example, in connection with the well-known Law of Corresponding States, if the reduced densities ρ/ρ^c of liquids (where ρ^c is the density at the critical temperature T^c), are plotted against T/T^c , a single curve results onto which a number of liquid substances fit very well. Reduced potential energy functions that apply to a range of related diatomic molecules are a second example and provide useful generalisations that provoke questions about what is common to molecules that can be so described. Quantities such as reduced mass and reduced energies feature as simplifying and unifying tools in the theory of rotational spectroscopy. In this article, we shall show that, for a wide range of non-covalent interactions $B \cdots A$ of Lewis bases B with Lewis acids A (namely, interactions through hydrogen, halogen, coinage-metal and alkali-metal bonds), we can define a reduced electrophilicity of A. We shall show that the reduced electrophilicity is an interesting general property of complexes $B \cdots YX$ in which the non-covalent interaction is one of several different

types. It is independent of the atom X in the molecule YX involved in the non-covalent interaction. Thus, the reduced electrophilicity is a property that describes the propensity of Y to form the appropriate non-covalent bond, independently of X.

The terms nucleophile and electrophile were introduced by Ingold¹ in 1933. In his book “Structure and Mechanism in Organic Chemistry”,² Ingold gave the following definitions: “Reagents which act by donating their electrons to, or sharing them with, a foreign atomic nucleus will be called nucleophilic reagents or sometimes nucleophiles” and “Reagents which act by acquiring electrons, or a share in electrons, which previously belonged exclusively to a foreign molecule will be called electrophilic reagents, or sometimes electrophiles.” Since 1933, there has been an extensive literature on the subject of nucleophilicity and electrophilicity. For example, using both words ‘nucleophilicity and electrophilicity’ together as a search topic in the Web Of Science returns 694 publications with an average citation of 30 per article. The great majority of the work is concerned with scales of nucleophilicity and electrophilicity based on rates of reaction, mainly involving organic molecules, as reviewed, for example, by Mayr and Patz³ in an article entitled: ‘Scales of Nucleophilicity and Electrophilicity: A System for Ordering Polar Organic and Organometallic Reactions.’ The definitions employed here, on the other hand, are based on the energy D_e of the (usually weak) non-covalent interaction of a pair of (initially isolated) molecules (a Lewis base B and a Lewis acid A) when they come together to form the isolated complex $B \cdots A$ without chemical reaction. Although these definitions are also consistent with those of Ingold quoted earlier, the subjects

^a Instituto de Química Médica (IQM-CSIC), Juan de la Cierva, 3, E-28006 Madrid, Spain. E-mail: ibon@iqm.csic.es; Tel: +34 915622900

^b School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK. E-mail: a.c.legon@bristol.ac.uk; Tel: +44 (0)117 331 7708

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d2cp00779g



of the present article might be more correctly called, gas-phase nucleophilicities and electrophilicities of molecules involved in non-covalent interactions.

Some years ago,⁴ a method was proposed for reproducing the intermolecular stretching force constants k_σ of simple hydrogen-bonded complexes $B \cdots HX$ by assigning dimensionless nucleophilicities N_B and electrophilicities E_{HX} to the Lewis base B and to the Lewis acids HX , respectively, according to eqn (1):

$$k_\sigma = cN_B E_{HX}, \quad (1)$$

in which c is a constant. The series for which $X = F, Cl, Br$ and CN and $B = N_2, OC, PH_3, H_2S, HCN, CH_3CN, H_2O$ and NH_3 was the main focus initially⁴ but this group was soon extended to include the π - and pseudo- π -Lewis bases acetylene, ethylene and cyclopropane.⁵ Later, it was shown that the approach could also be applied to several series of halogen-bonded complexes $B \cdots XY$, where XY is one of the Lewis acids $F_2, ClF, Cl_2, BrCl, Br_2$ ^{6,7} and ICl .⁸ In all these cases, the k_σ were experimental, zero-point state values, as determined from the centrifugal distortion constants⁹ available from analyses of the rotational spectra of such complexes. A detailed analysis^{10,11} of *ab initio* calculated, equilibrium dissociation energies D_e showed that experimental k_σ values are directly proportional to D_e through the simple equation:

$$D_e = C_\sigma k_\sigma, \quad (2)$$

and that the constant $C_\sigma = 1.50(3) \times 10^3 \text{ m}^2 \text{ mol}^{-1}$ was appropriate to the whole series $B \cdots HX$ ($B = N_2, CO, C_2H_2, C_2H_4, H_2S, H_3P, HCN, H_2O$ or NH_3 and $X = F, Cl, Br$ or I) and $B \cdots XY$ (with XY as listed earlier). This relationship was shown to hold also when both D_e and k_σ were equilibrium values, as calculated *ab initio* at the CCSD(T)CBS level (CBS = complete basis set extrapolation) for the restricted series $B \cdots HF, B \cdots HCl, B \cdots F_2, B \cdots Cl_2$ and $B \cdots ClF$ but involving the same set of Lewis bases B .¹² Plots of D_e versus k_σ yielded good straight lines having gradients in agreement with those established in ref. 10 and 11 for these series. In a further, more wide-ranging analysis,¹³ the *ab initio* calculated values of D_e for 250 complexes were considered. The complexes were formed from 11 Lewis bases B and 23 Lewis acids A , chosen to include several types of non-covalent interaction (hydrogen, halogen, tetrel, pnictogen and chalcogen bonds). The D_e values were fitted by a least-squares method to eqn (3) (with c' set as 1.00 kJ mol^{-1} for convenience) to yield N_B and E_A values.

$$D_e = c' N_B E_A \quad (3)$$

There was evidence that each of the five types of non-covalent interaction was fitted equally well and therefore it was concluded that eqn (3) is valid for all those interaction types. Thus the determined values of N_B and E_A were established as properties of the individual molecules.

Further investigations of complexes $B \cdots MX$ ($M = Cu, Ag$ or $Au, X = F$ or Cl) involving the coinage-metal bond¹⁴ and of alkali-metal bonded complexes $B \cdots LiX$ and $B \cdots NaX$ ($X = F, H$ or CH_3)¹⁵ have revealed that the *ab initio* calculated values of D_e

for the complexes involving these two types of non-covalent interaction can also be represented by eqn (3). However, when eqn (3) was fitted to D_e values by using the least-squares method described in ref. 13 although the fits were good, somewhat different N_B values from those obtained in ref. 13 resulted, perhaps not surprisingly in view of the much more polar character of the LiX, NaX and CuX, AgX and AuX diatomic molecules than of HX, XY , etc. For the $B \cdots MX$ coinage metal series¹⁴ it was discovered that graphs of D_e versus N_B were good straight lines through the origin and led to the following order of electrophilicities $E_{AuF} > E_{AuCl} > E_{CuF} > E_{CuCl} > E_{AgF} \approx E_{AgCl}$. It was also noted that, to good approximation, plots of D_e/σ_{\max} versus N_B , where σ_{\max} is the maximum positive value of the molecular electrostatic surface potential (MESP) on the $0.001 \text{ e bohr}^{-3}$ electron density iso-surface, fall on the same straight line for both MF and MCl for a given M. This suggests that a reduced electrophilicity might be defined as (E_{MX}/σ_{\max}) .

In the present article, we examine whether this definition of reduced electrophilicity is generally valid. To do so we present *ab initio* calculations of the equilibrium dissociation energy D_e accompanying the process $B \cdots IX = B + IX$ for the series of halogen-bonded complexes $B \cdots IX$ in which the Lewis base B is $N_2, CO, C_2H_2, C_2H_4, H_2S, HCN, H_2O$ or NH_3 and IX is variously IF, ICl, IBr or I_2 . The corresponding series $B \cdots BrX$ ($X = Br, Cl$ or F) and $B \cdots ClX$ ($X = Cl$ or F) are also investigated. The calculations of D_e were conducted at the CCSD(T)(F12c)/cc-pVDZ-F12 level of theory. A least-squares fit of the D_e values to eqn (3) was carried out using the set of known N_B values¹³ for the Lewis bases to determine the E_{YX} ($Y = I, Br, Cl$) for the Lewis acids involved. Not only can eqn (3) be tested for this extended series, but also we shall investigate whether the graph of D_e/σ_{\max} versus N_B is a single straight line for the full set of iodine-bonded complexes $B \cdots IX$ ($X = F, Cl, Br$ and I), and also for several other series. Put another way, we ask the question: Does E_{IX}/σ_{\max} provide a reduced electrophilicity for the set of Lewis acids IF, ICl, IBr and I_2 , *i.e.* one independent of the atom attached to I ? Some previously published D_e values for the series of hydrogen-bonded complexes $B \cdots HX$ ($X = F, Cl, Br$ and I)^{10,11} and the series of alkali-metal-bonded complexes $B \cdots NaX$ and $B \cdots LiX$ ($X = F, H$ and CH_3)¹⁵ will be treated in the same way to test whether the concept of reduced electrophilicity is generally applicable.

2. Theoretical methods

The geometry optimizations of the isolated monomers and complexes were carried out using the CCSD(T)(F12c) explicitly correlated, coupled-cluster method.^{16,17} The cc-pVDZ-F12 basis sets, which are optimised for use with the CCSD(T)-F12 method,¹⁸ were used for all of the first and second row atoms. Pseudo-potential (PP) basis sets were employed for the atoms Br and I in combination with the ECP-10-MDF¹⁹ and ECP-28-MDF²⁰ effective core potentials to account for scalar relativistic effects. The frozen-core approximation was used in all the calculations, which employed the MOLPRO package.^{21,22}



Dissociation energies, so computed at the CCSD(T)(F12c)/cc-pVDZ-F12 level, were corrected for basis set superposition error (BSSE)²³ by means of the counterpoise keyword in MOLPRO. D_e values for 132 new complexes and 12 monomers were calculated specifically for this article. The remaining D_e values employed here are from earlier publications: the data for coinage-metals complexes are from ref. 14, those for the alkali-metal complexes are from ref. 15 and some of the $B \cdots HX$ ($X = F, Cl, Br, I$) and $B \cdots XY$ (X and Y are halogen atoms) are from ref. 10 and 11.

Molecular electrostatic surface potentials (MESP) were calculated by using the MP2/aug-cc-pVTZ wavefunction in the GAUSSIAN program²⁴ and represented at the 0.001 e bohr⁻³ electron density iso-surface with the Jmol program²⁵ (an open-source Java viewer for chemical structures in 3D, available at <http://www.jmol.org/>). The extreme values (maxima and minima) on the iso-surface were calculated with the multiwfn program.²⁶

The results for 132 new complexes and 12 monomers are presented in this article for the first time and are recorded in the ESI.† Those for complexes involving the Li and Na non-covalent bonds were published previously in ref. 15) and those involving Cu, Ag and Au with $X = F$ and Cl were first published in ref. 14 but results for CuBr and CuI are new. All values of D_e for the coinage-metal complexes were calculated at the CCSD(T)/aug-cc-pVTZ, aug-cc-pVTZ-PP levels as appropriate and the binding energy D_e was obtained by using the CBS extrapolation method.^{27,28} In excess of 300 complexes are discussed.

3. Results

3.1 The iodine-bonded series $B \cdots IX$ ($X = F, Cl, Br, I$)

Fig. 1 shows a graph in which the equilibrium dissociation energies D_e of the iodine-bonded complexes $B \cdots IX$ ($X = F, Cl, Br$ and I) are plotted against the nucleophilicity N_B of the series of Lewis bases $B = N_2, CO, C_2H_2, C_2H_4, H_2S, HCN, H_2O$ or NH_3 , the latter quantities as determined in ref. 13. D_e is the energy required to move the components of $B \cdots IX$ from their equilibrium distance in the complex to become infinitely separated molecules B and IX relaxed to their hypothetical equilibrium states. The Lewis base PH_3 has been omitted from the list because it was shown previously that complexes such as $H_3P \cdots ICl$ ²⁹ and $H_3P \cdots ClF$ ³⁰ are of a different type from the remainder and have significant electric charge redistribution on complex formation. The linear-regression fits to the points (with the origin taken as a point under the assumption that the nucleophilicity $N_B = 0$ when $D_e = 0$) are shown as solid straight lines. The R^2 values shown in Fig. 1 indicate that the fits are satisfactory, with that for $B \cdots IF$ the worst at 0.95. Clearly, eqn (3) is obeyed by each of the $B \cdots IX$ series. The gradient of each straight line corresponds to the electrophilicity of the Lewis acid according to eqn (3) given that $c' = 1.00 \text{ kJ mol}^{-1}$ was used in the determination of the N_B values by the least squares fit to $250D_e$ values in ref. 13. The conclusion is therefore that




Fig. 1 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the four series of iodine bonded complexes $B \cdots IX$ ($X = F, Cl, Br$ and I). The continuous straight lines are linear regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit, as measured by R^2 .

$E_{IF} = 8.6(7) > E_{ICl} = 6.1(4) > E_{IBr} = 5.3(3) > E_{I_2} = 3.9(2)$, which order is consistent with electron withdrawing power of the X atom in the order $F > Cl > Br > I$.

The concept of reduced electrophilicity discussed in the Introduction can now be tested for the $B \cdots IX$ series. Dividing eqn (3) by the value of σ_{\max} (the maximum positive value of the MESP on the 0.001 e bohr⁻³ electron density iso-surface near to the I atom along the axis of the IX diatomic molecule), we have

$$D_e/\sigma_{\max} = c' N_B (E_A/\sigma_{\max}) = N_B \Xi_A, \quad (4)$$

in which $\Xi_A = c' E_A/\sigma_{\max}$ defines the reduced electrophilicity of the Lewis acid A and is dimensionless given that $c' = 1.0 \text{ kJ mol}^{-1}$, σ_{\max} is in kJ mol^{-1} , and E_A is dimensionless. Thus, a plot of D_e/σ_{\max} versus N_B should be a straight line through the origin and have gradient Ξ_A . Values of σ_{\max} calculated at the MP2/aug-cc-pVTZ level are available from Table 1. Fig. 2 displays these plots for the series $B \cdots IX$, when $X = F, Cl, Br$ and I .

Table 1 Maximum molecular electrostatic potentials $\sigma_{\max}/(\text{kJ mol}^{-1})$ on the 0.001 e bohr⁻³ electron-density iso-surface of various Lewis acids YX (on the axis at the Y end of the molecule)^a

Lewis acid YX	X = I	X = Br	X = Cl	X = F	X = CN	X = CP	X = CCH	X = H	X = CH ₃
IX	125.7	158.4	180.1	241.3	201.3	—	142.0	—	—
BrX	—	118.8	141.3	205.6	—	—	—	—	—
ClX	—	—	107.0	172.1	—	—	—	—	—
HX	119.4	160.1	190.2	287.9	216.9	126.3	134.8	—	—
CuX	342.8	366.1	379.8	405.1	—	—	—	—	—
AgX	296.0	305.1	308.0	294.5	—	—	—	—	—
AuX	270.1	314.6	346.7	404.8	—	—	—	—	—
LiX	—	—	—	764.2	—	—	—	809.5	793.1
NaX	—	—	—	566.8	—	—	—	510.0	471.0

^a Calculated at the MP2/aug-cc-pVTZ level of theory.






Fig. 2 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the four series of iodine-bonded complexes $B \cdots IX$ ($X = F, Cl, Br$ and I). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line and the quality of the fit as measured by R^2 .

Fig. 2 confirms that there is indeed a reduced electrophilicity Ξ_A having the value 0.034(2) for the iodine atom in the four diatomic molecules IF, ICl, IBr and I_2 . The scatter and R^2 are rather larger for IF, but the gradients of the four straight lines are identical within their quoted errors. In fact, some points are obscured by superposition of others. It is of interest to examine, in the same way, the situation when the Lewis acids are the pseudo-halogens ICN and ICCH. The graph of D_e versus N_B is shown in Fig. 3 while that when the ordinate is D_e/σ_{\max} is in




Fig. 3 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the two series of iodine-bonded complexes $B \cdots IX$ ($X = CN$ and CCH). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .




Fig. 4 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the two series of iodine-bonded complexes $B \cdots IX$ ($X = CN$ and CCH). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line and the quality of the fit as measured by R^2 .

Fig. 4. The latter diagram shows that the gradients in Fig. 4 are equal within the fitting error and hence both ICN and ICCH can be described by a reduced electrophilicity $\Xi = 0.0229(4)$. We note, however, that this value is different from that for the IX ($X = F, Cl, Br, I$) given in Fig. 2.

3.2 The bromine-bonded series $B \cdots BrX$ ($X = F, Cl, Br$)

Fig. 5 shows the graphs of D_e versus N_B for the three series $B \cdots BrX$ ($X = F, Cl, Br$). The same pattern as observed for the $B \cdots IX$ series is evident, with the largest scatter from its




Fig. 5 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the three series of bromine-bonded complexes $B \cdots BrX$ ($X = F, Cl$ and Br). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .



regression line again when $X = F$. Values of D_e are not given in ref. 10 and 11 for $N_2 \cdots Br_2$ and $HCN \cdots Br_2$ and these were calculated for the present work. Evidently, eqn (3) is obeyed by each series, as indicated by R^2 values close to unity. The electrophilicities of the BrX are in the order $X = F > Cl \geq Br$.

The corresponding graphs of D_e/σ_{max} versus N_B for the series $B \cdots BrF$, $B \cdots BrCl$ and $B \cdots Br_2$ are set out in Fig. 6. Note again that the gradients, which yield the reduced electrophilicity \mathcal{E}_A for the molecules BrX , are almost equal within the errors of the fits and suggest that, as for the $B \cdots IX$ series, the definition of reduced electrophilicity used here is reasonable.

3.3 The chlorine-bonded series $B \cdots ClX$ ($X = F, Cl$)

There are only two members of this series. Fig. 7 shows the graph of D_e versus N_B for both the $B \cdots ClF$ and $B \cdots Cl_2$ series. Eqn (3) appears to be obeyed by both. The D_e/σ_{max} versus N_B plots shown in Fig. 8 have gradients = 0.030(2) and 0.022(1) for the $B \cdots ClF$ and $B \cdots Cl_2$ series, respectively. Hence, we cannot unambiguously assign a reduced electrophilicity to the ClF and Cl_2 molecules for their interactions with the Lewis bases B , as was the case for the $B \cdots BrX$ and $B \cdots IX$ series. Nevertheless these values are much closer together than the gradients of 5.2(4) and 2.3(1) kJ mol^{-1} of the fitted points in the graphs of D_e versus N_B for the $B \cdots ClF$ and $B \cdots Cl_2$ series, respectively.

3.4 Is there a reduced electrophilicity \mathcal{E}_A appropriate to the hydrogen halides HX ?

The evidence presented in Sections 3.1–3.3 indicates that the quantity \mathcal{E}_A defined by means of eqn (3) and (4) might be used as a reduced electrophilicity for the Lewis acids A involved in series of halogen-bonded complexes. A reduced electrophilicity so defined is a measure of the propensity of, *e.g.* IX to form $B \cdots IX$ halogen bonds independently of the halogen atom X




Fig. 6 D_e/σ_{max} plotted against the nucleophilicity N_B of the Lewis base B for the three series of bromine-bonded complexes $B \cdots BrX$ ($X = F, Cl$ and Br). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line and the quality of the fit as measured by R^2 .




Fig. 7 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the two series of chlorine-bonded complexes $B \cdots ClX$ ($X = F$ and Cl). The continuous straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .




Fig. 8 D_e/σ_{max} plotted against the nucleophilicity N_B of the Lewis base B for the two series of chlorine-bonded complexes $B \cdots ClX$ ($X = F$ and Cl). The straight lines are regression fits to the points (solid dots) for each series, (origin as a point). The inset gives the gradient of each fitted line and the quality of the fit, as measured by R^2 .

attached to it. It is of interest to investigate whether this concept has generality. In this section, hydrogen-bonded complexes $B \cdots HX$ having $X = F, Cl, Br$ and I will be discussed first and then $X = CN, CP$ and CCH will be considered.

Dissociation energies D_e for the series $B \cdots HX$ (now including $B = PH_3$), calculated at the explicitly-correlated level CCSD(T)(F12c)/cc-pVDZ-F12 and corrected for basis set






Fig. 9 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the four series of hydrogen-bonded complexes $B \cdots HX$ ($X = F, Cl, Br$ and I). The straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .

superposition error (BSSE), have been published elsewhere.^{10,11} Using again N_B values determined in ref. 13, Fig. 9 was constructed. It shows clearly that the $B \cdots HX$ series obey eqn (3) with well-determined gradients and R^2 values, the latter close to 1. The gradients provide measures of the electrophilicities E_{HX} of the hydrogen halides, which have the values 7.0(4), 4.5(2), 3.9(2) and 2.8(1) for $X = F, Cl, Br$ and I , respectively, an order consistent with chemical intuition (the greater the slope the stronger the hydrogen bonds). Fig. 10 shows the




Fig. 10 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the four series of hydrogen-bonded complexes $B \cdots HX$ ($X = F, Cl, Br$ and I). The straight lines are regression fits to the points (solid dots) for each series, (origin as a point). The inset gives the gradient of each fit and the quality of the fit as measured by R^2 .

corresponding set of graphs in which D_e/σ_{\max} is now the ordinate. The gradients determined by linear regression fits are closely bunched and have the mean value of 0.0239(7), which is, according to eqn (4), the reduced electrophilicity \mathcal{E}_{HX} of the hydrogen halides. The narrow spread of values indicates that the reduced electrophilicity of the hydrogen halides appears to be a well-defined quantity.

Given the results for the hydrogen halides, it is of interest to consider hydrogen acids of the type $H CX$ in which H is now directly attached to a carbon atom. D_e values for $B \cdots HCN$, $B \cdots HCP$ and $B \cdots HCCH$ complexed with the same set of Lewis bases B were calculated at the CCSD(T)(F12c)/cc-pVTZ-F12 level of theory and corrected for BSSE. The graphs of D_e plotted against N_B for these three series are in Fig. 11. Note that the points for $B \cdots HCCH$ and $B \cdots HCP$ are close to coincident. Convergence in the geometry optimisation for $H_2S \cdots HCN$ and $H_2S \cdots HCCH$ was not achieved and so points for these are missing.

Although the scatter of the points is rather large for the HCN series, eqn (3) is obeyed by all three Lewis acids HCN, HCP and HCCH. Moreover, as can be seen in Fig. 12, the plots of D_e/σ_{\max} against N_B for all three series lead to the same gradient within the fitting error. Thus, the reduced electrophilicity appears to be a property that can be assigned to these three Lewis acids also. We note, however, that $\mathcal{E}_{HCX} = E_{HCX}/\sigma_{\max}$ is different from the value obtained for the hydrogen halides.

3.5 Reduced electrophilicity and the coinage-metal halides MX ($M = Cu, Ag, Au$; $X = F, Cl, Br, I$)

Complexes of the type $B \cdots MX$, where $M = Cu, Ag, Au$, $X = F, Cl$ were discussed in ref. 14, where the possibility of a reduced electrophilicity was first raised. The Lewis bases B are again the




Fig. 11 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for three series of hydrogen-bonded complexes $B \cdots HX$ ($X = CN, CP$ and CCH). The straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .






Fig. 12 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the three series of hydrogen-bonded complexes $B \cdots HX$ ($X = \text{CN}$, CP , and CCH). The straight lines are regression fits to the points (solid dots) for each series, (origin as a point). The inset gives the gradient of each fit and the quality of the fit as measured by R^2 .

nine simple molecules $B = \text{N}_2$, CO , C_2H_2 , C_2H_4 , H_2S , PH_3 , HCN , H_2O , and NH_3 . For the present article, calculations have been extended to include $X = \text{Br}$ and I . The graphs of D_e versus N_B for the series $B \cdots \text{AuX}$ ($X = \text{F}$, Cl , Br and I) are displayed in Fig. 13, while the plots with D_e/σ_{\max} as the ordinate are in Fig. 14. The corresponding pairs of diagrams for $M = \text{Cu}$ are available as Fig. S1 and S2 of the ESI[†] while for $M = \text{Ag}$ Fig. S3 and S4 (ESI[†]) are appropriate. The magnitude and order of the D_e values for the $B \cdots \text{MX}$ complexes (and therefore the values of the N_B of the Lewis bases when involved in such complexes, as published in




Fig. 13 Dissociation energy D_e plotted against the nucleophilicity of the Lewis base B for the four series of coinage-metal-bonded complexes $B \cdots \text{AuX}$ ($X = \text{F}$, Cl , Br , I). The straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .




Fig. 14 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the four series of coinage-metal-bonded complexes $B \cdots \text{AuX}$ ($X = \text{F}$, Cl , Br , I). The straight lines are regression fits to the points (solid dots) for each series, (origin as a point). The inset gives the gradient of each fit and the quality of the fit as measured by R^2 .

ref. 14) are both quite different from those used¹³ for the hydrogen- or halogen-bonded series. Nevertheless, it is clear from Fig. 13 and Fig. S1, S3 (ESI[†]) that eqn (3) is valid for coinage-metal bonds as well as hydrogen- and halogen bonds. Moreover, the order of the electrophilicities of the aurous halides is $E_{\text{AuF}} > E_{\text{AuCl}} > E_{\text{AuBr}} > E_{\text{AuI}}$. The order for the argentous and cuprous halides is the same.

Fig. 14 illustrates how division of the D_e value by σ_{\max} reduces the four straight lines in Fig. 13 to a single line. Thus, the gradients of the four lines in Fig. 14 are identical within the fitting error, with the mean value 0.0442(18). This is the dimensionless reduced electrophilicity $\bar{\epsilon}_{\text{AuX}}$ of the AuX diatomic molecules. Fig. S2 and S4 (ESI[†]) indicate that $\bar{\epsilon}_{\text{CuX}} = 0.0283(18)$ and $\bar{\epsilon}_{\text{AgX}} = 0.0287(22)$.

The coinage-metal diatomic molecules are much more polar than the Lewis acids considered in earlier sections. For example, the electric dipole moments of CuCl and AgCl are 5.74 D³¹ and 6.076(6) D,³² respectively, while those of BrCl , ICl and HBr are 0.519(4) D,³³ 1.24(2) D³⁴ and 0.8280(6) D,³⁵ respectively. CuCl and AgCl (and presumably AuCl) also have large ionicities i_c . The ionicity of a diatomic molecule is a measure of the fractional contribution of the structure M^+Cl^- to the valence-bond description of the molecule. It can be calculated³⁶ from the nuclear quadrupole coupling constant $\chi_z(\text{X})$ of the halogen atom X in the appropriate diatomic molecule by using eqn (5), in which $eQq_{n,1,0}$ is the quadrupole coupling constant arising from a single electron in a np_z orbital.

$$i_c = 1 + \{\chi_z(\text{X})/(eQq_{n,1,0})\} \quad (5)$$

The i_c values for CuCl and AgCl are 0.71 and 0.67,³⁷ while those for BrCl , ICl and HBr are 0.06, 0.25 and 0.30.³⁶ Despite their greater ionicity, the concepts of electrophilicity and reduced electrophilicity introduced by eqn (3) and (4), respectively,






Fig. 15 Dissociation energy D_e plotted against the nucleophilicity N_B of the Lewis base B for the six series of alkali-metal-bonded complexes $B \cdots LiX$ and $B \cdots NaX$ ($X = F, H, CH_3$). The straight lines are regression fits to the points (solid dots) for each series, with the origin taken as a point. The inset gives the gradient of each fitted line in kJ mol^{-1} and the quality of the fit as measured by R^2 .

appear to apply at least as well to the coinage-metal complexes $B \cdots MX$ as they do to the hydrogen-bonded $B \cdots HX$ and halogen-bonded $B \cdots YX$ discussed here.

3.6 Reduced electrophilicity of the alkali-metal compounds LiX and NaX ($X = F, H, CH_3$)

Application of eqn (5) to alkali-metal halides such as $NaCl$ and $LiCl$ shows values of i_c that are close to 1, indicating that these molecules are essentially ion pairs in the gas phase.³⁶ The question that then arises is: Does the concept of reduced




Fig. 16 D_e/σ_{\max} plotted against the nucleophilicity N_B of the Lewis base B for the six series of alkali-metal-bonded complexes $B \cdots LiX$ and $B \cdots NaX$ ($X = H, F, CH_3$). The six straight lines are regression fits to the points (solid dots) for each series, (origin as a point). The inset gives the gradient of each fit and the quality of the fit as measured by R^2 .

Table 2 Values of the reduced electrophilicity \mathcal{E}_A determined for various Lewis acids A^a

Lewis acid A	Reduced electrophilicity \mathcal{E}_A
IX ($X = F, Cl, Br, I$)	0.033(2)
IX ($X = CN, CCH$)	0.0229(4)
BrX ($X = F, Cl, Br$)	0.033(3)
ClX ($X = F, Cl$)	$\sim 0.026(4)$
HX ($X = F, Cl, Br, I$)	0.0239(7)
HX ($X = CN, CP, CCH$)	0.0164(13)
CuX ($X = F, Cl, Br, I$)	0.0283(18)
AgX ($X = F, Cl, Br, I$)	0.0287(22)
AuX ($X = F, Cl, Br, I$)	0.0442(18)
LiX ($X = H, F, CH_3$)	0.01114(2)
NaX ($X = H, F, CH_3$)	0.01007(10)

^a The reduced electrophilicity of a given Lewis acid A is defined as E_A/σ_{\max} and is the gradient of the straight line obtained by plotting D_e/σ_{\max} against N_B for a series of complexes $B \cdots A$, where the N_B and E_A are the nucleophilicity and electrophilicity of the Lewis bases B and the Lewis acid A, respectively. σ_{\max} is the maximum positive value of the MESP of A on the $0.001 e \text{ bohr}^{-3}$ iso-surface along its symmetry axis and near to the atom that is directly involved in the non-covalent interaction of A with B.

electrophilicity apply as well to alkali-metal bonded complexes such as $B \cdots NaX$ and $B \cdots LiX$ as it does to hydrogen-bonded, halogen-bonded and coinage-metal-bonded complexes? D_e and N_B values for $B \cdots MF$, $B \cdots MH$, $B \cdots MCH_3$ were recently published¹⁵ for $M = Na$ and Li , with Lewis bases $B = CO$, HCN , H_2O , NH_3 , H_2S and PH_3 . A set of N_B , E_{LiX} and E_{NaX} were derived by the least-squares method from 34 values of D_e for complexes that could be identified as isostructural with their hydrogen- and halogen-bonded counterparts. As for the coinage-metal bond series, the N_B scale had some significant differences from that used for the hydrogen- and halogen-bonded series. The variation of D_e with N_B for the alkali-metal bonded complexes (first published in ref. 15) is set out again in Fig. 15 (redrawn). Each series leads to an excellent straight line, which indicates eqn (3) is appropriate here also. The electrophilicities have the order $E_{LiH} > E_{LiF} \approx E_{LiCH_3} > E_{NaF} > E_{NaH} > E_{NaCH_3}$, as previously established.¹⁵

The corresponding graphs of D_e/σ_{\max} versus N_B are set out in Fig. 16. The gradients of the three straight lines for the $B \cdots LiX$ complexes in Fig. 16 are very nearly equal. However, a separate linear regression fit of all the $B \cdots LiX$ points alone gives $\mathcal{E}_{LiX} = E_{LiX}/\sigma_{\max} = 0.01114(2)$ for the reduced electrophilicity of the LiX molecules, while for the $B \cdots NaX$ complexes alone the value is $0.01007(10)$. It appears that the reduced electrophilicities of the LiX and NaX molecules are very similar, but not equal.

4. Conclusions and discussion

The main conclusion of this work is that it is possible to define a reduced electrophilicity \mathcal{E}_A for a Lewis acid A when involved in non-covalent complexes $B \cdots A$ formed with Lewis bases B. This is found to be true for a range of non-covalent interaction types. A summary of the values of \mathcal{E}_A determined here is available in Table 2.



Graphs of the equilibrium dissociation energy D_e for complexes $B \cdots IX$ plotted against the nucleophilicity N_B of the Lewis bases $B = N_2, CO, C_2H_2, C_2H_4, H_2S, HCN, H_2O,$ and NH_3 for the four series $X = I, Br, Cl, F$ have been used to show that eqn (3), $D_e = c'N_B E_A$, is obeyed by each of these series of iodine-bonded complexes. It is thereby established that the order of the gas-phase non-covalent electrophilicities E_{IX} of these dihalogen molecules is $E_{IF} > E_{ICl} > E_{IBr} > E_{I_2}$. Moreover, when D_e is divided by σ_{max} , (*i.e.* the value of the molecular electrostatic potential on the $0.001 \text{ e bohr}^{-3}$ electron density iso-surface on the molecular axis near to I), the plot of D_e/σ_{max} versus N_B reveals that the four distinct straight lines of different gradient in the D_e versus N_B plots have become conflated to give a single straight line to a good level of approximation. The gradient of the single straight line is referred to as the reduced electrophilicity $\Xi_{IX} = D_e/\sigma_{max}$ of the IX molecules ($X = I, Br, Cl$ and F). The same procedure followed for the series $B \cdots BrX$ and (less certainly) the series $B \cdots ClX$ allows similar conclusions and leads to the order of the reduced electrophilicities $\Xi_{IX} = \Xi_{BrX} > \Xi_{ClX}$. The series $B \cdots ICX$ ($X = N, CH$) also behaved in the same manner as the $B \cdots IX$ series, but the conflation of the two straight lines when D_e/σ_{max} is plotted against N_B yields an Ξ_{ICX} value smaller than that of the $B \cdots IX$ series.

It has also been possible to show that a similar conflation of four straight lines to a single straight line results when D_e/σ_{max} is plotted against N_B for the hydrogen-bonded series $B \cdots HX$ ($X = F, Cl, Br, I$) and hence provides a reduced electrophilicity Ξ_{HX} for the hydrogen halides. When HX is replaced by $H CX$, where $X = CN, CP$ and CCH , a value of Ξ_{HCX} is again established but, as for the corresponding IX and ICX series, it is found that $\Xi_{HX} > \Xi_{HCX}$.

When the Lewis acid is more polar, as for AuX, AgX and CuX in the three series of coinage-metal bonded complexes $B \cdots MX$ ($M = Cu, Ag, Au; X = F, Cl, Br, I$), it is still possible to establish well-determined reduced electrophilicities for the Lewis acids in the order $\Xi_{AuX} > \Xi_{AgX} \approx \Xi_{CuX}$. Even when the Lewis acid is close to being an ion pair, as in LiX and NaX ($X = F, H, CH_3$), the approach is again successful and leads to well-determined reduced electrophilicities in the order $\Xi_{NaX} \approx \Xi_{LiX}$.

It is important to discuss the significance of the newly-defined quantity referred to as reduced electrophilicity. Eqn (3) in Section 3.1 has been established to hold for a wide range of non-covalent interactions involved in axially symmetric complexes $B \cdots YX$, where Y is the atom of the Lewis acid directly involved in that interaction. An important quantity in this context is the most positive electrostatic potential σ_{max} at the $0.001 \text{ e bohr}^{-3}$ iso-surface on the symmetry axis in the vicinity of the atom Y of the Lewis acid ($0.001 \text{ e bohr}^{-3}$ is the conventional choice of electron density for such surfaces). It was shown in Section 3.1 that division of eqn (3) by σ_{max} to give eqn (4) and then a plot of D_e/σ_{max} versus N_B leads in all cases considered here to a straight line through the origin. According to eqn (4), the gradient of this line is E_{YX}/σ_{max} , that is the electrophilicity of the molecule YX per unit maximum positive electrostatic potential on the symmetry axis at Y . We have shown (by consideration of four different types of

non-covalent interaction and a range of axially symmetric Lewis acids YX) that the quantity E_{YX}/σ_{max} is an intrinsic property of the atom Y involved in the non-covalent interaction with the nucleophilic region of the Lewis base B . Thus, it is the same for all diatomic Lewis acids when the same atom Y is that involved in the non-covalent interaction with B . This result is made clear, for example, by Fig. 10, which involves the four hydrogen halides. When YX is a triatomic (or higher) molecule, as in Fig. 12 (which deals with the axially symmetric polyatomic Lewis acids HCN, HCP and $HCCH$), E_{YX}/σ_{max} is again an intrinsic property of the atom $Y = H$ for the group, but has a slightly different value from that of the hydrogen halides (Fig. 10). Therefore it seems justified to give the quantity E_{YX}/σ_{max} the name reduced electrophilicity and the special symbol Ξ_{YX} . This new reduced quantity clearly points to systematic family relationships among groups of Lewis acids involving different types of non-covalent interaction. Further investigation might reveal further aspects the relationship.

Thus, we conclude that the reduced electrophilicity of a Lewis acid YX is an intrinsic property of the atom Y and its axial electrostatic surface potential directly involved in the non-covalent interaction. It appears to be independent of the nature of the remainder of YX , certainly so when YX is a diatomic molecule and nearly so for more complex YX . It therefore provides a measure of the propensity of atom Y to participate in a wide range of hydrogen bonds (if $Y = H$), halogen bonds (if Y is a halogen atom), coinage-metal bonds (if Y is a Group 11 atom), alkali-metal bonds (if Y is an alkali-metal atom).

The vast majority of data on nucleophilicities and electrophilicities involves quantities that are derived from the rate constants of chemical reaction in solution between more complex molecules than those discussed here (see ref. 3, for example). The electrophilicities and their reduced values presented in the Results section refer to the limiting, gas-phase interactions between pairs of molecules otherwise isolated and therefore comparison of the two scales is unlikely to be useful. Two papers have been published,^{38,39} however, that make a theoretical interpretation of the gas-phase N and E values set out initially in ref. 4.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

ACL thanks the University of Bristol for the award of a University Senior Research Fellowship. IA thanks the Ministerio de Ciencia e Innovación of Spain (PGC2018-094644-B-C22 and 5931125495-125495-4-21) and Comunidad de Madrid (P2018/EMT-4329 AIRTEC-CM) for financial support.

References

- 1 C. K. Ingold, *J. Chem. Soc.*, 1933, 1120–1127.



- 2 C. K. Ingold, *Structure and Mechanism in Organic Chemistry*, Cornell University Press, Ithaca, New York, 1953, pp. 200–201.
- 3 H. Mayr and M. Patz, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 938–957.
- 4 A. C. Legon and D. J. Millen, *J. Am. Chem. Soc.*, 1987, **109**, 356–358.
- 5 A. C. Legon and D. J. Millen, *J. Chem. Soc., Chem. Commun.*, 1987, 986–987.
- 6 A. C. Legon, *J. Chem. Soc., Chem. Commun.*, 1998, 2585–2586.
- 7 A. C. Legon, *Angew. Chem., Int. Ed.*, 1999, **38**, 2686–2714.
- 8 A. C. Legon, in *Halogen Bonding: Fundamentals and Applications*, ed. P. Metrangolo and G. Resnati Structure and Bonding, Springer-Verlag, Berlin, 2008, ch. 2, vol. 126, pp. 17–64.
- 9 D. J. Millen, *Can. J. Chem.*, 1985, **63**, 1477–1479.
- 10 A. C. Legon, *Phys. Chem. Chem. Phys.*, 2014, **16**, 12415–12421.
- 11 A. C. Legon, *Phys. Chem. Chem. Phys.*, 2014, **16**, 25199.
- 12 I. Alkorta and A. C. Legon, *New J. Chem.*, 2018, **42**, 10548–10554.
- 13 I. Alkorta and A. C. Legon, *Molecules*, 2017, **22**, 1786–1799.
- 14 I. Alkorta, N. R. Walker and A. C. Legon, *Inorganics*, 2021, **9**, 13.
- 15 I. Alkorta, J. G. Hill and A. C. Legon, *Phys. Chem. Chem. Phys.*, 2020, **22**, 16421–16430.
- 16 C. Hättig, D. P. Tew and A. Köhn, *J. Chem. Phys.*, 2010, **132**, 231102.
- 17 C. Hättig, W. Klopper, A. Köhn and D. P. Tew, *Chem. Rev.*, 2012, **112**, 4–74.
- 18 J. G. Hill and K. A. Peterson, *J. Chem. Phys.*, 2014, **141**, 094106. See <http://bit.ly/ccBasis> for a repository of basis functions, including cc-pVDZ-F12 for Br and I, constructed by J. G. Hill.
- 19 M. Doig, U. Wedig, H. Stoll and H. Preuss, *J. Chem. Phys.*, 1987, **86**, 866–872.
- 20 I. S. Lin, H. Stoll and P. Schwertfeger, *J. Chem. Phys.*, 2006, **124**, 034107.
- 21 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schütz, 'Molpro: a general-purpose quantum chemistry program package', *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 242–253.
- 22 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schütz, *MOLPRO 2019.2, a package of ab initio programs*, 2019.
- 23 S. F. Boys and F. Bernardi, *Mol. Phys.*, 1970, **19**, 553–566.
- 24 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson and H. Nakatsuji, *et al.*, *Gaussian 16; Revision, A.03*, Gaussian, Inc., Wallingford, CT, USA, 2016.
- 25 Jmol: An Open-Source Java Viewer for Chemical Structures in 3D. Available online at: <http://www.jmol.org/>.
- 26 T. Lu and F. Chen, *Multiwfn: A multifunctional wavefunction analyzer*, *J. Comput. Chem.*, 2012, **33**, 580–592.
- 27 D. Feller, *J. Chem. Phys.*, 1993, **98**, 7059–7071.
- 28 A. Halkier, T. Helgaker, P. Jorgensen, W. Klopper and J. Olsen, *Chem. Phys. Lett.*, 1999, **302**, 437–446.
- 29 A. C. Legon, *Chem. Phys. Lett.*, 1999, **314**, 472–480.
- 30 R. A. Shaw, J. G. Hill and A. C. Legon, *J. Phys. Chem. A*, 2016, **120**, 8461–8468.
- 31 S. Hau and P. F. Bernath, *J. Phys. Chem. A*, 2015, **119**, 1435–1438.
- 32 K. P. R. Nair and J. Hoefft, *J. Phys. B: At. Mol. Phys.*, 1984, **17**, 735–738.
- 33 K. P. R. Nair, J. Hoefft and E. Tiemann, *Chem. Phys. Lett.*, 1978, **58**, 153–156.
- 34 E. Herbst and W. Steinmetz, *J. Chem. Phys.*, 1978, **56**, 5342–5347.
- 35 F. A. van Dijk and A. Dymanus, *Chem. Phys. Lett.*, 1970, **5**, 387–389.
- 36 W. Gordy and R. L. Cook, *Microwave Molecular Spectra*, John Wiley and sons, New York, 2nd edn, 1984, ch. 14, p. 752.
- 37 A. C. Legon and N. R. Walker, *Phys. Chem. Chem. Phys.*, 2018, **20**, 19332–19338.
- 38 A. Ebrahimi, H. Roohi, M. Habibi and M. Hasannejad, *Chem. Phys. Lett.*, 2006, **327**, 368–372.
- 39 A. Cedillo, R. Contreras, M. Galván, A. Aizman, J. Andrés and V. S. Safont, *J. Phys. Chem. A*, 2007, **111**, 2442–2447.

