

Showcasing research from Professor Martin Beyer's laboratory, Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Austria.

X-ray absorption spectroscopy of FeH^+ to aid its identification in astrochemical environments

The iron hydride diatomic molecular ion FeH^+ is expected to be present in the interstellar medium. Its X-ray absorption spectrum measured at the BESSY II synchrotron facility is the first laboratory spectrum that can be compared to astronomical observations.

As featured in:



See Norbert Przybilla, Martin K. Beyer *et al.*, *Chem. Commun.*, 2025, **61**, 2048.


 Cite this: *Chem. Commun.*, 2025, 61, 2048

 Received 11th September 2024,
 Accepted 16th December 2024

DOI: 10.1039/d4cc04702h

rsc.li/chemcomm

X-ray absorption spectroscopy of FeH⁺ to aid its identification in astrochemical environments†

 Shan Jin,^{id a} Max Flach,^{id bc} Alexander Ebenbichler,^{id d} Ethan M. Cunningham,^{id a}
 Vicente Zamudio-Bayer,^{id b} Konstantin Hirsch,^{id b} Christian van der Linde,^{id a}
 Norbert Przybilla,^{id *d} Milan Ončák,^{id a} J. Tobias Lau^{id bc} and Martin K. Beyer^{id *a}

We present the first absorption spectrum of the unperturbed diatomic molecular ion FeH⁺ in any wavelength range. The cryogenic X-ray absorption spectrum at the L₂ and L₃ edge is consistent with an iron 3d occupation of 6.24e. Comparison with the interstellar absorption spectrum of Cygnus X-1 indicates that FeH⁺ cannot be ruled out as a component of the absorbing medium.

With hydrogen as the most abundant element and iron the second most abundant metal in the Milky Way,¹ the diatomic molecular ion FeH⁺, among other iron containing molecular ions,² has long been suspected to be present in stellar atmospheres or in the interstellar medium (ISM).^{3,4} In the absence of laboratory data, however, it has not been identified so far in any astronomical environment. The only laboratory spectrum to date is the infrared photodissociation spectrum of Ar₂FeH⁺ in the Fe–H stretch region.⁵ Due to the shift induced by argon tagging, however, it is not suitable for comparison with data from astronomical observations.

Interstellar X-ray absorption spectroscopy in comparison with laboratory data has been used to assign the chemical form of iron in the ISM, and its fractionation into gas phase and dust particles.^{6–8} With no laboratory XAS data available, it has so far not been possible to consider iron containing molecular species like FeH⁺ in such studies. Here we report a spectrum of the iron L_{2,3} edges of FeH⁺ in the gas phase. It is the first laboratory spectrum of FeH⁺ in any spectral region that is suitable for comparison with data from astronomical observations. We also

briefly compare the results with recent spectra of diatomic iron-halide cations⁹ to discuss the implications of the spectrum for the electronic structure of FeH⁺, in particular the d-shell occupation, which so far has been studied only computationally.^{4,10,11}

X-ray absorption spectroscopy (XAS) was performed at the IonTrap station of the UE52-PGM beamline at the BESSY II electron storage ring operated by the Helmholtz-Zentrum Berlin für Materialien und Energie.¹² An electrospray ionization source with an ion funnel interface¹³ is used to produce FeH⁺ from dimethyl ferrocene in aqueous solution (see ESI† for details). This efficient way of making FeH⁺ was motivated by earlier work, which used electron ionization of dimethyl ferrocene.^{14,15} Mass selected FeH⁺ was trapped in a linear Paul trap and collisionally cooled to $T \approx 10$ K by using cryogenic helium buffer gas.¹⁶ X-ray absorption spectra with photon energy calibration with an uncertainty of less than 0.2 eV were recorded in partial ion yield mode,^{16,17} see ESI† for details. Supporting quantum chemical calculations were performed in Gaussian.¹⁸ Local 3d occupation was obtained by charge transfer multiplet (CTM) simulations using CTM4XAS,¹⁹ and compared with quantum chemical calculations. As usual, an empirical shift was applied to match the peak position of CTM4XAS calculations with experiment.

Fig. 1 summarizes the XAS spectra of FeH⁺, with the spectra of Fe⁺ shown for comparison.⁹ The L₃ edge of FeH⁺ exhibits less structure than the atomic ion, and the peak maximum is shifted to slightly smaller energies. However, the median L₃ energies at 708.1 eV are identical within the error bars, with a slightly higher uncertainty for FeH⁺ due to the smaller signal intensity for the molecular ion. The L₂ edge also stays at the same energy, with an overall similar peak shape. CTM simulations reproduce the overall spectral shape very well, with respect to peak broadening and substructure of the L₂ and L₃ edges. It should be noted that the CTM4XAS code and the underlying routines were developed for crystals in the solid state. Therefore the representation of the C_{∞v} point group of FeH⁺ by a crystal field in C₄ symmetry is not perfect.

In our earlier work on iron halides,⁹ we observed a strong correlation between the median position of the L₃ edge and the

^a Universität Innsbruck, Institut für Ionenphysik und Angewandte Physik, Technikerstraße 25/3, 6020 Innsbruck, Austria. E-mail: martin.beyer@uibk.ac.at

^b Abteilung für Hochempfindliche Röntgenspektroskopie, Helmholtz-Zentrum Berlin für Materialien und Energie, Albert-Einstein-Str. 15, 12489 Berlin, Germany

^c Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Hermann-Herder-Str. 3, 79104 Freiburg, Germany

^d Universität Innsbruck, Institut für Astro- und Teilchenphysik, Technikerstr. 25/8, 6020 Innsbruck, Austria. E-mail: norbert.prybilla@uibk.ac.at

† Electronic supplementary information (ESI) available: Tables S1 and S2 with results from quantum chemical calculations, further details on experiment, data evaluation, CTM simulations (with Table S3) and XMM-Newton data treatment (PDF); Experimental data shown in Fig. 1 and 2 (xlsx). See DOI: <https://doi.org/10.1039/d4cc04702h>






Fig. 1 Experimental iron $L_{3,2}$ edge XAS spectra of (a) FeH^+ (this work) and (b) Fe^+ (reproduced from Flach *et al.*, ref. 9), compared with CTM calculations (dashed lines). The black vertical lines in the figure are the median values of the experimental L_3 intensity distribution.

Mulliken electronegativity of the halogen^{9,21–24} as well as the 3d population of the iron center in FeX^+ , $X = \text{F}, \text{Cl}, \text{Br}, \text{I}$. Although both FeH^+ and FeX^+ contain a formal single bond and have $^5\Delta$ ground state, their electronic structure is completely different: one bonding σ molecular orbital is doubly occupied in FeH^+ , while doubly and singly occupied π , π^* and σ^* molecular orbitals contribute to the bonding in FeX^+ . To test whether the electronegativity correlation still holds despite these differences in electronic structure, we performed quantum chemical calculations. Tables S1 and S2 (ESI[†]) summarize Mulliken electronegativities and iron 3d and 4s populations obtained by a natural bond orbital analysis (NBO) with density functional theory (DFT). Obviously, the electronegativity correlation breaks down, while the 3d population prevails as the relevant property. CTM simulations yield an iron 3d population of 6.24, Table S3 (ESI[†]), in agreement with the values obtained from NBO analysis, Table S2 (ESI[†]).

In Fig. 2, we compare the Fe^+ and FeH^+ spectrum with observational data provided by the XMM-Newton Science Archive of the European Space Agency (ESA),²⁰ using spectra obtained *via* the reflecting grating spectrometer (RGS). Our




Fig. 2 Comparison of interstellar XAS spectrum with laboratory spectrum of Fe^+ and FeH^+ at the iron $L_{2,3}$ edges, after adding an empirical baseline and further processing (see text for details). The interstellar spectrum is taken from the XMM-Newton Science Archive.²⁰ The XMM-Newton RGS observations 0745250601 and 0745250501 were coadded.

laboratory spectra were converted to absorption spectra, convolved with a Gaussian kernel with the instrumental resolution of the RGS, binned to the RGS binning, and an arbitrary scaling and a simplified baseline were applied. The XMM-Newton data captures the total extinction, so both absorption and scattering, but scattering is negligible.⁶ This processing of the laboratory data more or less removes the subtle differences between the Fe^+ and FeH^+ laboratory spectra.

Since our model spectra agree in the onset and the lower-energy, but not in the higher-energy, part of the iron L_3 and L_2 edges of the interstellar spectrum, it is obvious that Fe^+ , FeH^+ , or any iron(II) species, alone cannot be the carrier of the $L_{2,3}$ edge absorption in the ISM along this sightline, as shown before,^{6,7} but that oxidation states of iron higher than +2 have to be present because the median excitation energy shifts to higher values with increasing oxidation state.²⁵ However, the interstellar X-ray absorption spectrum results from all iron compounds which are present in the interstellar medium. Since there is no feature in the FeH^+ spectrum that would not be covered by the absorptions in the Cygnus X-1 data, the presence of the FeH^+ molecular ion in the ISM cannot be ruled out. The similarity of the processed Fe^+ and FeH^+ spectrum in Fig. 2 underlines that FeH^+ may contribute to the gas-phase component of interstellar X-ray absorption at the iron $L_{2,3}$ edges.²⁶ However, the main carriers of the absorption obviously are other iron containing molecules or molecular ions, clusters or small particles.

With the X-ray absorption spectrum of FeH^+ we present the first laboratory data for this elusive species that can be directly compared with astronomical observations. Besides its astrochemical relevance, the spectrum carries information on the population of iron 3d orbitals in this molecular ion. Parameters for CTM simulations have been found that yield 3d occupations similar to quantum chemical NBO analysis. There is a clear need for X-ray reference data of iron species to identify further candidates for the carriers of interstellar X-ray spectra.

Shan Jin: data curation (supporting), investigation (equal), visualization (equal), writing – review and editing (supporting); Max Flach: conceptualization (equal), data curation (lead), formal analysis (lead), investigation (equal), writing – original draft (supporting), writing – review and editing (supporting); Alexander Ebenbichler: data curation (supporting), formal analysis (supporting), investigation (supporting), visualization (equal), writing – review and editing (supporting); Ethan M. Cunningham: funding acquisition (supporting), investigation (equal), project administration (supporting), resources (supporting), supervision (supporting); Vicente Zamudio-Bayer: investigation (supporting), writing – review and editing (supporting); Konstantin Hirsch: investigation (supporting), writing – review and editing (supporting); Christian van der Linde: methodology (supporting), writing – review and editing (supporting); Norbert Przybilla: data curation (supporting), funding acquisition (supporting), investigation (supporting), supervision (equal); Milan Ončák: data curation (supporting), formal analysis (supporting), investigation (equal), methodology (equal); J. Tobias Lau: conceptualization (equal), funding acquisition (equal), methodology (equal), writing – review and editing (supporting); Martin K.



Beyer: conceptualization (equal), formal analysis (supporting), funding acquisition (equal), supervision (equal), writing – original draft (lead).

Beamtime for this project was granted by HZB at beamline UE52-PGM Ion Trap of the BESSY II synchrotron radiation facility. JTL and MF acknowledge support by Deutsche Forschungsgemeinschaft (DFG) within RTG 2717. The computational results presented have been achieved using the HPC infrastructure LEO of the University of Innsbruck. This research was funded in part by the Austrian Science Fund (FWF), Grant DOIs 10.55776/W1259 (JS, AE, NP, MKB) and 10.55776/M3027 (EMC). For open access purposes, the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission. The authors thank Professor Frank de Groot for access to the CTM4XAS code and helpful advice on simulating iron $L_{2,3}$ edge XAS spectra.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Notes and references

- M.-F. Nieva and N. Przybilla, *Astron. Astrophys.*, 2012, **539**, A143.
- M. Lanza, A. Simon and N. Ben Amor, *J. Phys. Chem. A*, 2015, **119**, 6123.
- H. R. Johnson and A. J. Sauval, *Astron. Astrophys., Suppl. Ser.*, 1982, **49**, 77.
- Q. Cheng and N. J. DeYonker, *J. Chem. Phys.*, 2019, **150**, 234304.
- S. Jin, J. Heller, C. van der Linde, M. Ončák and M. K. Beyer, *J. Phys. Chem. Lett.*, 2022, **13**, 5867.
- L. Corrales, E. V. Gotthelf, E. Gatuzz, T. R. Kallman, J. C. Lee, M. Martins, F. Paerels, I. Psaradaki, S. Schippers and D. W. Savin, *Astrophys. J.*, 2024, **965**, 172.
- E. Costantini, C. Pinto, J. S. Kaastra, J. J. M. in't Zand, M. J. Freyberg, L. Kuiper, M. Méndez, C. P. de Vries and L. B. F. M. Waters, *Astron. Astrophys.*, 2012, **539**, A32.
- A. J. Westphal, A. L. Butterworth, J. A. Tomsick and Z. Gainsforth, *Astrophys. J.*, 2019, **872**, 66.
- M. Flach, K. Hirsch, M. Timm, O. S. Abyasova, M. da Silva Santos, M. Kubin, C. Bülow, T. Gitzinger, B. von Issendorff, J. T. Lau and V. Zamudio-Bayer, *Phys. Chem. Chem. Phys.*, 2022, **24**, 19890.
- M. Sodupe, J. M. Lluch, A. Oliva, F. Illas and J. Rubio, *J. Chem. Phys.*, 1989, **90**, 6436.
- S. R. Langhoff and C. W. Bauschlicher, Jr., *Astrophys. J.*, 1991, **375**, 843.
- K. Hirsch, J. T. Lau, P. Klar, A. Langenberg, J. Probst, J. Rittmann, M. Vogel, V. Zamudio-Bayer, T. Möller and B. von Issendorff, *J. Phys. B: At., Mol. Opt. Phys.*, 2009, **42**, 154029.
- S. Bari, D. Egorov, T. L. C. Jansen, R. Boll, R. Hoekstra, S. Techert, V. Zamudio-Bayer, C. Bülow, R. Lindblad, G. Leistner, A. Ławicki, K. Hirsch, P. S. Miedema, B. von Issendorff, J. T. Lau and T. Schlathöler, *Chem. – Eur. J.*, 2018, **24**, 7631.
- L. F. Halle, F. S. Klein and J. L. Beauchamp, *J. Am. Chem. Soc.*, 1984, **106**, 2543.
- E. I. Mysov, I. R. Lyatifov, R. B. Materikova and N. S. Kochetkova, *J. Organomet. Chem.*, 1979, **169**, 301.
- V. Zamudio-Bayer, R. Lindblad, C. Bülow, G. Leistner, A. Terasaki, B. von Issendorff and J. T. Lau, *J. Chem. Phys.*, 2016, **145**, 194302.
- J. T. Lau, J. Rittmann, V. Zamudio-Bayer, M. Vogel, K. Hirsch, P. Klar, F. Lofink, T. Möller and B. von Issendorff, *Phys. Rev. Lett.*, 2008, **101**, 153401.
- M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16 Revision A.03*, 2016.
- E. Stavitski and F. M. F. de Groot, *Micron*, 2010, **41**, 687.
- J. W. den Herder, A. C. Brinkman, S. M. Kahn, G. Branduardi-Raymont, K. Thomsen, H. Aarts, M. Audard, J. V. Bixler, A. J. den Boggende, J. Cottam, T. Decker, L. Dubbeldam, C. Erd, H. Goulooze, M. Güdel, P. Guttridge, C. J. Hailey, K. Al Janabi, J. S. Kaastra, P. A. J. de Korte, B. J. van Leeuwen, C. Mauche, A. J. McCalden, R. Mewe, A. Naber, F. B. Paerels, J. R. Peterson, A. P. Rasmussen, K. Rees, I. Sakelliou, M. Sako, J. Spodek, M. Stern, T. Tamura, J. Tandy, C. P. de Vries and A. Zehnder, *Astron. Astrophys.*, 2001, **365**, L7–L17.
- R. S. Mulliken, *J. Chem. Phys.*, 1934, **2**, 782.
- A. Kramida, Y. Ralchenko and J. Reader, *NIST ASD Team, NIST Atomic Spectra Database (ver. 5.8)*, [Online], available at: <https://physics.nist.gov/asd>, accessed 30 August 2021.
- R. C. Shiell, X. Hu, Q. J. Hu and J. W. Hepburn, *Faraday Discuss.*, 2000, **115**, 331–343.
- R. S. Berry and C. W. Reimann, *J. Chem. Phys.*, 1963, **38**, 1540.
- P. A. van Aken, B. Liebscher and V. J. Styrsky, *Phys. Chem. Miner.*, 1998, **25**, 323.
- E. Dwek, *Astrophys. J.*, 2016, **825**, 136.

