


# Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the RSC Publishing peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, which is prior to technical editing, formatting and proof reading. This free service from RSC Publishing allows authors to make their results available to the community, in citable form, before publication of the edited article. This *Accepted Manuscript* will be replaced by the edited and formatted *Advance Article* as soon as this is available.

To cite this manuscript please use its permanent Digital Object Identifier (DOI®), which is identical for all formats of publication.

More information about *Accepted Manuscripts* can be found in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics contained in the manuscript submitted by the author(s) which may alter content, and that the standard [Terms & Conditions](#) and the [ethical guidelines](#) that apply to the journal are still applicable. In no event shall the RSC be held responsible for any errors or omissions in these *Accepted Manuscript* manuscripts or any consequences arising from the use of any information contained in them.

## ARTICLE

# A combined solid- and solution-phase approach provides convenient access to analogues of the calcium-dependent lipopeptide antibiotics

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Peter 't Hart<sup>a,†</sup>, Laurens H. J. Kleijn<sup>a,†</sup>, Gerjan de Bruin<sup>b</sup>, Sabine F. Oppedijk<sup>a</sup>, Johan Kemmink<sup>a</sup>, and Nathaniel I. Martin<sup>a,\*</sup>


The calcium-dependent lipopeptide antibiotics represent a promising new class of antimicrobials for use in combatting drug-resistant bacteria. At present, daptomycin is the only such lipopeptide used clinically and displays potent antimicrobial activity against a number of pathogenic Gram-positive bacteria. Given the increasing need for new antibiotics, practical synthetic access to unnatural analogues of daptomycin and related antimicrobial lipopeptides is of value. We here report an efficient synthetic route combining solid- and solution-phase techniques that allows for the rapid preparation of daptomycin analogues. Using this approach, four such analogues, including two enantiomeric variants, were synthesized and their antimicrobial activities and hydrolytic stabilities evaluated.

## Introduction

The accelerated appearance of antibiotic resistance presents a serious and growing global health risk. Despite the increasing need for new antibacterial agents, only two mechanistically and structurally new antibiotics have reached the clinic in the past 40 years: linezolid and daptomycin.<sup>1,2</sup> While linezolid is of synthetic origin, daptomycin (Figure 1) is a natural product isolated from fermentations of *Streptomyces roseosporus*. Daptomycin is rapidly bactericidal against *Staphylococcus aureus*, including methicillin-resistant *S. aureus* (MRSA), vancomycin-intermediate *S. aureus* (VISA) and vancomycin-resistant *S. aureus* (VRSA) strains.<sup>3-5</sup> Marketed under the trade name Cubicin, daptomycin is the first lipopeptide antibiotic of its kind to be approved for clinical use. Structurally unique, daptomycin is a cyclic depsipeptide composed of 13 amino acids (including non-proteinogenic and D-amino acids) and bears an N-terminal 10-carbon lipophilic tail.

While the precise mechanistic details of daptomycin's antibacterial activity are unclear, it is known to disrupt aspects of bacterial cell membrane function.<sup>6,7</sup> In this regard, the current model for daptomycin's mode of action involves interaction with the bacterial membrane leading to rapid depolarization and a loss of membrane potential resulting in bacterial cell death without rupturing the cell.<sup>7</sup> Daptomycin's activity is calcium-dependent with serum levels of free calcium (45-55 µg/ml) sufficient to induce full antimicrobial activity.<sup>8</sup> Owing to the presence of four carboxylate side chains in the peptide, daptomycin is negatively charged at physiological pH and as such interacts with Ca<sup>2+</sup> ions.<sup>9</sup> Upon binding to Ca<sup>2+</sup> daptomycin is believed to oligomerize after which it is able to effectively disrupt the cell membranes of sensitive bacteria leading to defective cell division and cell wall synthesis.<sup>6,7</sup>

Given its role as the prototypical calcium-dependent lipopeptide antibiotic, daptomycin has garnered much attention from the synthetic community. Modified daptomycin analogues have been prepared via semi-synthesis<sup>10-13</sup> and the synthesis of daptomycin itself has been achieved using both chemo-enzymatic approaches or total synthesis. The Marahiel group demonstrated that linear precursor peptide thioesters could be converted to their corresponding daptomycin macrocycles by action of the appropriate recombinant cyclase enzyme.<sup>14</sup> In addition, the two total syntheses of daptomycin have also been reported to date.<sup>15,16</sup> In both cases the synthetic routes described are labor intensive and make use of multi-step synthetic strategies. For these reasons we were drawn towards developing a convenient synthetic route for the preparation of daptomycin analogues so as to provide more rapid access to structurally diverse derivatives of this important antibiotic.



**Figure 1.** Structure of daptomycin (**1**) with non-proteinogenic and D-amino acids indicated.


## Results and Discussion

In designing the daptomycin analogues to be prepared (Scheme 1) we opted to replace the L-threonine residue at position 4 with L-diaminopropionic acid. In doing so we were able to circumvent incorporation of the synthetically challenging<sup>15,16</sup> ester linkage between Thr4 and Kyn13 found in the natural daptomycin macrocycle by replacing it with the more accessible amide linkage. In this regard, we also speculated that the corresponding macrocyclic amide analogue of daptomycin might show improved hydrolytic stability (*vide infra*). Aside from the amide for ester modification, we also incorporated L-glutamic acid at position 12 in place of the (2*S*,3*R*)-3-methyl glutamate normally found in daptomycin. While published preparations of (2*S*,3*R*)-3-methyl glutamate are available,<sup>15-17</sup> they are rather lengthy (>10 steps) and time consuming. In addition, Marahiel and coworkers previously investigated the same Glu for 3-MeGlu substitution in their chemo-enzymatic approach to daptomycin analogues and found it to have a relatively small effect on antimicrobial activity (MIC values increased by ca. 7-fold).<sup>14</sup>

A variety of synthetic approaches were considered and explored in developing the route towards the daptomycin analogue illustrated in Scheme 1. Initial attempts at performing the entire synthesis on the solid phase by attachment to the resin via either of the aspartate side chains present in the macrocycle were unsuccessful and led us to explore the combined solid- and solution-phase approach indicated. To avoid any racemization in the later cyclization step, the peptide synthesis began at the glycine residue corresponding to position 5 in daptomycin. Employing an acid sensitive resin (2-chlorotrityl) and using standard SPPS techniques, the N-terminus of the peptide was first installed, including the C<sub>10</sub> fatty acid tail. At this point, the diaminopropionic acid side chain was deprotected providing an attachment point for the kynurenine residue (position 13) after which the remainder of the peptide was assembled without incident. Of particular note was the need to incorporate a rare Fmoc-Kyn building block which was obtained via a recently described methodology for the gram-scale production of kynurenine in either L- or D-stereochemistry.<sup>18</sup> In addition, DMB-protected glycine was introduced at position 10 to avoid aspartamide formation with the neighboring Asp residue. Upon completion, the intermediate protected peptide was cleaved from the resin using mild acid conditions and dissolved in CH<sub>2</sub>Cl<sub>2</sub> at high dilution (0.5 mM) followed by treatment with BOP/DIPEA which led to clean formation of the desired macrocycle (Scheme 1). Following deprotection and purification by RP-HPLC, daptomycin analogue **2** was obtained.

Based upon the concise route developed for the preparation of daptomycin analogue **2**, a second analogue, compound **3**, was also prepared wherein L-kynurenine at position 13 was replaced by the structurally similar L-tryptophan. The synthesis of **3** proceeded without incident after which the antimicrobial activity of analogues **2** and **3** was compared with that of authentic daptomycin (Table 1). Using a standard broth dilution assay employing *S. aureus* (ATCC 29213) as an indicator strain, analogues **2** and **3** were both found to exhibit calcium-dependent antimicrobial activity albeit at a significantly reduced level relative to that of daptomycin. The approximate 100-fold decrease in activity was somewhat expected given the structural modifications introduced in analogues **2** and **3**. Marahiel and coworkers reported similarly attenuated activities for daptomycin analogues obtained via their chemo-enzymatic approach.<sup>14</sup>

**Scheme 1.** Synthetic route developed for the preparation of daptomycin analogue **2**.



**Reagents and conditions:** a) Fmoc-Xaa-OH (or decanoic acid), BOP, DIPEA, DMF; b) Pd(PPh<sub>3</sub>)<sub>4</sub>, PhSiH<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>; c) Fmoc-Xaa-OH, BOP, DIPEA, DMF; d) HFIP/CH<sub>2</sub>Cl<sub>2</sub>; e) BOP, DIPEA, CH<sub>2</sub>Cl<sub>2</sub>, 40 h; f) TFA/TIS/H<sub>2</sub>O. Highlighted in blue are the structural differences between compound **2** and daptomycin

It is clear that replacement of the ester linkage in daptomycin with an amide dramatically alters antimicrobial activity. This effect may be due to conformational changes/restrictions in the macrocycle that result from incorporation of the amide. The amide for ester substitution is also expected to impart greater hydrolytic stability. Thus, the serum stability of analogues **2** and **3** was also evaluated and compared with that of daptomycin. Each peptide was incubated with human plasma serum at 37°C and sampled at specific time points. Under these conditions, daptomycin itself underwent significant degradation with an approximate 50% loss in the first 24 hours (Figure 2). This degradation is presumably due to

**Table 1.** Antimicrobial activity of daptomycin analogues against *S. aureus* (ATCC 29123)

Compound	MIC ( $\mu\text{M}$ ) <sup>a</sup>
<b>1 (Daptomycin)</b>	1.23
<b>2</b>	201.2
<b>3</b>	100.8
<b>ent-2</b>	not active <sup>b</sup>
<b>ent-3</b>	not active <sup>b</sup>

<sup>a</sup>Culture broth supplemented with  $\text{Ca}^{2+}$  (50 mg/L).<sup>b</sup>Not active at the highest concentration tested = 804  $\mu\text{M}$ .


hydrolytic opening of the macrocyclic lactone as evidenced by the appearance of a new  $\text{M}+\text{H}_2\text{O}$  species. By comparison, amide analogues **2** and **3** were much more stable under the same conditions with only minimal degradation detected over extended time periods of up to 48 hours.

The relative ease with which analogues **2** and **3** were assembled prompted us to also prepare the corresponding enantiomeric analogues *ent-2* and *ent-3*. The preparation of biologically active peptides in both enantiomeric forms is an established approach used in establishing the role of chiral biomolecular targets.<sup>19-22</sup> If as some have proposed, daptomycin kills bacteria via membrane disruption without invoking a specific chiral bacterial target biomolecule, one would expect the “mirror-image” enantiomeric form of daptomycin to show an equal antibiotic activity. Conversely, should the enantiomeric form of daptomycin not show antibacterial activity, it can be taken as an indication that a chiral interaction with a specific biomolecular target is integral to daptomycin’s mode of action. To this end the enantiomeric analogues *ent-2* and *ent-3* were assembled using the appropriate stereochemically-inverted amino acid building blocks. As expected, the enantiomeric analogues were shown to have identical retention times by analytical RP-HPLC (Figure 3) and exhibited optical rotations of equal magnitude but opposite sign. In addition, the circular dichroism spectra obtained for the daptomycin analogues further supported their enantiomeric nature (Supplemental Figure S3).

The antibacterial activities of *ent-2* and *ent-3* were next evaluated using the same assay as described above and revealed no detectable activity for both enantiomeric analogues (Table 1). These results indicate that a specific chiral interaction(s) is required for the activity of those analogues bearing the “native” daptomycin stereochemistry and may also support a similarly stereospecific mode of action for daptomycin itself. Plausible candidates for the requisite chiral target could include any number of membrane proteins or chiral phospholipids. In this regard, recent reports suggest that phosphatidylglycerol may play a role. Specifically, in daptomycin-resistant strains of enterococci and *S. aureus* levels of phosphatidylglycerol are significantly reduced.<sup>23</sup>


## Conclusions

In summary, we have developed a convenient approach employing both solid- and solution phase techniques for the preparation of analogues of the calcium-dependent lipopeptide family of antibiotics. While the antibacterial activity of the synthetic analogues was significantly reduced relative to that of



**Figure 2.** Serum stability of authentic daptomycin **1** (A) compared with analogues **2** (B) and **3** (C). Significant degradation of daptomycin is observed upon incubation at 37 °C while amide analogues **2** and **3** display increased stability.

daptomycin, their hydrolytic stability was greatly increased. The synthetic route here described should also be readily amenable to the production of new lipopeptide analogues with enhanced properties. Future work will be aimed at evaluating the effect of incorporating different amino acids in an attempt to increase antibacterial activity while maintaining the hydrolytic stability of these analogues. Support for this approach is evidenced by the observation that a tryptophan-for-kynurenine substitution as in compound **3** results to a more active analogue. In addition, our findings with the enantiomeric analogues implicate the involvement of a specific chiral target biomolecule(s) in bacterial strains that are sensitive to daptomycin.



**Figure 3.** Overlays of analytical RP-HPLC traces obtained for enantiomeric daptomycin analogues A) **2** and *ent-2* and B) **3** and *ent-3*.

## Experimental

**Reagents and General Methods.** All reagents employed were of American Chemical Society (ACS) grade or finer and were used without further purification unless otherwise stated. Both L- and D-kynurenine were synthesized according to a previously described procedure<sup>18</sup> and converted into the requisite Fmoc-building blocks based on established protocols.<sup>14</sup> A large scale preparation of Fmoc-(Dmb)Gly-OH was also developed based on existing literature procedures (see supporting information for details).<sup>24, 25</sup> All known compounds prepared had NMR spectra, mass spectra, and optical rotation values consistent with the assigned structures. Orthogonally protected Fmoc-L-Dap(Aloc)-OH and Fmoc-D-Dap(Aloc)-OH were obtained from Iris Biotech GmbH. All reagents employed were of American Chemical Society (ACS) grade or finer and were used without further purification unless otherwise stated. All reactions and fractions from column chromatography were monitored by thin layer chromatography (TLC) using plates with a UV fluorescent indicator (normal SiO<sub>2</sub>, Merck 60 F254). One or more of the following methods were used for visualization: UV absorption by fluorescence quenching; iodine staining; phosphomolybdic acid:ceric sulfate:sulfuric acid:H<sub>2</sub>O (10 g:1.25 g:12 mL:238 mL) spray; and ninhydrin staining. Flash chromatography was performed using Merck type 60, 230-400 mesh silica gel.

**Instrumentation for Compound Characterization.** NMR spectra were recorded at 300 or 500 MHz with chemical shifts

reported in parts per million (ppm) downfield relative to tetramethylsilane (TMS). <sup>1</sup>H NMR data are reported in the following order: multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; qn, quintet and m, multiplet), number of protons, coupling constant (*J*) in Hertz (Hz) and assignment. When appropriate, the multiplicity is preceded by br, indicating that the signal was broad. <sup>13</sup>C NMR spectra were recorded at 75.5 MHz with chemical shifts reported relative to CDCl<sub>3</sub> δ 77.0. 2D NMR experiments (TOCSY and HSQC) were performed on a 500 MHz instrument. High-resolution mass spectrometry (HRMS) analysis was performed using an ESI instrument. Circular dichroism spectra were recorded on a Jasco J-810 CD-spectrometer using a 2 mm cuvet.

**Peptide Synthesis.** Lipopeptide analogue **2** was synthesized beginning with Fmoc-Gly loaded 2-chlorotrityl resin (410 mg, 0.25 mmol). Fmoc groups were removed with 20% piperidine in DMF (10 mL, 1x1 min., 1x30 min.). Coupling reactions were done in DMF (10 mL) with 4 equivalents of Fmoc-amino acid (or capric acid), 4 equivalents of BOP and 8 equivalents of DIPEA (1 hour). For removal of the Aloc protecting group, the resin was first washed with CH<sub>2</sub>Cl<sub>2</sub> (2x10 mL) under argon after which PhSiH<sub>3</sub> (0.74 mL, 6.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) and Pd(PPh<sub>3</sub>)<sub>4</sub> (74 mg, 0.06 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (12 mL) were added and the mixture swirled under argon for 1 hour. The reaction mixture was drained and the procedure was repeated. To remove residual palladium catalyst, the resin was then washed with CH<sub>2</sub>Cl<sub>2</sub> (5x10 mL), a 0.5% solution of diethyldithiocarbamic acid trihydrate sodium salt in DMF (5x10 mL), and DMF (5x10 mL). After Aloc removal, the



remainder of the peptide was assembled on resin using standard SPPS approaches. The protected peptide was cleaved from the resin by addition of hexafluoroisopropanol (HFIP) in CH<sub>2</sub>Cl<sub>2</sub> (1:4, 6 mL, 1 hour) and the filtrate collected. The resin was rinsed with additional HFIP/CH<sub>2</sub>Cl<sub>2</sub> (2x3 mL) and the combined filtrates concentrated under vacuum. The crude linear peptide was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (500 ml, approximate peptide concentration of 0.5 mM) and BOP (0.22 g, 0.50 mmol, 2.0 equiv) and DIPEA (0.17 mL, 1.0 mmol, 4.0 equiv) were added. The cyclization reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) and was typically complete within 24 to 40 h. The reaction mixture was then concentrated, dissolved in EtOAc (250 mL) and washed with 1M KHSO<sub>4</sub> (2x200 mL) followed by saturated NaHCO<sub>3</sub> (2x200 mL). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under vacuum. The protected cyclic peptide was treated with a solution of TFA/TIS/H<sub>2</sub>O (95:2.5:2.5, 10 ml) for 1.5 hours. The mixture was then added to cold ether and the precipitated peptide collected by centrifugation and further was washed with cold ether (2x). Crude peptides were purified by RP-HPLC using a Maisch ReproSil-Pur C18-AQ column (250 x 22 mm, 10µm) and employing a gradient of 30% to 70% Buffer B with a flow of 12 ml/min (Buffer A: 95% H<sub>2</sub>O, 5% MeCN, 0.1% TFA; Buffer B: 5% H<sub>2</sub>O, 95% MeCN, 0.1% TFA). Product containing fractions were pooled and lyophilized to yield between 5-10 mg of pure peptide (1.2-2.4% overall yield). Table 1 summarizes the analytical data obtained for the four daptomycin analogues prepared as well as for authentic daptomycin.

**Biological activity assays.** The daptomycin analogues were tested against *S. aureus* (ATCC 29213) as an indicator strain. Two-fold serial dilutions of each compound were made in microtiter plates using Mueller-Hinton broth. After incubation at 37°C for 16 h, MIC values were determined visually. In these assays calcium- and magnesium-free Mueller-Hinton broth (Fluka) was supplemented with CaCl<sub>2</sub> (final Ca<sup>2+</sup> concentration 50 mg/L) and with MgCl<sub>2</sub> (final Mg<sup>2+</sup> concentration 10 mg/L) or alternatively with MgCl<sub>2</sub> alone to assess the effect of calcium on antibacterial activity.

**Serum stability assays.** Daptomycin and analogues **2** and **3** were added to human plasma serum to a final concentration of 150 µg/ml, followed by incubation at 37°C. At 0, 24 and 48 hours a 100 µl aliquot was taken and added to a 200 µl volume of methanol to precipitate plasma proteins (methanol also contained 0.075 mg/ml ethylparaben as an internal standard). The mixture was vortexed for 5 seconds and allowed to stand at room temperature for 15 minutes. After centrifugation at 13000 rpm for 5 minutes, the supernatant was removed and used for HPLC analysis. Analysis of the samples was performed by analytical RP-HPLC with a Maisch ReproSil-Pur 120 C18-AQ column (250 x 4.6 mm, 5µm) and a linear gradient of 0-100%

buffer B over 48 minutes at a flow rate of 1ml/min (Buffer A: 95% H<sub>2</sub>O, 5% MeCN, 0.1% TFA; Buffer B: 5% H<sub>2</sub>O, 95% MeCN, 0.1% TFA). Both the internal standard and peptides were detected at 220 nm allowing for the amount of intact peptide to be calculated based upon the ratio of peak areas corresponding to the internal standard and the intact peptides.

**Circular dichroism spectroscopy.** A 60 µM solution of each peptide was prepared by dissolving them in a 20 mM HEPES buffer (pH 7.4). Samples were measured at room temperature from 210-250 nm at a scan rate of 20 nm/min and a bandwidth of 0.1 nm. The spectra were converted to molar ellipticities in units of deg x cm<sup>2</sup>/dmol.

## Acknowledgements

We thank Eefjan Breukink for providing authentic daptomycin and Frederike Müskens for preparing L- and D-kynurenine. Javier Sastre Toraño is acknowledged for providing HRMS analysis and Jonas Dörr for assistance in acquiring the CD measurements. Linda Quarles van Ufford is kindly thanked for assisting with antibacterial assays. Financial support provided by The Netherlands Organization for Scientific Research (VIDI grant to NIM).

## Notes and References

<sup>a</sup>Medicinal Chemistry & Chemical Biology Group, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Universiteitsweg 99, 3584 CG Utrecht, The Netherlands. E-mail: n.i.martin@uu.nl  
<sup>b</sup>Bio-organic Synthesis Group, Leiden Institute of Chemistry, Einsteinweg 55, 2333 CC Leiden, The Netherlands.  
<sup>†</sup> Denotes equal contribution by P. t' Hart and L. H. J. Kleijn.

Electronic Supplementary Information (ESI) available: Preparation of Fmoc-(Dmb)Gly-OH and analytical RP-HPLC traces, CD spectra, and 2D-NMR spectra for peptides **2**, **3**, *ent-2* and *ent-3*. See DOI: 10.1039/b000000x/

1. T. Roemer and C. Boone, *Nat. Chem. Biol.*, 2013, **9**, 222.
2. G. D. Wright, *Nat. Rev. Microbiol.*, 2007, **5**, 175.
3. D. Patel, M. Husain, C. Vidailiac, M. E. Steed, M. J. Rybak, S. M. Seo and G. W. Kaatz, *Int. J. Antimicrob. Agents*, 2011, **38**, 442.
4. H. S. Sader and R. N. Jones, *Diagn. Microbiol. Infect. Dis.*, 2009, **65**, 158.
5. H. S. Sader, T. R. Fritsche and R. N. Jones, *Int. J. Infect. Dis.*, 2009, **13**, 291.
6. R. H. Baltz, *Curr. Opin. Chem. Biol.*, 2009, **13**, 144.
7. L. Robbel and M. A. Marahiel, *J. Biol. Chem.*, 2010, **285**, 27501.
8. G. M. Eliopoulos, C. Thauvin, B. Gerson and R. C. Moellering, Jr., *Antimicrob. Agents Chemother.*, 1985, **27**, 357.
9. R. H. Baltz, V. Miao and S. K. Wrigley, *Nat. Prod. Rep.*, 2005, **22**,

**Table 1:** Analytical data for compounds **1** (daptomycin), **2**, *ent-2*, **3**, and *ent-3*

Compound	R <sub>t</sub> (min)	[α] <sub>D</sub> <sup>25</sup>	Exact Mass (calc.)	Exact Mass (found)
<b>1</b> (Daptomycin)	30.2	N.D. <sup>a</sup>	1620.7182 [M+H] <sup>+</sup>	1620.7167 [M+H] <sup>+</sup>
<b>2</b>	29.3	-7.98 (0.28, H <sub>2</sub> O)	1591.7029 [M+H] <sup>+</sup>	1591.6997 [M+H] <sup>+</sup>
<i>ent-2</i>	29.3	+8.70 (0.28, H <sub>2</sub> O)	1591.7029 [M+H] <sup>+</sup>	1591.6995 [M+H] <sup>+</sup>
<b>3</b>	29.6	-9.45 (0.65, H <sub>2</sub> O)	1587.7080 [M+H] <sup>+</sup>	1587.7069 [M+H] <sup>+</sup>
<i>ent-3</i>	29.6	+9.24 (0.49, H <sub>2</sub> O)	1587.7080 [M+H] <sup>+</sup>	1587.7063 [M+H] <sup>+</sup>

<sup>a</sup>N.D. = not determined.

- 717.
10. J. Siedlecki, J. Hill, I. Parr, X. Yu, M. Morytko, Y. Zhang, J. Silverman, N. Controneo, V. Laganas, T. Li, J. Li, D. Keith, G. Shimer and J. Finn, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4245.
  11. J. Hill, J. Siedlecki, I. Parr, M. Morytko, X. Yu, Y. Zhang, J. Silverman, N. Controneo, V. Laganas, T. Li, J. J. Lai, D. Keith, G. Shimer and J. Finn, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4187.
  12. Y. He, J. Li, N. Yin, P. S. Herradura, L. Martel, Y. Zhang, A. L. Pearson, V. Kulkarni, C. Mascio, K. Howland, J. A. Silverman, D. D. Keith and C. A. Metcalf, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 6248.
  13. S. Yoganathan, N. Yin, Y. He, M. F. Mesleh, Y. Gui Gu and S. J. Miller, *Org. Biomol. Chem.*, 2013, **11**, 4680.
  14. J. Grunewald, S. A. Sieber, C. Mahlert, U. Linne and M. A. Marahiel, *J. Am. Chem. Soc.*, 2004, **126**, 17025.
  15. Cubist Pharmaceuticals Inc. (2007) Antiinfective Lipopeptides, WO Patent WO2006110185.
  16. H. Y. Lam, Y. Zhang, H. Liu, J. Xu, C. T. Wong, C. Xu and X. Li, *J. Am. Chem. Soc.*, 2013, **135**, 6272.
  17. C. Milne, A. Powell, J. Jim, M. Al Nakeeb, C. P. Smith and J. Micklefield, *J. Am. Chem. Soc.*, 2006, **128**, 11250.
  18. L. H. J. Kleijn, F. M. Muskens, S. F. Oppedijk, G. de Bruin and N. I. Martin, *Tetrahedron Lett.*, 2012, **53**, 6430.
  19. D. Wade, A. Boman, B. Wahlin, C. M. Drain, D. Andreu, H. G. Boman and R. B. Merrifield, *Proc. Natl. Acad. Sci. U. S. A.*, 1990, **87**, 4761.
  20. L. Z. Yan, A. C. Gibbs, M. E. Stiles, D. S. Wishart and J. C. Vederas, *J. Med. Chem.*, 2000, **43**, 4579.
  21. L. Sando, S. T. Henriques, F. Foley, S. M. Simonsen, N. L. Daly, K. N. Hall, K. R. Gustafson, M. I. Aguilar and D. J. Craik, *ChemBioChem*, 2011, **12**, 2456.
  22. F. Dettner, A. Hanchen, D. Schols, L. Toti, A. Nusser and R. D. Sussmuth, *Angew. Chem. Int. Ed. Engl.*, 2009, **48**, 1856.
  23. N. N. Mishra and A. S. Bayer, *Antimicrob. Agents Chemother.*, 2013, **57**, 1082.
  24. J. S. Yadav, S. Dhara, S. S. Hossain and D. K. Mohapatra, *J. Org. Chem.*, 2012, **77**, 9628.
  25. S. Zahariev, C. Guarnaccia, C. I. Pongor, L. Quaroni, M. Cemazar and S. Pongor, *Tetrahedron Lett.*, 2006, **47**, 4121.