

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Synthesis of Tetrabenzotriazacorrole μ -Oxo Dimer and Investigation of Its Stacking Effect

Cite this: DOI: 10.1039/x0xx00000x

Taniyuki Furuyama, Yusuke Sugiya and Nagao Kobayashi*

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

A μ -oxo dimer of phosphorus(V) tetrabenzotriazacorrole (PTBC) has been synthesized and characterized for the first time. Spectroscopic and theoretical analysis support the opinion that two PTBC chromophore groups interact with each other, while the low-symmetric PTBC structure is also efficient for generating singlet oxygen.

Phthalocyanine (Pc) and related tetrapyrrolic macrocycles are attractive target molecules in a variety of fields. These molecules have intense absorption bands in the UV-visible region, and their unique optical properties can be used as a key characteristic for constructing functional molecules. Since the optical properties of Pcs depend significantly on their structure, a number of Pc derivatives have been proposed and synthesized for over a century.¹ Among several methods in this regard, lowering of the molecular symmetry of Pcs is a good strategy.² However, separation of low-symmetry Pcs is generally not easy, and the yield is often low. Properties resulting from aggregation and/or oligomerization of Pcs are also important in various fields, such as material science and photochemistry.³ We focus on tetrabenzotriazacorrole (TBC) as a variant of low-symmetry Pc. TBC is known as a 18π aromatic Pc-corrole analogue, in which one *meso* nitrogen atom of Pc is missing. The direct pyrrole-pyrrole bond results in reduced symmetry (C_{2v}) compared to the D_{4h} of regular metalloPcs. TBCs can be synthesized from the corresponding symmetric Pcs in moderate yields with a simple procedure, and are easy to handle.⁴ In order to learn more about TBCs, we have chosen phosphorus(V) complexes which have two axial ligands.^{4a, c} Herein, we report the first synthesis and properties of a TBC μ -oxo dimer as a cofacial model of low-symmetry Pc oligomers. μ -oxo oligomers have been synthesized in Pc chemistry, and unique stacking effects observed.⁵

The synthetic route of the TBC μ -oxo dimer (**4**) is shown in Scheme 1a. Free-base β -(PhO)₈Pc **1** was reacted with excess phosphorus tribromide in pyridine, to give the TBC phosphorus complex **2**. The reaction was quenched with water, and the axial ligands of **2** were OH at this stage. The solution of **2** was heated in the presence of triphenylsilyl chloride, both the triphenylsiloxy-capped TBC monomer **3** and the desired dimer **4** were obtained. The bulkiness of the triphenylsilyl group appeared efficient for

stabilization of the dimer. The structure of **3** was finally elucidated by X-ray diffraction analysis of crystals (Scheme 1b and Fig. S1).[‡] The TBC macrocycle of **3** has a highly planar structure as previously reported phosphorus(V) corrorazine^{4c} and corrole⁶ complexes. The phosphorus ion is just the right size to fit into the center of the TBC ligand, while phosphorus(V) Pc has a ruffled structure⁷ due to mismatching between phosphorus and a central core of Pc.



Scheme 1 (a) Synthesis of tetrabenzotriazacorrole μ -oxo dimer. *Reagents and conditions:* (i) PBr₃ (20 eq), pyridine, reflux, 1 h, 55%; (ii) Ph₃SiCl, toluene, reflux, 3 h in the dark, 60% (**3**) and 3% (**4**). (b) X-ray crystal structure of **3**. The thermal ellipsoids were scaled to the 50% probability level. H atoms and axial ligands have been omitted for clarity.

¹H and ³¹P NMR spectra of **3** and **4** in CDCl₃ are shown in Fig. S2. Four sets of singlet assignable to the α -position of **3** are present at low magnetic field (9.09-9.00 and 8.32 ppm), reflecting the lower-symmetric TBC macrocycle. The ³¹P NMR spectrum of **3** exhibits only one peak, at -216 ppm, similar to porphyrin⁸ and phthalocyanine⁷ phosphorus(V) complexes. Both ¹H and ³¹P NMR spectra of **4** exhibit only one set of peaks for the TBC unit, and each TBC moiety could not be distinguished even at low temperature (Fig. S3). The peaks of **4** were observed at higher field than those of

3. A similar upfield shift was also observed in the case of the μ -oxo dimer of SiPc,^{5c} indicating that each TBC moiety is affected by the diatropic ring current generated in the other TBC moiety.

UV-vis and MCD spectra of compounds **3** and **4** in CHCl₃ are shown in Fig. 1. The absorption spectrum of capped monomer **3** exhibits two intense bands at 662 nm and 451 nm for the Q and Soret bands, respectively. Both the absorption coefficients and positions of the Q and Soret bands are close to those of the spectrum of **2** (Fig. S4). In the MCD spectrum, pseudo Faraday *A* terms of almost comparative intensity appeared corresponding to the Q and Soret absorption bands. On the other hand, the absorption spectrum of dimer **4** is completely different from that of **3**. Here, the absorption bands are blue-shifted to 635 and 425 nm with concomitant broadening, although the apparent intensity is close to that of the monomer. In the near-IR region (680–800 nm), a weak, broad band, monotonously decreasing toward longer wavelength, is seen, suggesting the presence of a low-energy forbidden transition for the dimer. In the MCD spectrum, an intense pseudo Faraday *A* term appeared, corresponding to the Q absorption peak, indicating that the MCD spectrum of cofacial dimers does not sensitively reflect the molecular symmetry of the constituent monomer.




Fig. 1 Magnetic circular dichroism (top), and electronic absorption (middle) spectra of **3** (black) and **4** (red) in CHCl₃, and theoretical absorption spectra (bottom) of **3'** (black) and **4'** (red). Calculations were carried out at the B3LYP/6-31G(d) level of theory.

The electrochemical properties of **3** and **4** were examined by cyclic voltammetry (Fig. 2). **3** showed one reversible oxidation couple at 0.19 V, while **4** showed two reversible oxidation couples at 0.44 and -0.02 V vs Fc^{+/0}. The first oxidation couple (E_{ox1}) shifted cathodically by ca. 0.2 V compared to that of the monomer **3**. Since the exact reduction potentials could not be estimated, their cathodic peak potentials, E_{pc} , were used as indexes of the first reduction potential. The close E_{pc} values clearly show that the LUMO energy levels of **3** and **4** are close. These voltammograms indicate that the gap between the HOMO and LUMO decreases after μ -oxo dimer formation of the TBC macrocycle.

In order to enhance the interpretation of the electronic structure of the TBC μ -oxo dimer and understand the origin of the blue-shifted absorption spectrum of **4** with narrower energy gap compared to

monomer **3**, model structures (**3'** and **4'**), whose peripheral substituents and axial triphenylsilyl groups are replaced by hydrogen, were considered and used in the calculations. The torsion angle (θ) of the optimized structure of **4'** is 41.4° (Fig. S5), which is larger than that of the crystallographic structure of the μ -oxo SiPc dimer ($\theta = 36.6^\circ$).^{5c} The optimized structures where θ was restricted were also obtained, but their energy difference was smaller than 4 kcal/mol (Fig. S6), thus allowing for the two TBC macrocycles to undergo rapid rotation, supporting the variable temperature ¹H NMR results.




Fig. 2 Cyclic voltammetry data for **3** (black), and **4** (red). Cyclic voltammograms were acquired from 1.0 mM solutions of analyte in 0.1 M ⁿBu₄NClO₄/o-DCB. Ferrocene was used as an internal standard and set to 0 V.

The calculated absorption spectra of **3'** and **4'** are shown at the bottom of Fig. 1, while partial MO energy diagrams are shown in Fig. S7, with the calculated transition energies, oscillator strengths (*f*), and configurations summarized in Table S1. As previously reported for the electronic structure of silicon(IV) corrolazine,⁹ the absorption spectrum of monomer **3'** can be explained by Gouterman's "four-orbital" theory.¹⁰ Both the calculated Q and Soret bands split into two due to the low-symmetry of the TBC chromophore. These bands are composed of transitions from the HOMO-1 and HOMO to the LUMO and LUMO+1, similarly to typical aromatic azaporphyrinoids. The frontier MOs of μ -oxo dimeric **4'** are delocalized over the entire complex. Here, the HOMO-LUMO gap of **4'** (1.93 eV) is narrower than that of **3'** (2.28 eV). On the other hand, the calculated HOMO-LUMO transition at 809 nm is very weak (*f* = 0.01). The main Q bands calculated at 583 and 565 nm are relatively strong, and the sum of the two *f* values (0.46) are close to that (0.41) of the calculated Q band for monomeric **3'** (595 and 564 nm), as is indeed observed experimentally for **3** and **4**. Based on the components of transitions for **4'**, the lowest-energy transition is forbidden, and the HOMO-3, HOMO-2, and HOMO-1 dominate in the intense Q absorption band for **4'**. Therefore, the transition component of the intense band in the Q band region of **4** is completely different from that of **3**, as a result of the orbital interaction between the two TBC chromophores. These spectroscopic features can also be interpreted conceptually by considering exciton coupling interactions between the co-facially arranged monomer units (H-type aggregate).¹¹ In the dimer, the orientation of two excitons derived from the monomer units are not perfectly parallel. However, as a first approximation, the band on the lower energy side of the Q band can be expressed as the difference of two electric dipole moments, and that on the higher energy side as the sum of the two moments; thus the band at higher energy becomes much stronger.

Finally, we demonstrate the photosensitising efficiency of TBCs¹² by determining their singlet oxygen quantum yields (Φ_{Δ}) by a steady-state method using 1,3-diphenylisobenzofuran (DPBF) as the chemical quencher. β -(4-*t*-BuOPh)₈Pc zinc complex (**Std-ZnPc**) was

used as a standard ($\Phi_{\Delta} = 0.73$ in CHCl_3).¹³ The photobleaching decays are shown in Fig. 3. TBC monomer **3** showed a high quantum yield ($\Phi_{\Delta} = 0.75$), while free-base Pc had a low Φ_{Δ} value. Compound **3** consists only of main-group elements above the third period, and therefore, its high Φ_{Δ} value is considered not to be caused by a heavy-metal effect, but due to the low-symmetry chromophore¹⁴ of TBC. On the other hand, the dimer **4** ($\Phi_{\Delta} = 0.25$) was much less efficient than **3**. In general, H-type aggregation in water reduces the lifetime of the excited state due to enhanced radiationless excited-state dissipation.¹⁵




Fig. 3 Comparison of the rate of decay of DPBF sensitized by **1** (red), **3** (blue), **4** (green) and **Std-ZnPc** (purple) in CHCl_3 as shown by the decrease in the absorbance at 415 nm. The absorption coefficient was normalized by the rates of light absorption coefficient over 600 nm.

In summary, a μ -oxo phosphorus(V) TBC dimer has been synthesized for the first time. In accordance with the narrower HOMO-LUMO gap derived from cyclic voltammograms, the dimer showed a weak Q absorption band with forbidden character at longer wavelength than the Q band of the monomer. The strong band of the dimer was observed to be blue-shifted than that of the constituting monomer. These spectral features could be explained by theoretical calculations, and conceptually by an exciton coupling model. The low-symmetry structure of TBC showed enhanced efficiency in generating singlet oxygen, so that the molecular symmetry appears to be important in the design of sensitizers. Further work is currently underway to tune the rotation properties by external stimuli.

This work was partly supported by a Grant-in-Aids for Scientific Research on Innovative Areas (25109502, ‘‘Stimuli-responsive Chemical Species’’), Scientific Research (B) (No. 23350095), Challenging Exploratory Research (No. 25620019) and Young Scientist (B) (No. 24750031) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT). The authors thank Dr. Eunsang Kwon (Tohoku University) for X-ray measurements. Some of the calculations were performed using supercomputing resources at the Cyberscience Center of Tohoku University.

Notes and references

Department of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan.

E-mail: nagaok@m.tohoku.ac.jp; Tel: +81 22 795 7719

‡ Crystallographic data for **3**: $\text{C}_{116}\text{H}_{78}\text{N}_7\text{O}_{10}\text{PSi}_2$, $M_w = 1817.00$, orthorhombic, space group $Pbca$ (no. 61), $a = 12.5583(2)$, $b = 31.0602(6)$, $c = 47.1556(8)$ Å, $V = 18393.7(6)$ Å³, $Z = 8$, $\rho_{\text{calcd}} = 1.312$ g cm⁻³, $T = -183^\circ\text{C}$, 202162 measured reflections, 16834 unique reflections ($R_{\text{int}} = 0.0934$), $R = 0.0974$ ($I > 2\sigma(I)$), $R_w = 0.3134$ (all data), goodness-of-fit on $F^2 = 1.035$, largest diff. peak/hole 1.007 and -0.317 e Å⁻³, CCDC 985980.

These data can be obtained free of charge from the Cambridge crystallographic data centre via <http://www.ccdc.cam.ac.uk/products/csd/request/>

† Additional spectroscopic data, full details of experimental and calculation procedures for all studied compounds. CCDC 985980. For ESI and crystallographic data in CIF or other electronic formats see 10.1039/c000000x/

- (a) *The Porphyrin Handbook*; K. M. Kadish, K. M. Smith and R. Guilard, Eds.; Academic Press: San Diego, United States, 2003; (b) *Handbook of Porphyrin Science*; K. M. Kadish, K. M. Smith and R. Guilard, Eds.; World Scientific Publishing, Singapore, 2010.
- (a) J. Mack and N. Kobayashi, *Chem. Rev.* 2011, **111**, 281; (b) J.-J. Cid, M. García-Iglesias, J.-H. Yum, A. Forneli, J. Albero, E. Martínez-Ferrero, P. Vázquez, M. K. Nazeeruddin, E. Palomares and T. Torres, *Chem. Eur. J.* 2009, **15**, 5130.
- (a) T. Fukuda, K. Matsumura and N. Ishikawa, *J. Phys. Chem. A* 2013, **117**, 10447; (b) M. García-Iglesias, K. Peuntinger, A. Kahnt, J. Krausmann, P. Vázquez, D. González-Rodríguez, D. M. Guldi and T. Torres, *J. Am. Chem. Soc.* 2013, **135**, 19311; (c) I. Sánchez-Molina, C. G. Claessens, B. Grimm, D. M. Guldi and T. Torres, *Chem. Sci.* 2013, **4**, 1338.
- (a) J. P. Fox and D. P. Goldberg, *Inorg. Chem.* 2003, **42**, 8181; (b) N. Kobayashi, F. Furuya, G.-C. Yug, H. Wakita, M. Yokomizo and N. Ishikawa, *Chem. Eur. J.* 2002, **8**, 1474; (c) B. Ramdhanie, C. L. Stern and D. P. Goldberg, *J. Am. Chem. Soc.* 2001, **123**, 9447; (d) J. Li, L. R. Subramanian and M. Hanack, *Chem. Commun.*, 1997, 679; (e) M. Fujiki, H. Tabei and K. Isa, *J. Am. Chem. Soc.*, 1986, **108**, 1532.
- (a) K. Oniwa, S. Shimizu, Y. Shiina, T. Fukuda and N. Kobayashi, *Chem. Commun.*, 2013, **49**, 8341; (b) J. Kleinwächter and M. Hanack, *J. Am. Chem. Soc.*, 1997, **119**, 10684; (c) D. W. DeWulf, J. K. Leland, B. L. Wheeler, A. J. Bard, D. A. Batzel, D. R. Dininny and M. E. Kenney, *Inorg. Chem.* 1987, **26**, 266; (d) B. L. Wheeler, G. Nagasubramanian, A. J. Bard, L. A. Schechtman, D. R. Dininny and M. E. Kenney, *J. Am. Chem. Soc.* 1984, **106**, 7404; (e) E. Ciliberto, K. A. Doris, W. J. Pietro, G. M. Reisner, D. E. Ellis, I. Fragala, F. H. Herstein, M. A. Ratner and T. J. Marks, *J. Am. Chem. Soc.* 1984, **106**, 7748.
- A. Ghosh and M. Ravikanth, *Chem. Eur. J.* 2012, **18**, 6386.
- T. Furuyama, K. Satoh, T. Kushiya and N. Kobayashi, *J. Am. Chem. Soc.* 2014, **136**, 765.
- K.-y. Akiba, R. Nadano, W. Satoh, Y. Yamamoto, S. Nagase, Z. Ou, X. Tan and K. M. Kadish, *Inorg. Chem.* 2001, **40**, 5553.
- N. Kobayashi, M. Yokoyama, A. Muranaka and A. Ceulemans, *Tetrahedron Lett.* 2004, **45**, 1755.
- M. Gouterman, *J. Mol. Spectrosc.* 1961, **6**, 138.
- M. Kasha, *Radiat. Res.*, 1963, **20**, 55.
- L. Huang, P. Zhao, Z. Li, F. Zhang and C.-H. Tung, *J. Phys. Chem. A*, 2008, **112**, 4165.
- S. E. Maree and T. Nyokong, *J. Porphyrin Phthalocyanines*, 2001, **5**, 782.
- K. Ishii, H. Itoya, H. Miwa, M. Fujitsuka, O. Ito and N. Kobayashi, *J. Phys. Chem. A* 2005, **109**, 5781.
- J. R. Darwent, P. Douglas, A. Harriman, G. Porter and M. C. Richoux, *Coord. Chem. Rev.* 1982, **44**, 83.