



**Cucurbiturils: From Synthesis to High-Affinity Binding and Catalysis**

Journal:	<i>Chemical Society Reviews</i>
Manuscript ID:	CS-REV-08-2014-000273.R1
Article Type:	Review Article
Date Submitted by the Author:	24-Sep-2014
Complete List of Authors:	Nau, Werner; Jacobs University, School of Engineering and Science Assaf, Khaleel; Jacobs University Bremen,

### Table of contents entry

Major developments in the synthesis of cucurbiturils and applications related to their high-affinity binding and catalysis have recently taken place.



## **Cucurbiturils: From Synthesis to High-Affinity Binding and Catalysis**

*Khaleel I. Assaf and Werner M. Nau*

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

## Cucurbiturils: From Synthesis to High-Affinity Binding and Catalysis


*Khaleel I. Assaf and Werner M. Nau*

Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

*Abstract.* In the wide area of supramolecular chemistry, cucurbit[*n*]urils (CB*n*) present themselves as a young family of molecular containers, able to form stable complexes with various guests, including drug molecules, amino acids and peptides, saccharides, dyes, hydrocarbons, perfluorinated hydrocarbons, and even high molecular weight guests such as proteins (*e.g.*, human insulin). Since the discovery of the first CB*n*, CB6, the field has seen tremendous growth with respect to the synthesis of new homologues and derivatives, the discovery of record binding affinities of guest molecules in their hydrophobic cavity, and associated applications ranging from sensing to drug delivery. In this review, we discuss in detail the fundamental properties of CB*n* homologues and their cyclic derivatives with a focus on their synthesis and their applications in catalysis.

## 1. Introduction


The chemistry of cucurbit[*n*]urils (CB*n*) is a rapidly developing field (Figure 1), which has been regularly reviewed from different perspectives.<sup>1-7</sup> Most recently, the synthesis of functionalized derivatives has progressed, and among the manifold applications, cucurbituril-catalyzed reactions are receiving increasing attention, which defines two focus areas of this review. The factors which govern their molecular recognition and high-affinity binding properties are continuously unfolding, such that we also update information pertinent in this fundamental area.<sup>8-11</sup> Biochemical and medicinal-chemical aspects<sup>12-32</sup> are excluded from this review, as are applications in chemosensing,<sup>33-48</sup> which in part have been recently reviewed elsewhere.<sup>49</sup>



**Fig. 1.** Molecular structures of the three smallest CB*n* homologues.


## 2. Synthesis

**Synthesis of Cucurbituril Homologues.** In 1905, the parent cucurbituril (CB6) was synthesized by Behrend and coworkers as a sparingly soluble “condensation product”.<sup>50</sup> Until today, cucurbiturils are produced by variations of the old synthesis, which involves acid-catalyzed condensation of glycoluril (**1**) and formaldehyde (Figure 2). The molecular structure of CB6 was uncovered by Mock and coworkers in 1981; Mock also coined the name “cucurbituril”, due to the resemblance of its structure to a pumpkin, which in turn belongs to the botanical family *cucurbitaceae*.<sup>51</sup> The group of Kim as well as that of Day varied the reaction conditions (e.g., 80-100°C, HCl or 9M H<sub>2</sub>SO<sub>4</sub>, 10-100h), which proved to be essential to successfully isolate other homologues, including CB5, CB6, CB7, CB8, and CB10•CB5; CB6 remained the major product.<sup>52-54</sup> The precise reaction mechanism of CB*n* was investigated in great detail by the group of Isaacs.<sup>55-58</sup>



**Fig. 2.** (Top) Synthesis of CB<sub>*n*</sub> homologues by condensation of glycoluril (**1**) and formaldehyde under acidic conditions. (Bottom) Different representations of the CB<sub>7</sub> structure.

Recently, the structure of the yet largest CB<sub>*n*</sub> member (CB<sub>14</sub>) has been reported with 14 normal glycoluril units linked by 28 methylene bridges (Figure 3).<sup>59</sup> The twisted CB<sub>14</sub> provided important information that larger CB<sub>*n*</sub> are actually being formed in the course of CB<sub>*n*</sub> synthesis.



**Fig. 3.** Top (left) and side (right) view of the X-ray crystal structure of the Eu<sup>3+</sup>CB<sub>14</sub> complex; hydrogen atoms removed for clarity.

It should be noted that CB<sub>*n*</sub> preparation and purification often introduces various impurities, always water and acid, frequently acetone and methanol, and, depending on the


isolated homologue, ammonium and alkali metal salts.<sup>53</sup> A number of techniques have been used to assess the purity (specifically the  $CB_n$  content) of  $CB_n$  samples, such as NMR, ITC, and TGA. Kaifer and coworkers have described a practical method to assay the purity of CB7 and CB8 based on UV-Vis titrations with cobaltocenium, which forms highly stable complexes with CB7 and CB8.<sup>60</sup>

**Synthesis of Cucurbituril Derivatives.** Functionalized  $CB_n$ , inverted- $CB_n$ , nor-seco- $CB_n$ , and various congeners have also been discovered (Table 1 and Figure 4), with essentially a new member reported in every year since 2000. Due to their enhanced cavity size and solubility, the new members display a range of novel applications, including their use for supramolecular vesicles, fluorescence sensing, drug delivery, catalysis, *etc.*<sup>22,25,31,44,48,61-65</sup>

**Table 1.** Timeline of the development of cucurbit[*n*]urils and related cyclic macrocycles.


Year	Compound name	Research group
1905	CB6 (not structurally identified)	Behrend <sup>50</sup>
1981	CB6	Mock <sup>51</sup>
1992	Me <sub>10</sub> CB5	Stoddart <sup>66</sup>
2000	CB5,7,8	Kim <sup>52</sup>
2001	CycH <sub>5</sub> CB5, CycH <sub>6</sub> CB6	Kim <sup>67</sup>
2002	Ph <sub>2</sub> CB6, CB10•CB5	Nakamura, Day <sup>54,68</sup>
2003	(HO) <sub>2n</sub> CB <sub>n</sub> , cucurbituril analogues, and partially substituted CB <sub>n</sub>	Isaacs, Kim, Day <sup>69-72</sup>
2004	Me <sub>4</sub> CB6, Me <sub>6</sub> CB6, hemicucurbit[ <i>n</i> ]uril ( <i>n</i> = 6 and 12)	Day, Tao, Keinan, Miyahara <sup>73-75</sup>
2005	CB10, <i>i</i> CB <sub>n</sub> ( <i>n</i> = 6 and 7), hemicucurbit[6]uril	Isaacs, Kim, Buschmann <sup>76-79</sup>
2006	<i>ns</i> -CB10	Isaacs <sup>80</sup>
2007	(±)-bis- <i>ns</i> -CB6, Me <sub>12</sub> CB6	Isaacs, Tao <sup>81,82</sup>
2008	Partially substituted CB <sub>n</sub> : Me <sub><i>n</i></sub> CB <sub>n</sub> ( <i>n</i> = 5 and 6), and CycH <sub><i>n</i></sub> CB6	Tao <sup>83-86</sup>
2009	Partially substituted CycP <sub><i>n</i></sub> CB6	Tao <sup>87</sup>
2010	Bambus[6]uril (Bu6)	Sindelar <sup>88</sup>
2011	(no new member reported)	
2012	Monofunctionalized CB6, monofunctionalized CB7	Scherman, Isaacs <sup>89-91</sup>
2013	CB14, norbornahemicucurbiturils, dimeric cucurbit[6]uril, and cyclohexylhemicucurbit[6]uril	Tao, Sindelar, Isaacs, Aav <sup>92,59,93,94</sup>
2014	Monosubstituted CB6 at methylene bridge (mPhCB6)	Sindelar <sup>95</sup>






**Fig. 4.** Molecular structures of different cucurbit[ $n$ ]urils and related cyclic macrocycles; colored parts indicate repetition units varying with  $n$ .

The first modified  $CB_n$  was reported by Stoddart in 1992, who synthesized the fully equatorially methylated  $CB_5$  ( $Me_{10}CB_5$ ).<sup>66</sup> Later, the group of Kim isolated the fully substituted cyclohexanecucurbit[ $n$ ]urils ( $CycH_5CB_5$  and  $CycH_6CB_6$ ), that are soluble in organic solvents and, surprisingly, also 170 times better soluble in water than parent  $CB_6$ .<sup>67,96</sup> Many more partially substituted  $CB_n$  and mixtures thereof have been obtained (Scheme 1 and Figure 5).<sup>53,68,69,71-73,82,84,85,87,96-101</sup> The reactions of substituted glycolurils so far reported favor invariably the formation of the smaller  $CB_n$  homologues,  $CB_5$  and  $CB_6$ . In fact, no fully substituted  $CB_7$  or  $CB_8$  have been isolated. Since alkylation of  $CB_n$  does not enable functionalization, Kim introduced in 2003 a direct method to produce fully hydroxylated  $CB_n$  by oxidation of unsubstituted  $CB_n$  under harsh conditions (hot  $K_2S_2O_8$ ). The resulting prehydroxyl- $CB_n$  could be easily modified to provide tailored  $CB_n$  derivatives with different functional groups, frequently accompanied by an enhanced solubility in common organic solvents.<sup>72</sup>




**Scheme 1.** Representative synthetic procedures for different cucurbituril derivatives; see Table 1 for references; dots mark connection points to the second identical half of the symmetric structures.




**Fig. 5.** Top views of the reported XRD molecular structures of substituted CB6 derivatives (codes taken from the CCDC data base); see Table 1 for references; dots mark connection points to the second identical half of the symmetric structures.

Isaacs *et al.* reported the isolation of inverted cucurbit[*n*]urils (*iCBn*, Figure 6), which contain a single inverted glycoluril unit directed towards the inside of the CB cavity.<sup>78</sup> *iCB6* and *iCB7* were isolated either by gel permeation chromatography or by fractional crystallization. Heating of *iCB6* or *iCB7* in concentrated HCl resulted in the conversion into mixtures of (regular) *CBn*.<sup>102</sup> An indication for the reverse process, interconversion of *CB7* to *iCB7*, has recently been obtained in the gas phase.<sup>103</sup> Until now, the inverted derivatives have not found any specific application.

Recently, different monofunctionalized *CB7* derivatives have been synthesized by Isaacs<sup>91</sup> through reaction of the acyclic glycoluril hexamer with a glycoluril bis(cyclic ether) (Scheme 2a). Further, the monofunctionalized *CB7* with a chloroalkyl tether can be transformed into the azide functional group, such that other substrates can be conveniently conjugated via click chemistry (Scheme 2b). The high water solubility of the monofunctionalized *CB7* (> 250 mM) and their host-guest recognition properties will render them solubilization agents for poorly soluble drug molecules and promising candidates for labeling and chemosensing applications as well as surface modification.




**Fig. 6.** Inverted cucurbiturils, *iCBn* ( $n = 6$  and  $7$ ); arrows indicate inverted glycoluril units; see Figure 4 for the molecular structure.




**Scheme 2.** a) Synthesis of Me<sub>2</sub>CB7, CycH<sub>2</sub>CB7, and MePhCB7. b) Synthesis of a monofunctionalized CB7 derivative; dots mark connection points to the second identical half of the symmetric structures.

A new related macrocycle, prepared by the acid-catalyzed condensation of 2,4-dimethylglycoluril and formaldehyde, was named bambus[6]uril (Figure 7), Bu6, which combines the structural features of both, cucurbiturils and hemicucurbiturils. Bu6 acts, however, as an anion receptor, e.g., it was shown to bind halide anions through C–H---X<sup>−</sup> interactions.<sup>88,104,105</sup>




**Fig. 7.** Side and top views of bambus[6]uril; see Figure 4 for the molecular structure.

By introducing new aldehydes to the reaction mixture of glycoluril and formaldehyde, Sindelar and coworkers have also synthesized the first example for a monosubstituted cucurbituril on the methylene-bridged position (Scheme 3). The most promising result was obtained with 3-phenylpropionaldehyde: (2-phenylethyl)-CB6 was isolated from the crude mixture in 0.2% yield.<sup>95</sup>



**Scheme 3.** Synthesis of the methylene bridge-monosubstituted cucurbituril mPheCB6.


Other prominent derivatives include bis-nor-seco-CB10<sup>80</sup> and nor-seco-CB6,<sup>81</sup> which were synthesized by the group of Isaacs. Bis-nor-seco-CB10 has a similar structure as CB10, but lacks two methylene bridges which results in two symmetrically interconnected cavities (Figure 8). Nor-seco-CB6 (Figure 4) is structurally comparable to CB6 but lacks a single methylene bridge. The removal of another methylene bridge results in ( $\pm$ )-bis-nor-sec-CB6 (Figure 4), the first reported chiral member of the CB $n$  family.



**Fig. 8.** Top view of bis-nor-seco-CB10; see Figure 4 for the molecular structure.

### 3. Physical Properties and Peculiarities of the Cucurbituril Cavity

The defining structural features of the  $CB_n$  family are their highly symmetric structure, with negatively charged carbonyl rims and a hydrophobic cavity. The electrostatic potential map of  $CB_n$  visualizes the high electron density at the carbonyl oxygens (Figure 9) and clearly illustrates their cation-receptor functionality. Their inner cavity, however, has neither functional groups nor electron pairs pointing towards the inside. It therefore cannot engage in hydrogen-bonding interactions, which – along with the non-dipolar nature of the macrocycle itself – provides an intuitive rationale for the high hydrophobicity of the cavity.



**Fig. 9.** Electrostatic potential map (top and side view) of CB7, revealing the negatively charged carbonyl portals (in red).

**Polarity and Polarizability.** Understanding the unique binding of cucurbiturils requires an understanding of the physical microenvironment of the inner cavity. When a guest is encapsulated within the inner cavity of CB, its absorption, fluorescence, and NMR spectra usually change. Such complexation-induced spectral shifts suggest changes in the surrounding electronic and/or magnetic environment. Several solvatochromic dyes have been utilized to



probe the cavity of  $CBn$ .<sup>36,106</sup> These probes are well known to display characteristic changes in their absorption, fluorescence, or phosphorescence spectra as the polarity of the (solvent) environment varies. For example, for CB7, the absorption spectra of Rhodamine 6G signal the immersion in a less polar environment when the dye is encapsulated. A polarity similar to that of alcohols (*n*-octanol,  $\epsilon = 10.3$ ) has been reported for CB7,<sup>107</sup> similar to the polarity determined for the cavities of other water-soluble macrocycles such as cyclodextrins or calixarenes. However, while  $CBn$  hosts have a nonpolar inner cavity and are themselves not dipolar, they display a very high quadrupole moment,<sup>8</sup> which allows higher-order electrostatic interactions. These have been implicated to account for the equatorial orientation of dipolar guests, such as ketones, in their cavity to achieve optimal quadrupole interactions.<sup>8, 108</sup> In contrast to the low polarity of the inner cavity of  $CBn$ , the portal regions of  $CBn$  have expectedly a more polar microenvironment.<sup>109</sup>

Beyond the relatively low polarity, we have shown that the inner cavity of  $CBn$  macrocycles has an extremely low polarizability/refractive index, which is closer to the gas-phase than to that of any other known solvent.<sup>110,111</sup> The polarizability was initially detected using DBO (2,3-diazabicyclo[2.2.2]oct-2-ene), a neutral azoalkane, as a spectrophotometric probe.<sup>110</sup> Its spherical shape and small size ( $111 \text{ \AA}^3$ ) allow it to be fully immersed inside the size-complementary cavity of CB7 ( $242 \text{ \AA}^3$ ) as well as that of other, similarly sized, water-soluble macrocycles, in particular  $\beta$ -cyclodextrin and *p*-sulfonatocalix[4]arene. A linear correlation between the inverse oscillator strength of the near-UV absorption band of DBO and the polarizability of different solvents was established, which could be subsequently used to interpolate or extrapolate the polarizability of several macrocyclic cavities, including CB7, cyclodextrins, and calixarenes (Table 2). The results showed that the CB7 interior has an extremely low polarizability (0.12), far below that of the two other macrocycles, which showed polarizabilities between those of water and benzene. The low polarizability of the  $CBn$  cavity can be rationalized along several lines: (i) There are no bonds or lone electron pairs pointing to the inside, (ii) the hosts are non-aromatic, (iii) any residual electron density at the nitrogens is mesomerically delocalized to the carbonyl oxygens, and, potentially, (iv) the packing is sufficiently loose to avoid close guest-to-wall contacts, e.g., the packing coefficient (*PC*) of DBO in CB7 is only 46% and therefore relatively loose, leaving some “vacuum”. For further discussion on *PC* arguments, see below and ref. 8.

**Table 2.** Refractive index and polarizability inside macrocyclic host molecules and different supramolecular environments, relative to those in selected solvents and the gas phase, taken from ref. 111.


Environment	Refractive index ( $n$ ) <sup>a</sup>	Polarizability ( $P$ ) <sup>b</sup>
Gas phase	1.000	0.000
CB7 <sup>c</sup>	1.17	0.11
Perfluorohexane	1.252	0.159
$\beta$ -Cyclodextrin	1.33	0.20
H <sub>2</sub> O	1.333	0.206
SDS micelles	1.34	0.21
<i>n</i> -Hexane	1.375	0.229
Triton XR-100 micelles	1.38	0.23
POPC liposomes	1.38	0.23
DMe- $\beta$ -CD (2:1 complex) <sup>d</sup>	1.39	0.24
Calixarene <sup>e</sup>	1.41	0.25
Benzene	1.501	0.295
Diiodomethane	1.742	0.404
Hemicarcerand <sup>f</sup>	1.86	0.45

<sup>a</sup> Refractive index, converted using the formula  $P = (n^2-1)/(n^2+2)$ . <sup>b</sup> From the empirical relationship ( $1/f = 3020-8320P$ ) by using the DBO chromophore as a solvatochromic probe. <sup>c</sup> For equatorial chromophore orientation; values for axial chromophore alignment are  $n = 1.19$  and  $P = 0.12$ . <sup>d</sup> Heptakis-(2,6-di-*O*-methyl)- $\beta$ -cyclodextrin. <sup>e</sup> *p*-Sulfonatocalix[4]arene, for equatorial chromophore orientation; values for axial chromophore alignment are  $n = 1.43$  and  $P = 0.26$ . <sup>f</sup> Value determined using biacetyl as solvatochromic probe, *cf.* ref. 110.

By exploiting DBO as a fluorescent probe, the low polarizability inside CB7 could be independently confirmed through very long radiative lifetimes, which are theoretically expected to increase as the refractive index of the environment, that is, its polarizability, decreases. Further independent evidence for the unique electronic environment offered by the interior of CB7 has recently been reported by Czar and Jockusch.<sup>112</sup> The authors used acridine orange (AOH<sup>+</sup>) as a probe to compare the gas-phase fluorescent properties of the free dye and its CB7 complex with that in aqueous solution. The excitation maxima of the dye in vacuum and in CB7 were the same, regardless of whether the CB7•AOH<sup>+</sup> complex is itself in the gas

phase or in aqueous solution. This result supports the finding that the cucurbituril interior is indeed “gas-phase-like”.

Most recently, we have involved NMR to further test the much debated hypothesis of a unique “phase of matter” inside cucurbiturils.<sup>45</sup> In detail, we took advantage of the fact that  $^{19}\text{F}$  chemical shifts do not strongly depend on magnetic shielding effects (such as  $^1\text{H}$  NMR shifts), but rather on the polarizability of the immediate surrounding of the F atoms.<sup>113, 114</sup> In an unconventional approach,  $\text{SF}_6$  and  $\text{CF}_4$  were used as “solvato-resonant” probes for the polarizability ( $P$ ) by establishing a linear empirical correlation between the  $^{19}\text{F}$  chemical shift and the polarizability of different solvents as well as the gas phase (Figure 10). The polarizability of the CB7 cavity, interpolated from the  $^{19}\text{F}$  chemical shift of the  $\text{CB7}\cdot\text{SF}_6$  complex, was found to be higher ( $P = 0.13$ ) than that of the gas phase ( $P \equiv 0.00$ ) but lower than that of water ( $P = 0.21$ ), a result which is well in line with previous results obtained by optical spectroscopy.<sup>36,106,110,112</sup>  $\text{CF}_4$  did not bind to CB7, but could be used as a solvato-resonant probe for CB6; in this case, the interpolated polarizability ( $P = 0.10$ ) was found to be even somewhat lower than the value obtained for  $\text{SF}_6$  inside CB7.<sup>115</sup>



**Fig. 10.** Linear correlation of the  $^{19}\text{F}$  chemical shift in the gas phase, different solvents, and in the complexes with CB6 and CB7 versus the polarizability ( $P$ ) of the environment a) for  $\text{SF}_6$  and b) for  $\text{CF}_4$  as solvato-resonant polarizability probes. The chemical shifts (down-field) are relative to the gas phase. The polarizabilities of the CB6 and CB7 cavities were interpolated from the correlations, see text and ref. 45.


**Structural features.** Table 3 lists pertinent structural parameters of common  $CB_n$  homologues and stereoisomers. In contrast to many other macrocyclic hosts,  $CB_n$  have a very rigid structure (borne out, for example, by small distortions in various crystal structures)<sup>116,117</sup> which renders the definition of the cavity parameters particularly informative. In detail, all  $CB_n$ , ( $n = 5-8$  and 10), have the same height ( $d = 9.1$  Å) but show large variations in cavity width (Table 3). It can be seen that the cavity width of  $CB_{10}$  is slightly too small to allow equatorial encapsulation of  $CB_5$ . Indeed, the x-ray structure of the  $CB_{10} \cdot CB_5$  complex showed that  $CB_5$  sits diagonally to the axis of  $CB_{10}$ .<sup>54</sup>

**Table 3.** Structural parameters of  $CB_n$  and  $iCB_n$ ; see Figure 2.

	Outer diameter	Inner cavity		Height
	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
$CB_5$	13.1	4.4	2.4	9.1
$CB_6$	14.4	5.8	3.9	9.1
$CB_7$	16.0	7.3	5.4	9.1
$CB_8$	17.5	8.8	6.9	9.1
$CB_9^a$	19.0	10.3	8.6	9.1
$CB_{10}^b$	20.0	11.7	10.0	9.1
$iCB_6^c$	10.7–14.4	3.8–5.8	4.3–3.9	9.1
$iCB_7^c$	11.2–16.0	5.4–7.3	6.7–5.4	9.1

<sup>a</sup> Values were calculated for the geometry of  $CB_9$  optimized by the B3LYP/6-311G\*\* method. <sup>b</sup> Average determined from the crystal structure with the CCDC code COQWOV, ref. 55. <sup>c</sup> Range defined by the shortest and longest wall-to-wall distances; determined from the crystal structures of  $iCB_6$  and  $iCB_7$  with the CCDC codes NEBDII and NEBDEE, respectively, from ref. 78.

The key-and-lock principle for macrocyclic hosts requires a complementarity between the guest size and the volume of the binding site to allow the formation of an inclusion complex. It has been shown that the “inner” cavity of  $CB_n$  (Figure 11) is best suited to assess this complementarity and to evaluate binding strengths on grounds of *PC* arguments (see next Section).<sup>8</sup> As can be seen from Table 4, the sizes of the inner  $CB_n$  cavities span a very large range from 68–691 Å<sup>3</sup>, those of the inverted  $iCB_n$  diastereoisomers are expectedly slightly smaller.



**Fig. 11.** a) Top and side views of different cavity definitions, taken for ref. 8. Copyright © 2011 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. b) Schematic illustration of an ideal cavity filling of 55% by encapsulation of a spherical guest.

**Table 4.** Calculated volumes ( $V/\text{Å}^3$ ) of the inner cavities of  $\text{CB}_n$  and  $i\text{CB}_n$ .<sup>8</sup>

Host	$V/\text{Å}^3$	Host	$V/\text{Å}^3$
CB5	68	CB7	242
$i\text{CB}_6^a$	121	CB8	367
CB6	142	CB9	515
$i\text{CB}_7^a$	221	CB10	691

<sup>a</sup> Values calculated in analogy to ref. 8.

**Packing Coefficient.** Rebek and Mecozzi introduced the packing coefficient ( $PC$ ), that is, the ratio of the guest size and the host cavity volume, as an estimator of the steric goodness of fit of host-guest inclusion complexes.<sup>118</sup> A value of ca. 55% was found to give the best binding affinities for the resulting host-guest complexes, while larger or smaller values were associated with lower affinities. Although the packing rule was originally derived for capsules as hosts, it has proven useful for several other macrocyclic receptors, particularly cucurbiturils.<sup>8</sup> It has not been written down yet that the number of 55% is actually a natural one as far as water-soluble host-guest systems are concerned: The packing coefficient of

water in water, for example, is also 55%, meaning that each water molecule ( $V = 17.0 \text{ \AA}^3$ ) has on average 45% void space at its disposition. The enthalpic minimum (estimated from the Lennard-Jones potential with  $\varepsilon_0 = 1.12\sigma$ ), lies at 71% for a spherical cavity, which is higher because of the neglect of entropy, the consideration of which lowers the ideal  $PC$  value.

We have previously demonstrated that  $PC$  arguments can be successfully applied to predict the stability of  $CB_n$  complexes.<sup>8</sup> The analysis of packing coefficients for representative sets of known guests with clearly defined hydrophobic binding motifs revealed average values of 47% for CB5, 58% for CB6, 52% for CB7, and 53% for CB8, which are well in line with the ideal packing (55% solution). In general, guests with associated packing coefficients of higher than 75% do not form stable inclusion complexes as a consequence of the associated entropic penalty unless additional interactions (such as ion-dipole ones) are at work. On the other hand, guests with packing coefficients lower than 30% also do not form stable complexes. Therefore, while the absolute  $PC$  values are only an empirical estimator, stable inclusion complexes result within the  $PC$  range of 45-65 %.

**Release of High-Energy Water Molecules.** The first guest molecules experimentally proven to be encapsulated in the hydrophobic cavity of a cucurbituril (CB6) were water molecules.<sup>52</sup> Depending on their cavity volume,  $CB_n$  can accommodate between 2 (for CB5) to 22 water molecules (for CB10). Estimates for the number of water molecules contained in the CB cavity are given in Table 5.<sup>8</sup> These water molecules are of high energy as a consequence of their greatly limited hydrogen bonding and due to weak dispersion interactions with the walls of the weakly polarizable CB cavity (see above).

In aqueous solution, the release of high-energy water from the cavity upon the complexation by a guest contributes a major part to the overall hydrophobic effect (Figure 12).<sup>8, 119-121</sup> Biedermann *et al.* used MD simulations to calculate the relative potential energy gain for the removal of all water molecules from  $CB_n$  cavities (Table 5).<sup>120</sup> The Results showed that the deficiency of hydrogen bonding of the encapsulated water molecules and their absolute number are the determinants for the overall gained energy. With increasing cavity size, the energy of each water molecule decreases due to improved hydrogen-bonding possibilities. The maximum calculated potential energy was obtained for CB7. Indeed, the  $242 \text{ \AA}^3$  large CB7 cavity does not allow the internal water molecules to arrange in a structurally stable H-bond network.


However, the energy of each individual internal water molecule is highest for small  $CB_n$ , such as CB5 and CB6, but the total energy for releasing all water from the inner cavity

of CB7 is higher due to the larger number of water molecules (more than twice) that are released upon the complexation process. On the other hand, the large cavity of CB8 allows the optimization of the H-bond network to a degree that is structurally similar to bulk water, with the result that the energy contribution for water release is less from CB8 than from CB7, despite its larger cavity. Kim and coworkers have reported another essential role for water molecules on host-guest recognition.<sup>122</sup> After the transfer of the CB $n$ -alkyldiammonium complexes from aqueous solution to the gas phase, it was found that the complex stabilities in the gas phase are different from those in aqueous solutions. The results were attributed to the fact that water molecules affect the ion-dipole interactions between the host and the guest. Indeed, in the absence of solvent, ion-dipole interactions are the main driving force for complexation. In contrast, in solution, the hydrophobic effect, and in particular the release of high-energy water, become dominant.

**Table 5.** Occupancy of the inner cavity of CB $n$  with water molecules.<sup>8</sup>

CB $n$	Number of water molecules in inner cavity		$\Delta E_{\text{pot}}^a$ (kJ mol <sup>-1</sup> )
	MD simulation	PC analysis	
CB5	2 [2.0] <sup>b</sup>	2	-41.6 ± 28.8
CB6	4 [3.3] <sup>b</sup>	4	-51.1 ± 29.0
CB7	7 [7.9] <sup>b</sup>	8	-102.4 ± 31.3
CB8	10 [13.1] <sup>b</sup>	12	-66.2 ± 10.7
CB9	14	16	----
CB10	20	22	----


<sup>a</sup> Difference in potential energy for the removal of all cavity water molecules and transfer of those to a spherical cavity in the aqueous bulk, see ref. 120. <sup>b</sup> Values from ref. 120 are given in square brackets.



**Fig. 12.** Schematic illustration of the release of high-energy water molecules from the CB7 cavity upon binding of a hydrophobic guest.


#### 4. Host-Guest Chemistry

The host-guest chemistry of  $CB_n$  has already been reviewed extensively.<sup>1,4,8,9,24,98,123-125</sup> The hydrophobic effect, with emphasis on the release of the high-energy water, as well as ion-dipole and dipole-dipole interactions have been addressed as the main driving forces for the binding of different guests by  $CB_n$ . In 1983, Mock and coworkers were the first to study the complexation of alkylammonium and alkyldiammonium ions with CB6 in aqueous formic acid and to determine their binding affinities.<sup>126</sup> Two years later, Freeman reported the first X-ray diffraction structure of a host-guest complex of a  $CB_n$ , that of the *p*-xylylenediammonium ion encapsulated by CB6 (Figure 13).<sup>127</sup>



**Fig. 13.** X-ray structure of the *p*-xylylenediammonium ion encapsulated by CB6, the first X-ray diffraction structure of a  $CB_n$  complex.<sup>127</sup>






**Fig. 14.** a) Relationship between the binding constant ( $\log K_a$ ) versus chain length  $m$  for  $\text{H}(\text{CH}_2)_m\text{NH}_3^+$  (●) and  $^+\text{H}_3\text{N}(\text{CH}_2)_m\text{NH}_3^+$  (▲). b) Dimensional comparison between CB6 and various  $\alpha,\omega$ -alkane diammonium ions.

The first structure-selectivity relationships between various alkyl- and aryl-substituted ammonium ions and CB6 were also established by Mock.<sup>128,129</sup> The extensive studies revealed that CB6 shows a chain length-dependent selectivity (Figure 14). Diammonium ions with a pentano or hexano bridge bind significantly stronger to CB6 than those with shorter or longer lengths. For 1,6-diammoniumhexane, the alkyl chain contracts to adopt a slightly folded conformation in order to optimize cavity filling and to additionally maximize ion-dipole interactions.<sup>130</sup> It was also found that CB6, with a cavity volume of  $142 \text{ \AA}^3$ , exhibits size and shape complementarity.<sup>44,128,129,131-134</sup> For example, CB6 can form an inclusion complex with *p*-tolylmethanamine, but not with its *ortho* or *meta* isomers. Similarly, we have shown that even though *n*-pentane, isopentane, and neopentane are equally large (all  $96 \text{ \AA}^3$ ,  $PC = 68\%$ ), only the first two guests showed sizable binding with CB6, while neopentane does not form

an inclusion complex with CB6, likely for kinetic reasons (slow ingress through the tighter portals).<sup>44</sup> A consistent result had been obtained by Mock for the neopentylammonium ion.<sup>126,128</sup> Moreover, CB6 differentiates between saturated and unsaturated hydrocarbons, with the latter displaying a weaker binding for the same carbon framework.<sup>44</sup>

As is the case for all macrocycles that possess a hydrophobic cavity, the guest hydrophobicity plays also an important role in the stability of CB complexes. For example, an alkylamine with a thioether linkage was found to bind more strongly than its oxygen counterpart, but less than the alkylamine itself. This result has been attributed to the hydrophobicity of an alkyl group being highest, that of an alkoxy group being lowest, and that of an alkylthio group being in between.<sup>128</sup> It should be noted that the driving force due to desolvation of the convex guest is mainly related to its surface area. It produces a “classical” hydrophobic effect with a favorable entropic component, which is different from the “nonclassical” effect due to the removal of high-energy water from concave cavities, which has a favorable enthalpic signature.<sup>121</sup>




**Fig. 15.** Representative structures illustrating the residue-selective binding of 1-(perfluorbutyl)pentane with a) CB6 and b) CB7.


Recently, we have reported a size-selective binding of alkyl *versus* perfluoroalkyl chains by CB $n$ .<sup>45</sup> CB7 shows preferential binding of perfluoroalkyl residues over normal alkyl groups, which characterizes this macrocycle as being at least as fluorophilic as lipophilic in nature. The <sup>1</sup>H and <sup>19</sup>F NMR of the *ditopic* guest 1-(perfluorbutyl)pentane clearly expose this binding preference (Figure 15). The preferential binding of the fluorinated residue can be reconciled in terms of the sufficiently large cavity size of CB7 (242 Å<sup>3</sup>), which snugly fits the perfluoroalkyl tail and results in ideal packing of 55%. In contrast, the smaller CB6 homologue (cavity size 142 Å<sup>3</sup>) showed a preferential complexation of the alkyl chain of the

same guest, because the volume of the perfluoralkyl group is slightly too large to be accommodated by the smaller host cavity.

**High-Affinity Binding.** CBs bind many size/shape complementary guests with very high affinity ( $10^3$ - $10^{10}$   $M^{-1}$ ). CB7, for example, forms stable 1:1 inclusion complexes with adamantanes<sup>135,136</sup> and diamantanes,<sup>137,138</sup> with bicyclooctanes,<sup>110,135,136</sup> and with ferrocene<sup>98,139-143</sup> as well as cobaltocene<sup>143-145</sup> derivatives (Table 6). The reported affinities of such residues can approach<sup>146,147</sup> or even exceed<sup>137</sup> those of the strongest non-covalent interactions found in nature, in the biotin-avidin pair. The ferrocene/CB7 complex was first reported by Kaifer<sup>143</sup> and Kim<sup>98</sup> in 2003. Figure 16 shows two crystallographically independent ferrocene/CB7 complexes that differ in their co-conformation, i.e., the relative alignment of the guest inside the host.<sup>148</sup> DFT calculations in the gas phase (B3LYP/6-31G\*) predict that the complex with perpendicular orientation should be favored.<sup>149</sup> We have attributed this to quadrupole interactions resulting from the high (negative) quadrupole moment of both, CB7 and ferrocene.<sup>8</sup> Further studies on ferrocene derivatives led to an extremely high affinity of  $10^{15}$   $M^{-1}$  by introducing an ammonium unit (Table 6).<sup>150</sup>




**Fig. 16.** Two different X-ray crystal structures of the complex between ferrocene and CB7 (side view for co-axial co-conformation and top view for orthogonal co-conformation), see ref. 139.



**Fig. 17.** Molecular structures of high-affinity binders to CB7.

Adamantylammonium was also found to form a very stable complex with CB7 ( $K_a = 4.2 \times 10^{12} \text{ M}^{-1}$ ).<sup>136</sup> Most recently, Isaacs and coworkers have reported the tightest binding that was ever measured for a monovalent molecular recognition event in water, between CB7 and diamantane diammonium ion, with  $K_a = 7.2 \times 10^{17} \text{ M}^{-1}$  in D<sub>2</sub>O and  $1.9 \times 10^{15} \text{ M}^{-1}$  in 50 mM NaO<sub>2</sub>CCD<sub>3</sub> by using <sup>1</sup>H NMR competition experiments.<sup>137</sup> The crystal structure of the CB7 complex, Figure 18, shows the complete inclusion of the diamantane residue in the CB7 cavity. The near perfect size as well as shape complementarity between the rigid CB7 cavity and the diamantane core led to an *attomolar* dissociation constant.



**Fig. 18.** Representations of the X-ray crystal structure of the CB7•diamantane diammonium complex. Taken from ref. 137. Copyright © 2014 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim.

The combined binding constants for CB7 in Table 6 can be approximately understood – considering only the composite hydrophobic effect and ion-dipole interactions, but ignoring differential dispersion interactions for the known reasons<sup>8</sup> – as follows: 1) The removal of all high-energy water molecules alone can account for a binding constant as high as  $10^8 \text{ M}^{-1}$ . 2) The desolvation of the guest contributes another factor of  $10^2$ - $10^4$ , depending on its

hydrophobic surface area. 3) Up to two suitably, ideally centro-symmetrically, placed trimethylammonium groups add another factor of  $10^3$  each. This results in the observed range from  $10^{10}$ - $10^{18}$ . Incidentally, the fact that a quarternary trimethylammonium group presents (when measured under identical conditions) a better binding motif than a ternary or primary ammonium group (Table 6) demonstrates nicely that hydrogen bonding interactions present no dominant driving force in the binding of guests to CB $n$  hosts. In fact, hydrogen bonding of guests with a host is in general unlikely to outperform hydrogen bonding with water molecules in aqueous solution.

**Table 6.** Binding constants for the complexation of adamantane (**A**), bicyclo[2.2.2]octane (**B**), ferrocene (**C**), and diamantane (**D**) derivatives with CB7 (see Figure 17 for molecular structures)

Guest	K / M <sup>-1</sup>	Guest	K / M <sup>-1</sup>
<b>A1</b> <sup>a</sup>	$7 \times 10^9$	<b>C1</b> <sup>d</sup>	$>10^6$
<b>A2</b> <sup>b</sup>	$2 \times 10^{10}$	<b>C2</b> <sup>e</sup>	$3 \times 10^9$
<b>A3</b> <sup>b</sup>	$2 \times 10^{14}$	<b>C3</b> <sup>b</sup>	$2 \times 10^{12}$
<b>A4</b> <sup>b</sup>	$1 \times 10^{14}$	<b>C4</b> <sup>e</sup>	$4 \times 10^{12}$
<b>A5</b> <sup>b</sup>	$8 \times 10^{14}$	<b>C5</b> <sup>e</sup>	$3 \times 10^{15}$
<b>A6</b> <sup>c</sup>	$2 \times 10^{12}$	<b>D1</b> <sup>a</sup>	$4 \times 10^9$
<b>A7</b> <sup>b</sup>	$5 \times 10^{15}$	<b>D2</b> <sup>f</sup>	$3 \times 10^{11}$
<b>B1</b> <sup>b</sup>	$6 \times 10^9$	<b>D3</b> <sup>f</sup>	$2 \times 10^{15}$
<b>B2</b> <sup>b</sup>	$2 \times 10^{14}$	<b>D3</b> <sup>f</sup>	$7 \times 10^{17g}$
<b>B3</b> <sup>b</sup>	$1 \times 10^{15}$		

<sup>a</sup> Unpublished results, by fluorescent indicator displacement. <sup>b</sup> From ref. 135. <sup>c</sup> From ref. 136.

<sup>d</sup> From ref. 143. <sup>e</sup> From ref. 150. <sup>f</sup> From ref. 137. <sup>g</sup> Measured in pure D<sub>2</sub>O.

Interestingly, diamantane in CB7 presents a more densely packed complex ( $PC = 79$  %) than adamantane in CB7 (61 %). The latter has an essentially perfect packing. Indeed, the binding constant of the diamantane derivative D2 is less than that of the adamantane derivative A3. The tight packing of D3 is structurally reflected in a significantly expanded cavity (increase in the diameter by  $\sim 0.3$  Å and increase in the volume from 242 to 258 Å<sup>3</sup>).<sup>137</sup> The fact that the D3 shows nevertheless the highest binding constant suggest that the  $PC$  value is over-compensated by the positive effect caused by the methylation of both amino groups; the latter causes a central positioning of the positive charge and, presumably,

increased Coulombic interactions due to a drop in the apparent dielectric permittivity between the interacting (partial) charges.

In contrast to other supramolecular recognition systems, the thermodynamic data for the complexation of  $CB_n$  with a variety of guest molecules overcome the enthalpy-entropy compensation trend.<sup>135,139,150</sup> The cancellation of enthalpic gain by commensurate entropic changes of opposite sign in the course of binding is common for tight non-covalent complexes, like protein-ligand binding; other macrocycles, e.g., cyclodextrins, also obey the enthalpy-entropy compensation principle.<sup>150-155</sup> This establishes a stand-alone feature of  $CB_n$  macrocycles, which can be attributed, among others, to the dominant role of the release of high-energy water as the enthalpic driving force for complexation.

Most recent, Vázquez *et al.* have reported a new method for identifying the quality of high affinity binders to CB7 and CB8 by indicator displacement, in which dyes with a macrocycle-specific optical fingerprint upon encapsulation by CB7 and CB8 were used. This method allows a rapid screening of strongly and selectively binding guest molecules at micromolar concentrations.<sup>156</sup>

## 5. Catalysis inside $CB_n$ Cavities


The potential of  $CB_n$  to control and catalyze chemical reactions through their distinctive inner cavity<sup>4,64,103,125,157-166</sup> has been explored to a considerably lesser extent than that of cyclodextrins as alternative macrocycles.<sup>167-173</sup> Nevertheless, several successful examples of  $CB_n$ -catalyzed or  $CB_n$ -promoted unimolecular and bimolecular reactions are nowadays available, and have been previously reviewed by Sivaguru.<sup>64</sup> Herein, we summarize chemical reactions catalyzed by  $CB_n$ , and highlight recent achievements (Table 7). Physical phenomena induced by complexation, such as the modulation of photoinduced electron transfer,<sup>174</sup> photostabilization,<sup>36</sup> complexation-induced  $pK_a$  shifts,<sup>16,18,175-177</sup> solubility enhancements of guests,<sup>14,178-182</sup> and adverse effects on chemical reactions, that is inhibition,<sup>183-187</sup> are not in the focus in this review, because they have been covered elsewhere.

**CB6-Catalyzed 1,3-Dipolar Cycloadditions.** The dipolar [3+2] cycloaddition between azides (**2**) and acetylenes (**3**) was the first reaction reported to be happening inside the CB cavity, particularly inside CB6.<sup>157</sup> In fact, it paved the way to a type of catalytic reactions which have later become famous as “click chemistry”.<sup>210,211</sup> As can be seen in Scheme 4, the formation of the [1,2,3]-triazole heterocycle **4** was accelerated by a factor of  $6 \times 10^4$  with

regiospecificity. Unambiguously, the strong ion-dipole interactions between the ammonium ions and the CB6 carbonyl rims stabilize the ternary complex. Mock explained the observed rate enhancement in terms of: 1) overcoming of entropic constraints and 2) a strain activation of the bound substrate.<sup>157</sup> Carlqvist and Maseras have used quantum-calculations (DFT, B3LYP/6-31G(d)) to provide more details on 1,3-dipolar cycloaddition inside CB6. The calculations, which lack consideration of dispersion interactions, showed no evidence for a transition state stabilization, which was interpreted in favour of the entropy constraints argument.<sup>212</sup> Be this as it may, the CB6-catalyzed click reaction has been subsequently extensively used for synthesis of rotaxanes, pseudorotaxanes, and polyrotaxanes.<sup>160,213-218</sup>

Table 7. Timeline for reports on chemical reactions mediated by CB $n$ .


Year	Reaction	Host (Catalyst)	Guest (Reactant)	Reference	Reaction conditions
1983	1,3-dipolar cycloaddition	CB6	<b>2</b> and <b>3</b>	157	Aq. formic acid, 40 °C
2001	Photodimeriz.	CB8	<b>10</b>	188	hv (300 nm)
2005,2006	Photodimeriz.	CB8	<b>20</b>	189, 190	hv (Hg lamp)
2005,2007,2014	Photodimeriz.	CB8	<b>13-15</b>	191-193	hv (Hg lamp)
2006	Photodimeriz.	CB7	<b>33</b>	194	hv (365 nm)
2008	Photodimeriz.	CB8	<b>29, 31</b>	195, 196	hv (Hg lamp)
2008,2010,2011	Photodimeriz.	CB8	<b>23</b>	197-200	hv (> 300 nm)
2008,2014	Photodimeriz.	CB8	<b>25</b>	201, 202	hv (> 300 nm)
2009	Hydrolysis	CB6, CB7	<b>7-9</b>	166	
2009,2011	Oxidation	CB8	alcohols	162, 203	IBX
2010	Oxidation	CB6	alkanes	204	Oxovanadium (IV)
2010	Desilylation	CB7	<b>44</b>	205	Ag(I) salts
2010	Hydrolysis	CB7	ArCOCl	184	
2011	Photolysis	CB7	<b>47</b> and <b>50</b>	206	Transition metal ions
2013	Oxidation	HemiCB6	<b>5</b>	207	
2013	Oxidation	HemiCB6	phenols	208	IBX
2013	Retro-Diels-Alder	CB6, CB7, CB8	azoalkanes, e.g., <b>47, 50</b>	103	Gas phase
2014	Photofragm.	CB8	<b>35</b> and <b>39</b>	209	hv (Xe lamp)



**Scheme 4.** Click reaction of azide (**2**) and alkyne (**3**) to form a [1,2,3]-triazole (**4**) catalyzed by CB6.

**Oxidation Reactions Catalyzed by CB $n$ .** The oxidation of alcohols into their corresponding aldehydes by *o*-iodoxylbenzoic acid (IBX) was investigated in the presence of CB8. The conversion of aryl and allyl alcohols was increased by 30-50% in the presence of CB8.<sup>162,203</sup> The pH effect on the conversion was also addressed; maximum conversion was obtained at neutral pH.<sup>203</sup> A similar catalytic effect was obtained when the same reaction was conducted in the presence of acetic acid.<sup>219</sup> The catalytic role of CB8 is not entirely understood in this case.

Reddy *et al.* studied a series of known reactions mediated by molecular halogens (iodine and bromine) and their solid-state CB6 complexes.<sup>220</sup> The solid state complexes were prepared by diffusing the halogen vapors through solid CB6. For example, the iodine-catalyzed Prins cyclization reaction was investigated by using free iodine *versus* the CB6•I<sub>2</sub> solid-state complex. The results revealed that, in the presence of the pre-complexed iodine, the cyclization proceeded in higher yields. It should be noted that the structures of the CB6•I<sub>2</sub> and CB6•Br<sub>2</sub> inclusion complexes are known,<sup>221</sup> while their use as reagents presents a novel application.<sup>220</sup>



**Scheme 5.** Aerobic oxidation of heterocyclic (**5**) mediated by hemicucurbit[6]uril.


Tao and coworker reported the aerobic oxidation of furan and thiophene (**5**) into their dione (**6**) conformation (Scheme 5). In aqueous solution, the oxidation was found to take place in the presence of HemiCB6, while no reaction happened in the absence of the host. The protonation of HemiCB6 appears to play a major role, because the oxidation was accelerated



at low pH.<sup>207</sup> The IBX oxidation of hydroxylbenzyl alcohols was investigated in the absence and presence of HemiCB6. The IBX oxidation of phenols is known to produce a mixture of aldehydes and o-quinones, while the presence of HemiCB6 directed the IBX oxidation towards the phenolic hydroxyl groups to produce aldehydes.<sup>208</sup>

**Hydrolysis Reactions Catalyzed by CBn.** García-Río and coworkers studied the hydrolysis of different substituted benzoyl chlorides in the presence of CB7 and DM- $\beta$ -CD (methylated  $\beta$ -cyclodextrin).<sup>184</sup> The reaction was catalyzed by CB7 and inhibited by DM- $\beta$ -CD when the benzoyl chloride was carrying electron-donating substituents. For the substrates with electron-withdrawing groups, the reaction was catalyzed by DM- $\beta$ -CD and inhibited by CB7. For example, the hydrolysis of 4-nitrobenzoyl chloride was inhibited in the presence of CB7 by a factor of 100, while for 4-methoxybenzoyl chloride the hydrolysis was accelerated 5-fold. The results were explained in terms of a differential stabilization of the acylium ion intermediate (in the transition state) by the CB7 carbonyl rim.<sup>184</sup>


Klöck *et al.* reported the hydrolysis of amides, carbamates (**7** and **8**), and oximes (**9**) in acidic aqueous solutions mediated by CBs (Figure 19).<sup>166</sup> In the presence of CB6 and CB7, the reaction showed a catalytic effect. The hydrolysis of carbamates with a cadaverine moiety, which is known to bind very strongly to CB6 and CB7, was accelerated by a factor of  $\sim 5$  ( $k_{\text{cat}} / k_{\text{uncat}}$ ) for CB6 and CB7 at pD 0.9 and by a factor of 11.6 for CB7 at pD 1.4. In the case of mono-*N*-(*tert*-butoxy)carbonyl (**8**), the deprotection rate constant was found to be  $\sim 30$  times faster with CB6 than the uncatalysed reaction. Furthermore, the oxime (**9**) hydrolysis in the presence of CB7 was studied. At high pD values (*e.g.* pD 5.8), the catalyzed reaction was very fast, with a peak acceleration factor of 285. The catalytic effect was attributed to a complexation-induced  $\text{p}K_{\text{a}}$  shift,<sup>15,17,159,160,181,182</sup> which assists the required protonation of the substrate.<sup>16,18,176,177,222,223</sup> Unfortunately, since the reaction product binds more strongly than the reactant (*e.g.*, benzaldehyde binds 6 times better than benzaldoxime), product inhibition was observed.



**Fig. 19.** Acidic hydrolysis of an amide (*N*-benzoyl-cadaverine, **7**), a carbamate (mono-*N*-(*tert*-butoxy)carbonyl cadaverine, **8**), and an oxime (benzaldoxime, **9**) mediated by *CBn*.<sup>166</sup>

**Photochemical Reactions Catalyzed or Templated by *CBn*.** Complexation by *CBn* greatly influences the photophysical properties of encapsulated guests<sup>34,36,42,175,224</sup> and their excited state processes<sup>4,53,183-187</sup> which can be used, in combination with their ability to control the spatial arrangement of two components in their ternary complexes, to design more efficient and more selective photochemical reactions.<sup>4,42,64,188,197,198,201</sup> Photochemical applications are facilitated by the fact that *CBn* are optically transparent in the near UV and the visible region, with the exception of some chromophore-modified derivatives.<sup>225-227</sup> In fact, other macrocyclic hosts (e.g., cyclodextrins,<sup>168-171</sup> calixarenes,<sup>228,229</sup> and the octaacid capsule<sup>230,231</sup>) have also been widely used as molecular containers for intermolecular photochemical reactions. Of interest in this context are particularly the larger *CBn* homologues, *CB7* and *CB8*, which are sufficiently large to accommodate two guests and, thus, to form ternary complexes, but which have only been reported in 2001.<sup>52</sup>

The first case of *CB*-catalyzed photoreactions with high stereoselectivity was reported by Kim and coworkers.<sup>188</sup> Two equivalents of (*E*)-diaminostilbene (**10**) were shown to form a stable 1:2 complex with *CB8* in aqueous solution, and upon UV-irradiation (at 300 nm, 30 min) the ternary complex underwent a [2+2] cycloaddition to yield the *syn*-adduct (**11a**) as major product and only a trace amount of the *anti*-adduct (**11b**). Interestingly, in the presence of *CB8*, no formation of the (*Z*)-isomer (**12**) was observed, while in the absence of *CB8* the (*Z*)-isomer (**12**) was the main product.<sup>232</sup> Other advantages offered by *CBn* were their ability to release the product from the cavity by increasing the pH (~9) and to be recycled after precipitation with excess methanol.<sup>188</sup>




Ramamurthy and coworkers carried out another impressive set of experiments.<sup>189-192</sup> By irradiation of different olefins (**13-15**) encapsulated in CB8 in aqueous solution, the *syn* dimer was obtained in 90% yield (**16a**), while irradiation of the 1:2 host-guest complex in the solid state showed no dimerization even after two days.<sup>191,192</sup> When these reactions were carried out without CB8, phenanthroline (**17**), the *cis*-isomer (**18**), and the hydration product (**19**, major product) were formed instead. Although photodimerization did not happen in the presence of CB7, the formation of **18** was promoted by this smaller host in higher yield (> 67%) than in the absence of CB7.<sup>191</sup> Symmetrical and unsymmetrical bispyridyl ethylenes, as well as stilbazole derivatives were employed in related photocycloaddition reactions; through a combination of best fit and a minimization of electrostatic repulsion within the CB8 cavity, new reactions could be designed.<sup>192</sup> Ramamurthy further extended this chemistry to include the photodimerization of neutral guests, such as trans-cinnamic acid (**20**).<sup>189,190</sup> Upon irradiation of **20** encapsulated by CB8 (in aqueous solution as well as in the solid state), the reaction afforded the *syn* head-to-head dimer (**21**), along with the corresponding *cis* isomer (**22**).<sup>189</sup> The latter was produced as dominant product in the presence of the smaller CB7. Zhang and coworkers have employed the same reaction type to construct covalently attached hyperbranched polymers, which present photosensitive supramolecular polymers with high molecular weight.<sup>193</sup>

Sivaguru reported the [2+2] cycloaddition of coumarin derivatives (**23**), both cationic and neutral, in the presence of CB8.<sup>197-199,233</sup> It was shown that upon UV excitation (> 300 nm) of the 1:2 complex of CB8 and two coumarins, the head-to-tail (HT) adduct was formed as the major product. The type of HT adduct (*syn* or *anti*) depended on the nature of substituents; coumarins with polar functional groups led to the *anti* photoadduct and those with nonpolar groups to the *syn* counterpart. Later, Sivaguru and coworkers showed that the formation of the 1:2 complex (ternary complex) is the rate determining step for the cycloaddition, in which the formation of the 1:1 complex is kinetically fast and that of the 1:2 complex is actually slow.<sup>200</sup> In the case of methylated coumarin (6-methylcoumarin), photodimerization led exclusively to the *syn* dimer (> 99%) with a *syn*-HH/*syn*-HT ratio of about 70 : 30.<sup>233</sup> The research groups of Inoue and Kim reported the effect of CB8 on the [4+4] cross-photodimerization of 2-anthracene carboxylate (**25**) and another 2-anthracene carboxylate linked to  $\alpha$ -CD (**27**).<sup>201</sup> In the case of the self-photodimerization of **25**, no significant effect on the *syn/anti* ratio or the head-to-head/head-to-tail ratio was observed in the presence and absence of CB8, e.g., the head-to-tail product remained the major isomer (**26a** and **26b**). In contrast, the cross-photodimerization of **27** inverted the HT/HH ratio form

98:2 to 1:99 by changing the templating host from  $\gamma$ -CD to CB8; accordingly, the products of the reaction with CB8 were **38a** and **38b**. The [4+4] photodimerization of anthracene in the presence of CB8 has recently been applied to polymer ligation and covalent network formation.<sup>202</sup> Another example, for the photodimerization of alkyl-2-naphthoate (**29**), was reported by Wu and coworkers, which afforded the cubane-type products **30**.<sup>195,196</sup> The photodimerization of methyl-2-naphthoate occurred efficiently in the presence of CB8 (60% conversion after 2 h), whereas in the free form methyl-2-naphthoate does not react. Similar results were obtained for 2-naphthalene-labeled poly(ethyl glycol) (**31**); the 1:1 complex was irradiated ( $> 280$  nm) for 12 min, sufficient to obtain more than 96% of the dimer (**32**); no product was formed in the absence of CB8.<sup>196</sup>

Macartney and coworkers demonstrated that photocycloadditions are not limited to the larger CB8 host, but that they can also be conducted with the intermediary sized CB7.<sup>194</sup> In the presence of 50 mol% of CB7, 2-aminopyridine (**33**, as hydrochloride salt) underwent a [4+4] photodimerization and afforded exclusively the *anti-trans* isomers (**34a**) in 90% yield after irradiation for 21 h. In the absence of CB7, a mixture of *anti-trans* (**34a**) and *syn-trans* (**34b**) was obtained in a 4:1 ratio. Importantly, CB7 was also found to stabilize the photoproduct and, thus, to protect it from thermal re-aromatization. Packing arguments suggest a stronger binding of the photoproduct in this case,<sup>8</sup> that is, product inhibition is expected, but actually desirable in this special case.




**Scheme 6.** CB8 modulates the phototransformation of a) benzimidazole (**35**) and b) thiabendazole (**39**); marked products were observed only in the presence of CB8.

The pre-assembled 1:2 host-guest complexes of CB8 and benzimidazole (**35**) underwent photohydrolysis to form a new major photoproduct, 2-aminofuranilide (**38**) (Scheme 6). When the reaction was carried out in the absence of CB8, the photoreaction of


BZ led to dehydrodimerization products (**36**) and (**37**). The phototransformation of free thiabendazole (**39**) also leads to the formation of **40**, **41**, and **42**, while in the presence of CB8 the coupling product **43** was formed as an additional product.<sup>209</sup>

**Catalysis Assisted by Metal Ions.** Due to their electronegative carbonyl rims,  $CB_n$  macrocycles are well known to bind metal cations at the portal regions.<sup>234-242</sup> This property has recently been employed to promote or catalyze organometallic reactions.<sup>204-206</sup> Demets and coworkers reported the oxidation of short unbranched alkanes, e.g. *n*-pentane, in the presence of CB6/oxovanadium (IV) (Figure 20).<sup>204</sup> Upon treatment of the complex with hydrogen peroxide or iodobenzene, a mixture of 2-pentanol, 2-pentanone, and 3-pentanone was obtained. The use of *Z*-cyclooctene, cyclohexane, and styrene as substrates showed no products formation, which was attributed to the fact that such guests do not form complexes with CB6, consistent with a detailed study of hydrocarbon binding to CB6.<sup>44</sup>




**Fig. 20.** The CB6/oxovanadium (IV) complex catalyzes *n*-pentane oxidation, but not that of larger alkanes.

Masson and Lu have recently reported a novel catalytic cycle for the desilylation of a trimethylsilylalkynyl derivative (**44**) assisted by  $CB_n$  in the presence of Ag(I) salts (Figure 21).<sup>205</sup> The formation of a ternary complex, e.g.,  $Ag^+ \cdot CB7 \cdot$ trimethylsilylalkynyl, was postulated, which would first lead to trimethylsilanol (**46**) complexed inside CB7 and an alkynylsilver organometallic complex. It was assumed that the latter is subsequently hydrolyzed to the desilylated alkyne (**45**) and  $Ag^+$ .



**Fig. 21.** CBn-catalyzed desilylation of trimethylsilylalkynyl derivatives (**44**) in the presence of Ag(I) salt; shown on top is the postulated ternary complex.

Our group documented a phase-selective photolysis of bicyclic azoalkanes (2,3-diazabicyclo[2.2.1]hept-2-ene (**47**) and 2,3-diazabicyclo[2.2.2]octa-2-ene (**50**)) promoted by transition-metal ions coordinated to the CB7 rim (Figure 22).<sup>206</sup> The reaction was selected to afford a photoproduct with a reduced affinity to the host and with a higher solubility in an organic phase than in water. In detail, irradiation of **47** and **50** in their near-UV  $n,\pi^*$  absorption band in water is known to cause nitrogen extrusion under formation of bicyclo[2.1.0]pentane (housane) (**48**) as exclusive photoproduct of **47** and of a 70:30 mixture of 1,5-hexadiene (**51**) and bicyclo[2.2.0]hexane (**52**) as photoproducts of **50**; the conversion can be conveniently monitored by gas chromatography. In the presence of CB7 and of specific transition metal ions ( $\text{Fe}^{3+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Ag}^+$ , also  $\text{Tl}^+$ ), the photolysis of DBO afforded a 5-fold to 10-fold excess of the diene product. The chemoselectivity was postulated to occur from the triplet-excited state, populated by heavy-atom-induced intersystem crossing. With the smaller azoalkane **46**, surprisingly, cyclopentene (**49**) was formed as a new photoproduct (41%), but only in the presence of both, CB7 and  $\text{Ag}^+$ . Because the formation of a ternary complex between CB7,  $\text{Ag}^+$ , and **50** could be crystallographically established, it is likely that a new mechanism, photoinduced electron-transfer, leads to cyclopentene. Such redox-active ternary complexes can be considered as metalloenzyme models.




**Fig. 22.** Chemoselective photoreactions of bicyclic azoalkanes (**54** and **57**) mediated by CB7 in the presence of transition metal ions.<sup>206</sup>

**Catalysis and Reactivity inside the Inner Phase of CB $n$  in the Gas Phase.** Lee *et al.* have investigated chemical reactions, retro-Diels-Alder reactions of bicyclic azoalkanes, inside CB $n$  molecular containers in the gas phase by using a mass spectrometric study combined with quantum-chemical calculations.<sup>103</sup> Ion mobility experiments as well as advanced DFT calculation were first performed to prove that the formed complexes between CB $n$  (CB6, CB7, and CB8) and the protonated azoalkanes (DBH, DBO, and DBN) are indeed inclusion complexes and not of the exclusion type. It was clearly shown that free CB7 exhibits the same drift time as the host-guest complex (Figure 23a), which depends directly on the collisional



cross section areas of the “empty” and the “filled” CB7. The fact that the cross sections are the same cannot be reconciled with an exclusion complex, because it would display a significantly larger cross section.



**Fig. 23.** A) Chemical structures of the bicyclic azoalkanes. b) Ion mobilitygrams of different CB7 complexes. c) Reaction and dissociation pathways of the inclusion complex of DBO inside CB7.

In a systematic approach, we have examined the fragmentation reaction of each viable complex of differently sized CB $n$  and guests (for example CB7, Figure 23b). After the mass selection, the protonated complexes, [CB $n$ •Guest•H]<sup>+</sup>, were subject to collision-induced dissociation (CID) experiments. This thermal activation of the CB $n$ -azoalkane complexes induced in some cases dominant chemical reactions of the encapsulated guests and the formation of [CB $n$ •fragment•H<sup>+</sup>] product complexes as opposed to irreversible dissociation of the host-guest complexes; the latter was expected and is observed for many supramolecular assemblies, including cyclodextrin complexes. For example, protonated DBH inside CB7 afforded elimination of ethylene with subsequent dissociation of the intermediary pyrazole complex upon CID. In contrast, when DBH was more tightly packed (inside CB6) or more loosely packed (with CB8), the simple dissociation pathway into guest and host dominated over the inner-phase chemical reaction. Most interestingly, thermal activation of DBO

complexed inside CB7 or CB8 (note that no complex was formed for DBO with CB6) led, predominantly, first to elimination of ethylene, followed by elimination of hydrogen cyanide, and, ultimately, dissociation of the protonated propenimine complex (Figure 23a).

The chemical reactivity of the gas phase complexes was interpreted in terms of three contributing factors: (1) the intrinsic activation energies for chemical reaction of the guest, (2) the constrictive binding displayed by the particular host, and (3) the void space inside the host-guest complex. Modelled Lennard-Jones 12-6 potentials of a spherical guest expanding inside the inner cavity of a rigid host, which included attractive (dispersion) and repulsive van der Waals forces, showed that the chemical reaction is “switched on” when the packing coefficients of the complexed guest falls within the range of 30-50%. In these cases, as shown by DFT calculations, the transition state experiences a dispersive stabilization, because the substrate must expand towards the cavity walls in order to eliminate the nitrogen molecule. These gas-phase reactions do therefore provide a genuine and puristic case of an inner-phase “catalysis” of a chemical reaction.

## 6. Conclusions and Outlook

Looking at only a single decade of active research on cucurbituril homologues, their potential in the areas of synthesis, high-affinity binding, and catalysis is still waiting to be fully explored. With respect to synthesis, several exciting developments have taken place, which included the emergence of monofunctionalized derivatives,<sup>90-92,96</sup> also through convergent synthesis routes (Table 1).<sup>87,95</sup> The monofunctionalized derivatives offer much potential for the labeling of surfaces and for selective conjugation to biomolecules. Also new homologues are still being reported, such as the recently reported largest member, CB14.<sup>59</sup> With respect to high-affinity binding, new host-guest combinations are still pushing the limits, and the highest value, attomolar binding, has just recently been reached.<sup>137</sup> Cucurbiturils are, among macrocycles, the hosts with highest (commonly 6 orders of magnitude higher) affinity, paving the way for *hitherto* unprecedented applications in biology, medicine, and material science. While recent studies point to the importance of the extrusion of high-energy water as an important denominator,<sup>119</sup> the quantitative understanding of the underlying factors responsible for the high-affinity binding of cucurbituril homologues remains challenging.<sup>243</sup> In contrast, dispersion interactions appear to be of very minor importance for guest binding by cucurbiturils in aqueous solution, which renders these hosts as interesting experimental and theoretical models to dissect the contributions of intermolecular interactions to host-guest binding, for example in the binding of perfluorinated compounds. Another mechanistic trend

in recent years involved the use of gas-phase studies to obtain fundamental information on the host-guest binding of cucurbiturils.<sup>103,112,122</sup> With respect to catalysis, to address the third thematic area of this review, several case studies are nowadays available (Table 7). Nevertheless, compelling application examples which fully exploit the high affinities of cucurbiturils, their peculiar binding properties, or the availability of derivatives (including monofunctionalized ones to explore heterogeneous conditions) are still waiting to be uncovered.

### Acknowledgements

The authors thank the Deutsche Forschungsgemeinschaft (DFG Grant NA-686/5), the Deutscher Akademischer Austausch Dienst (DAAD, graduate fellowships for KIA), and the COST Action CM1005 “Supramolecular Chemistry in Water” for their support.

### References

1. L. Isaacs, *Acc. Chem. Res.*, 2014, **47**, 2052.
2. *Special Issue: Cucurbiturils-consecutive reviews*, *Isr. J. Chem.*, 2011, **51**, 487.
3. N. Vallavoju and J. Sivaguru, *Chem. Soc. Rev.*, 2014, **43**, 4084.
4. E. Masson, X. X. Ling, R. Joseph, L. Kyeremeh-Mensah and X. Y. Lu, *RSC Advances*, 2012, **2**, 1213.
5. X.-L. Ni, X. Xiao, H. Cong, L.-L. Liang, K. Cheng, X.-J. Cheng, N.-N. Ji, Q.-J. Zhu, S.-F. Xue and Z. Tao, *Chem. Soc. Rev.*, 2013, **42**, 9480.
6. X. L. Ni, X. Xiao, H. Cong, Q. J. Zhu, S. F. Xue and Z. Tao, *Acc. Chem. Res.*, 2014, **47**, 1386.
7. J. Lü, J.-X. Lin, M.-N. Cao and R. Cao, *Coord. Chem. Rev.*, 2013, **257**, 1334.
8. W. M. Nau, M. Florea and K. I. Assaf, *Isr. J. Chem.*, 2011, **51**, 559.
9. Y. H. Ko, I. Hwang, D.-W. Lee and K. Kim, *Isr. J. Chem.*, 2011, **51**, 506.
10. E. Y. Chernikova, Y. V. Fedorov and O. A. Fedorova, *Russ. Chem. B+*, 2012, **61**, 1363.
11. A. E. Kaifer, *Acc. Chem. Res.*, 2014, **47**, 2160.
12. N. J. Wheate, D. P. Buck, A. I. Day and J. G. Collins, *Dalton Trans.*, 2006, 451.
13. H. D. Nguyen, D. T. Dang, J. L. J. van Dongen and L. Brunsveld, *Angew. Chem. Int. Ed.*, 2010, **49**, 895.
14. N. Dong, S.-F. Xue, Q.-J. Zhu, Z. Tao, Y. Zhao and L.-X. Yang, *Supramol. Chem.*, 2008, **20**, 659.
15. C. Kim, S. S. Agasti, Z. J. Zhu, L. Isaacs and V. M. Rotello, *Nat. Chem.*, 2010, **2**, 962.
16. I. Ghosh and W. M. Nau, *Adv. Drug Delivery Rev.*, 2012, **64**, 764.
17. A. L. Koner, I. Ghosh, N. Saleh and W. M. Nau, *Can. J. Chem.*, 2011, **89**, 139.
18. N. Saleh, A. L. Koner and W. M. Nau, *Angew. Chem. Int. Ed.*, 2008, **47**, 5398.
19. G. Ghale, V. Ramalingam, A. R. Urbach and W. M. Nau, *J. Am. Chem. Soc.*, 2011, **133**, 7528.
20. M. Florea and W. M. Nau, *Org. Biomol. Chem.*, 2010, **8**, 1033.
21. D. H. Macartney, *Future Med. Chem.*, 2013, **5**, 2075.
22. S. Walker, R. Oun, F. J. McInnes and N. J. Wheate, *Isr. J. Chem.*, 2011, **51**, 616.

23. N. Saleh, M. A. Meetani, L. Al-Kaabi, I. Ghosh and W. M. Nau, *Supramol. Chem.*, 2011, **23**, 654.
24. D. H. Macartney, *Isr. J. Chem.*, 2011, **51**, 600.
25. Y. Huang, Z. Tao, S.-F. Xue and Q.-J. Zhu, *Chem. J. Chinese. U.*, 2011, **32**, 2022.
26. N. J. Wheate, V. Vora, N. G. Anthony and F. J. McInnes, *J. Inclusion Phenom. Macrocyclic Chem.*, 2010, **68**, 359.
27. L. Goldoni, M. Grugni, S. De Munari, M. Cassin and R. Bernardini, *Chem. Lett.*, 2010, **39**, 676.
28. K. M. Park, K. Suh, H. Jung, D.-W. Lee, Y. Ahn, J. Kim, K. Baek and K. Kim, *Chem. Commun.*, 2009, 71.
29. Y. J. Jeon, S.-Y. Kim, Y. H. Ko, S. Sakamoto, K. Yamaguchi and K. Kim, *Org. Biomol. Chem.*, 2005, **3**, 2122.
30. S. Sun, F. Li, F. Liu, J. Wang and X. Peng, *Sci. Rep.*, 2014, **4**: 3570.
31. G. Hettiarachchi, D. Nguyen, J. Wu, D. Lucas, D. Ma, L. Isaacs and V. Briken, *PLoS One*, 2010, **5**, e10514.
32. V. D. Uzunova, C. Cullinane, K. Brix, W. M. Nau and A. I. Day, *Org. Biomol. Chem.*, 2010, **8**, 2037.
33. F. Biedermann, E. Elmalem, I. Ghosh, W. M. Nau and O. A. Scherman, *Angew. Chem. Int. Ed.*, 2012, **51**, 7739.
34. R. N. Dsouza, U. Pischel and W. M. Nau, *Chem. Rev.*, 2011, **111**, 7941.
35. M. Shaikh, S. D. Choudhury, J. Mohanty, A. C. Bhasikuttan and H. Pal, *Phys. Chem. Chem. Phys.*, 2010, **12**, 7050.
36. W. M. Nau and J. Mohanty, *Int. J. Photoenergy*, 2005, **7**, 133.
37. J. Mohanty, H. Pal, A. K. Ray, S. Kumar and W. M. Nau, *ChemPhysChem*, 2007, **8**, 54.
38. P. Montes-Navajas and H. Garcia, *J. Photochem. Photobiol. A*, 2009, **204**, 97.
39. N. Barooah, J. Mohanty, H. Pal and A. C. Bhasikuttan, *Org. Biomol. Chem.*, 2012, **10**, 5055.
40. H. Y. Zhang, L. L. Liu, C. Gao, R. Y. Sun and Q. C. Wang, *Dyes Pigm.*, 2012, **94**, 266.
41. E. Y. Chernikova, D. V. Berdnikova, Y. V. Fedorov and O. A. Fedorova, *Macroheterocycles*, 2013, **6**, 234.
42. G. Parvari, O. Reany and E. Keinan, *Isr. J. Chem.*, 2011, **51**, 646.
43. C. P. Carvalho, R. Ferreira, J. P. Da Silva and U. Pischel, *Supramol. Chem.*, 2013, **25**, 92.
44. M. Florea and W. M. Nau, *Angew. Chem. Int. Ed.*, 2011, **50**, 9338.
45. K. I. Assaf and W. M. Nau, *Supramol. Chem.*, 2014, **26**, 657.
46. R. N. Dsouza, A. Hennig and W. M. Nau, *Chem. Eur. J.*, 2012, **18**, 3444.
47. W. M. Nau, G. Ghale, A. Hennig, H. Bakirci and D. M. Bailey, *J. Am. Chem. Soc.*, 2009, **131**, 11558.
48. F. Biedermann and W. M. Nau, *Angew. Chem. Int. Ed.*, 2014, **53**, 5694.
49. G. Ghale and W. M. Nau, *Acc. Chem. Res.*, 2014, **47**, 2150.
50. R. Behrend, E. Meyer and F. Rusche, *Justus Liebigs Ann. Chem.*, 1905, **339**, 1.
51. W. A. Freeman, W. L. Mock and N.-Y. Shih, *J. Am. Chem. Soc.*, 1981, **103**, 7367.
52. J. Kim, I.-S. Jung, S.-Y. Kim, E. Lee, J.-K. Kang, S. Sakamoto, K. Yamaguchi and K. Kim, *J. Am. Chem. Soc.*, 2000, **122**, 540.
53. A. Day, A. P. Arnold, R. J. Blanch and B. Snushall, *J. Org. Chem.*, 2001, **66**, 8094.
54. A. I. Day, R. J. Blanch, A. P. Arnold, S. Lorenzo, G. R. Lewis and I. Dance, *Angew. Chem. Int. Ed.*, 2002, **41**, 275.
55. L. Isaacs, *Chem. Commun.*, 2009, 619.
56. W.-H. Huang, P. Y. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2008, **130**, 8446.

57. A. Chakraborty, A. X. Xu, D. Witt, J. Lagona, J. C. Fettinger and L. Isaacs, *J. Am. Chem. Soc.*, 2002, **124**, 8297.
58. L. Isaacs, *Isr. J. Chem.*, 2011, **51**, 578.
59. X.-J. Cheng, L.-L. Liang, K. Chen, N.-N. Ji, X. Xiao, J.-X. Zhang, Y.-Q. Zhang, S.-F. Xue, Q.-J. Zhu, X.-L. Ni and Z. Tao, *Angew. Chem. Int. Ed.*, 2013, **52**, 7252.
60. S. Yi and A. E. Kaifer, *J. Org. Chem.*, 2011, **76**, 10275.
61. H.-K. Lee, K. M. Park, Y. J. Jeon, D. Kim, D. H. Oh, H. S. Kim, C. K. Park and K. Kim, *J. Am. Chem. Soc.*, 2005, **127**, 5006.
62. M. del Pozo, E. Blanco, E. Fatás, P. Hernández and C. Quintana, *Analyst*, 2012, **137**, 4302.
63. F. Benyettou, I. Milosevic, Y. Lalatonne, F. Warmont, R. Assah, J.-C. Olsen, M. Jouaid, L. Motte, C. Platas-Iglesias and A. Trabolsi, *J. Mater. Chem. B.*, 2013, **1**, 5076.
64. B. C. Pemberton, R. Raghunathan, S. Volla and J. Sivaguru, *Chem. Eur. J.*, 2012, **18**, 12178.
65. S. Ghosh and L. Isaacs, *J. Am. Chem. Soc.*, 2010, **132**, 4445.
66. A. Flinn, G. C. Hough, J. F. Stoddart and D. J. Williams, *Angew. Chem. Int. Ed.*, 1992, **31**, 1475.
67. J. Z. Zhao, H.-J. Kim, J. Oh, S.-Y. Kim, J. W. Lee, S. Sakamoto, K. Yamaguchi and K. Kim, *Angew. Chem. Int. Ed.*, 2001, **40**, 4233.
68. H. Isobe, S. Sato and E. Nakamura, *Org. Lett.*, 2002, **4**, 1287.
69. C. A. Burnett, J. Lagona, A. X. Wu, J. A. Shaw, D. Coady, J. C. Fettinger, A. I. Day and L. Isaacs, *Tetrahedron*, 2003, **59**, 1961.
70. J. Lagona, J. C. Fettinger and L. Isaacs, *Org. Lett.*, 2003, **5**, 3745.
71. A. I. Day, A. P. Arnold and R. J. Blanch, *Molecules*, 2003, **8**, 74.
72. S. Y. Jon, N. Selvapalam, D. H. Oh, J.-K. Kang, S.-Y. Kim, Y. J. Jeon, J. W. Lee and K. Kim, *J. Am. Chem. Soc.*, 2003, **125**, 10186.
73. Y. J. Zhao, S. F. Xue, Q. J. Zhu, Z. Tao, J. X. Zhang, Z. B. Wei, L. S. Long, M. L. Hu, H. P. Xiao and A. I. Day, *Chin. Sci. Bull.*, 2004, **49**, 1111.
74. S. Sasmal, M. K. Sinha and E. Keinan, *Org. Lett.*, 2004, **6**, 1225.
75. Y. Miyahara, K. Goto, M. Oka and T. Inazu, *Angew. Chem. Int. Ed.*, 2004, **43**, 5019.
76. S. M. Liu, P. Y. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2005, **127**, 16798.
77. H.-J. Buschmann, A. Zielesny and E. Schollmeyer, *J. Inclusion Phenom. Macrocyclic Chem.*, 2006, **54**, 181.
78. L. Isaacs, S.-K. Park, S. M. Liu, Y. H. Ko, N. Selvapalam, Y. Kim, H. Kim, P. Y. Zavalij, G.-H. Kim, H.-S. Lee and K. Kim, *J. Am. Chem. Soc.*, 2005, **127**, 18000.
79. H.-J. Buschmann, E. Cleve and E. Schollmeyer, *Inorg. Chem. Commun.*, 2005, **8**, 125.
80. W.-H. Huang, S. M. Liu, P. Y. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2006, **128**, 14744.
81. W.-H. Huang, P. Y. Zavalij and L. Isaacs, *Angew. Chem. Int. Ed.*, 2007, **46**, 7425.
82. L.-B. Lu, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *Molecules*, 2007, **12**, 716.
83. J.-X. Lin, Y.-Q. Zhang, J.-X. Zhang, S.-F. Xue, Q.-J. Zhu and Z. Tao, *J. Mol. Struct.*, 2008, **875**, 442.
84. L.-B. Lu, D.-H. Yu, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *J. Mol. Struct.*, 2008, **885**, 70.
85. Z.-C. Tian, X.-L. Ni, X. Xiao, F. Wu, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *J. Mol. Struct.*, 2008, **888**, 48.
86. L. M. Zheng, J. N. Zhu, Y. Q. Zhang, Q. J. Zhu, S. F. Xue, Z. Tao, J. X. Zhang, X. Zhou, Z. B. Wei, L. S. Long and A. I. Day, *Supramol. Chem.*, 2008, **20**, 709.
87. L.-H. Wu, X.-L. Ni, F. Wu, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *J. Mol. Struct.*, 2009, **920**, 183.

88. J. Svec, M. Necas and V. Sindelar, *Angew. Chem. Int. Ed.*, 2010, **49**, 2378.
89. N. Zhao, G. O. Lloyd and O. A. Scherman, *Chem. Commun.*, 2012, **48**, 3070.
90. L. P. Cao and L. Isaacs, *Org. Lett.*, 2012, **14**, 3072.
91. B. Vinciguerra, L. P. Cao, J. R. Cannon, P. Y. Zavalij, C. Fenselau and L. Isaacs, *J. Am. Chem. Soc.*, 2012, **134**, 13133.
92. J. B. Wittenberg, P. Y. Zavalij and L. Isaacs, *Angew. Chem. Int. Ed.*, 2013, **52**, 3690.
93. T. Fiala and V. Sindelar, *Synlett*, 2013, **24**, 2443.
94. R. Aav, E. Shmatova, I. Reile, M. Borissova, F. Topić and K. Rissanen, *Org. Lett.*, 2013, **15**, 3786.
95. Laura Gilberg, M. S. A. Khan, M. Enderesova and V. Sindelar, *Org. Lett.*, 2014, **16**, 2446.
96. V. Lewin, J. Rivollier, S. Coudert, D.-A. Buisson, D. Baumann, B. Rousseau, F.-X. Legrand, H. Kouřilová, P. Berthault, J.-P. Dognon, M.-P. Heck and G. Huber, *Eur. J. Org. Chem.*, 2013, 3857.
97. A. X. Wu, A. Chakraborty, D. Witt, J. Lagona, F. Damkaci, M. A. Ofori, J. K. Chiles, J. C. Fettinger and L. Isaacs, *J. Org. Chem.*, 2002, **67**, 5817.
98. J. W. Lee, S. Samal, N. Selvapalam, H.-J. Kim and K. Kim, *Acc. Chem. Res.*, 2003, **36**, 621.
99. Y.-J. Zhao, S.-F. Xue, Y.-Q. Zhang, Q.-J. Zhu, Z. Tao, J.-X. Zhang, Z.-B. Wei and L.-S. Long, *Acta Chim. Sinica*, 2005, **63**, 913.
100. W.-J. Chen, J.-P. Zeng, Y.-Q. Zhang, Q.-J. Zhu, S. F. Xue, Z. Tao and G. Wei, *Chinese J. Inorg. Chem.*, 2010, **26**, 2018.
101. A. I. Day, X. Xiao, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *J. Inclusion Phenom. Macrocyclic Chem.*, 2011, **71**, 281.
102. S. M. Liu, K. Kim and L. Isaacs, *J. Org. Chem.*, 2007, **72**, 6840.
103. T. C. Lee, E. Kalenius, A. I. Lazar, K. I. Assaf, N. Kuhnert, C. H. Grun, J. Janis, O. A. Scherman and W. M. Nau, *Nat. Chem.*, 2013, **5**, 376.
104. V. Havel, J. Svec, M. Wimmerova, M. Dusek, M. Pojarova and V. Sindelar, *Org. Lett.*, 2011, **13**, 4000.
105. A. Révész, D. Schröder, J. Svec, M. Wimmerová and V. Sindelar, *J. Phys. Chem. A*, 2011, **115**, 11378.
106. M. A. Rankin and B. D. Wagner, *Supramol. Chem.*, 2004, **16**, 513.
107. J. Mohanty and W. M. Nau, *Angew. Chem. Int. Ed.*, 2005, **44**, 3750.
108. I. W. Wyman and D. H. Macartney, *Org. Biomol. Chem.*, 2008, **6**, 1796.
109. S. Senler, W. Li, M. H. Tootoonchi, S. Yi and A. E. Kaifer, *Supramol. Chem.*, 2014, **26**, 677.
110. C. Márquez and W. M. Nau, *Angew. Chem. Int. Ed.*, 2001, **40**, 4387.
111. W. M. Nau, A. Hennig and A. L. Koner, *Fluorescence of Supramolecules, Polymers, and Nanosystems*, 2008, **04**, 185.
112. M. F. Czar and R. A. Jockusch, *ChemPhysChem*, 2013, **14**, 1138.
113. R. J. Abraham and D. F. Wileman, *J. Chem. Soc., Perkin Trans. 2*, 1973, 1521.
114. D. F. Evans, *J. Chem. Soc.*, 1960, 877.
115. SF<sub>6</sub> as a solvato-resonant polarizability probe afforded a slightly higher polarizability for the CB6 cavity.
116. D. G. Samsonenko, A. V. Virovets, J. Lipkowski, O. A. Geras'ko and V. P. Fedin, *J. Struct. Chem.*, 2002, **43**, 664.
117. T. V. Mitkina, N. F. Zakharchuk, D. Y. Naumov, O. A. Gerasko, D. Fenske and V. P. Fedin, *Inorg. Chem.*, 2008, **47**, 6748.
118. S. Mecozzi and J. Rebek, Jr., *Chem. Eur. J.*, 1998, **4**, 1016.
119. F. Biedermann, M. Vendruscolo, O. A. Scherman, A. De Simone and W. M. Nau, *J. Am. Chem. Soc.*, 2013, **135**, 14879.

120. F. Biedermann, V. D. Uzunova, O. A. Scherman, W. M. Nau and A. De Simone, *J. Am. Chem. Soc.*, 2012, **134**, 15318.
121. F. Biedermann, W. M. Nau and H.-J. Schneider, *Angew. Chem. Int. Edit.*, 2014, DOI: 10.1002/anie.201310958.
122. S. J. C. Lee, J. W. Lee, H. H. Lee, J. Seo, D. H. Noh, Y. H. Ko, K. Kim and H. I. Kim, *J. Phys. Chem. B*, 2013, **117**, 8855.
123. P. Cintas, *J. Inclusion Phenom. Mol. Recognit. Chem.*, 1994, **17**, 205.
124. W. L. Mock, *Supramolecular Chemistry II - Host Design and Molecular Recognition*, 1995, **175**, 1.
125. J. Lagona, P. Mukhopadhyay, S. Chakrabarti and L. Isaacs, *Angew. Chem. Int. Ed.*, 2005, **44**, 4844.
126. W. L. Mock and N.-Y. Shih, *J. Org. Chem.*, 1983, **48**, 3618.
127. W. A. Freeman, *Acta Crystallogr. Sect. B-Struct. Sci.*, 1984, **40**, 382.
128. W. L. Mock and N.-Y. Shih, *J. Org. Chem.*, 1986, **51**, 4440.
129. W. L. Mock and N.-Y. Shih, *J. Am. Chem. Soc.*, 1988, **110**, 4706.
130. T. C. Krasia, S. Khodabakhsh, D. Tuncel and J. H. G. Steinke, *Cucurbituril: A Versatile "Bead" for Polyrotaxane Synthesis*, Springer-Verlag, Berlin - Heidelberg, 2004.
131. W. L. Mock and N.-Y. Shih, *J. Am. Chem. Soc.*, 1989, **111**, 2697.
132. W. L. Mock and J. Pierpont, *J. Chem. Soc., Chem. Commun.*, 1990, 1509.
133. C. Márquez, R. R. Hudgins and W. M. Nau, *J. Am. Chem. Soc.*, 2004, **126**, 5806.
134. A. V. Virovets, V. A. Blatov and A. P. Shevchenko, *Acta Crystallogr. Sect. B-Struct. Sci.*, 2004, **60**, 350.
135. S. Moghaddam, C. Yang, M. Rekharsky, Y. H. Ko, K. Kim, Y. Inoue and M. K. Gilson, *J. Am. Chem. Soc.*, 2011, **133**, 3570.
136. S. M. Liu, C. Ruspic, P. Mukhopadhyay, S. Chakrabarti, P. Y. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2005, **127**, 15959.
137. L. P. Cao, M. Šekutor, P. Y. Zavalij, K. Mlinarić-Majerski, R. Glaser and L. Isaacs, *Angew. Chem. Int. Ed.*, 2014, **53**, 988.
138. M. Šekutor, K. Molčanov, L. P. Cao, L. Isaacs, R. Glaser and K. Mlinarić-Majerski, *Eur. J. Org. Chem.*, 2014, **2014**, 2533.
139. W. S. Jeon, K. Moon, S. H. Park, H. Chun, Y. H. Ko, J. Y. Lee, E. S. Lee, S. Samal, N. Selvapalam, M. V. Rekharsky, V. Sindelar, D. Sobransingh, Y. Inoue, A. E. Kaifer and K. Kim, *J. Am. Chem. Soc.*, 2005, **127**, 12984.
140. Y. Chen, A. Klimczak, E. Galoppini and J. V. Lockard, *RSC Advances*, 2013, **3**, 1354.
141. I. Hwang, K. Baek, M. Jung, Y. Kim, K. M. Park, D.-W. Lee, N. Selvapalam and K. Kim, *J. Am. Chem. Soc.*, 2007, **129**, 4170.
142. A. E. Kaifer, W. Li and S. Yi, *Isr. J. Chem.*, 2011, **51**, 496.
143. W. Ong and A. E. Kaifer, *Organometallics*, 2003, **22**, 4181.
144. V. Logvinenko, T. Mitkina, V. Drebuschak and V. Fedin, *J. Therm. Anal. Calorim.*, 2011, **105**, 103.
145. D. Sobransingh and A. E. Kaifer, *Langmuir*, 2006, **22**, 10540.
146. N. M. Green, *Methods Enzymol.*, 1990, **184**, 51.
147. Y. Pazy, T. Kulik, E. A. Bayer, M. Wilchek and O. Livnah, *J. Biol. Chem.*, 2002, **277**, 30892.
148. X. Y. Zhang and W. M. Nau, *Angew. Chem. Int. Ed.*, 2000, **39**, 544.
149. R. V. Pinjari and S. P. Gejji, *J. Phys. Chem. A*, 2008, **112**, 12679.
150. M. V. Rekharsky, T. Mori, C. Yang, Y. H. Ko, N. Selvapalam, H. Kim, D. Sobransingh, A. E. Kaifer, S. M. Liu, L. Isaacs, W. Chen, S. Moghaddam, M. K. Gilson, K. M. Kim and Y. Inoue, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 20737.
151. M. V. Rekharsky and Y. Inoue, *Chem. Rev.*, 1998, **98**, 1875.

152. K. N. Houk, A. G. Leach, S. P. Kim and X. Y. Zhang, *Angew. Chem. Int. Ed.*, 2003, **42**, 4872.
153. Y. Inoue, Y. Liu, L.-H. Tong, B.-J. Shen and D.-S. Jin, *J. Am. Chem. Soc.*, 1993, **115**, 10637.
154. Y. Inoue, T. Hakushi, Y. Liu, L.-H. Tong, B.-J. Shen and D.-S. Jin, *J. Am. Chem. Soc.*, 1993, **115**, 475.
155. C.-E. Chang and M. K. Gilson, *J. Am. Chem. Soc.*, 2004, **126**, 13156.
156. J. Vázquez, P. Remón, R. N. Dsouza, A. I. Lazar, J. F. Arteaga, W. M. Nau and U. Pischel, *Chem. Eur. J.*, 2014, **20**, 9897.
157. W. L. Mock, T. A. Irra, J. P. Wepsiec and T. L. Manimaran, *J. Org. Chem.*, 1983, **48**, 3619.
158. T. C. Krasia and J. H. G. Steinke, *Chem. Commun.*, 2002, 22.
159. S. P. Gromov, A. I. Vedernikov, L. G. Kuz'mina, D. V. Kondratuk, S. K. Sazonov, Y. A. Strelenko, M. V. Alfimov and J. A. K. Howard, *Eur. J. Org. Chem.*, 2010, 2587.
160. D. Tuncel, O. Ünal and M. Artar, *Isr. J. Chem.*, 2011, **51**, 525.
161. Q.-S. Wang, H. Cong, J.-X. Zhang, Z. Tao and S.-F. Xue, *Chinese J. Org. Chem.*, 2011, **31**, 1049.
162. Y.-H. Wang, H. Cong, F.-F. Zhao, S.-F. Xue, Z. Tao, Q.-J. Zhu and G. Wei, *Catal. Commun.*, 2011, **12**, 1127.
163. H. Cong, Q.-S. Wang, J.-X. Zhang, Z. Tao and S.-F. Xue, *Chinese J. Org. Chem.*, 2012, **32**, 1093.
164. M. Wieland, J.-L. Mieusset and U. H. Brinker, *Tetrahedron Lett.*, 2012, **53**, 4351.
165. H. Cong, Z.-J. Li, Y.-H. Wang, Z. Tao, T. Yamato, S.-F. Xue and G. Wei, *J. Mol. Catal. A: Chem.*, 2013, **374**, 32.
166. C. Klöck, R. N. Dsouza and W. M. Nau, *Org. Lett.*, 2009, **11**, 2595.
167. R. Breslow, *Acc. Chem. Res.*, 1995, **28**, 146.
168. A. Nakamura and Y. Inoue, *J. Am. Chem. Soc.*, 2003, **125**, 966.
169. C. Yang, G. Fukuhara, A. Nakamura, Y. Origane, K. Fujita, D.-Q. Yuan, T. Mori, T. Wada and Y. Inoue, *J. Photochem. Photobiol. A*, 2005, **173**, 375.
170. A. Nakamura and Y. Inoue, *J. Am. Chem. Soc.*, 2005, **127**, 5338.
171. Q. A. Wang, C. Yang, G. Fukuhara, T. Mori, Y. Liu and Y. Inoue, *Beilstein J. Org. Chem.*, 2011, **7**, 1.
172. D. D. Sternbach and D. M. Rossana, *J. Am. Chem. Soc.*, 1982, **104**, 5853.
173. R. Breslow and S. D. Dong, *Chem. Rev.*, 1998, **98**, 1997.
174. S.-C. Cui, T. Tachikawa, M. Fujitsuka and T. Majima, *J. Phys. Chem. C*, 2011, **115**, 1824.
175. A. L. Koner and W. M. Nau, *Supramol. Chem.*, 2007, **19**, 55.
176. N. Saleh, Y. A. Al-Soud, L. Al-Kaabi, I. Ghosh and W. M. Nau, *Tetrahedron Lett.*, 2011, **52**, 5249.
177. M. Shaikh, J. Mohanty, P. K. Singh, W. M. Nau and H. Pal, *Photochem. Photobiol. Sci.*, 2008, **7**, 408.
178. Y. Huang, Q.-H. Hu, G.-X. Song, Z. Tao, S.-F. Xue, Q.-J. Zhu, Q.-D. Zhou and G. Wei, *RSC Advances*, 2014, **4**, 3348.
179. D. Ma, G. Hettiarachchi, D. Nguyen, B. Zhang, J. B. Wittenberg, P. Y. Zavalij, V. Briken and L. Isaacs, *Nat. Chem.*, 2012, **4**, 503.
180. N. Dong, X.-L. Wang, J.-P. Pan and Z. Tao, *Acta Chim. Sinica*, 2011, **69**, 1431.
181. D. Tuncel, M. Artar and S. B. Hanay, *J. Polym. Sci., Part A: Polym. Chem.*, 2010, **48**, 4894.
182. Y. Huang, S.-F. Xue, Z. Tao, Q.-J. Zhu, H. Zhang, J.-X. Lin and D.-H. Yu, *J. Inclusion Phenom. Macrocyclic Chem.*, 2008, **61**, 171.



183. Z. Miskolczy, M. Megyesi, G. Tárkányi, R. Mizsei and L. Biczók, *Org. Biomol. Chem.*, 2011, **9**, 1061.
184. N. Basilio, L. García-Río, J. A. Moreira and M. Pessêgo, *J. Org. Chem.*, 2010, **75**, 848.
185. R. B. Wang, L. N. Yuan and D. H. Macartney, *Chem. Commun.*, 2006, 2908.
186. L. S. Berbeci, W. Wang and A. E. Kaifer, *Org. Lett.*, 2008, **10**, 3721.
187. H. Cong, C.-R. Li, S.-F. Xue, Z. Tao, Q.-J. Zhu and G. Wei, *Org. Biomol. Chem.*, 2011, **9**, 1041.
188. S. Y. Jon, Y. H. Ko, S. H. Park, H.-J. Kim and K. Kim, *Chem. Commun.*, 2001, 1938.
189. M. Pattabiraman, A. Natarajan, L. S. Kaanumalle and V. Ramamurthy, *Org. Lett.*, 2005, **7**, 529.
190. M. Pattabiraman, L. S. Kaanumalle, A. Natarajan and V. Ramamurthy, *Langmuir*, 2006, **22**, 7605.
191. M. Pattabiraman, A. Natarajan, R. Kaliappan, J. T. Mague and V. Ramamurthy, *Chem. Commun.*, 2005, 4542.
192. M. V. S. N. Maddipatla, L. S. Kaanumalle, A. Natarajan, M. Pattabiraman and V. Ramamurthy, *Langmuir*, 2007, **23**, 7545.
193. H. Yang, Z. Ma, Z. Q. Wang and X. Zhang, *Polym. Chem.*, 2014, **5**, 1471.
194. R. B. Wang, L. N. Yuan and D. H. Macartney, *J. Org. Chem.*, 2006, **71**, 1237.
195. L. Lei, L. Luo, X.-L. Wu, G.-H. Liao, L.-Z. Wu and C.-H. Tung, *Tetrahedron Lett.*, 2008, **49**, 1502.
196. X.-L. Wu, L. Luo, L. Lei, G.-H. Liao, L.-Z. Wu and C.-H. Tung, *J. Org. Chem.*, 2008, **73**, 491.
197. N. Barooah, B. C. Pemberton and J. Sivaguru, *Org. Lett.*, 2008, **10**, 3339.
198. N. Barooah, B. C. Pemberton, A. C. Johnson and J. Sivaguru, *Photochem. Photobiol. Sci.*, 2008, **7**, 1473.
199. B. C. Pemberton, N. Barooah, D. K. Srivatsava and J. Sivaguru, *Chem. Commun.*, 2010, **46**, 225.
200. B. C. Pemberton, R. K. Singh, A. C. Johnson, S. Jockusch, J. P. Da Silva, A. Ugrinov, N. J. Turro, D. K. Srivastava and J. Sivaguru, *Chem. Commun.*, 2011, **47**, 6323.
201. C. Yang, T. Mori, Y. Origane, Y. H. Ko, N. Selvapalam, K. Kim and Y. Inoue, *J. Am. Chem. Soc.*, 2008, **130**, 8574.
202. F. Biedermann, I. Ross and O. A. Scherman, *Polym. Chem.*, 2014, **5**, 5375.
203. H. Cong, F.-F. Zhao, J. X. Zhang, X. Zeng, Z. Tao, S.-F. Xue and Q.-J. Zhu, *Catal. Commun.*, 2009, **11**, 167.
204. S. M. de Lima, J. A. Gómez, V. P. Barros, G. D. Vertuan, M. D. Assis, C. F. D. Graeff and G. J.-F. Demets, *Polyhedron*, 2010, **29**, 3008.
205. X. Y. Lu and E. Masson, *Org. Lett.*, 2010, **12**, 2310.
206. A. L. Koner, C. Márquez, M. H. Dickman and W. M. Nau, *Angew. Chem. Int. Ed.*, 2011, **50**, 545.
207. H. Cong, T. Yamato and Z. Tao, *J. Mol. Catal. A: Chem.*, 2013, **379**, 287.
208. H. Cong, T. Yamato and Z. Tao, *New J. Chem.*, 2013, **37**, 3778.
209. J. Smitka, A. Lemos, M. Porel, S. Jockusch, T. R. Belderrain, E. Tesarova and J. P. Da Silva, *Photochem. Photobiol. Sci.*, 2014, **13**, 310.
210. B. Schulze and U. S. Schubert, *Chem. Soc. Rev.*, 2014, **43**, 2522.
211. J. E. Moses and A. D. Moorhouse, *Chem. Soc. Rev.*, 2007, **36**, 1249.
212. P. Carlqvist and F. Maseras, *Chem. Commun.*, 2007, 748.
213. D. Tuncel and M. Katterle, *Chem. Eur. J.*, 2008, **14**, 4110.
214. D. Tuncel, Ö. Özsar, H. B. Tiftik and B. Salih, *Chem. Commun.*, 2007, 1369.
215. D. Tuncel, N. Cindir and U. Koldemir, *J. Inclusion Phenom. Macrocyclic Chem.*, 2006, **55**, 373.

216. D. Tuncel and J. H. G. Steinke, *Chem. Commun.*, 2002, 496.
217. D. Tuncel and J. H. G. Steinke, *Chem. Commun.*, 1999, 1509.
218. G. Celtek, M. Artar, O. A. Scherman and D. Tuncel, *Chem. Eur. J.*, 2009, **15**, 10360.
219. C.-K. Lin and T.-J. Lu, *Tetrahedron*, 2010, **66**, 9688.
220. K. R. K. Reddy, T. S. Cavallini, G. J. F. Demets and L. F. Silva, *New J. Chem.*, 2014, **38**, 2262.
221. H. S. El-Sheshtawy, B. S. Bassil, K. I. Assaf, U. Kortz and W. M. Nau, *J. Am. Chem. Soc.*, 2012, **134**, 19935.
222. A. Praetorius, D. M. Bailey, T. Schwarzlose and W. M. Nau, *Org. Lett.*, 2008, **10**, 4089.
223. J. Mohanty, A. C. Bhasikuttan, W. M. Nau and H. Pal, *J. Phys. Chem. B*, 2006, **110**, 5132.
224. H.-J. Buschmann and E. Schollmeyer, *J. Inclusion Phenom. Mol. Recognit. Chem.*, 1997, **29**, 167.
225. T. Minami, N. A. Esipenko, B. Zhang, L. Isaacs and P. Anzenbacher. Jr., *Chem. Commun.*, 2014, **50**, 61.
226. T. Minami, N. A. Esipenko, B. Zhang, L. Isaacs, R. Nishiyabu, Y. Kubo and P. Anzenbacher. Jr., *J. Am. Chem. Soc.*, 2012, **134**, 20021.
227. D. Lucas, T. Minami, G. Iannuzzi, L. P. Cao, J. B. Wittenberg, P. Anzenbacher. Jr. and L. Isaacs, *J. Am. Chem. Soc.*, 2011, **133**, 17966.
228. R. Kaliappan, L. S. Kaanumalle and V. Ramamurthy, *Chem. Commun.*, 2005, 4056.
229. R. Kaliappan, Y. H. Ling, A. E. Kaifer and V. Ramamurthy, *Langmuir*, 2009, **25**, 8982.
230. A. Parthasarathy, L. S. Kaanumalle and V. Ramamurthy, *Org. Lett.*, 2007, **9**, 5059.
231. A. Natarajan, L. S. Kaanumalle, S. Jockusch, C. L. D. Gibb, B. C. Gibb, N. J. Turro and V. Ramamurthy, *J. Am. Chem. Soc.*, 2007, **129**, 4132.
232. H. Meier, *Angew. Chem. Int. Ed.*, 1992, **31**, 1399.
233. B. C. Pemberton, E. Kumarasamy, S. Jockusch, D. K. Srivastava and J. Sivaguru, *Can. J. Chem.*, 2011, **89**, 310.
234. Y. Zhao, L.-L. Liang, K. Chen, N.-N. Ji, X.-J. Cheng, X. Xiao, Y.-Q. Zhang, S.-F. Xue, Q.-J. Zhu, N. Dong and Z. Tao, *Dalton Trans.*, 2014, **43**, 929.
235. W. H. Lei, G. Y. Jiang, Q. X. Zhou, Y. J. Hou, B. W. Zhang, X. X. Cheng and X. S. Wang, *ChemPhysChem*, 2013, **14**, 1003.
236. J.-X. Hu, Y.-F. Hu, X. Xiao, Y.-Q. Zhang, Z. Tao, S.-F. Xue, J.-X. Liu and Q.-J. Zhu, *Eur. J. Inorg. Chem.*, 2013, **2013**, 3632.
237. Z.-W. Gao, X. Feng, L. Mu, X.-L. Ni, L.-L. Liang, S.-F. Xue, Z. Tao, X. Zeng, B. E. Chapman, P. W. Kuchel, L. F. Lindoy and G. Wei, *Dalton Trans.*, 2013, **42**, 2608.
238. Y.-F. Hu, K. Chen, J.-X. Liu, R.-L. Lin, W.-Q. Sun, S.-F. Xue, Q.-J. Zhu and Z. Tao, *Polyhedron*, 2012, **31**, 632.
239. K. Chen, X. Feng, L.-L. Liang, Y.-Q. Zhang, Q.-J. Zhu, S.-F. Xue and Z. Tao, *Cryst. Growth Des.*, 2011, **11**, 5712.
240. X. X. Zhang, K. E. Krakowiak, G. P. Xue, J. S. Bradshaw and R. M. Izatt, *Ind. Eng. Chem. Res.*, 2000, **39**, 3516.
241. D. Whang, J. Heo, J. H. Park and K. Kim, *Angew. Chem. Int. Ed.*, 1998, **37**, 78.
242. R. Hoffmann, W. Knoche, C. Fenn and H.-J. Buschmann, *J. Chem. Soc. Faraday Trans.*, 1994, **90**, 1507.
243. C. N. Nguyen, T. K. Young and M. K. Gilson, *J. Chem. Phys.*, 2012, **137**.