

# **PAPER**

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2017, 7, 25753

# 2,3-Disubstituted-1,4-naphthoquinones containing an arylamine with trifluoromethyl group: synthesis, biological evaluation, and computational study†

Hatice Yıldırım, <sup>1</sup> \* Nilüfer Bayrak, <sup>1</sup> \* Amac Fatih Tuyun, <sup>1</sup> \* Emel Mataracı Kara, <sup>c</sup> Berna Özbek Çelik <sup>c</sup> and Girish Kumar Gupta <sup>1</sup> \* Emel Mataracı Kara, <sup>c</sup>

Antibacterial and antifungal organic compounds are becoming increasingly important for biomedical applications. This study deals with the synthesis, characterization of structures, in silico PASS prediction, and the discovery of antibacterial and antifungal properties based on new sulfanyl-1,4-naphthoguinone derivatives containing an arylamine with a trifluoromethyl group at different positions, which can be further applied in drug discovery and development. The in vitro antimicrobial potential of the newly synthesized compounds was evaluated in a panel of seven bacterial strains (three Gram-positive and four Gram-negative bacteria) and one yeast, with an additional study on antibiofilm activities. The compounds (5b and 5e) were identified as having strong antibacterial efficiency against the human-originated pathogen S. epidermidis, with minimal inhibitory concentration values (4.88 and 2.44 μg mL<sup>-1</sup>, respectively). The toxicity of both compounds (5b and 5e) was studied in detail to compare these compounds with Cefuroxime (a clinically proven drug). The antibacterial activity of the compound 5f was equal to that of Cefuroxime. Moreover, three compounds (5b, 5e, and 5f) exhibited excellent antibacterial activity, and 5b and 5e were two and four times more active than the reference antimicrobial compound (Cefuroxime), respectively. For this reason, these three compounds (5b, 5e, and 5f) are being considered as promising antibacterial agents. In addition, docking studies were used to better rationalize the action and prediction of the binding modes of these compounds.

Received 20th January 2017 Accepted 18th April 2017

DOI: 10.1039/c7ra00868f

rsc.li/rsc-advances

# Introduction

Although there has been a decline in the incidence of infectious diseases, there is no place in the world where infectious diseases are an insignificant cause of illness and death.<sup>1,2</sup> The number of deaths caused by pathogens and parasites have shown a minor decline of only 1% per year. Over 10 million people have died from infections since 1990.<sup>1a,2</sup> The World Health Organization (WHO) estimates 13 million deaths from infectious diseases in 2050.<sup>2</sup> People in low- and middle-income countries suffer from diseases such as HIV/AIDS, tuberculosis.

Infectious diseases are among the most serious threats of the "top global health risk factors", just like the high risk factors of chemical spills and nuclear accidents. <sup>1b,c,2,3</sup> WHO has, therefore, provided effective global strategies for the control of major infectious diseases that are being dubbed as the "infectious diseases control programs", which are necessary in order to help build these health systems.<sup>4</sup>

The fact that the resistance of bacterial and fungal pathogens leads to an ever-increasing global public health threat reflects the widespread and growing demand for new antibacterial and antifungal compounds, which are among the most prevalently used drugs. Unfortunately, very few new classes of antibacterial drugs have been registered for clinical practice in the past 50 years. Not surprisingly, this has recently caused a rapidly growing interest in the synthesis of new and efficient antibacterial and antifungal compounds that may be drug candidates for reducing infections caused by pathogens.

The antimicrobial activities of an increasing number of quinone compounds are being investigated for the discovery of new antibacterial and antifungal compounds.<sup>7,8</sup> A series of sulfanyl-1,4-naphthoquinone derivatives was synthesized and potent antifungal activity was exhibited *in vitro* by one

malaria, other pathogens, and infections that have been causing the majority of deaths since 2010. 14,2

<sup>&</sup>lt;sup>a</sup>Chemistry Department, Engineering Faculty, Istanbul University, Avcilar, 34320, Istanbul, Turkey. E-mail: hyildirim@istanbul.edu.tr; Fax: +90 212 473 7180; Tel: +90 212 473 7070

<sup>&</sup>lt;sup>b</sup>Engineering Sciences Department, Engineering Faculty, Istanbul University, Avcilar, 34320, Istanbul, Turkey. E-mail: aftuyun@gmail.com; aftuyun@istanbul.edu.tr; Fax: +90 212 473 7180: Tel: +90 212 473 7070

Pharmaceutical Microbiology Department, Pharmacy Faculty, Istanbul University, Beyazit, 34116, Istanbul, Turkey

<sup>&</sup>lt;sup>4</sup>Department of Pharmaceutical Chemistry, Maharishi Markandeshwar College of Pharmacy, Maharishi Markandeshwar University, Mullana, Ambala 133207, Haryana, India

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ra00868f

compound, compared to the clinically proven antifungal drug fluconazole, against S. schenckii, and the clinically proven drug amphotericin-B against T. mentagraphytes.9 1,4-Naphthoquinones containing an amino group have been used as antibacterials, antimalarials, antituberculars, larvicides and molluscicides, herbicides, and fungicides in many medical and biological applications.10-14 A series of phenylamino-1,4naphthoquinones was synthesized to investigate their antitumor effects against cancer cell lines and healthy fibroblasts, and it was reported that the presence of a chlorine atom in the acceptor quinone nucleus and/or the presence of a methyl group at the nitrogen atom of the donor phenylamino group induced changes in cytotoxic activity.15

Since the trifluoromethyl group (-CF<sub>3</sub>) is one of the most important substituents in organic chemistry because of its interesting stereoelectronic profile,16 our attention was turned to the naphthoquinone compounds containing the -CF<sub>3</sub> on the arylamines at the o-, m-, and p-positions, in light of previous studies. 17 The -CF3 has an effect based on the different electron density distributions on the reactivity of the molecule. The efficiency of -CF3 on the physiological activity is a very significant topic in pharmaceutical studies. The physiological profile is dependent on the position of -CF3 within the bioactive molecules.16

A series of naphthoguinone derivatives has been synthesized and tested for their biological activity against human African trypanosomiasis, normally known as sleeping sickness, a dangerous and often ignored parasitic disease. 17 The activities of some compounds have been compared to evaluate the importance of the position of the -CF3 group on the phenyl amine ring and its substitution that caused by the inhibition of T. brucei cell proliferation. Compounds shown in Fig. 1 with -CF<sub>3</sub> (electron withdrawing group) at different positions on the phenyl amine ring showed T. brucei inhibitory activity with average cytotoxicity. It was found that the presence of a chlorine atom instead of a second arylamine led to an increase in the activity. Additionally, changing the -CF<sub>3</sub> from the para position to the ortho position led to an increase in the activity. It was reported that the reason for this enhancement in T. brucei inhibitory activity might have been the strong electron withdrawing groups (-CF<sub>3</sub>, -NO<sub>2</sub>, and Cl) within the structure; replacing the -CF<sub>3</sub> with the methyl group at different positions led to activity loss. In this study, it can be clearly seen that the substitution patterns (2<sup>nd</sup> and 4<sup>th</sup> positions) on aniline and monoamino- or diamino-substituted compounds affect the T. brucei inhibitory activity.17

Sulfanyl aminonaphthoquinone and aminonaphthoquinone derivatives have important applications in medicinal chemistry as biologically active substances. Among them, some most striking examples are shown in Fig. 2. Tandon et al. reported the synthesis of some sulfanyl aminonaphthoquinone compounds (I-IV) and their biological evaluation. Some of these compounds showed significant antibacterial activity.7c,d Very recently, another research study on the synthesis of the sulfanyl arylamino quinone compounds was reported by Ryu et al. Some of the tested compounds (V and VI) completely inhibited fungal growth when compared to the Candida species.8a Tandon et al.

Nitro substituted aminonaphthoguinones containing -CF<sub>2</sub> group Nitro free aminonaphthoquinones containing -CF<sub>3</sub> group

Fig. 1 Structures of some nitro free and nitro substituted aminonaphthoquinones.17




Fig. 2 Structures of some important sulfanyl aminonaphthoguinones with biological activity in literature.

synthesized other series of sulfanyl aminonaphthoquinone and aminonaphthoquinone derivatives, and the synthesized compounds were tested for antibacterial and antifungal activities. Particularly, sulfanyl aminonaphthoquinone derivatives containing the arylsulfanyl group did not exhibit any significant activity (VII). In this study, the antifungal activity was observed to decrease when the 2,3-disubstituted moiety was altered by the phenyl group.8b

There are several factors that affect the biological evaluation of the quinone structures, which are listed as follows: (a) a naphthoquinoidal moiety; (b) an aromatic ring such as aniline; (c) the presence and position of substituents on the phenyl amine ring, and insertion of additional electronegative atoms (Cl, S, N etc.). 15,17,18 Encouraged by all these facts, and in Paper RSC Advances

Fig. 3 Structures of the sulfanyl derivatives of 1,4-naphthoquinone based on aryl amines containing -CF<sub>3</sub> group.

order to continue our previous studies, which were mainly about the synthesis of biologically active 1,4-naphthoquinone structures, we chose our compounds by (i) using sulfanyl derivatives of arylamino-1,4-naphthoquinone as the core structure; (ii) using arylamine ring containing the  $CF_3$  group at different positions (o-, m-, and p-); (iii) using different types of thiols, as shown in Fig. 3. From this point of view, profound pharmacological effects of related compounds in previous studies gave us clues regarding the synthesis, characterization, and evaluation of some new sulfanyl-1,4-naphthoquinone derivatives containing an arylamine with the  $-CF_3$  at different positions, as antimicrobial agents.

# 2. Results and discussion

## 2.1. The synthesis of substituted naphthoquinones

Herein, the synthesis of 3-arylamino-1,4-naphthoquinone derivatives (3a-c) was carried out via the reactions of 2,3dichloro-1,4-naphthoquinone (1) and primary arylamines (2a-c) containing an electron withdrawing substituent (-CF<sub>3</sub>) on the rings at the o-, m-, and p-positions as the starting materials in different mediums. 17,19,27 Unlike the synthesis of 3a and 3b reported previously in the scientific literature, compound 3c was obtained in low yields in DMF in the presence of K<sub>2</sub>CO<sub>3</sub> (10%). The synthesis of new sulfanyl 1,4-naphthoquinone derivatives containing an arylamine with the -CF<sub>3</sub> at different positions (5a-k) was achieved at room temperature via the reaction of 3arylamino-1,4-naphthoquinone derivatives (3a-c) with a variety of thiols (4a-d) in CH<sub>2</sub>Cl<sub>2</sub> in the presence of Et<sub>3</sub>N (1.1 eq.) as a base (shown in Scheme 1) in high yields.8 Additionally, compound 6 was obtained as a side-product from the reaction of 2-chloro-3-((4-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3a) with butane-2-thiol (4a). Among the synthesized compounds, the sulfanyl 1,4-naphthoquinone derivatives (5a-k) and the side product (6) are new, while the arylamino-1,4naphthoquinone derivatives (3a-c) have been reported in earlier studies. 17,19,27 New sulfanyl 1,4-naphthoquinone derivatives containing an arylamine with the -CF<sub>3</sub> at different positions (5a-k) were obtained in the range of 30-85% yield as orange and red powders or oils. Spectroscopic methods like one dimensional NMR techniques (1H, 13C), FTIR spectra and mass spectrometry were applied to confirm the structures of obtained compounds. The FTIR spectra of 5a-k and 6 showed some

characteristic absorptions around 3275–3346 cm<sup>-1</sup> (NH), 1657–1676 cm<sup>-1</sup> (C=O), 3419–3433 (OH) cm<sup>-1</sup>, 2851–2978 cm<sup>-1</sup> (CH<sub>aliphatic</sub>) and 3024–3076 cm<sup>-1</sup> (CH<sub>aromatic</sub>). The <sup>1</sup>H NMR spectra of **5a-k** and **6** (CDCl<sub>3</sub>-d<sub>1</sub>) exhibited the NH signal as a singlet between 7.64–7.93 ppm in the aromatic region and at 5.86–8.09 ppm corresponding to the aromatic protons. The signals associated with the other functional groups all appeared in the expected regions. Outstandingly, the presence of two peaks for the carbonyl groups (–C=O) in <sup>13</sup>C NMR spectra was the proof of the two substituents (amino and sulfanyl groups) attached to the naphthoquinone core. There was only one peak for the carbonyl group in the <sup>13</sup>C NMR spectrum of compound **6** because of the symmetry.

#### 2.2. The prediction of biological activity based on spectra

The prediction of biological activity based on the spectral analysis of the synthesized compounds in the present communication was easily carried out by using the freely available website with the internet version of PASS and PharmaExpert: http://www.pharmaexpert.ru/PASSOnline/. The activity spectrum for a molecule is a list of biological activity for which the probability of the compound being active  $(P_a)$  and the probability of the compound being inactive  $(P_i)$  are calculated.  $P_a$  and  $P_i$  values are independent and their values vary from 0 to 1. For example, the PASS prediction results for one of the compound (5e) are summarized in Table 1.

It was found that for antibacterial and anti-infective activity, the compounds had  $P_a$  nearly equal to 0.2. These findings also showed some good experimental results and can be considered to be valuable while planning the experiment, but some important factors should be taken into account such as the particular interest in some kind of activity, desirable novelty, the available facilities for experimental testing, *etc.* 

# 2.3. The antimicrobial activity

All the newly synthesized sulfanyl 1,4-naphthoquinone derivatives containing an arylamine with the –CF<sub>3</sub> at different positions (5a–k) and 2,3-disulfanyl 1,4-naphthoquinone (6) were evaluated for their *in vitro* antibacterial and antifungal activities. The antimicrobial assay results for all the newly synthesized 1,4-naphthoquinone structures (5a–k and 6) are given in Table 2.

*In vitro* activities of the studied molecules against planktonic cells are summarized in Table 2. Concerning the antibacterial activity, the results revealed that some compounds displayed varying effects on the growth of the tested Gram-positive and Gram-negative bacterial strains. As indicated in Table 2, among the synthesized compounds of **5b-g** and **6**, they generally showed potent antibacterial activity against some pathogenic strains that were tested. Actually, the activities of compounds **5b** and **5e** were superior as compared to that of Cefuroxime: **5b** and **5e** completely inhibited the growth of *S. epidermidis*, and were tested at the MIC level of 4.88 and 2.44 µg mL<sup>-1</sup>, respectively. In addition to these, the activity of compound **5f** was equal to that of Cefuroxime. In general, most of the tested compounds showed better antibacterial activity against the Gram-positive

**RSC Advances** Paper

Scheme 1 Synthetic pathway for the preparation of new sulfanyl 1,4-naphthoguinone derivatives substituted with arylamines containing the  $-CF_3$  at o-, m-, and p-positions.

bacteria compared to the Gram-negative ones. The results revealed that compounds 5d and 5g exhibited moderate activity against Gram-positive bacteria. The test-cultures P. aeruginosa, E. coli, and S. aureus appeared non-effective to most of the synthesized compounds. Some of the synthesized compounds (5d-e, 5g and 5j) possessed activity against E. faecalis, which had MIC values between 312.5-625  $\mu g \text{ mL}^{-1}$ . In addition to E. faecalis, the compounds (5d and 5g) possessed activity against S. epidermidis, which had the MIC value of 156.2  $\mu g \text{ mL}^{-1}$ . 2,3-Disulfanyl 1,4-naphthoquinone (6) was only able to induce appreciable growth inhibitory activity against S. epidermidis, which had the MIC value of 78.12 μg mL<sup>-1</sup>. The compounds

(5a-b, 5e-f, and 5h-k) exhibited no antibacterial activity against the Gram-negative bacteria. Regarding the antifungal activity of the tested compounds, only 2 compounds (5j and 5k) were able to induce appreciable growth inhibitory activity against C. albicans.

To evaluate the importance of the position of the -CF<sub>3</sub> on the phenyl amine ring with respect to the biological efficiency, the activities of all the synthesized compounds were compared (Scheme 1). The decrease in the antibacterial activities of the compounds with the -CF<sub>3</sub> at the p-positions on the phenylamine ring, (5a, 5d, 5g, and 5j) was the proof of an important factor for the antibacterial activity. In terms of the structure-

Table 1 PASS prediction results for compound (5e)

$P_{\rm a}$	$P_{\mathrm{i}}$	Activity
0.625	0.042	TP53 expression enhancer
0.589	0.094	Membrane permeability inhibitor
0.426	0.027	Pyruvate decarboxylase inhibitor
0.336	0.001	CDC25 phosphatase inhibitor
0.285	0.201	Aspartyltransferase inhibitor
0.233	0.173	Anti-infective
0.173	0.070	Protein synthesis inhibitor
0.179	0.144	Phosphatidylcholine-sterol O-acyltransferase inhibitor
0.158	0.045	Antibacterial, ophthalmic
0.138	0.025	Antibiotic anthracycline-like

activity relationship, the sulfanyl 1,4-naphthoguinone derivatives containing an arylamine with the -CF3 at the m-positions (5b, 5e, 5h, and 5k) showed, in general, more potent antibacterial activities than the other sulfanyl 1,4-naphthoquinone derivatives with an arylamine containing the -CF<sub>3</sub> at the p-positions (5a, 5d, 5g, and 5j). The sulfanyl 1,4-naphthoquinone derivatives containing an arylamine with the -CF<sub>3</sub> at the o-positions (5c, 5f, and 5i) showed, in general, more potent antibacterial activity than other sulfanyl 1,4-naphthoquinone derivatives with an arylamine containing the -CF<sub>3</sub> at the p-position, but unfortunately, not better than other sulfanyl 1,4-naphthoquinone derivatives with an arylamine containing the -CF3 at the m-positions. The replacement of a halogen with the different types of sulfanyl derivatives led to variable types of activities. However, aliphatic substitutions like sec-butylthio and 2-hydroxypropylthio moieties of the compounds (5b-f) in comparison to aralkyl or heteroaryl substitutions (5g, 5h, and 5k) may contribute partially toward biological potency with the additional effect of the position of -CF<sub>3</sub>. Moreover, this particular modification does not significantly affect the antifungal activity. Indeed, the aralkyl derivatives (5j and 5k) showed moderate inhibition effect against C. albicans; this leads to the conclusion that the moiety may be a factor affecting the antimicrobial activity of the synthesized sulfanyl 1,4-naphthoquinone derivatives. In this study, when the in vitro antimicrobial activities of all the newly synthesized sulfanyl 1,4-naphthoquinone derivatives were investigated, three molecules (5b, 5e, and 5f) had strong antibacterial efficiencies against S. epidermidis, which is usually considered as a common human originated pathogen.

The cytotoxic effects of the two compounds (5b and 5e) were obtained as the IC<sub>50</sub> values, which represent the molar drug concentrations required to cause 50% inhibition of cell viability. Consequently, all tested compounds significantly induced dose-dependent loss of viability in HepG2 cells after 24 h. The  $IC_{50}$  values (5b: 19.54 µg mL<sup>-1</sup>, 5e: 10.18 µg mL<sup>-1</sup>) for the compounds were four times higher than the MIC results shown in Fig. 4 and 5.

#### 2.4. Antibiofilm activity

A biofilm is a community of microbes embedded in a selfproduced polymeric matrix, which adheres to a surface.

		Microorganisms						
		Gram-negative bacteria	acteria			Gram-positive bacteria	eria	
		P. aeruginosa	E. coli	K. pneumoniae	P. mirabilis	S. aureus	S. epidermidis E. faecalis	E. faecalis
MIC values	5a	1	I	I	I	1	I	I
$(\mu \mathrm{g \ mL}^{-1})$	5 <b>b</b>	1	1	1	1	I	4.88	1
	5c	I	625	312.5	I	I	1250	I
	2 <b>d</b>	1	I	I	312.5	I	156.2	312.5
	5e	I	I	I	I	I	2.44	625
	5f	I	I	I	I	I	9.76	I
	5g	1	I	625	625	625	156.2	625
	5 <b>h</b>	I	I	I	I	I	I	I
	5i	1	I	I	I	I	1250	1
	J.	1	I	I	I	I	I	625
	5k	I	I	I	I	I	I	I
	9	I	I	I	I	I	78.12	I
	Reference	2.4	4.9	4.9	2.4	1.2	8.6	128
	antimicrobials Ceftazidime	Ceftazidime	Cefuroxime-Na	Cefuroxime-Na	Cefuroxime-Na	Cefuroxime-Na	Cefuroxime	Amikacin

C. albicans

Fungi

compounds

synthesized

for the

antifungal activity results

and

vitro antibacterial

7

~

RSC Advances Paper




Fig. 4 The effect of the compound **5b** on the % inhibition at different concentrations.




Fig. 5 The effect of the compound **5e** on the % inhibition at different concentrations.

Microbial cells that grow in biofilms behave quite differently from their planktonic counterparts. The bacteria on the biofilm structures are protected from the environmental conditions, antimicrobial agents, and host immune responses, and the biofilm organisms are far more resistant to antimicrobial agents than the organisms in the suspension (up to 1000-fold increase).<sup>22</sup>

When we considered the antibiofilm activities of the antibacterial molecules  $5\mathbf{b}$  and  $5\mathbf{e}$  against the *S. epidermidis* biofilms, the MBEC values were 5000 and 1250  $\mu g$  mL<sup>-1</sup>, respectively. The MBEC/MIC ratio, which is one of the important parameters for choosing antibiotics in the treatment of biofilm associated infections, was found to be 1024 and 512 fold, respectively, in contrast to many antibiotics (up to 10 000 fold).<sup>22a</sup>

Since the only biofilm susceptible to the studied active molecules was *S. epidermidis*, both the biofilm attachment and inhibition of the biofilm formation assays were performed. When these tests were carried out, **5b** inhibited the biofilm attachment according to the time, and it showed a significant inhibition activity against the biofilm formation at the 24<sup>th</sup> hour according to its varying concentrations (Fig. 6). In contrast to these findings, there was an inhibition in the biofilm attachment of **5e** at the end of the 2<sup>nd</sup> hour; following that, as the number of cells increased within the environment, this activity was decreased at the 4<sup>th</sup> hour.

The biofilms are not affected by "therapeutically achievable concentrations" of antimicrobial agents. The antibiofilm therapies are generally focused on the inhibition of the biofilm formation. For this purpose, the inhibitions of bacterial attachment to the surfaces were investigated; the inhibition of the biofilm production at the MIC or subMIC values of molecules were investigated as well. **5b** and **5e** were able to inhibit the attachment of bacteria at the MIC or subMIC values, and the biofilm formation after 24 hours was approximately 50% (at  $1/10 \times MIC\ p < 0.001$ ). Since inhibiting a mature biofilm can be very difficult, the inhibition of the biofilm formation in the early stages may seem to be more applicable and advantageous.

# 2.5. Docking studies

*In silico* approaches like docking studies are currently gaining much attention to pre-access the probable mechanism for the action and design of new templates towards more potential antibacterial agents.<sup>21</sup> The significant experimental




Fig. 6 The inhibition of *S. epidermidis*. (A) The surface attachment to the wells containing  $1/10 \times$  MIC of molecules and an inoculum of  $1 \times 10^7$  cfu per 200  $\mu$ L, incubated for 1, 2, or 4 h at 37 °C. (B) The biofilm formation in each well containing  $1 \times 1/10 \times$ , or  $1/100 \times$  MIC of molecules and an inoculum of  $5 \times 10^5$  cfu per 200  $\mu$ L, incubated for 24 h at 37 °C. The control bars indicating bacteria without the molecule of interest are accepted as 100%. Six wells were used for each molecule. Each experiment is representative of the two independent tests, and the error bars indicate the standard deviations. All the differences between the control and molecule-treated biofilms are statistically significant (p < 0.001).

Paper

antibacterial activities of the compounds 5b and 5e gave us a hint to perform molecular docking studies to understand the protein-ligand interactions. The docking was performed via the Molegro Virtual Docker 2010.4.1.0 program<sup>23</sup> for the active site of Staphylococcus epidermidis mevalonate diphosphate decarboxylase (PDB ID: 4DPT) in order to predict the binding mode and support the biological results.24

Mevalonate diphosphate decarboxylase (MDD) catalyzed the formation of isopentenyl 5-diphosphate in an ATP-dependent irreversible reaction and was therefore an attractive target for the inhibitor development that could lead to the new antimicrobial agents.24b In the case of compound 5b, it revealed the MolDock score of -72.77 and formed three interactions, shown as green dotted lines (Fig. 7), which were expressed as hydrogen bonds formed with the oxygen atom of the C=O (naphthoquinone) moiety at the position 4 with Arg 144 and Ala 14, with distances of 2.59 Å and 3.19 Å, respectively and one hydrogen bond formed between the -NH- moiety at the position of 3 and Asn 12 of distance 3.35 Å.

In the case of the compound 5e, it revealed the MolDock score of -78.46 and formed the main interactions as depicted in Fig. 8, expressed as one hydrogen bond formed with the oxygen atom of the C=O (naphthoquinone) moiety at the position 4, and bonds with the oxygen atom of the 2-hydroxypropylthio group with Ala 284 and Ser 141, with distances of 3.14 Å and 2.93 Å, respectively.

Compound 6 revealed the MolDock score of −56.46. The supposed Moldock score was found to be positively correlated with the experimental result values, where the antibacterial activities against S. epidermidis of the compounds of 5e, 5b and 6 decrease from left to right. Interactions with the Arg 144




Fig. 7 The binding mode of compound 5b in S. epidermidis mevalonate diphosphate decarboxylase [PDB ID: 4DPT].




Fig. 8 The binding mode of compound 5e in S. epidermidis mevalonate diphosphate decarboxylase [PDB ID: 4DPT].

(catalytic site residue, ligand to mevalonate C1 carboxyl) and Ser 141 (ATP binding motif residue) polar active side chains indicate the binding affinity of the compounds with the protein. This could be further investigated under the topic of MDD inhibitors.25

#### 3. Conclusion

The sulfanyl 1,4-naphthoquinone derivatives, containing an arylamine with the -CF<sub>3</sub> at the different positions (5a-k), and 2,3-disulfanyl 1,4-naphthoquinone (6) were synthesized via the nucleophilic substitution of aminonaphthoquinones (3a-c) with 1.1 eq. of aryl- and alkylthiols (4a-d) in CH<sub>2</sub>Cl<sub>2</sub> in the presence of Et<sub>3</sub>N (1.1 eq.) as a base at the room temperature.8 All of the newly synthesized compounds (5a-k and 6) were tested for their antimicrobial activities. Based on the testing data for the presented compounds, the *in vitro* antimicrobial activities were evaluated against the different Gram-positive and Gramnegative bacterial strains in addition to the antifungal activities. The antibacterial profile of the synthesized compounds (5a-k) indicated that compounds (5b-k) had potent antibacterial activities. Among the most promising antibacterial compounds, 5b and 5e exhibited better antibacterial activity than clinically prevalent antibacterial drug Cefuroxime against S. epidermidis, while 5f had similar activity. The compounds (5b, 5e, and 5f) are therefore lead compounds as potent antibacterial agents for further studies. The tested compounds (5b and 5e) significantly induced dose-dependent loss of viability in HepG2 cells after 24 h. The  $IC_{50}$  values (5b: 19.54 µg mL<sup>-1</sup>, 5e: 10.18 µg  $mL^{-1}$ ) for the compounds were four times higher, compared to the MIC results. These results suggest that the sec-butylthio and 2-hydroxypropylthio moieties with the additional effect of the position of -CF3 are promising leads for the development of antibacterial agents. 5b and 5e were able to inhibit the attachment of bacteria at the MIC or subMIC values and the biofilm formation after 24 hours was approximately 50%, at  $1/10 \times$  MIC (p < 0.001). The molecular docking studies mentioned above further helped in supporting the experimental results. These experimental findings and structures with the additional efforts will pave the way for more potent biologically active derivatives containing the quinone chromophore.

#### Experimental 4.

## Chemistry

All of the reagents were obtained from the commercial suppliers and were used without further purification unless otherwise noted. Petroleum ether had a boiling range of 40-60 °C. Analytical thin layer chromatography (TLC) plates were purchased from Merck KGaA (silica gel 60 F254) based on Merck DC-plates (aluminum based). Visualization of the chromatogram was performed by UV light (254 nm). Column chromatographic separations were carried out using silica gel 60 (Merck, 63-200 μm particle size, 60-230 mesh). <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded with the VarianUNITY INOVA spectrometers with 500 MHz frequency for <sup>1</sup>H and 125 MHz frequency for  $^{13}$ C NMR in ppm ( $\delta$ ).  $^{1}$ H NMR spectra and  $^{13}$ C NMR spectra in

 $CDCl_3$  were referenced to the solvent signals centered at  $\delta$  7.19 and  $\delta$  76.0 ppm, respectively. Standard abbreviations indicating multiplicity were used as follows: s (singlet), br s (broad singlet), d (doublet), t (triplet), and m (multiplet). Coupling constants I were given in Hz. FTIR spectra were recorded as ATR on either a Thermo Scientific Nicolet 6700 spectrometer or Alpha T FTIR spectrometer. Mass spectra were obtained on either a Thermo Finnigan LCQ Advantage MAX MS/MS spectrometer equipped with ESI (electrospray ionization) sources. The purity of the sulfanyl 1,4-naphthoquinone derivatives was obtained by HPLC analysis. HPLC was performed on a Shimadzu/DGU-20A5 HPLC apparatus fitted with a 25 cm Chiralpac AD-H chiral column. The melting points (mp) were determined with a Buchi B-540 melting point apparatus, but were uncorrected. The 3arylamino-1,4-naphthoguinone derivatives prepared via the reaction between the 2,3-dichloro-1,4naphthoquinone (1) and trifluoromethyl substituted aryl amines (2a-c) according to the literature reported previously and the references cited therein. 17,19,26,27 All of the synthesized sulfanyl 1,4-naphthoquinone derivatives (5a-k) and side product (6) are new.

**4.1.1.** General procedure for synthesis of the sulfanyl 1,4-naphthoquinone derivatives (5a–k). The 3-arylamino-1,4-naphthoquinone derivatives (3a–c) and thiols in  $CH_2Cl_2$  were stirred at room temperature using  $Et_3N$ . The resulting solution was extracted with 100 mL chloroform, and then washed with water (4  $\times$  100 mL) and dried over calcium chloride. The solvent was removed *in vacuo*. The residue was subjected to column chromatography on silica gel using suitable solvents to give the products.

4.1.1.1. 2-(Sec-butylthio)-3-(4-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5a). This compound was synthesized from 2-chloro-3-((4-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3a) and butane-2-thiol (4a) as a red oil by using the general procedure. Yield: 0.030 g, 52%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3323 (-NH), 3030 (CH<sub>arom</sub>), 2967, 2929, 2875 (CH<sub>aliphatic</sub>), 1659 (C=O). <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.10–8.08 dd, *J: 7.81*, 0.98 Hz, 1H (-CH<sub>arom</sub>); 8.03-8.01 dd, J: 7.81, 0.98 Hz, 1H (-CH<sub>arom</sub>); 7.75 bs, 1H (-NH); 7.70-7.66 td, J: 7.33, 1.47 Hz, 1H (-CH<sub>arom</sub>); 7.63-7.60 td, *J:* 7.81, 1.47 Hz, 1H (-CH<sub>arom</sub>); 7.51-7.49 d, *J:* 8.30 Hz, 2H (-CH<sub>arom</sub>); 6.99-6.97 d, J: 8.30 Hz, 2H (-CH<sub>arom</sub>); 3.02-2.98 q, J: 6.83 Hz, 1H (-CH-); 1.38-1.28 m, 2H (-CH<sub>2</sub>-); 0.99-0.97 d, J: 6.83 Hz, 3H (-CH<sub>3</sub>); 0.71-0.68 t, J: 7.32 Hz, 3H (-CH<sub>3</sub>). <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 180.2, 179.1 (C=O), 143.4, 140.5 (C<sub>a</sub>), 133.6, 132.1, 126.0, 125.7, 120.5 (CH<sub>arom</sub>), 132.2, 129.7, 124.7 (C<sub>q</sub>), 43.3 (-CH-); 28.8 (-CH<sub>2</sub>-); 19.4, 10.1 (-CH<sub>3</sub>). MS (ESI-) m/z (%): 402 (100,  $[M - 3H]^+$ ), 403 (22,  $[M - 2H]^+$ ). Anal. calcd for  $C_{21}H_{18}F_3NO_2S$  (405.43).

Additionally, **6** was also obtained from 2-chloro-3-((4-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (**3a**) and butane-2-thiol (**4a**) by using the general procedure.

4.1.1.1.1 2,3-Bis(sec-butylthio)naphthalene-1,4-dione (6). Red oil, yield: 0.035 g, 37%. FTIR (ATR)  $\nu$  (cm $^{-1}$ ): 2967, 2925, 2875 (-CH<sub>aliphatic</sub>), 1661 (C=O).  $^{1}$ H NMR (500 MHz,  $CDCl_{3}$ )  $\delta$  (ppm): 7.99–7.98 dd, J: 5.37, 2.93 Hz, 2H (-CH<sub>arom</sub>); 7.62–7.61 dd, J: 5.86, 2.93 Hz, 2H (-CH<sub>arom</sub>); 4.04–4.00 q, J: 6.83 Hz, 2H (-CH-);

1.61–1.48 m, 4H (–CH<sub>2</sub>–); 1.24–1.22 dd, *J*: 6.84, 2.93 *Hz*, 6H (–CH<sub>3</sub>); 0.96–0.93 td, *J*: 7.32, 2.44 *Hz*, 6H, (–CH<sub>3</sub>). <sup>13</sup>C NMR (125 MHz, *CDCl*<sub>3</sub>)  $\delta$  (ppm): 178.3 (–C=O); 148.1 (C<sub>q</sub>); 132.4, 125.9 (–CH<sub>arom</sub>); 132.1 (C<sub>q</sub>); 44.5 (–CH–); 29.8 (–CH<sub>2</sub>–); 20.2, 20.1, 10.4, 10.3 (–CH<sub>3</sub>). MS (ESI+) m/z (%): 334 (100, [M]<sup>+</sup>), 335 (71, [M + H]<sup>+</sup>). Anal. calcd for C<sub>18</sub>H<sub>22</sub>O<sub>2</sub>S<sub>2</sub> (334.50).

4.1.1.2. 2-(Sec-butylthio)-3-(3-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5b). This compound was synthesized from 2-chloro-3-((3-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3b) and butane-2-thiol (4a) as a red powder by using the general procedure. Yield: 0.045 g, mp: 109-110 °C, 39%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3275 (-NH), 3076 (-CH<sub>arom</sub>), 2967, 2929 (-CH<sub>aliphatic</sub>), 1662 (-C=O). <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.09-8.08 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 8.01-8.00 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.78 s, 1H (-NH); 7.69-7.65 td, J: 7.32, 0.97 Hz, 1H (-CH<sub>arom</sub>); 7.62-7.59 t, *J. 7.32 Hz*, 2H, (-CH<sub>arom</sub>); 7.38-7.35 t, *J.* 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.32-7.30 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.09-7.08 d, J: 7.81 Hz, 1H, (-CH<sub>arom</sub>); 2.98-2.93 m, 1H (-CH-); 1.36-1.23 m, 2H (-CH<sub>2</sub>-); 0.97-0.95 d, J: 6.83 Hz, 3H (-CH<sub>3</sub>); 0.69-0.66 t, J: 7.32 Hz, 3H (-CH<sub>3</sub>).  $^{13}$ C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  (ppm): 180.2, 179.1 (-C=O); 143.7, 137.8, 132.3, 129.7, 127.8, 124.4, 119.9, 118.9, 118.1 (C<sub>q</sub>); 133.6, 132.0, 126.0, 125.7 (-CH<sub>arom</sub>); 43.2 (-CH-); 28.7 (-CH<sub>2</sub>-); 19.3, 10.1 (-CH<sub>3</sub>). MS (ESI+) m/z (%): 406  $(100, [M + H]^{+})$ . Anal. calcd for  $C_{21}H_{18}F_{3}NO_{2}S$  (405.43).

4.1.1.3. 2-(Sec-butylthio)-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (5c). This compound was synthesized from 2-chloro-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3c) and butane-2-thiol (4a) as a red oil by using the general procedure. Yield: 0.039 g, 70%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3328 (-NH), 3072 (-CH<sub>arom</sub>), 2963, 2922, 2872 (-CH<sub>aliphatic</sub>), 1664 (-C= O). <sup>1</sup>H NMR (500 MHz, *CDCl*<sub>3</sub>) δ (ppm): 8.08–8.07 dd, *J: 7.32*, 1.46 Hz, 1H (-CH<sub>arom</sub>); 8.00-7.98 dd, J: 7.32, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.91 s, 1H (-NH); 7.68-7.65 td, J: 7.81, 0.98 Hz, 1H (-CH<sub>arom</sub>); 7.61-7.58 m, 2H, (-CH<sub>arom</sub>); 7.43-7.40 t, *J: 7.81 Hz*, 1H (-CH<sub>arom</sub>); 7.19-7.16 t, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 6.90-6.88 d, J: 8.30 Hz, 1H, (-CH<sub>arom</sub>); 3.07-3.02 m, 1H (-CH-); 1.42-1.26 m, 2H (-CH<sub>2</sub>-); 0.99-0.98 d, J: 6.34 Hz, 3H (-CH<sub>3</sub>); 0.73-0.70 t, J: 7.32 Hz, 3H  $(-CH_3)$ . <sup>13</sup>C NMR (125 MHz, *CDCl*<sub>3</sub>)  $\delta$  (ppm): 180.1, 178.8 (-C=O); 144.9, 132.2, 130.8, 129.8, 125.3, 118.9 (C<sub>a</sub>); 133.5, 125.9, 125.7, 124.0, 123.4 (-CH<sub>arom</sub>); 43.3 (-CH-); 28.9 (-CH<sub>2</sub>-); 19.3, 10.0  $(-CH_3)$ . MS (ESI+) m/z (%): 406 (100,  $[M + H]^+$ ). Anal. calcd for C<sub>21</sub>H<sub>18</sub>F<sub>3</sub>NO<sub>2</sub>S (405.43).

4.1.1.4. 2-(2-Hydroxypropylthio)-3-(4-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5d). This compound was synthesized from 2-chloro-3-((4-(trifluoromethyl)phenyl)amino) naphthalene-1,4-dione (3a) and 3-mercaptobutan-2-ol (4b) as an orange powder by using the general procedure. Yield: 0.085 g, mp: 139–140 °C, 73%. FTIR (ATR)  $\nu$  (cm $^{-1}$ ): 3419 (–OH), 3336 (–NH), 3030 (–CH $_{\rm arom}$ ), 2978, 2961, 2929 (–CH $_{\rm aliphatic}$ ), 1669 (–C=O).  $^{1}$ H NMR (500 MHz, *CDCl*<sub>3</sub>)  $\delta$  (ppm): 8.10–8.08 dd, *J:* 7.81, 1.46 Hz, 1H (–CH $_{\rm arom}$ ); 8.03–8.01 dd, *J:* 7.32, 0.98 Hz, 1H (–CH $_{\rm arom}$ ); 7.86 s, 1H (–NH); 7.71–7.68 td, *J:* 7.32, 1.47 Hz, 1H (–CH $_{\rm arom}$ ); 7.64–7.61 td, *J:* 7.32, 1.46 Hz, 1H (–CH $_{\rm arom}$ ); 7.53–7.51 d, *J:* 8.78 Hz, 2H (–CH $_{\rm arom}$ ); 7.05–7.03 d, *J:* 8.79 Hz, 2H (–CH $_{\rm arom}$ ); 3.64–3.60 m, 1H (–CH–); 2.77–2.73 dd, *J:* 13.18, 2.93 Hz, 1H (–CH $_{\rm 2}$ –); 2.48–2.43 dd, *J:* 13.66, 8.78 Hz, 1H (–CH $_{\rm 2}$ –); 1.06–1.05 d, *J:* 6.34 Hz, 3H (–CH $_{\rm 3}$ ).

Paper

<sup>13</sup>C NMR (125 MHz,  $CDCl_3$ ) δ (ppm): 181.4, 180.3 (-C=O); 135.1, 133.5, 127.4, 127.1 (-CH<sub>arom</sub>); 146.2, 142.1, 133.5, 130.8, 126.2, 122.5 (C<sub>q</sub>); 66.5 (-CH-); 43.6 (-CH<sub>2</sub>-); 22.1 (-CH<sub>3</sub>). MS (ESI+) m/z (%): 429 (100, [M + Na]<sup>+</sup>), 407 (52, [M]<sup>+</sup>). Anal. calcd for  $C_{20}H_{16}F_3NO_3S$  (407.41).

4.1.1.5. 2-(2-Hydroxypropylthio)-3-(3-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5e). This compound was synthesized from 2-chloro-3-((3-(trifluoromethyl)phenyl)amino) naphthalene-1,4-dione (3b) and 3-mercaptobutan-2-ol (4b) as a red oil by using the general procedure. Yield: 0.076 g, 65%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3433 (-OH), 3346 (-NH), 3050 (-CH<sub>arom</sub>), 2980, 2930 (-CH<sub>aliphatic</sub>), 1664 (-C=O). <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.09-8.07 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 8.01-7.99 d, J: 7.32 Hz, 1H (-CH<sub>arom</sub>); 7.91 s, 1H (-NH); 7.70-7.66 td, J: 7.81, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.62-7.59 td, J: 7.32, 1.47 Hz, 1H (-CH<sub>arom</sub>); 7.41-7.35 t, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.37-7.35 t, *J: 7.81 Hz*, 1H (-CH<sub>arom</sub>); 7.25 s, 1H (-CH<sub>arom</sub>); 7.16-7.15 d, *J: 7.32 Hz*, 1H (-CH<sub>arom</sub>); 3.62-3.58 m, 1H (-CH-); 2.71-2.68 dd, J: 13.67, 3.42 Hz, 1H (-CH<sub>2</sub>-); 2.43-2.39 dd, J: 13.67, 8.79 Hz, 1H (-CH<sub>2</sub>-); 1.05-1.03 d, J: 6.35 Hz, 3H (-CH<sub>3</sub>).  $^{13}$ C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 181.5, 180.4 (-C=O); 146.7, 139.6, 133.5, 130.7, 129.4, 126.4, 122.0, 120.1, 117.4 (C<sub>a</sub>); 135.1, 133.3, 127.4, 127.1 (-CH<sub>arom</sub>); 66.4 (-CH-); 43.8 (-CH<sub>2</sub>-); 22.1 (-CH<sub>3</sub>). MS (ESI-) m/z (%): 407 (16, [M]<sup>+</sup>), 406 (35,  $[M - H]^+$ ), 405 (100,  $[M - 2H]^+$ ). Anal. calcd for  $C_{20}H_{16}F_3NO_3S$  (407.41).

4.1.1.6. 2-((2-Hydroxypropyl)thio)-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (5f). This compound was synthesized from 2-chloro-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3c) and 3-mercaptobutan-2-ol (4b) as a red powder by using the general procedure. Yield: 0.059 g, mp: 85-87 °C, 66%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3423 (-OH), 3328 (-NH), 3075 (-CH<sub>arom</sub>), 2966, 2919 (-CH<sub>aliphatic</sub>), 1667 (-C=O). <sup>1</sup>H NMR (500 MHz, *CDCl*<sub>3</sub>)  $\delta$  (ppm): 8.09–8.07 dd, *J:* 7.32, 0.98 Hz, 1H (-CH<sub>arom</sub>); 8.01–7.99 dd, J: 7.81, 0.97 Hz, 1H (-CH<sub>arom</sub>); 7.93 s, 1H (-NH); 7.70-7.66 td, J: 7.81, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.62-7.60 m, 2H (-CH<sub>arom</sub>); 7.47-7.44 t, J: 7.80 Hz, 1H (-CH<sub>arom</sub>); 7.26-7.23 t, J: 7.80 Hz, 1H (-CH<sub>arom</sub>); 7.00-6.99 d, J: 7.80 Hz, 1H (-CH<sub>arom</sub>); 3.62-3.54 m, 1H (-CH-); 2.96 s, 1H (-OH); 2.78-2.75 dd, J: 13.67, 2.96 Hz, 1H (-CH<sub>2</sub>-); 2.42-2.38 dd, J: 13.67, 8.79 Hz, 1H (-CH<sub>2</sub>-); 1.04-1.03 d, J: 6.35 Hz, 3H (-CH<sub>3</sub>). <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 180.3, 178.8 (-C=O); 146.2, 132.2, 131.1, 129.5, 125.6 (C<sub>q</sub>); 133.8, 132.1, 126.2, 125.9, 124.9, 124.4 (-CH<sub>arom</sub>); 65.0 (-CH-); 42.7 (-CH<sub>2</sub>-); 20.7 (-CH<sub>3</sub>). MS (ESI+) m/z (%): 408 (100,  $[M + H]^+$ ), 430 (54,  $[M + Na]^+$ ). Anal. calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>NO<sub>3</sub>S (407.41).

4.1.1.7. 2-(Furan-2-ylmethylthio)-3-(4-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5g). This compound was synthesized from 2-chloro-3-((4-(trifluoromethyl)phenyl)amino) naphthalene-1,4-dione (3a) and furan-2-ylmethanethiol (4c) as a red powder by using the general procedure. Yield: 0.037 g, mp: 91–92 °C, 30%. FTIR (ATR)  $\nu$  (cm $^{-1}$ ): 3334 (–NH), 3104 (–C-O-ether), 2964, 2918, 2851 (–CH<sub>aliphatic</sub>), 1659 (–C=O).  $^{1}$ H NMR (500 MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.10–8.08 dd, J: 7.81, 1.46 Hz, 1H (–CH<sub>arom</sub>); 7.99–7.97 dd, J: 7.81, 1.46 Hz, 1H (–CH<sub>arom</sub>); 7.65 td, J: 7.32, 1.46 Hz, 1H (–CH<sub>arom</sub>); 7.66 bs, 1H (–NH); 7.62–7.59 td, J: 7.81, 1.47 Hz, 1H (–CH<sub>arom</sub>); 7.48–7.47 d, J: 8.30 Hz, 2H (–CH<sub>arom</sub>); 7.12–7.11 t, J: 0.98 Hz, 1H, (–CH<sub>arom</sub>); 6.93–6.91 d, J:

8.29 Hz, 2H (-CH<sub>arom</sub>); 6.09–6.08 dd, J: 2.93, 1.95 Hz, 1H (-CH<sub>arom</sub>); 5.88–5.87 d, J: 3.42 Hz, 1H (-CH<sub>arom</sub>); 3.91 s, 2H (-CH<sub>2</sub>-).  $^{13}$ C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 179.7, 178.9 (-C=0); 149.7, 144.8, 141.2, 140.9 (C<sub>q</sub>); 133.7, 132.3, 132.0, 129.6, 125.9, 125.8, 121.0 (-CH<sub>arom</sub>); 124.8, 117.4, 109.5, 107.2 (C<sub>q</sub>); 29.4 (-CH<sub>2</sub>-). MS (ESI+) m/z (%): 452 (27, [M+Na]+), 451 (100, [M+Na-1]+). Anal. calcd for  $C_{22}H_{14}F_3NO_3S$  (429.41).

4.1.1.8. 2-(Furan-2-ylmethylthio)-3-(3-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5h). This compound was synthe-2-chloro-3-((3-(trifluoromethyl)phenyl)amino) naphthalene-1,4-dione (3b) and furan-2-ylmethanethiol (4c) as a red oil by using the general procedure. Yield: 0.08 g, 66%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3308 (-NH), 3072 (-C-O<sub>ether</sub>), 2928 (-CH<sub>aliphatic</sub>), 1668 (-C=O). <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.11-8.10 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.99-7.97 d, J: 7.80 Hz, 1H (-CH<sub>arom</sub>); 7.71-7.67 td, *J: 7.81*, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.66 s, 1H (-NH); 7.62-7.59 td, J: 7.81, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.35-7.34 m, 2H (-CH<sub>arom</sub>); 7.13-7.12 m, 2H (-CH<sub>arom</sub>); 7.03-7.01 m, 1H (-CH<sub>arom</sub>); 6.10-6.09 dd, J: 3.41, 1.47 Hz, 1H (-CH<sub>arom</sub>); 5.87-5.86 d, J: 3.41 Hz, 1H (-CH<sub>arom</sub>); 3.88 s, 2H (-CH<sub>2</sub>-). <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 179.6, 179.0 (-C=O); 149.8, 145.7, 141.2, 138.5, 132.4, 129.6, 128.1, 125.0, 120.5, 118.7, 115.9 (C<sub>q</sub>);133.7, 131.9, 126.0, 125.8, 109.5, 107.2 (-CH<sub>arom</sub>); 29.6 (-CH<sub>2</sub>-). MS (ESI-) m/z (%): 428 (100,  $[M - H]^+$ ). Anal. calcd for C<sub>22</sub>H<sub>14</sub>F<sub>3</sub>NO<sub>3</sub>S (429.41).

4.1.1.9. 2-((Furan-2-ylmethyl)thio)-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (5i). This compound was synthesized from 2-chloro-3-((2-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3c) and furan-2-ylmethanethiol (4c) as a red oil by using the general procedure. Yield: 0.07 g, 85%. FTIR (ATR)  $\nu$  $(cm^{-1})$ : 3294 (-NH), 2922 (-CH<sub>aliphatic</sub>), 1676 (-C=O). <sup>1</sup>H NMR (500) MHz,  $CDCl_3$ )  $\delta$  (ppm): 8.09–8.07 dd,  $J: 7.81, 0.98 Hz, 1H (-CH_{arom});$ 7.97-7.95 dd, J: 7.81, 0.98 Hz, 1H (-CH<sub>arom</sub>); 7.75 s, 1H (-NH); 7.68-7.65 td, J: 7.32, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.60-7.56 m, 2H (-CH<sub>arom</sub>); 7.39-7.36 t, J: 7.80 Hz, 1H (-CH<sub>arom</sub>); 7.21-7.17 q, J: 7.32 Hz, 1H (-CH<sub>arom</sub>); 7.12–7.10 d, *J: 2.44 Hz*, 1H (-CH<sub>arom</sub>); 6.82–6.80 d, *J: 7.81* Hz, 1H (-CH<sub>arom</sub>); 6.07-6.06 dd, J: 3.44, 1.96 Hz, 1H (-CH<sub>arom</sub>); 5.86-5.85 d, J: 3.90 Hz, 1H (-CH<sub>arom</sub>); 3.91 s, 2H (-CH<sub>2</sub>-). <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 179.6, 178.7 (-C=O); 149.7, 145.9, 141.2, 132.4, 130.9, 129.5, 125.5, 116.6, 116.1 (C<sub>a</sub>); 133.7, 131.9, 131.8, 125.9, 125.7, 124.5, 124.0 (-CH<sub>arom</sub>); 29.4 (-CH<sub>2</sub>-). MS (ESI+) m/z (%): 452 (100, [M + Na]<sup>+</sup>), 430 (22, [M + H]<sup>+</sup>). Anal. calcd for C<sub>22</sub>H<sub>14</sub>F<sub>3</sub>NO<sub>3</sub>S (429.41).

4.1.1.10. 2-(Phenethylthio)-3-(4-(trifluoromethyl)phenylamino)-naphthalene-1,4-dione (5j). This compound was synthesized from 2-chloro-3-((4-(trifluoromethyl)phenyl)amino)naphthalene-1,4-dione (3a) and 2-phenylethanethiol (4d) as a red powder by using the general procedure. Yield: 0.074 g, mp: 85–87 °C, 57%. FTIR (ATR) ν (cm $^{-1}$ ): 3321 (–NH), 3272, 3060, 3025 (–CH<sub>arom</sub>); 2930 (–CH<sub>aliphatic</sub>), 1662 (–C=O).  $^{1}$ H NMR (500 MHz, *CDCl*<sub>3</sub>) δ (ppm): 8.07–8.05 dd, *J:* 7.33, 1.46 Hz, 1H (–CH<sub>arom</sub>); 8.00–7.98 dd, *J:* 7.32, 1.47 Hz, 1H (–CH<sub>arom</sub>); 7.64 bs, 1H (–NH); 7.63–7.66 td, *J:* 7.33, 1.46 Hz, 1H (–CH<sub>arom</sub>); 7.50–7.48 d, *J:* 8.78 Hz, 2H (–CH<sub>arom</sub>); 7.06–7.03 t, *J:* 7.33 Hz, 2H (–CH<sub>arom</sub>); 6.98–6.96 d, *J:* 7.32 Hz, 1H (–CH<sub>arom</sub>); 6.94–6.92 d, *J:* 8.30 Hz, 4H (–CH<sub>arom</sub>); 2.88–2.85 t, *J:* 7.32 Hz, 2H (–CH<sub>2</sub>–); 2.66–2.63 t, *J:* 7.81 Hz, 2H (–CH<sub>2</sub>–).  $^{13}$ C NMR (125 MHz,

 $CDCl_3$ )  $\delta$  (ppm): 180.0, 178.9 (-C=O); 142.5, 140.5, 138.5, 134.2 (C<sub>0</sub>); 133.5, 132.3, 132.0, 127.5, 127.3, 125.9, 125.7 (-CH<sub>arom</sub>); 129.6, 125.4, 124.7, 120.2 (C<sub>a</sub>); 35.3, 33.5 (-CH<sub>2</sub>-). MS (ESI-) m/ z (%): 453 (20, [M]<sup>+</sup>), 452 (48, [M - H]<sup>+</sup>), 451 (100, [M - 2H]<sup>+</sup>). Anal. calcd for  $C_{25}H_{18}F_3NO_2S$  (453.48).

4.1.1.11. 2-(Phenethylthio)-3-(3-(trifluoromethyl)phenylamino)naphthalene-1,4-dione (5k). This compound was synthesized from 2-chloro-3-((3-(trifluoromethyl)phenyl)amino)naphthalene-1,4dione (3b) and 2-phenylethanethiol (4d) as a red powder by using the general procedure. Yield: 0.05 g, mp: 93-95 °C, 39%. FTIR (ATR)  $\nu$  (cm<sup>-1</sup>): 3308 (-NH), 3063, 3025 (-CH<sub>arom</sub>), 2931 (-CH<sub>aliphatic</sub>), 1657 (-C=O).  $^{1}$ H NMR (500 MHz,  $CDCl_{3}$ )  $\delta$  (ppm): 8.06-8.05 dd, J: 7.32, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.99-7.97 dd, J: 7.32, 0.97 Hz, 1H (-CH<sub>arom</sub>); 7.68-7.65 td, J: 7.32, 1.47 Hz, 1H (-CH<sub>arom</sub>); 7.66 s, 1H (-NH); 7.61-7.58 td, J: 7.80, 1.46 Hz, 1H (-CH<sub>arom</sub>); 7.36-7.33 t, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.31-7.29 d, J: 7.81 Hz, 1H (-CH<sub>arom</sub>); 7.15 s, 1H, (-CH<sub>arom</sub>); 7.06-7.03 m, 3H (-CH<sub>arom</sub>); 6.98-6.93 m, 3H (-CH<sub>arom</sub>); 2.86-2.83 t, J: 7.33 Hz, 2H (-CH<sub>2</sub>-); 2.63-2.60 t, *J. 7.32 Hz*, 2H (-CH<sub>2</sub>-). <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ )  $\delta$  (ppm): 179.9, 179.0 (-C=O); 143.1, 138.6, 138.0, 132.4, 129.6, 127.9, 125.4, 124.0, 119.9, 118.6, 117.8 (C<sub>q</sub>); 133.6, 132.0, 127.5, 127.3, 125.9, 125.7 (-CH $_{arom}$ ); 35.2, 33.6 (-CH $_{2}$ -). MS (ESI-) m/z (%): 453 (25, [M]<sup>+</sup>), 452 (100, [M - H]<sup>+</sup>). Anal. calcd for  $C_{25}H_{18}F_3NO_2S$  (453.48).

#### 4.2. In vitro antimicrobial activity

4.2.1. Determination of minimum inhibitory concentrations (MIC). Antibacterial activities against the three Grampositive bacteria (Staphylococcus aureus ATCC 29213, Staphylococcus epidermidis ATCC 12228, and Enterococcus faecalis ATCC 29212) and four Gram-negative bacteria (Pseudomonas aeruginosa ATCC 27853, Escherichia coli ATCC 25922, Klebsiella pneumoniae ATCC 4352, and Proteus mirabilis ATCC 14153) and the antifungal activity against a yeast Candida albicans ATCC 10231 were determined by the microbroth dilutions technique using the Clinical Laboratory Standards Institute (CLSI) recommendations.<sup>28,29</sup> Mueller-Hinton broth for bacteria and RPMI-1640 medium for the yeast strain were used as the test media. Serial two-fold dilutions ranging from 2500  $\mu g$  mL<sup>-1</sup> to 0.61  $\mu g$ mL<sup>-1</sup> were prepared in the medium. Inoculum was prepared using a 4-6 h broth culture for each bacteria type and 24 hours of culture of yeast strains were adjusted to a turbidity equivalent to the 0.5 McFarland standard, diluted in broth media to give a final concentration of  $5 \times 10^5$  cfu mL<sup>-1</sup> for the bacteria and 5  $\times$  10<sup>3</sup> cfu mL<sup>-1</sup> for the yeast in the test tray. The trays were covered and placed into plastic bags to prevent evaporation. The trays containing Mueller-Hinton broth were incubated at 35 °C for 18-20 h, while the trays containing RPMI-1640 medium were incubated at 35 °C for 46-50 h. The MIC was defined as the lowest concentration of compound, where the complete inhibition of visible growth can be seen. As the control, the antimicrobial effects of the solvents were investigated against the test microorganisms. According to these values of the control groups, the results were evaluated. The MIC values of the compounds are given in Table 2.

4.2.2. Determination of antibiofilm activity. Measurements of the antimicrobial susceptibilities of the bacterial and C. albicans biofilms were assessed using a minimum biofilm eradication concentration (MBEC) assay, which was performed with the stated mentioned techniques as in the following modifications.<sup>30</sup> The 24 h biofilms were grown in a 96 well tissue culture, the microtiter plates were washed three times with 250 μL PBS solutions and air-dried. Serial two-fold dilutions ranging from 10 000 to 625  $\mu g \text{ mL}^{-1}$  for the molecules were prepared in cation adjusted Mueller-Hinton broth (CAMHB). Following that, 200 µL of each sample concentration were added to each corresponding well and the plates were incubated for 24 h at 37 °C. After the incubation, the antibiotics were gently aspirated, the plates were washed, thoroughly scraped, and the contents of each well were incubated in a sonicating water bath for 5 minutes to disrupt the biofilms. 100 μL samples were plated on TSA and the colonies were counted after 24 h of incubation at 37 °C. MBEC was defined as the lowest concentration of molecules where the microorganism failed to regrow after the exposure.

Biofilm attachment and inhibition of the biofilm formation assays were performed as in the previously described method with some further modifications.30 1/10× MIC of molecules were added to the 24 h biofilm and the plates were incubated for 1, 2 and 4 h at 37 °C; molecules at  $1\times$ ,  $1/10\times$  and  $1/100\times$  MIC concentrations were added to the 24 h biofilm, and plates were incubated for 24 h at 37 °C, respectively. Six wells were used for each molecule. The positive controls were microorganisms in TSB-glucose without the molecules of interest. After the incubation, wells were washed with PBS solutions and measured at  $OD_{595}$  nm.

4.2.3. Statistical analysis. All of the experiments were performed in two independent assays. In the determination of MIC and MBEC values, when the results were different in the both experiments, we made another test for the final result. In the assays of biofilm attachment and the inhibition of biofilm formation, results were presented as mean  $\pm$  standard deviation of two independent experiments. One way ANOVA-Bonferroni's multiple comparison test was used to compare the differences between the control and the antimicrobial treated biofilms. P value < 0.001 was considered as a statistically significant parameter.

## 4.3. Cell culture and cytotoxicity assays

Compounds 5b and 5e were assayed for in vitro cytotoxicity against the hepatocellular carcinoma (HepG2) cell line, which was kindly provided by the Department of Pharmaceutical Toxicology in the Faculty of Pharmacy at Istanbul University. HepG2 cells were grown in Eagle's Minimum Essential Medium (EMEM) (Gibco), and were supplemented with 10% FBS (Gibco), and 100 units per mL penicillin G and 100 units per mL streptomycin under the humidified, 5% CO<sub>2</sub> atmosphere at 37 °C. For the assays, the cells were seeded as  $1 \times 10^4$  cells per well in 96 well plates and were allowed 24 h to ensure the attachment. Following the 24 h, the cells were incubated either in the absence or presence of serial dilutions of test compounds for a further 24 h. After the 24 h, cell viability was determined by XTT (Xenometrix) (2,3-bis-(2-methoxy-4-nitro-5-sulfophenyl)-2H-tetrazolium-5-carboxanilide) assay in accordance with the manufacturer's protocol.<sup>31</sup> In each assay, three replicates for each concentration were used and this process, which used three replicates was repeated four times. The cytotoxic effects of each compound were obtained as IC<sub>50</sub> values, which represent the molar drug concentrations required to cause 50% inhibition

#### 4.4. Docking studies

of cell viability.

Paper

The molecules were built using ChemSketch 12.0.1 and converted to 3D structure from the 2D structures, and further 3D structures were energetically minimized and saved as MDL Mol File (\*.mol2). The docking was performed by Molegro Virtual Docker 2010.4.1.0 program.<sup>23</sup> Staphylococcus epidermidis mevalonate diphosphate decarboxylase [PDB ID: 4DPT] was taken as the target enzyme for the rationalization of the antibacterial activity.<sup>24</sup> The polar hydrogen atoms, partial atomic charges and solvation parameters were added. The MolDock optimization search algorithm with a maximum of 100 runs was used for the calculations, while all the other parameters were kept as default. All the spatial structures were examined manually, and the best one was attained for further use.

# Acknowledgements

This work was financially supported by the Scientific Research Projects Coordination Unit of Istanbul University (Project number: 53374) in the supply of the equipment and materials. We also gratefully thank to Prof. Dr Buket Alpertunga and Res. Assist. Ayşe Tarbın Jannuzzi for providing us the hepatocellular carcinoma (HepG2) cell lines and valuable suggestions from the Department of Pharmaceutical Toxicology in the Faculty of Pharmacy at Istanbul University.

# References

- 1 (a) C. Dye, *Philos. Trans. R. Soc., B*, 2014, 369; (b) M. L. Cohen, *Nature*, 2000, **406**, 762–767; (c) A. S. Fauci, *Clin. Infect. Dis.*, 2001, **32**, 675–685.
- 2 World Health Organization, *Mortality and Global Health Estimates*, Geneva, Switzerland, 2013.
- 3 K. F. Smith, M. Goldberg, S. Rosenthal, L. Carlson, J. Chen, C. Chen and S. Ramachandran, *J. R. Soc., Interface*, 2014, **11**, 20140950.
- 4 World Health Organization, *Global Tuberculosis Control*, Geneva, Switzerland, 2010.
- 5 (a) M. N. Gwynn, A. Portnoy, S. F. Rittenhouse and D. J. Payne, Ann. N. Y. Acad. Sci., 2010, 1213, 5–19; (b) D. Sanglard, Curr. Opin. Microbiol., 2002, 5, 379–385; (c) M. L. Cohen, Science, 1992, 257, 1050–1055; (d) H. S. Gold and R. C. Moellering Jr, N. Engl. J. Med., 1996, 335, 1445–1453.
- 6 (a) A. R. M. Coates and Y. Hu, Br. J. Pharmacol., 2007, **152**, 1147–1154; (b) K. Lewis, Nature, 2012, **485**, 439–440.

- 7 (a) A. Riffel, L. F. Medina, V. Stefani, R. C. Santos, D. Bizani and A. Brandelli, Braz. J. Med. Biol. Res., 2002, 35, 811–818;
  (b) C. K. Ryu, R. E. Park, M. Y. Ma and J. H. Nho, Bioorg. Med. Chem. Lett., 2007, 17, 2577–2580;
  (c) V. K. Tandon, H. K. Maurya, D. B. Yadav, A. Tripathi, M. Kumar and P. K. Shukla, Bioorg. Med. Chem. Lett., 2006, 16, 5883–5887;
  (d) V. K. Tandon, H. K. Maurya, M. K. Verma, R. Kumar and P. K. Shukla, Eur. J. Med. Chem., 2010, 45, 2418–2426.
- 8 (a) C. K. Ryu, J. A. Choi and S. H. Kim, Arch. Pharmacal Res.,
  1998, 21, 440–444; (b) V. K. Tandon, H. K. Maurya,
  N. N. Mishra and P. K. Shukla, Eur. J. Med. Chem., 2009,
  44, 3130–3137; (c) V. K. Tandon, D. B. Yadav, H. K. Maurya,
  A. K. Chaturvedi and P. K. Shukla, Bioorg. Med. Chem.,
  2006, 14, 6120–6126.
- 9 V. K. Tandon, H. K. Maurya, A. Tripathi, G. B. ShivaKeshava, P. K. Shukla, P. Srivastava and D. Panda, *Eur. J. Med. Chem.*, 2009, 44, 1086–1092.
- 10 R. F. Silver and H. L. Holmes, Can. J. Chem., 1968, 46, 1859– 1864.
- 11 I. Oeriu, Biokhimiia, 1963, 28, 380-383.
- 12 B. Prescott, J. Med. Chem., 1969, 12, 181-182.
- 13 E. M. Hodnett, C. Wongwiechintana, W. J. Dunn 3rd and P. Marrs, *J. Med. Chem.*, 1983, **26**, 570–574.
- 14 N. G. Clark, Pestic. Sci., 1984, 15, 235-240.
- 15 J. Benites, J. A. Valderrama, K. Bettega, R. C. Pedrosa, P. B. Calderon and J. Verrax, Eur. J. Med. Chem., 2010, 45, 6052–6057.
- 16 S. V. Druzhinin, E. S. Balenkova and V. G. Nenajdenko, *Tetrahedron*, 2007, **63**, 7753–7808.
- 17 B. S. Samant and C. Chakaingesu, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 1420–1423.
- 18 V. K. Tandon, H. K. Maurya, D. B. Yadav, A. Tripathi, M. Kumar and P. K. Shukla, *Bioorg. Med. Chem. Lett.*, 2006, 16, 5883–5887.
- 19 A. F. Tuyun, N. Bayrak, H. Yildirim, N. Onul, E. M. Kara and B. O. Celik, *J. Chem*, 2015, 645902.
- 20 Website: http://www.ibmc.msk.ru/PASS.
- 21 (a) V. V. Poroikov, D. A. Filimonov, Y. V. Borodina, A. A. Lagunin and A. Kos, J. Chem. Inf. Comput. Sci., 2000, **40**, 1349–1355; (b) V. V. Poroikov and D. A. Filimonov, J. Des., 2002, **16**, 819–824; (c) Comput.-Aided Mol. A. Geronikaki, E. Babaev, J. Dearden, W. Dehaen, D. Filimonov, I. Galaeva, V. Krajneva, A. Lagunin, F. Macaev, G. Molodavkin, V. Poroikov, S. Pogrebnoi, V. Saloutin, A. Stepanchikova, E. Stingaci, N. Tkach, L. Vlad and T. Voronina, Bioorg. Med. Chem., 2004, 12, 6559-6568; (d) A. Geronikaki, A. Lagunin, V. Poroikov, D. Filimonov, D. Hadjipavlou-Litina and P. Vicini, SAR QSAR Environ. Res., 2002, 13, 457-471; (e) R. K. Goel, V. Kumar and M. P. Mahajan, Bioorg. Med. Chem. Lett., 2005, **15**, 2145–2148; (f) G. K. Gupta, V. Saini, R. Khare and V. Kumar, Med. Chem. Res., 2014, 23, 4209-4220; (g) A. Kumar, P. Lohan, D. K. Aneja, G. K. Gupta, D. Kaushik and O. Prakash, Eur. J. Med. Chem., 2012, 50, 81-89.
- 22 (a) R. M. Donlan and J. W. Costerton, Clin. Microbiol. Rev., 2002, 15, 167; (b) P. Jorge, A. Lourenco and M. O. Pereira, Biofouling, 2012, 28, 1033-1061.

23 R. Thomsen and M. H. Christensen, *J. Med. Chem.*, 2006, **49**, 3315–3321.

**RSC Advances** 

- 24 (a) M. L. Barta, D. A. Skaff, W. J. McWhorter, T. J. Herdendorf, H. M. Miziorko and B. V. Geisbrecht, J. Biol. Chem., 2011, 286, 23900–23910; (b) D. A. Skaff, W. J. McWhorter, B. V. Geisbrecht, G. J. Wyckoff and H. M. Miziorko, Arch. Biochem. Biophys., 2015, 566, 1–6.
- 25 (a) D. V. Krepkiy and H. M. Miziorko, *Biochemistry*, 2005, 44, 2671–2677; (b) D. Krepkiy and H. M. Miziorko, *Protein Sci.*, 2004, 13, 1875–1881; (c) N. E. Voynova, Z. J. Fu, K. P. Battaile, T. J. Herdendorf, J. J. P. Kim and H. M. Miziorko, *Arch. Biochem. Biophys.*, 2008, 480, 58–67.
- 26 N. P. Buu-Hoi, R. Royer and M. Hubert-Habart, *Recl. Trav. Chim. Pays-Bas*, 1954, 73, 188–192.

- 27 A. Mital, M. Sonawane, S. Bindal, S. Mahlavat and V. Negi, *Der Pharma Chem.*, 2010, 2(3), 63–73.
- 28 Clinical and Laboratory Standards Institute, *Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts*, CLSI, Wayne, PA, 2th edn, 1997, Approved Standard M27-A2.
- 29 Clinical and Laboratory Standards Institute, *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically*, CLSI, Wayne, PA, 7th edn, 2006, Approved standard M7-A7.
- 30 E. Mataraci and S. Dosler, Antimicrob. Agents Chemother., 2012, 56, 6366-6371.
- 31 http://www.aniara.com/Resources/Xenometrix-Literature.aspx.