


 Cite this: *Chem. Commun.*, 2024, 60, 12417

 Received 9th July 2024,  
 Accepted 2nd October 2024

DOI: 10.1039/d4cc03427a

rsc.li/chemcomm

# Characterizing excited-state intramolecular proton transfer in 3-hydroxyflavone with ultrafast transient infrared spectroscopy†

 Valerie S. Winkler and Joseph A. Fournier \*


**Vibrational dynamics associated with excited-state intramolecular proton transfer in 3-hydroxyflavone and 3-hydroxy-2-(thiophen-2-yl)chromen-4-one are characterized with ultrafast transient infrared spectroscopy. The spectroscopic data reveal rapid (<100 fs) proton transfer dynamics in both species, followed by vibrational relaxation of the tautomer products within a few picoseconds. Coherent coupling of the shared proton to low-frequency modes along the hydrogen bond coordinate are also observed in the ground electronic states.**

Excited-state proton transfer reactions are a crucial class of reactions that are ubiquitous in biological systems, chemical catalysis, and photoredox chemistry.<sup>1,2</sup> Many model systems have been investigated to characterize the fundamental dynamics and mechanisms of proton transfer processes. In particular, 3-hydroxyflavone (3HF) is a prototypical system for the study of excited-state intramolecular proton transfer (ESIPT) reactions.<sup>3,4</sup> Upon UV excitation to the first electronic excited state ( $S_1$ ), the “normal” enol isomer undergoes ESIPT to form the keto tautomer (referred to as the  $S_1'$  state), as depicted in Fig. 1. The  $S_0$  to  $S_1$  excitation is a  $\pi$ - $\pi^*$  transition where significant electron density is removed from the enol OH group, resulting in a highly acidic proton where transfer to the tautomer  $S_1'$  structure becomes energetically favourable. Dual fluorescence from both the normal and tautomer states have been observed, with their ratio being highly dependent on solvent and temperature.<sup>3–9</sup> Due to ease of chemical modification of the 3HF scaffold, flavonoids and their derivatives have been widely explored as tuneable chemical probes and sensors.<sup>10–14</sup>

Time-resolved fluorescence<sup>15–17</sup> and UV/Vis transient absorption<sup>9,18</sup> spectroscopic investigations of 3HF have reported ultrafast (<100 fs) ESIPT timescales occurring within the instrument

response. In polar solvents such as ethanol and acetonitrile, hydrogen bonding between the solvent and enol group disrupts the intramolecular hydrogen bond. Consequently, slower (few ps) proton transfer dynamics have been measured that involve breaking of the enol-solvent hydrogen bond.<sup>16,17</sup> Numerous computational studies have also been performed on 3HF to investigate the ESIPT process.<sup>19–23</sup> Molecular dynamics simulations, in general, corroborate both the ultrafast proton transfer dynamics in gas phase or nonpolar solvents and the slower solvent-inhibited picosecond dynamics in polar solvents. The rapid transfer dynamics arise from the relatively small barrier (700–1400  $\text{cm}^{-1}$ , 2–4  $\text{kcal mol}^{-1}$ ) between the  $S_1$  and  $S_1'$  states, indicating a near-barrierless proton transfer reaction upon excitation into the Franck–Condon region.<sup>23</sup>

The multiple broad features observed in the transient UV/Vis spectra of 3HF, however, have led to interpretational difficulties in determining the molecular-level mechanism of ESIPT.<sup>9</sup> To



**Fig. 1** Calculated energy level diagrams for 3HF (black) and 3HTC (purple). Barrier heights relative to the  $S_0$  and  $S_1$  normal states are shown for the transition state levels. Calculations were performed at the B3LYP-D3BJ/aug-cc-pVDZ level of theory and basis set.

Washington University in St. Louis, One Brookings Dr, St. Louis, MO 63130, USA.

E-mail: jfournier@wustl.edu; Tel: +1-314-935-8183

 † Electronic supplementary information (ESI) available: Experimental and computational methods, computational results, UV/Vis spectra, IR pump-IR probe transient absorption spectra. See DOI: <https://doi.org/10.1039/d4cc03427a>


uncover the vibrational dynamics underlying the ESIPT process, Chevalier *et al.* reported transient infrared spectra of 3HF in acetonitrile following UV excitation at 360 nm.<sup>24</sup> Global fitting analysis of the transient infrared spectra resulted in a  $\sim 3$  ps spectral decay component assigned to the solvent-inhibited ESIPT process. A slower  $\sim 10$  ps spectral component was assigned to vibrational relaxation of the tautomer  $S_1'$  state following proton transfer. The 500 fs experimental resolution, however, did not allow the faster ESIPT component unperturbed by solvent to be measured.

To better characterize ESIPT and provide improved experimental benchmarks for computational predictions, we have measured transient infrared spectra of 3HF and the derivative 3-hydroxy-2-(thiophen-2-yl)chromen-4-one (3HTC)<sup>25</sup> with approximately 100 fs resolution in a non-perturbing solvent (chloroform). Detailed experimental, computational, and data analysis methods are provided in the ESI.<sup>†</sup>

The calculated (B3LYP-D3BJ/aug-cc-pVDZ) energy levels for isolated structures of the normal, transition state, and tautomer species in the ground and first excited electronic states for 3HF and 3HTC are shown in Fig. 1. Optimized structures for each species are given in ESI,<sup>†</sup> Fig. S1. Small barriers between the normal and tautomer structures are predicted in the excited state, with 3HF calculated to have a barrier of about  $1200\text{ cm}^{-1}$  and 3HTC about  $1700\text{ cm}^{-1}$ , consistent with previous computational studies.<sup>19,26</sup>

Transient infrared spectra following UV excitation are shown in Fig. 2a for 3HF (excitation wavelength 363 nm) and Fig. 2b

for 3HTC (excitation wavelength 379 nm). The UV/Vis spectra of 3HF and 3HTC are provided in Fig. S2 (ESI<sup>†</sup>). In both species, there are two dominant negative signals corresponding to bleaching of ground-state  $S_0$  vibrational transitions: the ketone carbonyl stretch near  $1630\text{ cm}^{-1}$  and the enol OH bend near  $1440\text{ cm}^{-1}$  (FTIR spectra are given by the grey inverted traces in Fig. 2a and b). Computational predictions of the vibrational spectra in the  $S_0$  states of the isolated molecules are provided in Fig. S3 (ESI<sup>†</sup>). The positive signals correspond to vibrational transitions in excited electronic states. In both species, the transient spectra show minor evolution after about 200 fs, indicating that the excited-state features beyond 200 fs derive from the  $S_1'$  tautomer following a rapid proton transfer process. For 3HF, the most intense excited-state features appear near  $1300$  and  $1350\text{ cm}^{-1}$ . Based on the spectral components and computational assignments made by Chevalier *et al.*<sup>24</sup> and our calculated  $S_1'$  spectrum (Fig. S4, ESI<sup>†</sup>), these features are both expected to have mixed C–C and C–O stretch character. Two weaker excited-state features appear near  $1450\text{ cm}^{-1}$ , which are assigned to transitions with mixed CH and OH bend character based on the calculated  $S_1'$  spectrum. The calculations by Chevalier *et al.*<sup>24</sup> (B3LYP/TZVP), to the best of our knowledge, are the only computed excited-state vibrational spectra reported for 3HF, and were shown to be strongly influenced by inclusion of either a single explicit acetonitrile molecule or a polarizable continuum model (PCM) for the solvent. While our excited-state calculations are in close agreement to those reported by Chevalier *et al.*<sup>24</sup> on isolated 3HF, both sets of calculations




Fig. 2 Transient infrared spectra of (a) 3HF and (b) 3HTC at selected delay times following UV excitation. FTIR spectra are presented as inverted grey traces in both panels. Assignment of the OH bend transition in the  $S_1$  state and integration area for the time traces shown in Fig. 3 are given by the green boxes. Global fit analysis of the transient infrared spectra between 0.2–10 ps for (c) 3HF and (d) 3HTC. The  $A_0$  components are static contributions representing the ground-state bleach and excited-state  $S_1'$  spectra. The  $A_1$  components follow a monoexponential decay (time constants given inset) and are assigned to vibrational relaxation of the  $S_1'$  states following proton transfer.



show only modest overall agreement with the experimental excited-state spectra. A more thorough computational characterization of the excited-state vibrational spectra of 3HF and its derivatives is clearly warranted.

Transient IR spectra for 3HTC have not been previously reported, but show similar patterns to 3HF. An intense, closely spaced excited-state doublet feature appears near  $1325\text{ cm}^{-1}$ . The other strong excited-state features occur at  $1375$  and  $1450\text{ cm}^{-1}$ . Based on the similarities of the experimental spectra, the strongest 3HTC  $S_1'$  transitions also likely derive from normal mode vibrations that involve C–C stretch, C–O stretch, CH bend, and OH bend motions. The calculated  $S_1'$  spectrum of 3HTC is presented in Fig. S4 (ESI†).

Global fit analysis of the transient spectra was performed starting at 200 fs, following proton transfer. Two dominant decay-associated spectral components were recovered for both 3HF and 3HTC, which are presented in Fig. 2c and d, respectively. The largest component for each species, labelled  $A_0$ , is constant over the 10 ps experimental window and correspond to the  $S_0$  ground-state bleach and the excited-state  $S_1'$  spectra. The constant nature of the  $A_0$  components indicate that the  $S_1'$  tautomers form within 200 fs following electronic excitation. A weaker second profile observed for both compounds, labelled  $A_1$ , have monoexponential decay constants of about 1 ps for 3HF and 3 ps for 3HTC. The 3HF  $A_1$  spectrum is similar in appearance to the longer-lived spectral component identified by Chevalier *et al.* in acetonitrile.<sup>24</sup> We likewise assign the  $A_1$  decay-associated spectra of 3HF and 3HTC to changing intensities and frequencies of the  $S_1'$  states resulting from vibrational relaxation following ESIPT. With 3HF and 3HTC having very similar structures, it is expected that intermolecular relaxation to the solvent will occur on similar timescales for both species. The timescale difference between 3HF and 3HTC, therefore, is attributed to differences in the efficiency of intramolecular vibrational relaxation between the two species following the proton transfer event.

There are no unambiguous signatures, however, of the normal  $S_1$  structure for either species. Based on the few-ps solvent-inhibited  $S_1$  spectral component of 3HF determined by Chevalier *et al.* in acetonitrile,<sup>24</sup> the most intense  $S_1$  transitions are masked by the strong and broad  $S_1'$  background near  $1350\text{ cm}^{-1}$  (computational predictions of the  $S_1$  state vibrational spectra are provided in Fig. S5, ESI†). The extracted  $S_1$  spectra by Chevalier *et al.*, however, showed a relatively isolated transition near  $1275\text{ cm}^{-1}$  that falls below the strong  $S_1'$  background and was assigned as the OH bend transition. This transition, therefore, acts as a direct reporter of the proton transfer coordinate. The transient spectra at 100 fs delay time does show larger intensity in this spectral region compared to later spectra, indicating the presence of the  $S_1$  species at the earliest delay times measured. The decay dynamics of the putative  $S_1$  OH bend transitions near  $1275\text{ cm}^{-1}$  for both 3HF and 3HTC are presented in Fig. 3. While the measured signals at  $>50$  fs show intensity above the instrument response (grey traces in Fig. 3), it is difficult to deconvolve the responses at the current signal-to-noise levels. Therefore, we characterize the decay of the  $S_1$




Fig. 3 Spectral dynamics of (a) 3HF and (b) 3HTC for the  $S_1$  OH bend transitions near  $1275\text{ cm}^{-1}$ . The data were fit to a monoexponential decay function,  $f(t) = Ae^{-t/\tau} + b$  (solid lines). The exponential decay time constants ( $24 \pm 3$  fs for 3HF and  $29 \pm 8$  fs for 3HTC) are convolved with the instrument response (grey trace) and, therefore, are characterized as being sub-100 fs.

states and the ESIPT timescales in 3HF and 3HTC as being sub-100 fs, consistent with those measured by Ameer-Beg *et al.* for 3HF in nonpolar solvent by transient UV/Vis spectroscopy.<sup>9</sup> Time profiles in the  $1300$ – $1350\text{ cm}^{-1}$  regions are provided in Fig. S6 for 3HF and S7 for 3HTC (ESI†). These kinetics display similarly fast initial decays with time constants  $<100$  fs, indicating that strong  $S_1$  transitions are present at the earliest delay times. In particular, time traces and fits of isolated  $S_1'$  transitions near  $1460\text{ cm}^{-1}$  in Fig. S6c and S7c (ESI†) have an initial sub-100 fs increase in signal, consistent with the rapid decay of the  $S_1$  features. These  $S_1'$  features also have slower exponential growth time constants of about 1.2 ps and 2.5 ps for 3HF and 3HTC, respectively, in reasonable agreement with the  $A_1$  decay time constants from the global fit analysis. Although more accurate ESIPT timescales cannot be currently measured, these measurements provide a more direct experimental estimate for the ESIPT timescales in 3HF and 3HTC.

The intramolecular hydrogen bond in the ground  $S_0$  state in both species is quite weak, with proton donor–acceptor distances between the sharing oxygen atoms calculated to be approximately  $2.6\text{ \AA}$ . This distance decreases significantly in the  $S_1$  state, with optimized values close to  $2.5\text{ \AA}$ . The large decrease in the O–O distance suggests that strong anharmonic coupling should exist between the proton degrees of freedom (OH stretch and bend) and low-frequency vibrational modes that alter the OH–O hydrogen bond interaction (*e.g.*, the O–O stretch). Interestingly, the OH bend ground-state transitions for 3HF and 3HTC exhibit highly damped oscillatory behaviour at early pump–probe delay times (Fig. 4). Identical oscillations are also present in the ground-state IR pump–IR probe transient absorption spectra (Fig. S8 and S9, ESI†). The presence of these oscillations in the






Fig. 4 Spectral dynamics of the  $S_0$  ground-state OH bend transitions for 3HF (black dots) and 3HTC (purple dots). Fits to a damped oscillatory function,  $f(t) = A \sin\left(\frac{2\pi t}{T} + b\right)e^{-t/\tau} + c$ , are given by solid lines. The exponential decay ( $\tau$ ) and oscillation period ( $T$ ) fitting parameters and their standard deviations are shown inset. The oscillation periods are consistent with coupling between the shared proton's degrees of freedom and low-frequency vibrational modes.

ground electronic state is consistent with impulsive excitation and coherent coupling dynamics between the OH bend and low-frequency hydrogen bond motions within the  $S_0$  state.<sup>19,23,27–29</sup> The oscillation periods of 264 fs in 3HF and 295 fs in 3HTC correspond to vibrational frequencies of about  $125\text{ cm}^{-1}$  and  $110\text{ cm}^{-1}$ , respectively. The  $S_0$  calculations predict several modes in this low-frequency region that involve the proton donor and acceptor groups that are most likely involved in the coupling: in-plane bending motions that bring the donor and acceptor oxygen atoms towards each other ( $118\text{ cm}^{-1}$  in 3HF) and out-of-plane bending motions ( $102\text{ cm}^{-1}$  in 3HF,  $102\text{ cm}^{-1}$  and  $107\text{ cm}^{-1}$  in 3HTC) perpendicular to the OH–O coordinate (Fig. S10, ESI†). Such motions that directly impact the proton transfer coordinate are expected to play prominent roles in the ESIPT process upon electronic excitation.

The observed rapid ESIPT dynamics and wavepacket dynamics on the ground state provide new experimental benchmarks for advanced theoretical methods. Higher-order experiments with better time resolution that resolve the UV excitation axis (namely, two-dimensional electronic-vibrational spectroscopy) are necessary to more accurately measure the ESIPT dynamics and identify the nuclear coordinates that drive ESIPT through the measurement of vibrational wavepacket dynamics on the excited electronic surfaces.<sup>30</sup> Experiments on the deuterated species to measure kinetic isotope effects would further validate the ESIPT mechanism. Temperature-dependent studies could also provide an experimental measurement of the ESIPT barrier. Improved fundamental understanding of ESIPT and vibrational relaxation mechanisms will aid in the development of more tuneable and sensitive molecular probes and sensors for material and biological applications.

This work was supported by the US National Science Foundation under grant CHE-2044927.

## Data availability

The following data are provided in ESI†: Cartesian coordinates of the optimized  $S_0$ ,  $S_1$  and  $S_1'$  structures; transient infrared spectra at the pump–probe waiting times and the global fit analysis presented in Fig. 2. Data from all other pump–probe times are available on the Harvard Dataverse at <https://doi.org/10.7910/DVN/Q71QIH>.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- 1 A. P. Demchenko, K. C. Tang and P. T. Chou, *Chem. Soc. Rev.*, 2013, **42**, 1379–1408.
- 2 T. Kumpulainen, B. Lang, A. Rosspeintner and E. Vauthey, *Chem. Rev.*, 2017, **117**, 10826–10939.
- 3 P. K. Sengupta and M. Kasha, *Chem. Phys. Lett.*, 1979, **68**, 382.
- 4 G. J. Wolfe and P. J. Thistlethwaite, *J. Am. Chem. Soc.*, 1981, **103**, 6919.
- 5 M. Itoh, K. Tokumura, Y. Tanimoto, Y. Okada, H. Takeuchi, K. Obi and I. Tanaka, *J. Am. Chem. Soc.*, 1982, **104**, 4146.
- 6 A. J. G. Strandjord, S. H. Courtney, D. M. Friedrich and P. F. Barbara, *J. Phys. Chem.*, 1983, **87**, 1125–1133.
- 7 D. McMorro and M. Kasha, *J. Phys. Chem.*, 1984, **88**, 2235.
- 8 S. Protti and A. Mezzetti, *J. Mol. Liq.*, 2015, **205**, 110–114.
- 9 S. Ameer-Beg, S. M. Ormson, R. G. Brown, P. Matousek, M. Towrie, E. T. J. Nibbering, P. Foggi and F. V. R. Neuwahl, *J. Phys. Chem. A*, 2001, **105**, 3709–3718.
- 10 M. Sarkar and P. K. Sengupta, *Chem. Phys. Lett.*, 1991, **179**, 68.
- 11 A. Sytnik, D. Gormin and M. Kasha, *Proc. Natl. Acad. Sci. U. S. A.*, 1994, **91**, 11968–11972.
- 12 J. Guharay, B. Sengupta and P. K. Sengupta, *Proteins*, 2001, **43**, 75–81.
- 13 A. S. Klymchenko and A. P. Demchenko, *Langmuir*, 2002, **18**, 5637.
- 14 S. Chaudhuri, K. Basu, B. Sengupta, A. Banerjee and P. K. Sengupta, *Luminescence*, 2008, **23**, 397–403.
- 15 A. N. Bader, V. G. Pivovarenko, A. P. Demchenko, F. Ariese and C. Gooijer, *J. Phys. Chem. B*, 2004, **108**, 10589.
- 16 T. C. Swinney and D. F. Kelley, *J. Phys. Chem.*, 1991, **95**, 10369.
- 17 G. A. Brucker, T. C. Swinney and D. F. Kelley, *J. Phys. Chem.*, 1991, **95**, 3190.
- 18 B. J. Schwartz, L. A. Peteanu and C. B. J. Harris, *Phys. Chem.*, 1992, **96**, 3591.
- 19 M. Nottoli, M. Bondanza, F. Lipparini and B. J. Mennucci, *Chem. Phys.*, 2021, **154**, 184107.
- 20 Y. Li, F. Siddique, A. J. Aquino and H. Lischka, *J. Phys. Chem. A*, 2021, **125**, 5765–5778.
- 21 X. P. Chang, F. R. Fan, G. Zhao, X. Ma and T. S. Zhang, *Chem. Phys.*, 2023, **575**, 112056.
- 22 M. J. Colin, M. Á. Aguilar and M. E. Martín, *ACS Omega*, 2023, **8**, 19939–19949.
- 23 M. A. Bellucci and D. F. Coker, *J. Chem. Phys.*, 2012, **136**, 194505.
- 24 K. Chevalier, M. M. N. Wolf, A. Funk, M. Andres, M. Gerhards and R. Diller, *Phys. Chem. Chem. Phys.*, 2012, **14**, 15007–15020.
- 25 D. Dziuba, I. A. Karpenko, N. P. F. Barthes, B. Y. Michel, A. S. Klymchenko, R. Benhida, A. P. Demchenko, Y. Mély and A. Burger, *Chem. – Eur. J.*, 2014, **20**, 1998–2009.
- 26 H. Ma and J. Huang, *RSC Adv.*, 2016, **6**, 96147.
- 27 J. Stenger, D. Madsen, J. Dreyer, E. T. J. Nibbering, P. Hamm and T. Elsaesser, *J. Phys. Chem. A*, 2001, **105**, 2929–2932.
- 28 C. Chudoba, E. Riedle, M. Pfeiffer and T. Elsaesser, *Chem. Phys. Lett.*, 1996, **263**, 622.
- 29 M. Balasubramanian, A. Reynolds, T. J. Blair and M. Khalil, *Chem. Phys.*, 2019, **519**, 38–44.
- 30 J. D. Gaynor and M. Khalil, *J. Chem. Phys.*, 2017, **147**, 094202.

