

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

*Activation of C-H and B-H bonds through agostic bonding: an ELF/QTAIM insight*Emilie-Laure Zins,^{a,b} Bernard Silvi,^{c,d} M. Esmail Alikhani,^{a,b,*}

a-Sorbonne Universités, UPMC Univ. Paris 06, MONARIS, UMR 8233, Université Pierre et Marie Curie, 4 Place Jussieu, case courrier 49, F-75252 Paris Cedex 05, France

b-CNRS, MONARIS, UMR 8233, Université Pierre et Marie Curie, 4 Place Jussieu, case courrier 49, F-75252 Paris Cedex 05, France

c-Sorbonne Universités, UPMC Univ. Paris 06, Laboratoire de Chimie Théorique (LCT), UMR 7616, Université Pierre et Marie Curie, 4 place Jussieu, case courrier 137, F-75252 Paris Cedex 05, France

d-CNRS, Laboratoire de Chimie Théorique (LCT), UMR7616, Université Pierre et Marie Curie, 4 place Jussieu, case courrier 137, F-75252 Paris Cedex 05, France

Abstract

Agostic bondings are of paramount importance in C-H bond activation processes. The reactivity of the σ C-H bond thus activated will depend on the nature of the metallic center, the nature of the ligand involved in the interaction and co-ligands, as well as on geometric parameters. Because of their importance in organometallic chemistry, a qualitative classification of agostic bondings could be very much helpful. Herein we propose descriptors of the agostic character of bondings based on the electron localization function (ELF) and Quantum Theory of Atoms in Molecules (QTAIM) topological analysis. A set of 31 metallic complexes taken, or derived, from the literature was chosen to illustrate our methodology. First, some criteria should prove that an interaction between a metallic center and a σ X-H bond can indeed be described as an "agostic" bonding. Then, the contribution of the metallic center in the protonated agostic basin, in the ELF topological description, may be used to evaluate the agostic character of bondings. A σ X-H bond is in agostic interaction with a metal center when the protonated X-H basin is a trisynaptic basin with a metal contribution strictly larger than the numerical uncertainty, i.e. 0.01 e. In addition, it was shown that the weakening of the electron density at the X – H_{agostic} bond critical point with respect to that of X–H_{free} well correlates with the lengthening of the agostic X – H bond distance as well as with the shift of the vibrational frequency associated with the ν_{X-H} stretching mode. Furthermore, the use of a normalized parameter that takes into account the total population of the protonated basin, allows to compare the agostic character of bondings involved in different complexes.

Keywords

Topological Description, Electron Localization Function (ELF), Agostic Bonding, C-H Bond Activation, Trisynaptic Basin, Covariance Matrix

* Corresponding author: esmail.alikhani@upmc.fr

I. Introduction

In the context of catalytic processes, understanding the molecular mechanism involving transition metals is of paramount importance. Indeed, since the pioneering works of Chatt and Davidson¹ from the one hand, and Bergman² and Jones³ from the other hand, a considerable number of investigations have demonstrated that transition metal complexes may be involved in C-H bond activation. A thorough study of structures and reactivity of alkyl transition metal complexes has led to the introduction of a new concept: the term "agostic" was proposed to characterize the formation of a 3 centers - 2 electrons (3c-2e) interaction leading to the activation of a C-H bond around transition metal centers.^{4,5,6,7} The choice of a specific term to underline the importance of the activation of C-H bond proved to be remarkably appropriate. Indeed, thorough investigations in various catalytic processes further lead to the development of numerous reactions involving agostic interactions, such as the alkane oxidative addition,⁸ C-H bond elimination,⁹ transcyclometallation,¹⁰ cyclometallation of benzoquinoline,¹¹ and Ziegler-Natta polymerization.^{12,13} In processes such as cyclometallation using d-block transition metals, it was proposed that agostic interactions may play a decisive role. Indeed, these interactions may either favor or prevent some cyclometallations.¹⁴ A mechanistic study has shown how agostic bondings may lead to deactivation processes of the Grubb catalyst.¹⁵ Thus, a thorough understanding of the agostic bondings may help in designing new reagents that may react in a specific way.

Despite their paramount importance in catalytic processes, the identification of an agostic interaction is far from obvious. Indeed, even if some approaches were proposed to experimentally or theoretically characterize these 3c-2e interactions, a consensual tool that may qualitatively describe the strength of every kind of agostic bondings is still missing.

The formation of an agostic bonding comes from an interaction between a C-H σ bond and an unoccupied orbital of an hypovalent transition metal center. Criteria were proposed to determine whether an organometallic complex contains an agostic interaction. More specifically, the following geometric parameters were established: a C-H agostic bonding should be characterized by a distance between the metallic center and the hydrogen atom in the 1.8 - 2.3 Å range, as well as an M-H-C angle in the 90 - 140° range.⁷ On the other hand, the presence of sterically-constrained ligands or pincer ligands, for instance, may lead to profound geometric distortions.¹⁶ Furthermore, close to the threshold values, it may be difficult to conclude whether a complex contains an agostic interaction or not. Thus, these geometric criteria are not unambiguous, without mentioning the difficulty in determining them in some cases, specifically in dynamic systems.

As far as other "weak" interactions are concerned, criteria derived from topological studies were proposed to classify hydrogen bonding into three categories, and to propose a quantitative scale for these types of interactions.¹⁷ Since there are some common points between agostic and hydrogen bondings,¹⁶ the use of a similar approach to characterize agostic bonding comes naturally.

Several reviews aimed at presenting experimental and theoretical tools to identify agostic bonding.^{18,19,20} In the present article, we will first briefly recall the different definitions that were / are considered for agostic bonding. Their theoretical studies are then briefly summarized, with a particular emphasis on topological approaches. The following part of this article is dedicated to the presentation of a new methodological approach aiming at estimating the agostic character of a bonding. Representative examples of several kinds of agostic bonding taken from the recent literature on the topic are used to illustrate this approach. The topological description of different types of $M \cdots H-X$, interactions with $X = B$ or C , will be presented. A classification of agostic bondings based on the strength of the interaction, is then proposed. To this end, the use statistical descriptors to qualitatively evaluate the strength of an agostic bonding is validated by comparison with experimental parameters.

II. Agostic bonding: several definitions and nuances

Despite the fundamental importance of the concept of agostic bonding in terms of reactivity and chemical activation of σ bonds, the definition of these interactions is neither unique nor unambiguous. The main definitions associated to these interactions and the context in which they were introduced, are presented herein.

1. Historical definition and evolution

The "agostic" term was initially introduced to describe a specific intramolecular 3c-2e interaction in which a σ C-H bond "could act as a ligand to a transition metal center".⁷ Thus, a specific term was coined for interactions involving C-H bond to underline their importance in catalytic processes. First of all, β agostic interactions were characterized and thoroughly investigated both theoretically and experimentally. α C-H agostic bonding appears to be more rare. Even if in the initial definition, the term "agostic" was specifically devoted to C-H bond, intramolecular 3c-2e interactions between a metallic center and a σ bond involving H bonded to a heteroatom are now almost always called "agostic". The agostic bondings that will be discussed in the present article are presented in Figure 1.




Figure 1: Types of agostic bondings that will be investigated in the present study.

From an experimental point of view, the characterization of an agostic bonding is mainly based on four criteria:¹⁸

- crystallographic data,
- NMR chemical shifts to high field $\delta = -5$ to -15 ppm,
- reduced NMR coupling constants $^1J = 75$ to 100 Hz,
- low vibrational frequencies $\nu_{\text{C-H}} = 2700 - 2300 \text{ cm}^{-1}$.

As far as geometric properties are concerned, Brookhart and Green proposed in their seminal article,⁷ the criteria summarized in the Table 1 may be used to distinguish between an agostic and an anagostic interaction:

	M...H-C agostic	M...H-C anagostic
Distance M...H (Å)	1.8 - 2.3	2.3 - 2.9
Angle MHC (°)	90 - 140	110 - 170

Table 1: Geometric criteria used in the literature to distinguish between an agostic and an anagostic interaction.⁷

Based on QTAIM calculations, five topological criteria were proposed to characterize an agostic bonding¹⁸:

- a triplet of concomitant topological objects: a bond critical point, a bond path and an interatomic surface,
- a ring critical point, that is a signature of a structural instability,

- a Laplacien of the electron density of the bond critical point $\nabla^2\rho_{\text{BCP}}$ in the 0.15 - 0.25 a.u. range,
- a negative net charge for an hydrogen atom involved in an agostic bonding,
- a dipolar polarization that is 15 - 30% larger for an agostic hydrogen compared with a non-agostic one.

However, these criteria are not unambiguous since they cannot be applied to any type of agostic bondings, as it will be further discussed in the part III.

2. Other definitions

In addition with the rigorous initial definition, the term "agostic" is also sometimes used in a broader definition²¹, to characterize intermolecular interactions, especially in the context of sigma alkane,^{22,23} silane^{24,25} and sigma boranes.^{25,26}

In an attempt to fully characterize some systems potentially containing weak agostic interactions, experimental and theoretical approaches were applied to numerous and various organometallic complexes characterized by geometric distortions. While in some cases clear agostic bondings were identified, in some other systems the approaches used failed to characterize such interactions. The term "anagostic" was proposed to define systems where an interaction between a σ bond and a metallic center leads to a geometric distortion of the structure whereas some considered criteria are not meet to label this interaction as "agostic".^{27,28,29}

In most of the above-mentioned studies, the agostic interaction occurs in an organometallic complex centered on a metallic atom. In a wider sense however, this concept was employed to qualify situations in which organic molecules interact with a metallic surfaces during catalytic processes in heterogeneous phase. For instance, mechanisms involving agostic interactions were proposed in the chemisorption of hydrogen, hydrocarbons and intermediates on Pt(111).^{30,31} The formation of a particularly rare agostic interaction was even proposed in the context of the water dissociation of a Pt(111) surface, in combination with hydrogen bondings.³²

III. Theoretical characterizations of agostic bondings

1. Equilibrium geometry

Since agostic interactions are generally defined on the basis of geometric criteria, geometry optimization may appear as a suitable tool to investigate this type of interactions. Such an approach may complement X-ray structures analyses. In some cases however, the complexes cannot be isolated in crystalline form (especially in the case of dynamic systems), and a theoretical approach may be the only way to determine whether an agostic interaction is involved in a given reaction pathway. In such a case, a particular care should be taken in the choice of the level of theory. In the case of density functional theory calculations, the choice of the functional, the description of the metallic atom as well as the basis set are crucial in the geometrical parameters that will be obtained. Furthermore, in a rigorous theoretical characterization of agostic bondings, such geometry optimizations would only correspond to the first step of a more complete investigation involving either a description in terms of molecular orbitals, or a topological study. As an example, H.G. Cho has investigated the agostic structure of titanium methylidene hydride at 15 different levels of theories using Hartree-Fock, Density Functional Theory and Post-Hartree-Fock calculations. He has shown the importance of the choice of the basis set on the agostic distortions, and on the delocalization energies as calculated with the Natural Bond Orbital (NBO) method (vide infra).³³ The influence of the basis set superposition error (BSSE) was also underlined by several studies. As an example, the α -agostic interaction involved in $\text{Cp}^*\text{Rh}(\text{CO})(\eta^2\text{-alkane})$ was investigated at various levels of theories,^{34,35} and the lack of BSSE

correction in the conventional *ab initio* methods leads to an overestimation of the agostic bonding, while some DFT approaches underestimate this interaction.³⁶ Multi-configurational quantum chemical methods (complete active space self-consistent field CASSCF)/second-order perturbation theory (CASPT2)) is a powerful approach to compare the agosticity of several metallic center. Such an approach was used by Roos et al. to explain experimental results in the case of methyldiene metal dihydride complexes.³⁷

2. NBO

Agostic interactions may be seen as donor-acceptor systems, with an electron transfer from a σ bond of the ligand to an unoccupied orbital of the metallic atom. In this respect, techniques for the studies of molecular orbitals may shed some light onto agostic bondings. Natural bond orbital theory is a powerful approach to characterize donor and acceptor groups in a molecule, and to quantify delocalization energies. In the natural bond orbital approach, a suite of mathematical algorithms are used to diagonalize the density matrix generated during Hartree-Fock, Density Functional Theory or Post Hartree-Fock calculations. These mathematical transformations allow the generation of orbitals in terms of effective atom-like constituents within the molecular environment.^{38,39,40} This approach allows the study of delocalization from atomic orbitals. Toward the natural population analysis, the NBO approach offers a quantitative way to characterize the donor-acceptor relationship. An interesting example of the use of NBO for a quantitative study of agostic bondings is given by the work of Cho et al.^{33,41} Based on a systematical study, a methodology was proposed to analyze the C-H...M agostic bondings in terms of donor-acceptor analysis as defined in the NBO theory.⁴² Thus, NBO method provides a valuable way to evaluate the agostic character of interactions.⁴³

3. QTAIM

In addition with an orbital description of agostic interactions, topological tools were applied in an attempt to qualitatively and quantitatively investigate these bondings. Two topological approaches were particularly applied to the study of agostic bonding: the Bader's quantum theory of atom in molecules (QTAIM) and the electron localization function (ELF). These topological descriptions are based on the dynamical analysis of gradients, from the electronic density, or from functions of electronic density. A vector field is derived from gradients of a potential, and critical points are defined on the basis of the vector fields.

The QTAIM approach is based on a partition of the molecular space into non-overlapping regions within which the local virial theorem is fulfilled. This implies, that the kinetic energy of each of these regions has a definite value what is achieved if and only if the integral of the Laplacian of the electron density, $\nabla^2\rho(r)$, over each region vanishes, a condition which is fulfilled for boundaries which are zero-flux surfaces for the gradient of the density.⁴⁴ In this method, gradient fields of the electron density $\nabla\rho(r)$ are studied. Since an atomic center corresponds to a local maximum of electron density, each atomic center acts as an attractor, and field lines define a basin. Basins are separated from each other by zero-flux surfaces. On each zero-flux surface, gradient lines converge toward a critical point called a bond critical point (BCP). The presence of such a BCP is one of the criterion that was proposed by Popelier et al. to characterize an agostic bonding in the QTAIM framework.^{18,45} On the other hand, the identification of BCP may be difficult in the case of weak agostic interactions. In their systematic study of a set of 20 crystal structures potentially characterized by agostic bonding, Thakur and Desiraju noticed that NBO may be more relevant than QTAIM in describing these weak interactions.³³ Numerous studies have now proven that QTAIM can indeed describe β C-H agostic interactions,^{46, 47} but is not suitable for the study of α C-H agostic interactions or weaker interactions

such as C-C agostic bondings.⁴⁸ Indeed, the use of new local criteria seem to be the only way to detect weak agostic bonding by means of the QTAIM approach.⁴⁹

4. ELF

The ELF approach^{50,51,52} intends to provide a partition into basins of attractors which closely match the VSEPR electronic domains, or in other words Lewis's representation. The assumption that groups of electrons can be localized within space filling non-overlapping domains implies that the variances (the squared standard deviation) of the domain populations are minimal. It was pointed out that it is convenient to define a localization function $\eta(\mathbf{r})$ such as:⁵³

$$\frac{\delta \sigma^2(\bar{N})}{\delta V} = \oint n \cdot \nabla \eta(\mathbf{r}) ds = 0 \quad (1)$$

enabling to carry out the partition into basins of the attractors of the gradient dynamical system of $\eta(\mathbf{r})$. It was further proposed⁵⁴ to minimize the Frobenius norm of the covariance matrix:

$$\|\sigma^2(\Omega_A, \Omega_B)\|_F = (\sum_{A,B} (\sigma^2(\Omega_A, \Omega_B))^2)^{1/2} \quad (2)$$

where the matrix element $\sigma^2(\Omega_A, \Omega_B)$ is the covariance of the populations of the basins labelled Ω_A and Ω_B . The covariance operator is defined as:

$$\widehat{\sigma^2}(\Omega_A, \Omega_B) = \hat{N}(\Omega_A)\hat{N}(\Omega_B) - \bar{N}(\Omega_A)\bar{N}(\Omega_B) \quad (3)$$

with $\hat{N}(\Omega_A) = \sum_i^N \hat{y}(r_i)$ ($\hat{y}(r_i) = 1, r_i \in \Omega_A$, and $\hat{y}(r_i) = 0, r_i \notin \Omega_A$) where $\hat{N}(\Omega_A)$ is the population operator introduced by Diner and Claverie⁵⁵ and $\hat{N}(\Omega_A) = \int_{\Omega_A} \rho(\mathbf{r}) d\mathbf{r}$ is the basin population. The minimization is carried out with the help of the local covariance measure for σ spin electrons:

$$\xi_\sigma^h[\rho; r] = \rho_\sigma(r) \int (h_{\sigma,\sigma}(r, r'))^2 dr' \quad (4)$$

where $h_{\sigma,\sigma}(\mathbf{r}, \mathbf{r}')$ is the exchange hole, and can be approximated by an expression proportional to the kernel of the Becke and Edgecombe function.⁵⁰

The covariance can be written as the sum of four spin components, i.e.:

$$\sigma^2(\Omega_A, \Omega_B) = \sigma_{\alpha\alpha}^2(\Omega_A, \Omega_B) + \sigma_{\alpha\beta}^2(\Omega_A, \Omega_B) + \sigma_{\beta\alpha}^2(\Omega_A, \Omega_B) + \sigma_{\beta\beta}^2(\Omega_A, \Omega_B) \quad (5)$$

With

$$\sigma_{\alpha\alpha}^2(\Omega_A, \Omega_B) = \bar{\Pi}_{\alpha\alpha}(\Omega_A, \Omega_B) - \bar{N}_\alpha(\Omega_A)\bar{N}_\alpha(\Omega_B) \quad (6)$$

$$\sigma_{\alpha\beta}^2(\Omega_A, \Omega_B) = \bar{\Pi}_{\alpha\beta}(\Omega_A, \Omega_B) - \bar{N}_\alpha(\Omega_A)\bar{N}_\beta(\Omega_B) \quad (7)$$

$$\sigma_{\beta\alpha}^2(\Omega_A, \Omega_B) = \bar{\Pi}_{\beta\alpha}(\Omega_A, \Omega_B) - \bar{N}_\beta(\Omega_A)\bar{N}_\alpha(\Omega_B) \quad (8)$$

$$\sigma_{\beta\beta}^2(\Omega_A, \Omega_B) = \bar{\Pi}_{\beta\beta}(\Omega_A, \Omega_B) - \bar{N}_\beta(\Omega_A)\bar{N}_\beta(\Omega_B) \quad (9)$$

where $\bar{\Pi}_{\sigma\sigma'}(\Omega_A, \Omega_B) = \int_{\Omega_A} \int_{\Omega_B} \Pi_{\sigma\sigma'}(r_1; r_2) dr_1 dr_2$ is the number of $\sigma\sigma'$ electron pairs between Ω_A and Ω_B . The opposite spin contributions are expected to be negligible with respect to the same spin ones, and, in fact they are null in the Hartree-Fock approximation. Except for specific delocalized bonding situations such as charge-shift bonds, the covariance between non adjacent basins is negligible. The zero flux surfaces of the gradient field of the spin pair composition for which ELF is

an excellent approximation, provide a partition which nearly minimizes the covariance between basins limited a same separatrix.⁵⁶

The ELF partition yields basins of attractors clearly related to Lewis's model: core and valence basins. A core basin surrounds a nucleus with atomic $Z > 2$, it is a single basin for the elements of the second period or the union of the basins belonging to the inner shells for heavier elements. It is labelled $C(A)$ where A is the element symbol. In the study of systems involving transition metal elements it is often useful to consider independently the basins of the metal external core (subvalence) shell. The valence basins are characterized by the number of atomic valence shells to which they participate, or in other words by the number of core basins with which they share a boundary. This number is called the synaptic order. Thus, there are monosynaptic, disynaptic, trisynaptic basins, and so on. Monosynaptic basins, labeled $V(A)$, correspond to the lone pairs of the Lewis model, and polysynaptic basins to the shared pairs of the Lewis model. In particular, disynaptic basins, labeled $V(A, X)$ correspond to two-centre bonds, trisynaptic basins, labeled $V(A, X, Y)$ to three-centre bonds, and so on. The valence shell of a molecule is the union of its valence basins. As hydrogen nuclei are located within the valence shell they are counted as a formal core in the synaptic order because hydrogen atoms have a valence shell. For example, the valence basin accounting for a C–H bond is labeled $V(C, H)$ and called protonated disynaptic. The valence shell of an atom, say A , in a molecule is the union of the valence basins whose label lists contain the element symbol A .

The ELF population analysis provides not only the basin populations and the associated covariance matrix but also the probability of finding n electron in a given basin and the contribution of the QTAIM basins to the ELF basins. The contribution of the atomic basin of A to the ELF disynaptic basin $V(A,B)$ which is denoted by $N[V(A,B)|A]$ is evaluated by integrating the electron density over the intersection of the $V(A,B)$ basin and of the atomic basin of A . Raub and Jansen⁵⁷ have introduced a bond polarity index defined as:

$$p_{AB} = \frac{\bar{N}[V(A,B)|A] - \bar{N}[V(A,B)|B]}{\bar{N}[V(A,B)|A] + \bar{N}[V(A,B)|B]} \quad (10)$$

As an example, Fig. 2 displays the QTAIM and ELF basins as well as the ELF/QTAIM projection in the case of a simple molecule that is involved as a ligand in the following study, namely the NCCH_3 molecule. To obtain an optimal clarity, only two QTAIM atomic surfaces (H and N) are shown. It is interesting to note that there no ELF core basin for a hydrogen atom. We reported the ELF valence basin populations as well as the corresponding atomic contributions.




Figure 2: QTAIM and ELF basins for free NCCH_3 molecule.

IV. Topological descriptions, probabilistic approach and qualitative estimators

1. Context

ELF has proven to be a valuable tool for characterizing all kinds of chemical bonds. In the case of weak interactions such as 2c-3e and hydrogen bondings, the ELF analysis enables us to describe and classify them in terms of topological properties very close and similar to the traditional chemical concepts.

The analysis of basin population was helpful to distinguish between weak, medium and strong hydrogen bondings. It was shown that the core-valence bifurcation index is a suitable criterion to quantitatively describe these interactions.^{17,58}

In the case of 2c-3e bondings, no disynaptic basins are found. However, this is not the sole criterion to determine whether a bonding exists or not, and once again, core-valence bifurcation index has proved to be a very powerful tool. In addition, in this specific case, a topological delocalization index was helpful in quantifying the electron fluctuation.^{59,60}

These examples prove that, beyond the BCP, topological approaches can be used in a quantitative way to thoroughly characterize electron localization / delocalization involved in different types of interactions. Below we will present a similar approach to characterize agostic interactions.

2. Presentation of the approach

a. A statistical approach

The bonding in many molecular systems could not be fully described by considering only a perfect bonding electron localization following Lewis structure. It is principally due to the indiscernibility of the electron which implies an always present delocalization. This electronic delocalization is not observable, and therefore there is no experimental property that allows its *direct measurement*.

The multivariate analysis is a basic statistical method enabling one to reveal the correlations between different groups of data. It relies upon the construction of the covariance matrix elements defined by

$$\langle cov(i, j) \rangle = \langle ij \rangle - \langle i \rangle \langle j \rangle \quad (11)$$

where $\langle i \rangle$, $\langle j \rangle$, and $\langle ij \rangle$ are the averages of the data values and of their product. The diagonal elements of the covariance matrix are the variances $\sigma^2(i) = \langle i^2 \rangle - \langle i \rangle^2$ which measure the statistical dispersion of the data among the group. The standard deviations $\sigma(i)$ are the square root of the variances. The off-diagonal matrix elements indicate which data are involved in the delocalization.

The bonding in most molecular systems can be described by a strict localization of electron pairs. A more realistic picture, closely related to the concept of resonance, is provided by the superposition of electron multiplets distributed among the basins and therefore accounting for the electron delocalization. The population of a given basin Ω_A appears accordingly as the average of such n-tuplets weighted by the probability, $P_n(\Omega_A)$ of finding n electron in Ω_A :

$$\bar{N}(\Omega_A) = \sum_{n=1}^N n P_n(\Omega_A) \quad (12)$$

A measure of the delocalization is given by the covariance matrix whose diagonal matrix elements are the square of the standard deviations of the basin populations whereas the off matrix elements indicate which basins are involved in the delocalization.⁶¹

In our search of several parameters to qualitatively estimate the strength of agostic bondings, we studied a large number of systems. From this study, it was found that four parameters from QTAIM

and ELF taken together may allow a comparison of agostic bondings present in different systems between X-H bond (X = C, B) and a metallic center M:

- First of all, we will consider only hydrogen atoms as potentially involved in agostic interactions. In the Brookhart and Green's definition,^{6,7} even if this is not the only criterion, the presence of three atomic centers sharing two electrons is a fundamental aspect of an agostic bonding. Thus, in the topologic description, the total population of the protonated potentially agostic basin $V(H)$ is an obvious important parameter.
- The projection of ELF on AIM basins gives some information on the atomic contribution in the agostic protonated basin. Hereafter in this paper, this quantity will be labeled as $M/X/H$ in the case of X-H agostic bond (X = C or B). Hereafter, this information will be used as a clear indicator of the trisynaptic character of a protonated basin. It is worthy to note that an atomic contribution only makes sense if its value is larger than the numerical error, i.e. 0.01 electron.
- The covariance calculated from the ratio between the basin's population of the potentially agostic H atom and the population of the metallic center core basin $C(M)$ is an important parameter to characterize the interaction between these two basins $Cov(V(H)/C(M))$. The covariance thus obtained from ELF topological analysis gives some insights on the delocalization of electrons between the two atoms. Obviously this value depends on the theoretical description of the metallic center: the covariance is smaller when a pseudo potential is used for the metallic center. Taken together with the previous parameters, a covariance larger than 0.03 (in absolute value) is a proof of an agostic interaction between X-H σ bond and M.
- Furthermore, in the context of comparison between several levels of calculations, the σ^2 and the variance calculated with the ELF topological approach are an indication of deviation from perfect localization coming from inter-population. Similar values of variances for a same system calculated at different levels of theories thus allows to ensure that all the theoretical descriptions are consistent with each other.
- Furthermore, it was shown that the electron density (ρ) of the bond critical point (BCP) could be related to the bond order and thus the bond strength.⁶² We propose the use of three $\rho(\text{BCP})$ to gain some insight on the strength of the agostic interaction. In line with Popelier and Logothetis,¹⁸ we suggest the use of the $\rho(\text{BCP})$ of the $M - H_\beta$ bonding, when it exists. Additionally, the comparison of the $\rho(\text{BCP})$ values associated with the $C - H_{\text{agostic}}$ and $C - H_{\text{free}}$ allows to estimate the weakening of the $C - H_{\text{agostic}}$ bond caused by the agostic interaction. Two conditions are necessary for the use of the $\rho(\text{BCP}(C - H_{\text{agostic}}))$ and $\rho(\text{BCP}(C - H_{\text{free}}))$ values. First, the carbon bearing the hydrogen atom potentially involved in an agostic interaction should also bear an additional non-agostic hydrogen atom. This condition is often fulfilled. Furthermore, neither of the H_{agostic} and H_{free} atoms should be involved in another non-covalent interaction. Provided that these conditions are satisfied, we can propose the following reference values. In the alkyl complexes, the $\rho(\text{BCP}(C - H_{\text{free}}))$ are characterized by

$$0.28 \leq \rho(\text{BCP}(C - H_{\text{free}})) \leq 0.29$$

whereas the $C - H_{\text{agostic}}$ are characterized by smaller electron densities at the BCP, with

$$0.200 \leq \rho(\text{BCP}(C - H_{\text{agostic}})) \leq 0.27$$

From the ELF point of view, a protonated basin is considered as a trisynaptic basin when its population originates from three atomic centers. An H-agostic bond is a protonated trisynaptic basin whose population is around of $2 e^-$. Consequently, it corresponds to the traditional 3c-2e interaction in chemistry.

These parameters were not only chosen for their ability to describe an agostic interaction, but also for their phenomenological significances. Indeed, when the covariance $\text{Cov}(V(H)/C(M))$ increases, this signifies that the delocalization of electrons between the metallic center and the agostic protonated basin increases, concomitantly with a weakening of the bond between the hydrogen and other atom X. The H-X bond is thus activated and the agostic character of the interaction between H and M is stronger. In a limit case, when the agostic bonding becomes stronger and stronger, the ELF basin evolves from a trisynaptic to a disynaptic basin, and the covariance increases till the formation of a covalent M-H bond. This corresponds to a metallic hydride.

Moreover, the strength of an agostic H – X bond may be estimated from the QTAIM properties calculated at the H – X bond critical point (BCP). Interestingly, the charge density at the agostic H – X BCP compared to that of a non-agostic H – X bond in the complex (and/or compared to the charge density at the H – X BCP of free ligand) provides an indication to the strength of the agostic interaction.

b. Choice of a representative set of agostic bondings

When Popelier and Logothetis carried out their seminal work on the topological characterization of agostic bondings, they have chosen TiCl_2 -alkyl complexes as models, and they investigated agostic bondings by means of QTAIM approach¹⁸. Thus, these simple models already cover four types of potentially agostic bondings. The QTAIM method was helpful in characterizing an agostic bonding in the case of β C-H, γ C-H and C-C interactions with the metallic center, but not in the case of an interaction with an α C-H. These systems therefore appear as references in the context of setting a new theoretical approach for the characterization of agostic bondings.

From this study as well as from further theoretical investigations, it was found that QTAIM suitably describe β C-H and γ C-H agostic bondings, but not the α C-H ones, even in cases for which experimental data tend to prove that α C-H agostic bondings were indeed formed.

Some compounds were observed under different agostomers. We will choose a few examples of agostomers based on experimental investigations carried out by Baird et al.^{63,64} In the case of $[\text{Cp}_2\text{TiCH}_2\text{CHMe t-Bu}]^+$, they observed that the α -agostic isomer is preferentially formed although a β -agostic isomer could have been formed. Such a situation is relatively rare: when both α - and β -agostic isomers may be formed, generally the β -agostic form is more stable. Another alkyl-titanium complex that may exist as β - and γ -agostomers, will be considered.

The complex formed between acetonitrile and zero-valent nickel that may lead to the formation of a C-H agostic bonding⁶⁵ will be investigated.

In an attempt to understand the parameters that influence the formation of agostic bondings, the effects of co-ligands, small changes in the structures and the nature of the metallic center, will be investigated. The effect of the presence of a co-ligand will be topologically investigated based on the example of rhodium thiophosphoryl pincer complex studied by Milstein et al.⁶⁶

Different titanium complexes will be compared, based on the compounds studied by Popelier,¹⁸ Baird^{63,64} and Mc Grady⁶⁷. The comparison of complexes studied by Mc Grady⁶⁷ and Forster⁶⁸ will allow to study the influence of the nature of the metallic center. Further examples derived from the model compounds of Popelier¹⁸ as well as from Sabo-Etienne⁶⁹ will also be discussed.

Lastly, interactions involving heteroatoms will be considered. Indeed, such interactions are sometimes considered as "agostic" whereas some authors consider that all the donation from a σ bond and an unoccupied orbital of an hypovalent transition metal center cannot be classified under an unique appellation. Our aim here is to determine whether the methodology above presented is able to differentiate between these types of interactions, or whether these interactions are due to a similar effect. For this purpose, titanocene and zirconocene amidoborane complexes^{67,68},

dimethylaminoborane complexes,⁶⁹ and mesitylborane complexes,⁷⁰ will be considered. The last examples corresponds to intermolecular interactions, whereas in the other systems, the weak interaction between a σ X – H bond and the metal center should be considered as an intramolecular interaction.

V. Application to a representative set of examples

1. Influence of the level of theory

To begin with, the influence of the level of theory on the parameters calculated with the ELF topology will be checked. To this end, DFT calculations were carried out, and three different types of functionals were selected:

- B3LYP because this is one of the most popular hybrid functional,
- PBE0 because this is a non-empirical hybrid functional⁷¹ that was widely employed in the context of agostic interactions,
- TPSSh because this Meta-GGA hybrid functional can be used for reference calculations, when combined with a suitable basis set.⁷²

In combination with these functionals, six different basis sets were selected:

- the 6-311++G(2d, 2p) basis set, in combination with a pseudo-potential LANL-2TZ-f including a triple ξ and an additional diffuse f function for the metallic center,
- the 6-311++G(2d, 2p) basis set, in combination with a pseudo-potential LANL-2TZ-p including a triple ξ and an additional diffuse p function for the metallic center,
- the 6-311++G(2d, 2p) basis set, in combination with a pseudo-potential LANL-2DZ including a double ξ function for the metallic center,
- the 6-31++G(2d, 2p) basis set, in combination with a pseudo-potential LANL-2DZ including a double ξ function for the metallic center,
- the 6-311++G(2d, 2p) basis set, without any pseudo potential for the metallic center,
- the 6-31++G(3df, 3pd) basis set, without any pseudo potential for the metallic center.

For these tests of the influence of the levels of theories, a simple model molecule taken from the study of Popelier and Logothetis was selected (see below, $[\text{TiCl}_2\text{CH}_2\text{CH}_3]^+$).

In a first series of tests, the molecule was re-optimized at each level of theory prior to the topological investigation. In a second series of tests, the complex was optimized using the highest level of theory, namely B3LYP, PBE0, or TPSSh/6-31++g(3df, 3pd), and ELF calculations were then carried out on the wave function of the single point geometries using the LanL2DZ as basis set.

Optimized geometries are compared in Table 2 and the topological parameters are summed up in Table 3.


	Level of theory	d(Ti-C) (Å)	d(Ti-H _{agostic}) (Å)	θ (°)
	B3LYP/6-311++G(2d,2p)	2.008	2.034	113.8
B3LYP/6-311++G(3df,3pd)	2.008	2.030	113.8	
PBE0/6-311++G(3df,3pd)	1.986	2.015	114.3	
TPSSh/6-311++G(3df,3pd)	2.009	2.008	113.5	

Table 2: Distances (in Å) and angles (in °) for the $[\text{TiCl}_2\text{CH}_2\text{CH}_3]^+$ complex optimized at different levels of theory.

The values presented in the Table 2 clearly show that the geometry of the complex is correctly described even with the B3LYP functional, when used in combination with a relatively large basis set including 2d and 2p polarization functions. Indeed, if we compare the distances and the angles calculated at the B3LYP/6-311++G(2d,2p) level of theory with the reference values (calculations at the TPSSh/6-311++G(3df,3pd) level of theory), the errors are 0.001 Å, 0.026 Å and 0.3° for the d(Ti-C), d(Ti-H_{agostic}), and θ , respectively.

Level of theory	$V(H_{\beta})$	$Ti/C_{\beta}/H_{\beta}$	Cov (V(H _β)/C(Ti))		σ^2	d(Ti-H _{agostic}), θ , d(Ti-C)
B	Lan12DZ	1.89	0.04/0.82/1.03	-0.08	0.72	2.116, 115.6, 1.982
3	6-31++G(2d,2p)	1.95	0.08/0.78/1.09	-0.09	0.74	2.034, 113.8, 2.003
L	6-311++G(2d,2p)	1.93	0.08/0.76/1.09	-0.09	0.73	2.034, 113.8, 2.008
Y	6-31++G(2d,2p)/LanL2DZ	1.92	0.03/0.78/1.11	-0.09	0.72	2.025, 113.7, 1.994
P	6-311++G(2d,2p)/LanL2DZ	1.92	0.04/0.75/1.13	-0.09	0.73	2.005, 113.4, 1.996
	6-311++G(2d,2p)/LanL2TZ(P)	1.90	0.04/0.73/1.13	-0.10	0.72	2.009, 113.6, 2.004
	6-311++G(2d,2p)/LanL2TZ(F)	1.90	0.05/0.72/1.13	-0.10	0.72	2.005, 113.4, 2.002
P	6-31++G(2d,2p)	1.93	0.07/0.78/1.08	-0.09	0.74	2.019, 114.4, 1.983
B	6-311++G(2d,2p)	1.92	0.07/0.77/1.08	-0.10	0.74	2.018, 114.3, 1.986
E	6-31++G(2d,2p)/LanL2DZ	1.91	0.04/0.75/1.12	-0.10	0.73	2.002, 117.0, 1.975
0	6-311++G(2d,2p)/LanL2DZ	1.91	0.04/0.75/1.12	-0.10	0.73	1.985, 113.9, 1.977
	6-311++G(2d,2p)/LanL2TZ(P)	1.90	0.04/0.74/1.12	-0.10	0.74	1.990, 114.1, 1.985
	6-311++G(2d,2p)/LanL2TZ(F)	1.90	0.05/0.73/1.12	-0.10	0.73	1.989, 114.1, 1.982
T	6-31++G(2d,2p)	1.93	0.08/0.74/1.11	-0.10	0.73	2.016, 113.6, 2.006
P	6-311++G(2d,2p)	1.93	0.08/0.74/1.11	-0.10	0.72	2.011, 113.5, 2.009
S	6-31++G(2d,2p)/LanL2DZ	1.93	0.03/0.77/1.13	-0.10	0.73	2.001, 113.4, 1.995
S	6-311++G(2d,2p)/LanL2DZ	1.90	0.03/0.72/1.15	-0.09	0.72	1.980, 113.1, 1.996
H	6-311++G(2d,2p)/LanL2TZ(P)	1.91	0.04/0.72/1.15	-0.10	0.73	1.985, 113.3, 2.005
	6-311++G(2d,2p)/LanL2tz(F)	1.91	0.04/0.72/1.15	-0.10	0.72	1.985, 113.3, 2.002
S	B3LYP/LanL2DZ//	1.83	0.03/0.73/1.07	-0.08	0.72	Single-Point calculations
P	B3LYP/6-311++G(3df,3pd)					
	PBE0/LanL2DZ//	1.89	0.03/0.82/1.04	-0.08	0.73	
	PBE0/6-311++G(3df,3pd)					
	TPSSh/LanL2DZ//	1.89	0.02/0.81/1.06	-0.08	0.72	

Table 3: Topological and geometrical parameters obtained at several levels of theories. The total population of the agostic protonated basin $V(H_{\beta})$ and the contribution from Ti, C and H ($Ti/C_{\beta}/H_{\beta}$), the covariance $Cov(V(H_{\beta})/C(Ti))$ and the variance (σ^2) are given. Distances (in Å) and angles (in degrees) are given for the optimized structures.

Table 3 clearly shows that, all the selected levels of theory identify an agostic bonding in this simple TiCl₂-alkyl system. Indeed,

- the total population of the valence basin of H_{β} is slightly smaller than $2e^-$,
- the variance value (σ^2) does not depend on the level of theory which indicates the stability of the ELF topological procedure in partitioning of the molecular space,
- in all the cases, the contribution of the titanium atom to this basin is in the range of 0.07 – 0.08 e^- provided to use an explicit triple zeta quality basis set for metallic center,
- the calculated covariance is close to -0.1, clearly indicating a delocalization of the electrons between agostic hydrogen and metallic center.

These four criteria should be considered all together, the presence or the absence of one of the criterion is not sufficient to drive any conclusion.

We would like to emphasize the fact that, whatever the level of theory selected, the topological analysis of the agostic bonding leads to similar results. Thus, the criteria selected are robust toward the level of theory.

This study thus show that the B3LYP/6-311++G(2d,2p) can be used for a topological investigation of agostic interactions. This level of theory was selected for all the further studies presented below.

2. Study of different agostomers

Let's first consider alkyl titanocene compounds $[\text{Cp}_2\text{TiCH}_2\text{CHMe}(\text{t-Bu})]^+$ studied by Baird et al.⁶³ Two agostic isomers were characterized: α - and β -agostomers (Figure 3).




Figure 3: α - and β -agostomers observed for an alkyl titanocene compound.⁶³

The experimental study clearly show that, between two diastereoisomers α - and β -agostic, the α -agostic one is more stable. The Table 4 shows, without any surprise, that a BCP is indeed obtained in the case of the β -agostic isomer that is not the case for the α -agostic one. Nevertheless, it is worth noting that the electron density at the C-H bond slightly decreases by 0.05 *a.u.* due to the agostic deformation for both α - and β -agostic isomers. Simultaneously, the Laplacian of charge density ($\nabla^2\rho$) and the energy density (H) at the BCP increase which leading to the reduction of the covalent character of the C-H bond.

α -agostomer	β -agostomer
BCP(Ti-H α): does not exist	BCP(Ti-H β): $\rho=0.03$, $\nabla^2\rho=+0.11$, $H=0.00$ $\epsilon=0.37$
BCP(Ti-C α): $\rho=0.10$, $\nabla^2\rho=+0.07$, $H=-0.04$	RCP(Ti-H β -C β): $\rho=0.03$, $\nabla^2\rho=+0.13$
BCP(C α -H α): $\rho=0.24$, $\nabla^2\rho=-0.70$, $H=-0.23$	BCP(Ti-C α): $\rho=0.10$, $\nabla^2\rho=+0.03$, $H=-0.04$
BCP(C α -H): $\rho=0.29$, $\nabla^2\rho=-1.05$, $H=-0.31$	BCP(C α -H): $\rho=0.29$, $\nabla^2\rho=-1.04$, $H=-0.31$
BCP(C α -C β): $\rho=0.24$, $\nabla^2\rho=-0.51$, $H=-0.19$	BCP(C α -C β): $\rho=0.24$, $\nabla^2\rho=-0.49$, $H=-0.19$
BCP(C β -H): $\rho=0.29$, $\nabla^2\rho=-1.03$, $H=-0.30$	BCP(C β -H β): $\rho=0.24$, $\nabla^2\rho=-0.69$, $H=-0.22$

Table 4: Search for BCP's and RCP's in the α - and β -agostic alkyl titanocene compounds studied by Baird et al.⁶³

To further characterize these two isomers, the methodological approach above proposed was applied, and the Table 5 summarizes the ELF investigation of these two isomers.

Isomer		
	Ti-Alpha-Baird	Ti-Beta-Baird
V(H)	1.89	1.9
M/X/H	0.04/0.78/1.07	0.04/0.80/1.11
Cov(V(H)/ C(M))	-0.09	-0.06
d(Ti-H _{agostic}) (Å)	2.05	2.01
d(C-H _{agostic}) (Å)	1.14	1.17

Table 5: Topological characterization of the α - and β -agostic alkyl titanocene (Cp = Cyclopentadienyl) compounds studied by Baird et al.⁶³

The results presented in Table 5 clearly show that, despite the absence of BCP in the alpha agostomer, the agostic bonding is indeed described by the combined ELF/QTAIM studies. Indeed, the total population of the valence basin of H_α is below $2 e^-$, the contribution of the Ti atom to the protonated valence basin of H_α is 0.04, which is not negligible, and the covariance between two basins V_{H_α}/C_M is close to -0.1. In comparison, the total population of the agostic H atom is closer to $2 e^-$ and the covariance is slightly smaller (in absolute value) in the case of the β -agostic isomer. As a conclusion, the agostic character is slightly more pronounced in the α -agostic isomer compared with the β -agostic one, for this specific case. Furthermore, these two agostic bondings are relatively weak.

Baird et al.⁶⁴ also studied an alkyl-titanium complex that exist under two isomeric forms, one presenting a β -agostic bonding, and another presenting a γ -agostic bonding. Table 6 summarizes the ELF/QTAIM characteristics of these two isomers. As in the previous case, the statistical parameters of ELF clearly describe both agostic bondings. These bondings are characterized by BCP's and RCP's in the QTAIM description. The quantitative study of these bondings shows that the σ C-H interactions are relatively weak, as it was the case in the previous α - and β -agostic isomers.

Isomer		
	Ti-Beta-Si-Baird	Ti-Gamma-Si-Baird
V(H)	1.93	1.95
M/X/H	0.06/0.77/1.10	0.05/0.81/1.09
Cov(V(H)/ C(M))	-0.08	-0.07
d(Ti-H _{agostic}) (Å)	2.09	2.00
d(C-H _{agostic}) (Å)	1.16	1.15

Table 6: Topological characterization of the β - and γ -agostic alkyl titanium complexes studied by Baird et al.⁶⁴

As a conclusion, these examples show that the topological tools of the ELF approach indeed allows to characterize α -, β - and γ -agostic bondings.

3. Topological characterization of a representative set of C-H agostic bondings

A set of complexes was used to probe the agostic character of different features of the $M \cdots H-C$ bonding by means of the topological criteria previously presented, and the selected topological criteria are summarized in the Table 7.








Compound	 Ti-Popelier ¹⁸	 Ti-Alpha-Baird ⁶³	 Ti-Beta-Baird ⁶³	 EtTiCl ₃ (dmpe)-McGrady ⁶⁷
V(H) M/X/H Cov(V(H)/ C(M))	1.93 0.08/0.76/1.09 -0.09	1.90 0.04/0.79/1.07 -0.09	1.96 0.04/0.81/1.11 -0.06	1.97 0.03/0.94/1.00 -0.04
d(Ti-H _{agostic}) (Å) d(C-H _{agostic}) (Å) ω(C-H _{agostic}) (cm ⁻¹)	2.034 1.146 2534	2.049 1.140 2589	2.157 1.151 2429	2.183 1.115 2815
d(C _α -H _{free}) (Å) ω(C-H _{free}) (cm ⁻¹)	1.0874, 1.0874 3087 – 3172	1.085 3037 – 3134	1.083 3034 – 3197	1.083 3067 – 3177
QTAIM topological parameters for agostic compound: ρ, ∇ ² ρ, H(ρ) in a.u.				
BCP(C-H _β) BCP(C-H _{free}) BCP(Ti-H _β)	0.233, -0.67, -0.22 0.287, -1.07, -0.30 0.045, +0.14, 0.00	0.238, -0.70, -0.30 0.288, -1.05, -0.31 Does not exist	0.235, -0.69, -0.22 0.288, -1.05, -0.31 0.029, +0.11, 0.00	0.262, -0.86, -0.26 0.285, -1.03, -0.31 Does not exist
Some relevant parameters for free ligand:				
d(C-H) (Å) ω(C-H) (cm ⁻¹) BCP(C-H _β)	1.091 3034 - 3101 0.281, -1.01, -0.30	1.096 2991 - 3110 0.282, -1.01, -0.30		1.0908 3034 - 3101 0.281, -1.01, -0.30
Electron density at BCP(C-H) in function of d(C-H)				

Table 7: to be continued







Compound	 CpTiNiPr ₂ Cl ₂ -McGrady ⁶⁷	 Rh-butene-Milstein ⁶⁶	 Rh-H ₂ CO-Milstein ⁶⁶	 Ni-Jones ⁶⁵
V(H) M/X/H Cov(V(H)/ C(M))	2.06 0.01/1.07/0.98 -0.02	2.01 0.05/1.02/0.95 -0.15	2.12 0.23/0.93/0.96 -0.38	2.01 0.07/0.98/0.96 -0.13
d(Ti-H _{agostic}) d(C-H _{agostic}) ω(C-H _{agostic})	2.363 1.095 3006	1.939 1.131 2612	1.665 1.222 1950	1.811 1.126 2665
d(C _α -H _{free}) ω(C-H _{free})	1.088, 1.090, 1.090 3040 – 3130	1.088 3174 - 3192	1.088 3174 - 3182	1.088 3070 – 3119
QTAIM topological parameters for agostic compound: ρ, ∇ ² ρ, H(ρ) in a.u.				
BCP(C-H _{agostic}) BCP(C-H _{free}) BCP(M-H _{agostic})	0.287, -1.05, -0.30 0.284, -1.03, -0.30 Does not exist	0.250, -0.77, -0.24 0.281, -0.97, -0.28 0.055, +0.20, -0.01	0.202, -0.47, -0.16 0.282, -0.98, -0.85 0.107, +0.26, -0.05	0.256, -0.82, -0.25 0.285, -1.04, -0.30 0.057, +0.23, -0.01
Some relevant parameters for free ligand:				
d(C-H) ω(C-H) BCP(C-H _β)	1.093 3020 - 3040 0.288, -1.06, -0.31	1.092 3050 0.277, -0.94, -0.28	1.089 3060 - 3128 0.283, -1.03, -0.30	
Electron density at BCP(C-H) in function of d(C-H)	Borderline case: If there was an agostic case, It should correspond to a very weak interaction!			

Table 7: Topological characterization of a set of M···H-C complexes reported in the literature as agostic compounds. Optimized geometries have been found at the B3LYP/6-311++G(2d,2p) level of theory.

For all the species the total population of the valence basin of the agostic hydrogen atom is in the $2 \pm 0.15 e^-$ range. Both the covariance $\text{Cov}(V(H)/C(M))$ and the atomic contributions in the valence basin of H (M, X, H) fluctuate depending on the complex.

Table 7 presents complexes containing intramolecular σ C-H agostic bonding. However, the Rh-H₂CO-Milstein⁶⁶ and Rh-butene-Milstein⁶⁶ complexes could not be easily classified as β agostic species. We note in passing with these two examples that a change in the nature of a co-ligand may strongly affect the agostic character of a C-H bonding. Indeed a weak agostic character is topologically predicted in the case of the Rh-butene-Milstein⁶⁶ complex, whereas a strong one is predicted for the analogous compound containing H₂CO instead of butene as co-ligand. These two cases will be further discussed hereafter.

A close look of the data reported in table 7 some trends which be summarized as follows:

- In view of the metal contribution in the protonated basin, we can classify the complexes in four categories: (1) $M = 0.01 e^-$ corresponding to an undefined case, (2) $0.01 < M < 0.05$ for weak - medium agostic bonding, (3) $0.05 < M < 0.20$ for medium - strong agostic interaction, and (4) $M > 0.20$ for almost pre-dissociated C - H_{agostic} or pre-hydride H_{agostic} - M.
- One can note that the calculated harmonic vibrational frequency of the C - H_{agostic} oscillator is always red-shifted with respect to that of C - H_{free}. To a certain extent, the amount of this red-shift reflects the strength of the agostic interaction. It is interesting to note that the harmonic vibrational frequency of C - H_{free} is sometimes blue-shifted with respect to the C - H frequency in the free ligand. However, on the ground of C-H vibrational frequency red-shift one can easily distinguish three categories of agostic species: red-shift $\approx 2\%$ for the weakest agostic compound (CpTiNiPr₂Cl₂-McGrady), red-shift $\approx 40\%$ for the strongest agostic compound (Rh-H₂CO-Milstein), and $5\% < \text{red-shift} < 40\%$ for the other compounds going from weak to strong cases.
- Case of CpTiNiPr₂Cl₂-McGrady compound: This compound has been previously considered as an agostic case by McGrady et al.⁶⁷ and also Scherer and coworkers.⁷³ We would like to emphasize that the very low metal contribution (0.01 e) in the population of V(H) makes actually impossible to decide the presence or absence of an agostic interaction, because of the numerical uncertainty of our ELF analysis which is just equal to 0.01 e.

This is also consistent with the geometrical properties of the complex: if we refer to the criteria summarized in the Table 1, the CpTiNiPr₂Cl₂-McGrady compound is anagostic ($d(M-H) > 2.3 \text{ \AA}$).

In order to check the possible effect of the dispersion contribution in the electronic structure, we also optimized the studied structure using two hybrid functionals (wB97XD and B2PLYPD3) which are suitable to treat the very weak non-covalent interactions.

As shown by the results reported in Table 8, topological differences between the C _{β} -H _{β} so-called agostic bond and C _{β} -H_{free} within the same compound are minor so that we can confidently exclude a dominant agostic interaction within this complex. This description is also supported by a very small vibrational $\omega(\text{C-H})$ frequency shift (less than 2%) with respect to free $\omega(\text{C-H})$. Furthermore, we found a BCP between the so-called agostic H atom and one of the two chlorine atoms. At this latter BCP, the electron density is equal to 0.015 e belonging to the hydrogen bonded range.⁷⁴

	wB97XD	B2PLYPD3	B3LYP
$d(\text{C}_{\beta}\text{-H}_{\beta})$ (Å)	1.096	1.095	1.095
$\omega(\text{C-H})$ (cm ⁻¹)	3035		3006
BCP(C _{β} -H _{β}): $\rho, \nabla^2\rho$ in a.u.	0.286, -1.04	0.287, -1.05	0.284, -1.03

$d(C_{\beta}-H_{\text{free}})$ (Å)	1.091	1.090	1.090
$\omega(C-H)$ (cm^{-1})	3077		3040
BCP($C_{\beta}-H_{\text{free}}$): ρ , $\nabla^2\rho$ in a.u.	0.290, -1.067	0.291, -1.08	0.287, -1.05

Table 8: Effect of the density functional on the characterization of the CpTiNiPr₂Cl₂-McGrady complex using the 6311++G(2d,2p) basis set. No BCP(Ti-H _{β}) was found for any of the three functional.

- Agostic bonding and QTAIM bond critical point: we note that the presence of a BCP(H-Ti) in the case of EtTiCl₃(dmpe)-McGrady compound actually depend on the level of theory. Indeed, we found a BCP only at BP86/6-311++G(d,p) level, whereas there is no BCP(H-Ti) when we use B3LYP, PBE0 or BP86 with 6-311++G(2d,2p) as basis set.
- Concerning the strength of an agostic bonding, we note that the electron density at the BCP(C-H agostic) decreases when the agosticity increases. This trend is graphically shown for each compound in Table 7. A global linear regression graph for all the species will be discussed in section VI.

4. Topological characterization of a representative set of B-H agostic bondings

In Table 9 are gathered five complexes containing intramolecular σ B-H agostic bondings. These complexes have been subject of experimental and/or theoretical study.^{65,67,68}







Compound	 ZrCINH ₂ BH ₃ -a-Forster ⁶⁸	 ZrCINH ₂ BH ₃ -b-Forster ⁶⁸	 TiHNBH ₂ BH ₃ -a-McGrady ⁶⁷	 TiHNBH ₂ BH ₃ -b-McGrady ⁶⁷	 TiCp ₂ NH ₂ BH ₃ -McGrady ⁶⁷
V(H)	1.89	1.92	1.89	1.85	1.92
M/X/H	0.07/0.22/1.59	0.07/0.27/1.58	0.07/0.24/1.58	0.07/0.26/1.52	0.12/0.23/1.57
Cov(V(H)/C(M))	-0.13	-0.13	-0.14	-0.14	-0.15
dM-H _{agostic}	2.030	2.039	1.892	1.855	1.892
d(B-H _{agostic})	1.302	1.276	1.300	1.310	1.299
ω (B-H _{agostic})	1932	2084	1959	1814	1895
d(B-H _{free})	1.203	1.203	1.205	1.204	1.204
ω (B-H _{free})	2488 – 2542	2487 – 2539	2474 – 2526	2480 – 2530	2477 – 2524
QTAIM topological parameters for agostic compound: ρ , $\nabla^2\rho$, H(ρ) in a.u.					
BCP(B-H _{agostic})	0.122, +0.02, -0.11	0.132, 0.00, -0.13	0.123, +0.02, -0.11	0.122, -0.01, -0.11	0.126, -0.01, -0.12
BCP(B-H _{free})	0.175, -0.27, -0.20	0.174, -0.26, -0.19	0.170, -0.21, -0.18	0.171, -0.21, -0.18	0.174, -0.26, -0.19
BCP(M-H _{agostic})	0.057, +0.11, -0.01	0.056, +0.11, -0.01	0.059, +0.11, -0.01	0.062, +0.13, -0.01	0.057, +0.16, -0.01
Some relevant parameters for free ligand:					
d(B-H) (Å)	1.208				
BCP(B-H _{β})	0.171, -0.24, -0.19				
Electron density at BCP(B-H) in function of d(B-H)					

Table 9: Topological characterization of a set of M...H-B complexes reported in the literature as agostic compounds. Optimized geometries have been found at the B3LYP/Def2-TZVPD level of theory.


For all the amidoborane titanocene or zirconocene complexes, where the nitrogen atom is at the α -position and the boron atom at β -position, the presence of a B-H protonated basin containing a

metallic contribution ranging from 3% to 6% of the $V(H)$ population is a clear indicator for the existence of a β -agostic bonding. This is supported by the bond lengthening and frequency red-shift of the $B-H_{\text{agostic}}$ bond. Compared to the $C-H$ agostic bonding, the $B-H$ agosticity should be considered as medium to strong interaction. This consideration is naturally in line with the decrease of the electron density at the $B-H_{\text{agostic}}$ bond critical point. It is graphically evidenced on the linear regression graph (Table 9).

Globally, the ELF/QTAIM criteria lead to an homogeneous and consistent description of the bondings thus supporting the use of "agostic" for both $\sigma C-H \cdots M$ and $\sigma B-H \cdots M$ intramolecular bonding.

5. Parameters influencing the agostic character of bondings

To further investigate $\sigma C-H$ β -agostic interactions and the parameters affecting such bondings, an interesting case-study of rhodium thiophosphoryl pincer investigated by Milstein et al. will be detailed below. Indeed, the authors reported the formation of identical agostic bondings with Rh that also interacts either with but-2-ene or with formaldehyde. The agostic hydrogen atoms are characterized by largely different geometrical parameters, and the topological investigation proves that the nature of the co-ligand R is of paramount importance on the agostic interaction (See Scheme 1).



Scheme 1: Rhodium thiophosphoryl pincer studied by Milstein et al, R = but-2-ene or formaldehyde.⁶⁶

To better understand the co-ligand influence we will carefully analyze and compare the topological properties of the Rh-butene-Milstein compound to those of Rh- H_2CO -Milstein. In Tables 10 and 11 are reported the most relevant topological properties of both Rh-butene-Milstein and Rh- H_2CO -Milstein complexes.

H-agostic compound	QTAIM properties ⁽¹⁾	ELF properties ⁽²⁾
Pincer (S-C-S)-R: R = Cis-2-butene	BCP($H\beta$, Rh): Yes BCP($C\alpha$, Rh): Yes BCP($C\beta$, Rh): No RCP(Rh, $H\beta$, $C\beta$, $C\alpha$): Yes	$V(Rh, C\beta, H\beta)$: Yes $V(Rh, C\alpha)$: No $V(Rh, C\beta)$: No $V(C\alpha, C\beta, Rh)$: Yes
	BCP(C(Ligand), Rh): Yes BCP(C(Ligand), Rh): Yes RCP(Rh, C, C): Yes	$V(C(Ligand), Rh)$: Yes $V(C(Ligand), Rh)$: Yes
Pincer (S-C-S)-R: R = OCH_2	BCP($H\beta$, Rh): Yes BCP($C\alpha$, Rh): No BCP($C\beta$, Rh): No RCP(Rh, $H\beta$, $C\beta$, $C\alpha$): No	$V(Rh, C\beta, H\beta)$: Yes $V(Rh, C\alpha)$: No $V(Rh, C\beta)$: No $V(C\alpha, C\beta, Rh)$: Yes

	BCP(O(Ligand), Rh): Yes	V1(O(Ligand), Rh): Yes V2(O(Ligand), Rh): No
--	-------------------------	---

Table 10: ELF and QTAIM topological features of the agostic bondings for the Rh-H₂CO-Milstein and Rh-butene-Milstein compounds.⁶⁶

(1) BCP and RCP are the bond and ring critical points corresponding to the (3,-1) and (3,+1) critical points. (2) V(X,Y) and V(X,Y,Z) stand for the disynaptic and trisynaptic basins which share two and three core basins, respectively.

Agostic compound	QTAIM properties ⁽¹⁾ $\rho, \nabla^2\rho, H, \epsilon$	ELF properties ⁽²⁾ M, C and H contributions
Pincer (S-C-S)-R: R = Cis-2-butene Rh-butene-Milstein ⁶⁶	BCP(H β , Rh): 0.055; +0.204, -0.008, 0.29	V(Rh, C β , H β): 0.05/1.02/0.95
	BCP(C α , Rh): 0.050, +0.141, -0.008, 0.12	V(C α , C β , Rh): 1.25/0.91/0.06
	RCP(Rh, H β , C β , C α): 0.045, +0.172, -0.004, No ϵ	
Pincer (S-C-S)-R: R = OCH ₂ Rh-H ₂ CO-Milstein ⁶⁶	BCP(C(Ligand), Rh): 0.102, +0.192, -0.034, 1.13	V(C(Ligand), Rh): 0.54/0.23
	BCP(C(Ligand), Rh): 0.111, +0.158, -0.043, 0.37	V(C(Ligand), Rh): 0.63/0.21
	RCP(Rh, C, C): 0.100, +0.297, -0.027, No ϵ	
Pincer (S-C-S)-R: R = OCH ₂ Rh-H ₂ CO-Milstein ⁶⁶	BCP(H β , Rh): 0.107, +0.255, -0.046, 0.26	V(Rh, C β , H β): 0.23/0.93/0.96
	BCP(O(Ligand), Rh): 0.086, +0.488, -0.011, 0.46	V1(O(Ligand), Rh): 2.31/0.03 V2(O(Ligand), Rh): 2.81/0.00

Table 11: The quantitative topological characteristics of different types of agostic bondings.

(1) The four QTAIM characteristics at a critical point are given by ρ (the charge density), $\nabla^2\rho$ the Laplacian of charge density), H (the energy density) and ϵ (the ellipticity). Note that we have only the first three characteristics at a RCP. These quantities are given in atomic units. (2) The X/Y/Z contributions are the atomic contributions in the averaged population of the V(X, Y, Z) basin. These numbers are in electrons.

A close look of the data reported in tables 10 and 11 allows us to summarize the similarities and differences between the titled complexes as follows:

- As it concerns the weak interactions, despite overall agreement between the topologies of ELF and QTAIM, few differences however have been underlined.^{49, 75,76,77,78} Nevertheless, we would like to emphasize that we believe in the complementarity of these two methods, rather than mutual exclusion. But having said that, we remember the readers that there is no bond critical point between H α and metallic center, while we have a trisynaptic protonated basin accounting for the α -agostic interaction. Topological analysis of the C-C bonding - bond between the carbon of methyl and that of aryl - obtained from both QTAIM and ELF methods are likewise complementary and often clarify each other. In both Rh-butene-Milstein and Rh-H₂CO-Milstein pincer complexes the C-C bond valence basin is indeed a trisynaptic basin with a non-negligible contribution from the metallic center (0.06 and 0.09 e⁻) which is a clear indication on the η^3 -C-C-

H agostic compound. This conclusion supports the analysis advanced in the paper of Milstein and coworkers.⁶⁶ As for the QTAIM analysis, it gives a BCP between the rhodium and C atom of aryl group in the case of the Milstein pincer bonded with cis-2-butene, while there no such a BCP for the other species. This means that the agostic interaction in the Rh-butene-Milstein could be referred to as traditional β -agostic compound, while this not the case for the Rh-H₂CO-Milstein pincer complex.

- The ligand effect is another striking feature of these pincer-R (R = cis-2-butene or OCH₂) agostic compounds. It is worth noting that both QTAIM and ELF topologies provide the same analysis for the ligand effect.
 - In the case of the cis-2-butene ligand, rhodium thiophosphoryl pincer cation involves in the formation of two non-equivalent disynaptic basins labeled as V(C(Ligand), Rh) in Table 10 and 11. The metal atom contribution amounts to 33% and 42% of the total averaged population of these metal-ligand bonds. These basins clearly are indicative of the formation of two metal-carbon coordinate covalent bonds. In parallel, we found two bond critical points for two Rh-C bonding and a RCP in the center of C-Rh-C triangle. The non-negligible negative values of the energy density at the BCPs (-0.034 and -0.043) clearly indicate the non-negligible covalency of these bonds. As a consequence, the formation of these coordinate covalent bonds enriches the valence shell of metal leading to lower acidity. Accordingly, the agostic interaction is relatively weak (as shown by values presented in table 7, with a small covariance V(H)/C(M), and a small contribution of the metal in the valence basin of the agostic hydrogen atom). This weak agostic interaction is geometrically confirmed by a small H_{agost} – C distance and a large H_{agost} – Rh distance.
 - Contrarily, in the case of the formaldehyde-rhodium thiophosphoryl pincer cation, the interaction between oxygen and rhodium atoms is almost weak – manifested by small contribution of Rh in the one of the two valence basins of oxygen – and thus without noticeable change in the valence shell of metal. Thus the electron-deficiency of the Rh center is not counterbalanced by a notable electron transfer. As a consequence, the agostic bonding between Rh and H will be enhanced. Geometrically, the H_{agost} – Rh distance will be smaller than in the previous case, and the H_{agost} – C distance will be larger, thus leading to a more pronounced activation of the H – C bond.

Thus, the presence of a co-ligand can be of paramount importance in the activation of a C-H bond by means of the formation of an agostic bonding, and a topological description of the systems may help in understanding these differences.

On the other hand, a change in the nature of the co-ligand do not necessary lead to a fundamental change in the agostic character of the bonding. As an example, Table 12 allows to follow the geometry and the topological characterization of the valence basin of the agostic hydrogen atom in β - and γ -agostic alkyl titanium complexes studied by Baird et al.⁶⁴ The initial agostomers contain two cyclopentadienyl ligands. We suggest the substitution of a cyclopentadienyl ligand either by formaldehyde or by a chlorine atom to not lead to a huge distortion of the agostic bonding: H_{agost} – Ti and H_{agost} – C distances are almost unchanged (see Table 11). As far as the topological description of the valence basin of the agostic hydrogen atom is concerned, almost no change is observed, neither in the contribution of the metal center to the protonated valence basin nor in the covariance values. Thus, in this case, the substitution of cyclopentadienyl ligand by formaldehyde or by chlorine atom do not affect the agostic character of the bonding.



Isomer	Characterization		
R ¹ = R ² = Cp	V(H)	1.93	1.95
	Ti/C/H	0.06/0.77/1.10	0.05/0.81/1.09
	Cov(V(H)/ C(M))	-0.08	-0.07
	$d_{Ti-H}^{agostic}$ (Å)	2.09	2.00
	$d_{H-C}^{agostic}$ (Å)	1.16	1.15
R ¹ = Cp R ² = OCH ₂ (TRANS)	V(H)	1.98	1.99
	Ti/C/H	0.04/0.86/1.08	0.03/0.88/1.07
	Cov(V(H)/ C(M))	-0.07 (AIM : -0.06)	-0.06 (AIM : -0.05)
	$d_{Ti-H}^{agostic}$ (Å)	2.12	2.10
	$d_{H-C}^{agostic}$ (Å)	1.13	1.12
R ¹ = R ² = OCH ₂	V(H)	Starting from β isomer, it coverges to γ agostomer.	1.99
	Ti/C/H		0.06/0.85/1.07
	Cov(V(H)/ C(M))		-0.09 (AIM : -0.07)
	$d_{Ti-H}^{agostic}$ (Å)		2.00
	$d_{H-C}^{agostic}$ (Å)		1.13
R ¹ = Cp R ² = Cl (TRANS)	V(H)	1.98	1.97
	Ti/C/H	0.08/0.80/1.09	0.06/0.84/1.08
	Cov(V(H)/ C(M))	-0.09 (AIM : -0.07)	-0.08 (AIM : -0.06)
	$d_{Ti-H}^{agostic}$ (Å)	2.06	2.01
	$d_{H-C}^{agostic}$ (Å)	1.14	1.14
R ¹ = R ² = Cl	V(H)	Starting from β isomer, it coverges to γ agostomer.	1.98
	Ti/C/H		0.08/0.84/1.06
	Cov(V(H)/ C(M))		-0.09 (AIM : -0.07)
	$d_{Ti-H}^{agostic}$ (Å)		1.99
	$d_{H-C}^{agostic}$ (Å)		1.15

Table 12: Topological characterization of the β - and γ -agostic alkyl titanium complexes as studied by Baird et al.⁶⁴ and differently substituted.

To further investigate the parameters influencing the formation of agostic bondings, we studied the influence of the metallic center on the agosticity. The β model compound of Popelier and Logothetis was chosen. Table 13 shows that the substitution of the titanium atom by either Zr or Hf do not affect the geometry of the agostic bonding, and the topological description of the valence basin of the agostic hydrogen atom remains similar. Thus, in some cases, and specifically in the case of titanocene compounds, the substitution of the metallic center by an atom belonging to the same chemical family, does not affect the agostic character of the bonding.



	 β model compound (Popelier & Logothetis)	Topological description	
	Geometry	ELF	AIM
M = Ti	$d(\text{H}\beta\text{-C}\beta) = 1.152$ $d(\text{M-C}\alpha) = 2.010$ $d(\text{M-C}\beta) = 2.393$ $d(\text{M-H}\beta) = 2.009$ $a(\text{M-C}\alpha\text{-C}\beta) = 84.3$ $a(\text{C}\alpha\text{-C}\beta\text{-H}\beta) = 113.5$	$V(\text{C}\beta\text{-H}\beta) = 1.93$ Ti(0.08)/C(0.75)/H(1.10) $\text{Cov}(V(\text{H}\beta)/\text{C}(\text{M})) = -0.10$	$Q(\text{H}\beta) = -0.10$ $Q(\text{Ti}) = +1.9$ $\text{Cov}(\text{Ti}/\text{H}\beta) = -0.08$
M = Zr	$d(\text{H}\beta\text{-C}\beta) = 1.156$ $d(\text{M-C}\alpha) = 2.148$ $d(\text{M-C}\beta) = 2.553$ $d(\text{M-H}\beta) = 2.149$ $a(\text{M-C}\alpha\text{-C}\beta) = 86.3$ $a(\text{C}\alpha\text{-C}\beta\text{-H}\beta) = 113.9$	$V(\text{C}\beta\text{-H}\beta) = 1.94$ Zr(0.05)/C(0.74)/H(1.15) $\text{Cov}(V(\text{H}\beta)/\text{C}(\text{M})) = -0.10$	$Q(\text{H}\beta) = -0.15$ $Q(\text{Zr}) = +2.6$ $\text{Cov}(\text{Zr}/\text{H}\beta) = -0.08$
M = Hf	$d(\text{H}\beta\text{-C}\beta) = 1.157$ $d(\text{M-C}\alpha) = 2.140$ $d(\text{M-C}\beta) = 2.545$ $d(\text{M-H}\beta) = 2.155$ $a(\text{M-C}\alpha\text{-C}\beta) = 85.9$ $a(\text{C}\alpha\text{-C}\beta\text{-H}\beta) = 114.5$	$V(\text{C}\beta\text{-H}\beta) = 1.93$ Hf(0.04)/C(0.71)/H(1.18) $\text{Cov}(V(\text{H}\beta)/\text{C}(\text{M})) = -0.09$	$Q(\text{H}\beta) = -0.18$ $Q(\text{Hf}) = +2.8$ $\text{Cov}(\text{Hf}/\text{H}\beta) = -0.07$

Table 13: Influence of the substitution of the metallic center in the case of the β model compound of Popelier and Logothetis.¹⁸

6. Characterization of double $\sigma\text{-BH}_2$ and $\sigma\text{-CH}_2$ interactions with a metallic center

The term "agostic bonding" was also used in the literature to describe situations in which a small molecule is in interaction with a metallic complex by means of two simultaneous weak interactions.^{26, 69,70} Sabo-Etienne, Alcaraz et al. are particularly active in the study of such intermolecular interactions. To complete our topological study of agostic bondings, we applied our methodology to intermolecular interactions involving dimethylaminoborane⁶⁹ and mesitylborane.⁷⁰

The complexes involved intermolecular interactions involving dimethylaminoborane and studied by Sabo-Etienne et al. are presented in Table 14.

				
Nature of M and X	Ru, H	Ru, Cl	Os, H	Os, Cl
$d_{\text{H1-M}}^{\text{agostic}}$ (Å)	1.85	1.62	1.81	1.62
$d_{\text{H1-B}}^{\text{agostic}}$ (Å)	1.30	1.55	1.37	1.95
$d_{\text{H2-M}}^{\text{agostic}}$ (Å)	1.85	2.00	1.81	1.97
$d_{\text{H2-B}}^{\text{agostic}}$ (Å)	1.30	1.27	1.37	1.29
Agostic character	agostic	H ₁ : strongly agostic	agostic	H ₁ : hydride

		H ₂ : weakly agostic		H ₂ : agostic
--	--	---------------------------------	--	--------------------------

Table 14: Characterization of the intermolecular interactions between ruthenium- and osmium-containing complexes and dimethylaminoborane investigated by Etienne-Sabo et al.⁶⁹

In the case of the osmium-containing complexes, an interaction is observed between the σ B-H bond and the metallic center when X = H. The substitution of this hydride by X = Cl dramatically affect the intermolecular interaction. The H atom *trans* to the chlorine atom leads to the formation of an hydride and the bond between H₁ and B is broken. On the other hand the σ B – H agostic interaction with the metal is maintained for the H₂ atom.

In the case of the ruthenium-containing complexes, an interaction is similarly observed between the σ B-H bond and the metallic center when X = H. As in the case of the osmium-containing complex, the substitution of the X = H atom by a Cl atom causes a distinction between the H₁ and H₂ atoms: The H atom *trans* to the Cl atom leads to the formation of a stronger interaction with the metal, whereas the interaction between H₂ and B becomes weaker.

For the present study, we selected the osmium-containing complexes for a topological investigation. Indeed, the two osmium-containing complexes allowed us to compare our quantitative approach with the strength of the interaction experimentally observed. Furthermore, these examples gave the opportunity that the topological descriptors herein chosen correctly discriminate a strong σ bond interaction and the formation of a hydride. For a sake of comparison, we also investigate the same system with M = Fe. Table 15 summarizes the topological data obtained for the intermolecular interactions of the four complexes.

In the case of the osmium-containing complexes, the topological criteria selected for the present study are indeed consistent with the formation of an agostic bonding, even if the total population of the valence basin of the hydrogen atom is a little bit high for a 3c-2e interaction.

As expected, H₁ and H₂ atoms are equivalent, and the agostic interaction is relatively strong compared with what was expected in the case of intramolecular interactions.

When the X = H atom is substituted by a chlorine atom, the topological description is fully consistent with what is expected from the data available in the literature. In the case of the H₁ atom, the total population of the basin is 1.67 e⁻ and the boron atom is not involved in this basin, which signifies that an hydride is formed, as already reported in the literature⁶⁹ for this case. On the other hand, the topological description of the H₂ atom is consistent with the formation of an agostic interaction. The total population of the valence basin of the hydrogen atom is 2.01 and both Os and B are involved in this basin. The covariance Cov(V(H)/C(M)) as well as the small contribution of the metal in the valence basin of H₂ is clearly consistent with a weaker agostic interaction compared with what is observed for the complex with X = H.

Thus, the topological description herein proposed is qualitatively and quantitatively consisted with what is already reported for these osmium-containing complexes.

When the osmium atom is replaced by Fe, the total population of the valence basin of the hydrogen atoms interacting with the metallic center increases: a total population of 2.35 is calculated when X = H, whereas slightly smaller populations are obtained when X = Cl (2.33 and 2.29 e⁻).

For the " Fe₂H- MeB" compound, both H₁ and H₂ atoms are identical, and the contribution of the metallic center to the valence basin of the hydrogen atom is quite large (0.30 e⁻), concomitantly with a large contribution of the boron atom (0.54). Furthermore, the covariance Cov(V(H)/ C(M)) is large, thus suggesting that the agostic character of this compound is quite large.

When X = H is substituted by X = Cl, the two H₁ and H₂ atoms are characterized by quantitatively different interactions with the metallic center. Contrary to what was observed in the case of the osmium, the agostic interaction of the two H₁ and H₂ atoms is conserved. The agostic character of the

hydrogen atom *trans* to the chloride atom is slightly reinforced, whereas the agostic character of the hydrogen atom *cis* to the chloride atom is slightly weakened.

In the case of a ruthenium-containing dimethylaminoborane compound,⁷⁰ a similar topological description is calculated. Once again, the total population of the valence basin of the hydrogen atom is relatively high, but the contribution of the metallic atom in this basin and the covariance $\text{Cov}(V(\text{H})/C(\text{M}))$ are fully consistent with the description of an agostic bonding.

During the discussions it is clear that the chlorine substituted compounds leads to the formation of a hydridic bond only when the metallic atom is an osmium atom, whereas a strong agostic interaction remains the final product with iron and ruthenium atoms.

Thus, from a topological point of view, these intermolecular interactions are similar to the intramolecular, agostic interactions, and there is no topological reason for not using the same name, "agostic".

Compound					
	Os,H-MeB-Alcaraz ⁶⁹	Os,Cl-MeB-Alcaraz ⁶⁹	Fe,H-MeB	Fe,Cl-MeB	Ru,dimethyl amino-borane-Alcaraz ⁷⁰
V(H)	2.25	H ₁ : 1.67 H ₂ : 2.00	2.35	H ₁ : 2.32 H ₂ : 2.28	2.16
M/B/H	0.18/0.59/1.48	H ₁ : 0.39/0.01/1.27 H ₂ : 0.04/0.42/1.54	0.30/0.54/1.51	H ₁ : 0.35/0.46/1.51 H ₂ : 0.22/0.52/1.54	0.18/0.59/1.37
Cov(V(H)/C(M))	-0.31	H ₁ : -0.40 H ₂ : -0.17	-0.31	H ₁ : -0.37 H ₂ : -0.29	-0.33
$d_{H1-M}^{agostic}$ (Å)	1.81	1.62	1.69	1.62	1.87
$d_{H1-B}^{agostic}$ (Å)	1.37	1.95	1.27	1.29	1.28
$d_{H2-M}^{agostic}$ (Å)	1.81	1.97	1.69	1.78	1.87
$d_{H2-B}^{agostic}$ (Å)	1.37	1.29	1.27	1.24	1.28

Table 15: Intermolecular interactions involving dimethylaminoborane⁶⁹ and mesitylborane⁷⁰

In an attempt to better understand the formation of such intermolecular agostic bondings, model systems were studied:

1. BH₂Cl forming simultaneously two interactions with Ru(PH₃)Cp,
2. an isomer of the previous model system in which the BH₂Cl molecule only forms one σ B-H interaction with the metallic center,
3. an analog of the first model system in which BH₂Cl is substituted by CH₂Cl₂. The formation of such complexes was proposed in the literature^{70,79} and was reported with Li as a metallic center,⁸⁰ but not, to our knowledge, in the case of transition metal complexes. On the other hand, cases in which the three hydrogen atoms of a CH₃-R group are simultaneously involved in agostic bondings with a same metallic center, were reported or proposed in the literature.^{81,82} Thus, a topological description of multiple σ C-H intermolecular agostic bondings is needed.

The topological criteria obtained for these three systems are presented in the Table 16.




Compound			
	Model system - 1	Model system - 2	Model system - 3
V(H)	2.26	2.07	2.16
M/B/H	0.22/0.56/1.45	0.19/0.43/1.44	0.04/0.19/0.93
Cov(V(H)/ C(M))	-0.41	-0.35	-0.14
$d_{H-M}^{agostic}$ (Å)	1.73	1.73	2.03
$d_{H-B \text{ or } C}^{agostic}$ (Å)	1.34	1.35	1.12

Table 16: Model compounds used for the study of intermolecular interactions


As already noted in the case of compounds reported in the Table 15, relatively large values of total population of the hydrogen basins are observed in the case of the model systems 1 and 3. This corresponds to cases for which the molecule forms simultaneously two σ interactions with the metallic center. In the case of the Model system - 1, the topological criteria suggest the formation of two identical and relatively strong agostic bondings. A similarly strong interaction is calculated in the isomeric system forming only one σ B-H interaction with the metallic center (Model system 2). On the other hand, in this case the total population of the valence basin is not particularly high, thus suggesting that the relatively large values of V(H) reported in tables 15 and 16 are a specific signature of double intermolecular interactions.

In the case of the Model system - 3, both of the hydrogen atoms involved in a σ interaction with the metallic center are identical. They are characterized by a lower total population of the valence basin of the hydrogen atoms, in comparison with the Model system - 1. Furthermore, the contribution of the metallic center in these valence basins are particularly weak. This may explain why the experimental formation, the isolation and the characterization such systems may be difficult.

VI. Discussion

1. Identification of the existence or non-existence of an agostic bonding

To begin with, the Scheme 2 summarizes the conditions that should be fulfilled to conclude that an agostic bonding exists between the C-H bond and the metallic center.



Scheme 2: Determination of the existence or non-existence of an agostic bonding between a C-H bond and a metallic center.

2. A qualitative comparison of different agostic bondings

To further evaluate the capability of the above presented statistical approach to qualitatively characterize the agostic bondings, our theoretical approach will be compared with experimental data. The first necessary step for such a comparison was obviously to find an experimental criteria that correctly describe the agostic character of the bonding:

- around different metallic centers,
- involving totally different compounds, and not focused in a well-defined chemical family of compounds,
- for α -, β -, γ -agostomers,
- involving σ C-H and σ B-H agostic bonding, with the possible presence of heteroatoms,
- with constrained geometries in the case of pincer ligands and bi-agostic compounds.

a. Which kinds of experimental data are available to classify the agostic bondings depending on the strength of the interaction?

Prior to introduce the parameters that we chose as an "universal" measurement of the agostic character of the bonding, we would like to briefly overview the experimental approaches currently available of the characterization of such interactions.

The experimental approaches generally used to characterize agostic bondings include NMR shifts, vibrational $\nu(\text{C-H})$ shifts (IR spectroscopy) and X-ray structures.^{7, 18,19, 20,83} Obviously, the changes in

reactivity due to the formation of an agostic bonding, and more precisely the activation of the σ bond involved in the interaction with the metallic center, is in itself an evidence that such an interaction exists. More rarely EPR spectroscopy and visible/UV spectroscopy also proved, in some specific cases, to be able to describe the formation of an agostic bonding.^{84,85} These methods will be shortly described below and in the Table 17, in the context of the study of agostic bondings.

Experimentally, the formation of an agostic bonding is distinguished first and foremost by an activation of the σ bond interacting with the metallic center.⁸⁶ In a same chemical family of compounds, it may be possible to qualitatively estimate the agostic character of the bonding, by comparing their reactivity toward a same reagent. On the other hand, such a qualitative approach will be limited to a specific chemical family of compounds and may depend on the reagents. As a conclusion, such an approach, if fully relevant in a screening approach for the most suitable complex in a given reaction process, will not allow to determine "absolute" agostic character of the bonding.

Spectroscopic methods that were applied to the study of agostic bondings almost cover the whole electromagnetic spectrum, from NMR to X ray spectroscopy.

The comparison of NMR shifts will not allow to compare agostic bondings of different chemical nature. Furthermore, only the strongest agostic bondings can be characterized by "classical" one-dimensional NMR experiments. Indeed, all the agostic bondings except the strongest ones are flexible in solution in the timescale of the NMR experiment, thus obscuring the effect.^{87,88}

Complementarily to the NMR spectroscopy, EPR spectroscopy was also successfully applied to the characterization of agostic bondings.⁸⁵ However, since this method is restricted to the study paramagnetic compounds, EPR spectroscopy is far from an "universal" experimental method that may be used to compare the agostic bonding of a wide set of complexes.

Another spectroscopic approach that was used to experimentally characterize agostic bonding is infrared (IR) spectroscopy.⁸⁸ Once again, the characterization of weak interactions in mixtures that may contain agostic and anagostic isomers may be difficult. An elegant procedure to characterize agostic interactions by means of IR spectroscopy, particularly applied by Andrews et al.,⁸⁹ consists in a coupling with the matrix isolation technique. Species to be analyzed are diluted in an inert medium in the gas phase, and further condensed on surface maintained at cryogenic temperatures. However, such a characterization is far from an easy tool that may be routinely used, without speaking on the experimental difficulty to generate a cryogenic matrix in which an agostic isomer is isolated. On the other hand, vibrational frequencies may be calculated by ab initio or DFT procedures once the geometry of the agostic compound is accurately determined.

Complexes of the transition metals are often colored. When the formation of an agostic bonding affects the energies of the d orbitals of the metallic center (especially the HOMO and the LUMO levels), visible / UV spectroscopy may be used to further characterize these interactions. Molybdenum β - and γ -agostomers were recently characterized by this technique.⁸⁴ However the characterization of agostic bondings by visible / UV spectroscopy can only be applied in very specific cases for which the interactions between the metallic center and the σ bond affect the color of the species.

Since an agostic bonding should lead to a distortion of the geometry of a compound, the crystalline X-ray structure is an obvious characterization tool.⁹⁰ The low scattering factor for hydrogen atoms and the difficulty to localize hydrogen in the vicinity of a metallic center are some of the limits of the method, in addition with the fact that agostic bondings may exist as intermediate species in liquid solution during reaction processes. In these latter cases, agostomers may not be obtained in crystalline form. Even if limited to agostomers that may be isolated as crystals, and despite its limitations X-ray diffraction structures were often reported in the literature for compounds involving agostic bondings. Furthermore, other experimental techniques provide data that may be further used for theoretical studies of the species. Thus geometries of agostomers can be obtained from DFT calculations and post-Hartree-Fock approaches. Theoretical calculations also allow to determine spectroscopic data that may be compared with experimental values. It is then possible to determine structures for these compounds. From a geometrical point of view, the formation of an agostic bonding will affect:

- the angles between the atoms,
- the $H_{\text{agost}}-M$ distance,
- the $H_{\text{agost}}-X$ distance.

Method	Effect leading to a possible characterization of an agostic bonding	General values expected in the case of an agostic bonding (and "normal" values expected for all other cases)	Limitations of the method
Reactivity	Activation of the C-H (or X-H, X = heteroatom such as B) bond.	Change in the reactivity. ^{14, 86}	The reactivity will obviously depend on co-ligands, geometry and the exact chemical nature of agostic bonding.
NMR	Redistribution of bonding-electron density in the formation of an agostic bonding.	$\delta = -5$ to -15 ppm, $^1J_{\text{C-H}_{\text{agostic}}} = 75$ to 100 Hz for C-H agostic bonding ($^1J_{\text{C}_{\text{sp}^3}-\text{H}_{\text{free}}} = 128$ MHz) ^{7-18,91, 83} Downfield paramagnetic shifts in the 700 - 1100 ppm range for axial ligands. ⁹²	NMR spectra cannot be obtained for dynamical systems.
EPR spectroscopy	Influence on the Zeeman electronic effect of the distortion of the geometry.	Variations in the values of g's and determination of ΔH° and ΔS° . ^{85, 93}	Limited to paramagnetic compounds for which the single electron is involved in the agostic bonding.
IR spectroscopy	Weakening of the C-H bond leading to a reduction in frequency for $\nu(\text{C-H})$ vibrational mode.	$\nu(\text{C-H}) = 2300 - 2700 \text{ cm}^{-1}$ (to be compared with the $\approx 2700 - 3000 \text{ cm}^{-1}$ range for free ligands). ^{18,19, 20}	Difficulty to identify a small weak signal that may overlap with other signals.
Visible / UV spectroscopy	Valence electron transition that may be affected by the formation of the M...H agostic bonding.	Depends on the crystal field perturbation induced by the agostic bonding. ⁸⁴	Limited to cases for which the agostic bondings affect the color of the compound
X-ray diffraction	Geometrical distortion.	Distance $H_{\text{agost}}-M = 1.8-2.3 \text{ \AA}$ ($2.3-2.9 \text{ \AA}$). ^{7,94} Distance $H_{\text{agost}}-X$ larger than the corresponding value	Difficulty to localize hydrogen atoms. Limited to agostomers that may be isolated as crystals.

		for the free ligand. Change in the angles between the atoms. ⁷	
--	--	---	--

Table 17: Comparison of different experimental approaches that may be used to characterize agostic bondings.

b. Choice of representative experimental criteria that may describe the strength of agostic bondings

Among above-mentioned characterizations, only vibrational frequencies and geometrical parameters can easily be obtained for all the types of agostic bondings, either directly experimentally, or by means of a combined experimental and theoretical investigation.

The (harmonic) vibrational shift of the H-X stretching mode, $\Delta \omega(\text{H-X})$, caused by the agostic bonding, is an obvious tool to investigate the strength of the interaction. This shift is defined as follows:

$$\Delta \omega(\text{H-X}) = \omega(\text{H-X})^{\text{free ligand}} - \omega(\text{H-X})^{\text{agostic compound}}$$

The shift may depend on the nature of the X atom. In order to compare agostic bondings involving chemically different X atoms, we suggest to use a normalized parameter:

$$\frac{\Delta \omega(\text{H-X})}{\omega(\text{H-X})^{\text{free ligand}}}$$

Additionally, geometrical parameters such as angles and distances were already proposed to characterize agostic bondings.⁷ The angles between the atoms will obviously depend on the system, and will not be directly linked to the agostic character of the bonding.

As already mentioned, the values of the angles cannot be used to compare the agostic character of bondings, in the set of compounds chosen for the present study, because their initial geometries are too much different from each other (Figure 1). The $H_{\text{agost}}\text{-M}$ distance will decrease, concomitantly with the increase of the $H_{\text{agost}}\text{-M}$ distance, during the formation of an agostic bonding. On the other hand, these two distances will be strongly affected by the nature of the X atom. In a first approximation, we can suggest that the ratio between $d_{\text{H-X}}^{\text{agostic}}$ and $d_{\text{H-M}}^{\text{agostic}}$, $\frac{d_{\text{H-X}}}{d_{\text{H-M}}}$ will not strongly depend on the nature of X.

Thus, two normalized experimental parameters may be used to compare the strength of agostic bondings, $\frac{d_{\text{H-X}}}{d_{\text{H-M}}}$ and $\frac{\Delta \omega(\text{H-X})}{\omega(\text{H-X})^{\text{free ligand}}}$. A simpler way to compare agostic bondings in a same family of complexes involving only one chemical type of H-X as agostic bondings, is to directly use the $d_{\text{H-X}}^{\text{agostic}}$ and $\Delta \omega(\text{H-X})$ experimental parameters.

c. May statistical parameters obtained from the ELF/QTAIM study be used to qualitatively estimate the agostic bonding strength?

Once it was checked that the criteria summarized in the Scheme 2 are consistent with the existence of an agostic bonding, several statistical parameters appear to be particularly relevant to qualitatively compare the strength of agostic bondings.

First of all, the contribution of the metallic center in the protonated agostic $V(\text{H})$ basin, $M_{V(\text{H})}$, is an obvious important parameter.

On the other hand, this parameter depends on the whole structure of the metallic complex. As a consequence, a direct comparison of the $M_{V(\text{H})}$ values will not allow to rigorously quantify the agostic character of bondings. To overcome this limit, we propose to introduce a normalized parameter, the

normalized metal contribution in the V(H) basin ($N_{M_{V(H)}}$). Its value is calculated by the ratio between the metal contribution in the V(H) basin and the total population of the V(H) basin:

$$N_{M_{V(H)}} = \frac{M_{V(H)}}{V(H)} \times 100 \quad (13)$$

The main advantage of this normalized parameter toward the $M_{V(H)}$ value, is to take into account the variation of the number of electrons in the V(H) basin that may be induced by other atoms in the vicinity of H.

Figure 4 shows the variations of $M_{V(H)}$ and $N_{M_{V(H)}}$ as a function of the $\left(\frac{d_{H-X}}{d_{H-M}}\right)$ ratio. The last point in the plots of the metal contribution corresponds to the hydride atom. This point was not included in the curve of the covariance. For all the other studied systems, all the values were taken into account in the plots.

As already pointed out, the complexes selected for the present study cover a wide range of chemical families, owing to the nature of the metallic center, the nature of the ligand involved in the agostic bonding, the nature of co-ligands and the geometry of the complexes. Because of these chemical differences, a relative spread of the of the $M_{V(H)}$ and $N_{M_{V(H)}}$ was expected. Given this point, a satisfactory linear correlation is observed for $M_{V(H)}$ and $N_{M_{V(H)}}$ with the ratio of the distances. The coefficient of the linear regression clearly confirm that $N_{M_{V(H)}}$ is a suitable criterion to compare agostic bondings in compounds belonging to different chemical families. Furthermore, the linear dependence observed demonstrates that $N_{M_{V(H)}}$ can indeed be used to qualitatively characterize the agostic character of the bonding.




Figure 4: Variation of the $M_{V(H)}$ and $N_{M_{V(H)}}$ parameters as a function of the $\frac{d_{H-X}}{d_{H-M}}$ ratio, with X = B or C, and $N_{M_{V(H)}} = \frac{M_{V(H)}}{V(H)} \times 100$.

As mentioned earlier, the electron density at the bond critical point of the X-H bond is also an indication of the σ -donation to the metallic center and thus of the strength of the agostic bonding. Once again, if directly used, this parameter will depend on the system under investigation. As a consequence, the direct comparison of the $\rho(\text{BCP}_{X-H})$ may be a suitable way to classify the strength of agostic bondings for a series of complexes belonging to a same chemical family of compounds, whereas the definition of a normalized value may be useful to compare agostic bonding in a set of complexes belonging to different chemical families. To this extent, we suggest to use the ratio between the $\rho(\text{BCP})$ for the X-H bond in the agostic complex and the $\rho(\text{BCP})$ for the X-H bond in the free ligand:

$$\frac{\rho(\text{BCP}(X - H_{\text{agostic}}))}{\rho(\text{BCP}(X - H_{\text{free}}))}$$

Figure 5 illustrates the possibility of using:

- direct H-X distances to characterize the strength of agostic bondings inside a same chemical family of compounds;
- reduced frequencies of the H-X stretching mode to characterize the strength of agostic bondings bonding in a set of complexes belonging to different chemical families;
- reduced $\rho(\text{BCP})$ for the H-X bond to classify the interaction with a metallic center.




Figure 5: Variation of the reduced $\frac{\rho(\text{BCP}(X - H_{\text{agostic}}))}{\rho(\text{BCP}(X - H_{\text{free}}))}$ QTAIM parameter as a function of the: A) d_{H-X} distance, and B) reduced frequencies of the H-X stretching mode. The strength of the agostic bonding increases from left to right and from top to bottom in both of the graphs.

Finally, the X-H bonds can be divided into four categories, depending on the strength of the M--H agostic bonding probably formed with a metallic center. Once the existence of the agostic bonding is ascertained based on criteria presented in the Scheme 2, four main estimators can be used to evaluate the strength of the interaction:

- two experimental parameters, $\frac{d_{H-X}}{d_{H-M}}$ and $\frac{\Delta \omega(H-X)}{\omega(H-X)_{\text{free ligand}}}$; alternatively these parameters may be obtained by geometry optimization and theoretical calculations;
- the $N_{M_{V(H)}}$ normalized parameter, based on ELF/QTAIM calculations;
- the $\frac{\rho(\text{BCP}(X-H_{\text{agostic}}))}{\rho(\text{BCP}(X-H_{\text{free}}))}$ normalized QTAIM parameter.

In Table 18, a global classification is proposed to estimate the strength of an agostic bonding, based on these four parameters.

Characterization of the M--H interaction Criterion	Anagostic	Weak-Medium agostic	Medium-Strong agostic	Pre-hydride
$\frac{d_{H-X}}{d_{H-M}}$	< 0.5	0.5 - 0.7	0.65 - 0.8	> 1
$\frac{\Delta \omega(H-X)}{\omega(H-X)_{\text{free ligand}}}$	≥ 0.90	0.90 - 0.70		< 0.65
$N_{M_{V(H)}}$	< 1	1 - 5	5 - 15	> 20
$\frac{\rho(\text{BCP}(X-H_{\text{agostic}}))}{\rho(\text{BCP}(X-H_{\text{free}}))}$	≥ 0.90	0.90 - 0.70		< 0.65

Table 18: A qualitative classification of the strength of agostic bonding based on three criteria.

VII. Conclusion

Herein we presented a topological quantitative investigation of σ C-H interactions. It was shown that the agostic character of the bonding can be compared even in the case of

- complexes around a different metallic center,
- complexes containing different ligands and co-ligands,
- α -, β -, γ -, and η^3 -CCH agostomers,
- mono- and bis-agostic species,
- inter- and intramolecular agostic bonding.

Thus the present method makes it possible to compare the agostic character of interactions that take place in systems that do not belong to a same chemical family, by means of a simple topological approach. The large set of compounds presented in this article shows that the methodology proposed here can lead to an unambiguously determination of the agostic character, both qualitatively and quantitatively of a wide range of chemical compounds.

Furthermore, the topological description of intramolecular σ C-H and σ B-H interactions, as well as intermolecular σ B-H interactions, is similar, thus suggesting that the use of the same "agostic" term for all these interactions, is appropriate.

It has been evidenced that any X-H...M agostic bond could be fully analyzed using three topological descriptors obtained easily with ELF method within the free TopMod code. We have to pay a particular attention on the stability and quality of the wave function derived from an optimization procedure. Indeed, the ELF topological analysis is actually an a posteriori approach whose reliability depend on that of wave function in hand.

The most relevant descriptors are:

1. The existence of a protonated trisynaptic basin, labeled as $V(H_{ag})$,
2. Three contributors participate on the averaged population of this basin: M/X/H,
3. The first and second descriptors are actually similar. We suggest them explicitly in order to underline the trisynaptic character of this basin which accounts for the 3c-2e interactions. Nevertheless, we also suggest the use of dimensionless descriptors to qualitatively estimate the strength of agostic bondings:
 - the ratio between the metal contribution and the total population of the protonated basin, in order to emphasize the importance of the relative metal contribution. This $\mathcal{N}_{M_{V(H)}} = \frac{MV(H)}{V(H)} \times 100$ ratio allows to quantitatively classify the agostic character of bondings;
 - the ratio of the electron densities at the BCP, $\frac{\rho(\text{BCP}(X - H_{agostic}))}{\rho(\text{BCP}(X - H_{free}))}$ in order to take into account the weakening of the X-H bond due to the interaction with the metallic center.
4. We suggest to use the above-mentioned descriptors in combination with two other criteria, ideally derived from experimental data. Indeed, experimental or, if not available, theoretical data, may be used to determine two additional reduced parameters that suitably describe the strength of agostic bondings:
 - the weakening of the H-X bond due to the agostic interaction is characterized by a shift in the frequencies of the vibrational stretching mode of the H-X bond. The $\frac{\Delta \omega(H-X)}{\omega(H-X)_{\text{free ligand}}}$ reduced frequency parameter may be obtained from experimental IR spectra, or from (harmonic) calculations;
 - the $\frac{d_{H-X}}{d_{H-M}}$ reduced distance may be obtained or derived from X-ray structures. This reduced distance is increasing within the strength of the agostic bonding.
5. The covariance between two basins: the agostic protonated basin and the metallic core basin $C(M)$. This quantity provide a measure of association between two quantities: $C(M)$ and $V(H_{ag})$. For agostic bondings for which the ratio $\mathcal{N}_{M_{V(H)}}$ is similar, the agostic character of the bondings increase with the $|C(M)|$.

Finally, we would like to emphasize the mutually complementary aspect of the topological results obtained with QTAIM and ELF analysis.

VIII. Computational details

The ELF calculations have been done using TopMod software.²⁴ ⁹⁵ Furthermore, the AIMAll software was used for the quantitative study of the topological QTAIM data.⁹⁶

All the necessary wfn files for the topological investigations have been obtained using the Gaussian 09 RevD.01 quantum chemical package. Calculations have been performed using different density functionals (B3LYP, PBE0 and TPSSh) as well as several basis sets (6-31++G(2d,2p), 6-311++G(2d,2p), TZVP, and LanL2DZ) as implemented in the Gaussian 09 package. In addition, the Def2-TZVP, Def2-TZVPD, LanL2TZ(P), and LanL2TZ(F) basis sets of Ahlrichs et al.,⁹⁷ obtained from “EMSL Basis Set Exchange Library”.⁹⁸ Furthermore, in order to take into account the dispersion

contribution in some model compounds (mono- and bi-agostic model structures), geometry optimizations have been done using wB97XD range separated hybrid functional⁹⁹ (Table 16) and also with B2PLYPD3 double hybrid functional including the D3 version of Grimme's dispersion with the original D3 damping function.¹⁰⁰ In order to ensure that an optimized structure corresponds to a true minimum, a frequency analysis was performed. Concerning the geometrical structures of the studied complexes, additional data on the optimized geometries are available in the Supporting Information.

¹ J. Chatt and J. M. J. Davidson, The tautomerism of arene and ditertiary phosphine complexes of ruthenium(0), and the preparation of new types of hydrido-complexes of ruthenium(II). *J. Chem. Soc.*, 1965, 843-855.

² A. H. Janowicz and R. G. Bergman, Carbon-hydrogen activation in saturated hydrocarbons: direct observation of $M + R-H \rightarrow M(R)(H)$. *J. Chem. Soc.*, 1982, **104**, 352-354.

³ W. D. Jones, A. J. Vetter, D. D. Wick and T. O. Northcutt, Alkane Complexes as Intermediates in C-H Bond Activation Reactions. In *Activation and Functionalization of C-H Bonds*, ed. K. I. Goldberg and A. S. Goldman, ACS Symposium Series 885, Washington, DC, 2004, ch. 3, pp 56-69.

⁴ M. Brookhart, M. L. H. Green and R. B. A. Parry, Two-electron, three-centre carbon-hydrogen-cobalt bonds in the compounds $[Co(\eta-C_5Me_4R)(\eta-C_2H_4)(\eta-C_2H_4-\mu-H)]BF_4$, R= Me and Et. *J. Chem. Soc. Chem. Commun.*, 1983, 691-693.

⁵ M. Brookhart and M. L. H. Green, Carbon-hydrogen-transition metal bonds. *Journal of Organometallic Chemistry*, 1983, **250**, 395-408.

⁶ M. Brookhart, M. L. H. Green and L. L. Wong, Carbon-Hydrogen-Transition Metal Bonds. *Prog. Inorg. Chem.*, 1988, **36**, 1-124.

⁷ M. Brookhart, M. L. H. Green and G. Parkin, Agostic interactions in transition metal compounds. *P.N.A.S.*, 2007, **104**, 6908-6914.

⁸ R. H. Crabtree, Organometallic alkane CH activation. *Journal of Organometallic Chemistry*, 2004, **689**, 4083-4091.

⁹ B. Rybtchinsk, L. Konstantinovskiy, L. J. W. Shimon, A. Vigalok and D. Milstein, Solvent-Stabilized Alkyrhodium(III) Hydride Complexes: A Special Mode of Reversible C-H Bond Elimination Involving an Agostic Intermediate. *Chemistry - A European Journal*, 2000, **6**, 3287-3291.

¹⁰ I. Omae, Agostic bonds in cyclometallation. *Journal of Organic Chemistry*, 2011, **696**, 1128-1145.

¹¹ E. Clot, O. Eisenstein, T. Dubé, J.W. Faller and R.H. Crabtree, Interplay of Weak Interactions: An Iridium(III) System with an Agostic tert-Butyl but a Nonagostic Isopropyl Group. *Organometallics*, 2002, **21**, 575-580.

¹² T. Ziegler, The 1994 Alcan Award Lecture Density functional theory as a practical tool in studies of organometallic energetics and kinetics. Beating the heavy metal blues with DFT. *Can. J. Chem.*, 1995, **73**, 743-761.

¹³ H. H. Brintzinger, D. Fischer, R. Mrlhaupt, B. Rieger, R. M. Waymouth, Stereospecific Olefin Polymerization with Chiral Metallocene Catalysts. *Angew. Chem. Int. Ed. Engl.*, 1995, **34**, 1143-1170.

¹⁴ M. Albrecht, Cyclometalation Using d-Block Transition Metals: Fundamental Aspects and Recent Trends. *Chem. Rev.*, 2010, **110**, 576-623.

¹⁵ J. Mathew, N. Koga and C.H. Suresh, C-H Bond Activation through σ -Bond Metathesis and Agostic Interactions: Deactivation Pathway of a Grubbs Second-Generation Catalyst. *Organometallics*, 2008, **27**, 4666-4670.

¹⁶ M. Lein, J. A. Harison and A. J. Nielson, Identification of non-classical C-H...M interactions in early and late transition metal complexes containing the $CH(ArO)_3$ ligand. *Dalton Trans.*, 2013, **42**, 10939-10951.

¹⁷ F. Fuster and B. Silvi, Does the topological approach characterize the hydrogen bond? *Theor. Chem. Acc.*, 2000, **104**, 13-21.

¹⁸ P. L. A. Popelier and G. Logothetis, Characterization of an agostic bond on the basis of the electron density. *Journal of Organometallic Chemistry*, 1998, **555**, 101-111.

- ¹⁹ W. Scherer and G.S. McGrady, Agostic Interactions in d⁰ metal alkyl complexes. *Angew. Chem. Int. Ed.*, 2004, **43**, 1782-1806.
- ²⁰ M. Lein, Characterization of agostic interactions in theory and computation. *Coordination Chemistry Reviews*, 2009, **253**, 625-634.
- ²¹ M. Etienne and A.S. Weller, Intramolecular C-C agostic complexes: C-C sigma interactions by another name. *Chem. Soc. Rev.*, 2014, **43**, 242-259.
- ²² W. H. Bernskoetter, C. K. Schauer, K. I. Goldberg and M. Brookhart, Characterization of a rhodium(I) σ -methane complex in solution. *Science*, 2009, **326**, 553-556.
- ²³ J. A. Calladine, S. B. Duckett, M. W. George, S. L. Matthews, R. N. Perutz, O. Torres and K. Q. Vuong, Manganese alkane complexes: an IR and NMR spectroscopic investigation. *J. Am. Chem. Soc.*, 2011, **133**, 2303-2310.
- ²⁴ S. Lachaize and S. Sabo-Etienne, σ -Silane Ruthenium Complexes: The Crucial Role of Secondary Interactions. *Eur. J. Inorg. Chem.*, 2006, 2115-2127.
- ²⁵ G. Alcaraz and S. Sabo-Etienne, NMR: A good tool to ascertain σ -silane or σ -borane formulations? *Coord. Chem. Rev.*, 2008, **252**, 2395-2409.
- ²⁶ J. F. Hartwig, C. N. Muhoro and X. He, Catecholborane bound to titanocene. Unusual coordination of ligand σ bonds. *J. Am. Chem. Soc.*, 1996, **118**, 10936-10937.
- ²⁷ A. T. Çolak, O. Z. Yeşilel and O. Büyükgüngör, A novel anagostic Csingle bondH...Cu interaction and clustered water molecules in copper(II)-pyridine-2,5-dicarboxylate complex. *Journal of Molecular Structure*, 2011, **991**, 68-72.
- ²⁸ H. V. Huynh, L. R. Wong and P. S. Ng, Anagostic Interactions and Catalytic Activities of Sterically Bulky Benzannulated N-Heterocyclic Carbene Complexes of Nickel(II). *Organometallics*, 2008, **27**, 2231-2237.
- ²⁹ M. G. D. Holaday, G. Tarafdar, A. Kumar, M. L. P. Reddy and A. Srinivasan, Exploring anagostic interactions in 5,15-porphodimethene metal complexes. *Dalton Trans.*, 2014, **43**, 7699-7703.
- ³⁰ M. L. Yang, Y. A. Zhu, C. Fan, Z. J. Sui, D. Chen and X. G. Zhou, Density functional study of the chemisorption of C1, C2 and C3 intermediates in propane dissociation on Pt(111). *Journal of molecular catalysis A: Chemical*, 2010, **321**, 42-49.
- ³¹ G. Papoian, J.K. Nørskov and R. Hoffmann, A comparative theoretical study of the hydrogen, methyl, and ethyl chemisorption on the Pt(111) surface. *J. Am. Chem. Soc.*, 2000, **122**, 4129-4144.
- ³² T. Jacob and W. A. Goddard III, Agostic Interactions and Dissociation in the First Layer of Water on Pt(111). *J. Am. Chem. Soc.*, 2004, **126**, 9360-9368.
- ³³ H. G. Cho, Agostic Structure of Titanium Methylidene Hydride (CH₂=TiH₂): A Theoretical Investigation for the Elusive Intramolecular Interaction. *Bulletin of the Korean Chemical Society*, 2009, **30**, 1631-1633.
- ³⁴ S. Niu and M. B. Hall, Theoretical Studies on Reactions of Transition-Metal Complexes. *Chem. Rev.*, 2000, **100**, 353-406.
- ³⁵ M. D. Su and S. Y. Chu, Theoretical Model for Insertion of the 16-Electron Species (η^5 -C₅H₅)M(L) into Saturated Hydrocarbons. A (η^5 -C₅H₅)M(CO) + CH₄ (M = Ru⁻, Os⁻, Rh, Ir, Pd⁺, Pt⁺). *Organometallics*, 1997, **16**, 1621-1627.
- ³⁶ S. Zaric and M. B. Hall, Ab Initio Calculations of the Geometries and Bonding Energies of Alkane and Fluoroalkane Complexes with Tungsten Pentacarbonyl. *J. Phys. Chem. A*, 1997, **101**, 4646-4652.
- ³⁷ B. O. Roos, R. Lindh, H. G. Cho and L. Andrews, Agostic Interaction in the Methylidene Metal Dihydride Complexes H₂MCH₂ (M = Y, Zr, Nb, Mo, Ru, Th, or U). *J. Phys. Chem. A*, 2007, **111**, 6420-6424.
- ³⁸ E. D. Glendening, C. R. Landis and F. Weinhold, Natural bond orbital methods. *Wiley Interdisciplinary Reviews: Computational Molecular Science*, 2012, **2**, 1-42.
- ³⁹ F. Weinhold and C. R. Landis, Natural bond orbitals and extensions of localized bonding concepts. *Chemistry education: Research and practice in Europe*, 2001, **2**, 91-104.
- ⁴⁰ F. Weinhold and C. R. Landis, Valency and Bonding - A natural bond donor-acceptor perspective, ed Cambridge University Press, 2005.
- ⁴¹ H. G. Cho and B. S. Cheong, Agostic Interaction of the Smallest Zirconium Methylidene Hydride: Reproduction of the Distorted Structure Experimentally Observed in Matrix Infrared Spectra. *Bull. Korean Chem. Soc.*, 2009, **30**, 470-481.

- ⁴² T. S. Thakur and G. R. Desiraju, Theoretical investigation of C–H...M interactions in organometallic complexes: A natural bond orbital (NBO) study. *Journal of Molecular Structure: THEOCHEM*, 2007, **810**, 143-154.
- ⁴³ M. Kaupp, The Unusual Fluxional Structure of Tetramethyloxotungsten: Quantum Chemical Structure Predictions for the d0 and d1 Complexes [MOR₄] (M = W, Re; R = H, CH₃). *Chem. Eur. J.*, 1998, **4**, 2059-2071.
- ⁴⁴ R. F. Bader, Atoms in Molecules: A Quantum Theory, ed Oxford University Press, Oxford, 1994.
- ⁴⁵ S. Pillet, G. Wu, V. Kulsomphob, B. G. Harvey, R. D. Ernst and P. Coppens, Investigation of Zr–C, Zr–N, and Potential Agostic Interactions in an Organozirconium Complex by Experimental Electron Density Analysis. *J. Am. Chem. Soc.*, 2003, **125**, 1937-1949.
- ⁴⁶ J. Saßmannshausen, Agostic or not? Detailed Density Functional Theory studies of the compounds. *Dalton Trans.*, 2011, **40**, 136-141.
- ⁴⁷ W. Scherer, V. Herz, A. Brück, C. Hau, F. Reiner, S. Altmannshofer, D. Leusser and D. Stalke, The Nature of β -Agostic Bonding in Late-Transition-Metal Alkyl Complexes. *Angew. Chem. Int. Ed.*, 2011, **50**, 2845-2849.
- ⁴⁸ R. F. W. Bader and C. F. Matta, Bonding to titanium. *Inorg. Chem.*, 2001, **40**, 5603-5611.
- ⁴⁹ V. Tognetti, L. Joubert, R. Raucoles, T. de Bruin and C. Adamo, Characterizing agosticity using the Quantum Theory of Atoms in Molecules: Bond Critical Points and their local properties. *J. Phys. Chem. A*, 2012, **116**, 5472-5479.
- ⁵⁰ A. D. Becke and K. E. Edgecombe, A simple measure of electron localization in atomic and molecular system. *J. Chem. Phys.*, 1990, **92**, 5397-5403.
- ⁵¹ B. Silvi and A. Savin, Classification of chemical bonds based on topological analysis of electron localization functions. *Nature*, 1994, **371**, 683-686.
- ⁵² A. Savin, B. Silvi and F. Colonna, Topological analysis of the electron localization function applied to delocalized bonds. *Can. J. Chem.*, 1996, **74**, 1088-1096.
- ⁵³ B. Silvi, I. Fourré and M. E. Alikhani, The topological analysis of the electron localization function. A key for a position space representation of chemical bonds. *Monatshefte für Chemie/Chemical Monthly*, 2005, **136**, 855-858.
- ⁵⁴ P. W. Ayers, Electron localization functions and local measures of the covariance. *J. Chem. Sci.*, 2005, **117**, 441-454.
- ⁵⁵ S. Diner and P. Claverie, in *Localization and Delocalization in Quantum Chemistry*. eds O. Chalvet, R. Daudel, S. Diner, R. Malrieu, Dordrecht, 1976, vol II, pp 395-448.
- ⁵⁶ B. Silvi, The Spin-Pair Compositions as Local Indicators of the Nature of the Bonding. *J. Phys. Chem. A*, 2003, **107**, 3081-3085.
- ⁵⁷ S. Raub and G. Jansen, A quantitative measure of bond polarity from the electron localization function and the theory of atoms in molecules. *Theor Chem Acc*, 2001, **106**, 223-232.
- ⁵⁸ F. Fuster and S. Grabowski, Intramolecular hydrogen bonds: the QTAIM and ELF characteristics. *J. Phys. Chem. A*, 2011, **115**, 10078-10086.
- ⁵⁹ I. Fourré and J. Bergès, Structural and topological characterization of the three-electron bond: the SO radical. *J. Phys. Chem A*, 2004, **108**, 898-906.
- ⁶⁰ I. Fourré and B. Silvi, What can we learn from two-center three-electron bonding with the topological analysis of ELF? Heteroatom Chemistry, 2007, **18**, 135-160.
- ⁶¹ J. Poater, M. Duran, M. Solà and B. Silvi, Theoretical evaluation of electron delocalization in aromatic molecules by means of Atoms in Molecules (AIM) and Electron Localization Function (ELF) topological approaches. *Chem. Rev.*, 2005, **105**, 3911-3947.
- ⁶² K. B. Wiberg, R. F. W. Bader, Clement D. H. Lau, Theoretical analysis of hydrocarbon properties. 2. Additivity of group properties and the origin of strain energy. *J. Am. Chem. Soc.*, 1987, **109**, 1001-1012.
- ⁶³ A. F. Dunlop-Brière, P.H.M. Budzelaar and M.C. Baird, α - and β -agostic alkyl-titanocene complexes. *Organometallics*, 2012, **31**, 1591-1594.
- ⁶⁴ A. F. Dunlop-Brière, M.C. Baird and P.H.M. Budzelaar, [Cp₂TiCH₂CHMe(SiMe₃)]⁺, an Alkyl–Titanium Complex Which (a) Exists in Equilibrium between a β -Agostic and a Lower Energy γ -Agostic Isomer and (b) Undergoes Hydrogen Atom Exchange between α -, β -, and γ -Sites via a

Combination of Conventional β -Hydrogen Elimination–Reinsertion and a Nonconventional CH Bond Activation Process Which Involves Proton Tunnelling. *J. Am. Chem. Soc.*, 2013, **135**, 17514-17527.

⁶⁵ T. A. Atesin, T. Li, S. Lachaize, W. W. Brennessel, J. J. Garcia and W. D. Jones, Experimental and theoretical examination of C-CN and C-H bond activation of acetonitrile using zerovalent nickel. *J. Am. Chem. Soc.*, 2007, **129**, 7562-7569.

⁶⁶ M. Montag, I. Efremenko, Y. Diskin-Posner, Y. Ben-David, J. M. L. Martin and D. Milstein, Exclusive C-C oxidative addition in a rhodium thiophosphoryl pincer complex and computational evidence for an η^3 -C-C-H agostic intermediate. *Organometallics*, 2012, **31**, 505-512.

⁶⁷ D. J. Wolstenholme, K. T. Trabousee, A. Decken and G. S. McGrady, Structure and Bonding of Titanocene Amidoborane Complexes: A Common Bonding Motif with Their β -Agostic Organometallic Counterparts. *Organometallics*, 2010, **29**, 5769-5772.

⁶⁸ T. D. Forster, H. M. Tuononen, M. Parvez and R. Roesler, Characterization of β -B-Agostic Isomers in Zirconocene Amidoborane Complexes. *J. Am. Chem. Soc.*, 2009, **131**, 6689-6691.

⁶⁹ G. Bénac-Lestrille, U. Helmstedt, L. Vendier, G. Alcaraz, E. Clot and S. Sabo-Etienne. Dimethylaminoborane coordination to late transition metal centers: Snapshots of the B-H oxidative addition process. *Inorganic Chemistry*, 2011, **50**, 11039-11045.

⁷⁰ G. Alcaraz, E. Clot, S. Sabo-Etienne, L. Vendier and U. Helmstedt, Mesitylborane as a bis (σ -B-H) ligand: an unprecedented bonding mode to a metal center. *J. Am. Chem. Soc.*, 2007, **129**, 8704-8705

⁷¹ C. Adamo and V. Barone, Toward reliable density functional methods without adjustable parameters: The PBE0 model. *J. Chem. Phys.*, 1999, **110**, 6158-69.

⁷² J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, Climbing the density functional ladder: Nonempirical meta-generalized gradient approximation designed for molecules and solids. *Physical Review Letters*, 2003, **146401**, 1-4.

⁷³ W. Scherer, D. J. Wolstenholme, V. Herz, G. Eickerling, A. Brück, P. Benndorf and P. W. Roesky, On the nature of agostic interaction in transition-metal amido complexes. *Angew. Chem. Int. Ed.* 2010, **49**, 2242–2246.

⁷⁴ U. Koch and P.L.A. Popelier, Characterization of C-H-O Hydrogen Bonds on the Basis of the Charge Density. *J. Phys. Chem.* 1995, **99**, 9747-9754.

⁷⁵ I. Vidal, S. Melchor, I. Alkorta, J. E Iguero, M. R. Sundberg, and J. A. Dobado, On the Existence of r -Agostic Bonds: Bonding Analyses of Titanium Alkyl Complexes. *Organometallics*, 2006, **25**, 5638-5647.

⁷⁶ B. Silvi, The synaptic order: a key concept to understand multicenter bonding. *J. Mol. Struct.*, 2002, **614**, 3-10.

⁷⁷ N. Berkaine, P. Reinhardt, M. E. Alikhani, Metal (Ti, Zr, Hf) insertion in the C–H bond of methane : Manifestation of an agostic interaction. *Chem. Phys.*, 2008, **343**, 241-249.

⁷⁸ P. L. A. Popelier, On the full topology of the Laplacian of the electron density. *Coord. Chem. Rev.*, 2000, **197**, 169-189.

⁷⁹ J. C. S. da Silva, W. B. de Almeida and W. R. Rocha, Theoretical investigation of the structure and nature of the interaction in metal-alkane σ -complexes of the type $[M(CO)_5(C_2H_6)]$ (M = Cr, Mo, and W). *Chem. Phys.*, 2009, **365**, 85-93.

⁸⁰ D. Braga, F. Grepioni, K. Biradha and G.R. Desiraju, Agostic interactions in organic metal compounds. A Cambridge Structural database study. *J. Chem. Soc., Dalton Trans.* 1996, 3925-3930.

⁸¹ J. Pinkas, V. Varga b, I. Cisarovac, J. Kubistaa, M. Horaceka, K. Mach, The first thermally stable half-sandwich titanium zwitterionic complex. *Journal of Organometallic Chemistry*, 2007, **692**, 2064-2070.

⁸² W. E. Billups, S. C. Chang, R. H. Hauge, and J. L. Margrave, Detection of a σ -Complex in the Reaction of Cobalt Atoms with Methane. *J. Am. Chem. Soc.*, 1993, **115**, 2041-2042.

⁸³ R. H. Crabtree, The organometallic chemistry of the transition metals. Fifth Edition, p307. John Wiley and Sons, Inc, 2009.

⁸⁴ E. F. van der Eide, P. Yang and R.M. Bullock, Isolation of two agostic isomers of an organometallic cation: different structures and colors. *Angew. Chem.*, 2013, **125**, 10380-10384.

⁸⁵ W. W. Lukens, M. R. Smith III and R. A. Andersen, A π -Donor Spectrochemical Series for X in $(Me_5C_5)_2TiX$, and β -Agostic Interactions in X) Et and N(Me)Ph. *J. Am. Chem. Soc.*, 1996, **118**, 1719-1728.

- ⁸⁶ See for instance: a) R. S. Anju, D. K. Roy, B. Mondal, K. Yuvaraj, C. Arivazhagan, K. Saha, B. Varghese, and S. Ghosh, Reactivity of Diruthenium and Dirhodium Analogues of Pentaborane - Agostic versus Boratrane Complexes. *Angew. Chem. Int. Ed.*, 2014, **53**, 2873-2877; b) M. A. Alvarez, D. Garcia-Vivo, M. E. Garcia, M. E. Martinez, A. Ramos, and M. A. Ruiz, Reactivity of the r-Agostic Methyl Bridge in the Unsaturated Complex $[\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\mu\text{-}\eta^1\text{:}\eta^2\text{-CH}_3)(\mu\text{-PCy}_2)(\text{CO})_2]$: Migratory Behavior and Methylidyne Derivatives. *Organometallics*, 2008, **27**, 1973-1975.
- ⁸⁷ G. Alcaraz, S. Sabo-Etienne, NMR: A good tool to ascertain σ -silane or σ -borane formulations? *Coordination Chemistry Reviews*, 2008, **252**, 2395-2409.
- ⁸⁸ See for instance: a) J. A. Calladine, S. B. Duckett, M. W. George, S. L. Matthews, R. N. Perutz, O. Torres and K. Q. Vuong, Manganese Alkane Complexes: An IR and NMR Spectroscopic investigation. *J. Am. Chem. Soc.*, 2011, **133**, 2303-2310; b) O. Torres, J. A. Calladine, S. B. Duckett, M. W. George and R. N. Perutz, Detection of s-alkane complexes of manganese by NMR and IR spectroscopy in solution: $(\eta^5\text{-C}_5\text{H}_5)\text{Mn}(\text{CO})_2(\text{ethane})$ and $(\eta^5\text{-C}_5\text{H}_5)\text{-Mn}(\text{CO})_2(\text{isopentane})$. *Chem. Sci.*, 2015, **6**, 418-424.
- ⁸⁹ See for instance H. G. Cho, X. Wang, L. Andrews, The C-H Activation of Methane by Laser-Ablated Zirconium Atoms: CH_2ZrH_2 , the Simplest Carbene Hydride Complex, Agostic Bonding, and $(\text{CH}_3)_2\text{ZrH}_2$. *J. Am. Chem. Soc.*, 2005, **127**, 465-473.
- ⁹⁰ See for instance F. M. Conroy-Lawis, L. Mole, A. D. Redhouse, S. A. Litster and J. L. Spencer, Synthesis of Coordinatively Unsaturated Diphosphine Nickel(II) and Palladium(II) β -Agostic Ethyl Cations: X-Ray Crystal Structure of $[\text{Ni}\{\text{Bu}^t\text{P}(\text{CH}_2)_2\text{PBu}^t\}_2(\text{C}_2\text{H}_5)][\text{BF}_4]$. *J. Am. Chem. Soc.*, 1991, 1601-1603.
- ⁹¹ C. Elschenbroich, in *Organometallics: a concise introduction*. Second Edition, p168. Weinheim New York Basel, 1992.
- ⁹² K. Rachlewicz, S.L. Wang, C.H. Peng, C.H. Hung and L. Latos-Grazynski. Remarkable paramagnetically shifted 1H and 2H NMR spectra of Iron(II) complexes of 2-aza-21-carbaporphyrin: an evidence for agostic interaction. *Inorg. Chem.*, 2003, **42**, 7348-7350.
- ⁹³ G. A. Luinstra, L. C. Ten Cate, H. J. Heeres, J. W. Pattiasina, A. Meetsmaand and J. H. Teuben, Synthesis and Reactivity of Tervalent Paramagnetic Titanium Compounds $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}$: Molecular Structure of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiCH}_2\text{CMe}_3$. *Organometallics*, 1991, **10**, 3227-3237.
- ⁹⁴ T. S. Thakur and G. R. Desiraju, Misassigned C-HCu agostic interaction in a copper(II) ephedrine derivative is actually a weak, multicentred hydrogen bond. *Chem. Commun.*, 2006, 552-554.
- ⁹⁵ S. Noury, X. Krokidis, F. Fuster and B. Silvi, Computational tools for the electron localization function topological analysis. *Comput. Chem.*, 1999, **23**, 597-604.
- ⁹⁶ AIMAll (Version 14.10.27), Todd A. Keith, TK Gristmill Software, Overland Park KS, USA, 2014 (aim.tkgristmill.com)
- ⁹⁷ F. Weigend and R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297-3307.
- ⁹⁸ a) D. Feller, The role of databases in support of computational chemistry calculations. *J. Comput. Chem.*, 1996, **17**, 1571-1586; b) K. L. Schuchardt, B. T. Didier, T. Elsethagen, L. Sun, V. Gurumoorthi, J. Chase, J. Li and T. L. Windus, Basis Set Exchange: A Community Database for Computational Sciences. *J. Chem. Inf. Model.*, 2007, **47**, 1045-1052.
- ⁹⁹ J.-D. Chai and M. Head-Gordon, Long-range corrected hybrid density functionals with damped atom-atom dispersion corrections, *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615-6620.
- ¹⁰⁰ L. Goerigk and S. Grimme, Efficient and Accurate Double-Hybrid-Meta-GGA Density Functionals—Evaluation with the Extended GMTKN30 Database for General Main Group Thermochemistry, Kinetics, and Noncovalent Interactions, *J. Chem. Theory Comput.*, 2011, **7**, 291-309.