

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## A family of low molecular-weight, organic catalyst for reductive C-C bond formation

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

Saad Shaaban,<sup>a</sup> Anaïs Jolit,<sup>a</sup> Desislava Petkova,<sup>a</sup> and Nuno Maulide<sup>a</sup>

Hydrazines form a new family of low molecular-weight reducing agents for diazonium salts. Using only small amounts of a hydrazine catalyst, the coupling of diazonium salts to a variety of reactive partners has been achieved, without the requirement for either metal adjuvants or irradiation with visible or ultraviolet light. The generality of the concept proposed herein as well as its advantages in preparative scale are outlined and discussed.


### Introduction

Diazonium salts are well-known highly reactive species in organic synthesis.<sup>1</sup> Despite their commercially important application in the dye industry,<sup>2</sup> diazonium salts function as ideal sources of organic radicals that engage in several important transformations such as the Meerwein arylation<sup>3</sup> or the Sandmeyer<sup>4</sup> and Pschorr reaction families.<sup>5</sup> Since the advent<sup>6</sup> and renaissance of photoredox catalysis,<sup>7</sup> many photocatalysts have been shown to catalyse the single electron transfer reduction of diazonium salts to organic radicals. In particular, ruthenium(II) and iridium(III) complexes have been successfully applied in many reactions that exploit their remarkable reactivity in either reductive or oxidative quenching pathways upon visible light (440nm) excitation.<sup>8</sup> The relatively high price of those noble/precious metal complexes has subsequently led to the development of alternatives. Namely, several organic dyes proved to be efficient photocatalysts upon excitation with visible light.<sup>9</sup> For instance, Eosin Y,<sup>10</sup> acridinium salts<sup>9a,11</sup> and perylene diimides (PDI)<sup>12</sup> were all successfully used in photocatalytic organic transformations. A recurrent structural feature of these organic dye catalysts is the presence of extended  $\pi$ -surfaces (mandated by the need to effectively absorb visible light) and high molecular-weight frameworks.<sup>13</sup> Recently, Carrillo et al. described the use of ascorbic acid as an initiator for a C-H arylation of anilines.<sup>14</sup> Herein we describe a family of low-molecular weight organic catalyst that promote a range of C-C bond forming reactions of diazonium salts, and achieve so

without the need for any metal adjuvant or irradiation with light.

### Results and discussion

We were inspired by the work of Shevlin *et al.*<sup>15</sup> who aimed to isolate stable tetrazenes by reacting phenyldiazonium tetrafluoroborate with phenylhydrazine (Scheme 1). However, those authors were not able to isolate the tetrazene intermediate but instead observed a complex mixture that featured benzene as major component in 54% yield and several other minor compounds. The formation of most of the components of the mixture was attributed to the intermediacy of free radicals, as supported by spectroscopic studies (CIDNP).



**Scheme 1** Reaction of phenyldiazonium salt with phenylhydrazine as reported by Shevlin *et al.*<sup>15</sup>

Arylated heteroarenes have proved to be of interest for various applications, such as material sciences and biomedicine.<sup>16</sup> One efficient route to install the aryl-heteroaryl bond is the Meerwein arylation, a cross-coupling reaction between heteroarenes and diazonium salts that is assumed to proceed by a free radical mechanism.<sup>17</sup>

We hypothesized that a hydrazine such as phenylhydrazine (employed in the work of Shevlin) could function as a novel


<sup>a</sup> University of Vienna, Institute of Organic Chemistry, Währinger Strasse 38, 1090 Vienna, Austria.

[nuno.maulide@univie.ac.at](mailto:nuno.maulide@univie.ac.at); <http://maulide.univie.ac.at>

† Electronic Supplementary Information (ESI) available: characterisation data for new compounds and ancillary experiments. See DOI: 10.1039/x0xx00000x

reducing agent towards aryl diazonium salts. However, it was crucial to generate the radical intermediate under conditions which would (a) avoid the occurrence of either homocoupling or hydrogen abstraction (as seen by Shevlin), (b) enable its efficient trapping by the heteroarene partner, (c) prevent unwanted disproportionation reactions (such as, e.g. leading to an azide and an aniline, as originally also observed by Shevlin) and, most importantly, (d) allow rebound electron-transfer from the coupling adduct to generate the final, neutral biaryl product.


From the outset of our studies on the Meerwein arylation of thiophene, it soon became apparent that hydrazines might have the potential to function as catalyst in such a coupling, thus addressing all the above concerns in a fell swoop (detailed optimisation is included as Table S1 of the ESI). Strikingly, yields of coupling reactions were unchanged when the reaction was performed in the dark, showing that *light plays no appreciable role in this transformation*. We chose 4-aminomorpholine **3d** to pursue our studies due to both practical considerations and its low price. Importantly, the catalyst proved to be essential for the formation of **4a**, as only traces of the desired product were obtained under similar conditions if it was absent from the reaction mixture.<sup>18</sup>



**Scheme 2.** Scope of the  $\alpha$ -arylation of heteroarenes with various diazonium salts using the optimised conditions detailed in the supporting information. The yields are given for the isolated products.


The scope of this hydrazine-catalyzed C-H arylation of heteroarenes with various aryl diazonium salts was then examined using the best conditions found. As shown in Scheme 2, a range of functional groups, such as nitro, cyano, ester, and trifluoromethane are compatible with the reaction

conditions. Notably, the carbon-halogen bonds in **4g** and **4i** were not affected and could serve as reactive handles for further elaboration.<sup>19</sup> A further potential advantage of this system is highlighted in Scheme 3. *N,N*-dimethylhydrazine **3c**, which has a very low molecular weight (MW = 60 g/mol, more than 11 times lower than eosin Y), was used to scale up the coupling reaction between nitrobenzene diazonium salt **1a** and furan **2b** to yield **4b** in ca. 85% yield in 15 min. Remarkably, only 7.6 mg of catalyst was used for 12.6 mol of substrate (3g scale, 1 mol% catalyst). For the sake of comparison, to achieve the same reaction König's elegant photochemical method would require 87 mg of eosin Y (1 mol%).<sup>10b</sup>



**Scheme 3.** Scale up of the  $\alpha$ -arylation of furan with *p*-nitrobenzene diazonium salt using *N,N*-dimethylhydrazine **3c** as catalyst.

At this juncture, we inquired whether this mode of redox catalysis could be extended to other related transformations. The synthesis of  $\alpha$ -aryl ketones is a challenging significant reaction.<sup>20</sup> It was recently shown that the capture of diazonium salts with enolacetates led to this motif.<sup>21</sup> Nevertheless, the reported methodologies mandate the use of transition metal catalysts,<sup>21a</sup> transition metal photocatalysts<sup>21b</sup> or stoichiometric amounts of base.<sup>21c</sup> We thus were intrigued by the possibility of employing hydrazine catalysis for this process (Scheme 4).




**Scheme 4.** Preparation of  $\alpha$ -aryl ketones via reduction of diazonium salts with hydrazine **3d**. Yields refer to isolated products.

Several aryl diazonium salts **1** were therefore treated with alkenyl acetates **5** in presence of 10 mol% of hydrazine **3d** in dimethylformamide (DMF) to prepare  $\alpha$ -arylketone products **6**. A series of  $\alpha$ -aryl methylketones and  $\alpha$ -aryl arylketones could be synthesized in moderate to good yield. The reaction was


tolerant of several functional groups, such as nitro, ester and cyano substituents.

We further applied this catalytic system to the preparation of 2-substituted thiophenes **8** from the corresponding *ortho*-(methylthio) diazonium tetrafluoroborate **1h**. This transformation has been previously reported using transition metal catalysis<sup>22a</sup> or visible light photoredox catalysis with eosin Y.<sup>22b</sup> 4-Aminomorpholine **3d** was able to catalyse this transformation furnishing the corresponding benzothiophenes in good to moderate yields (Scheme 5).




**Scheme 5.** Preparation of 2-substituted benzothiophenes *via* hydrazine-catalysed coupling of diazonium salt **1h** and alkynes **7**. Yields refer to isolated products.

Furthermore, the arylation of quinones with diazonium salts is an interesting transformation due to the important role of functionalized quinones in medicinal chemistry and biologically active compounds. Bearing this in mind, this reaction was also achieved employing 4-aminomorpholine **3d** as catalyst and the results are summarized in Scheme 6.<sup>23</sup>




**Scheme 6.** Hydrazine-catalysed C-H arylation of quinones with diazonium salts. Yields refer to isolated products.



**Scheme 7.** Trapping the aryl radical intermediate with TEMPO.

Finally, we attempted to trap the putative aryl radical intermediate, presumably formed upon diazonium salt reduction, with TEMPO **11**. As shown in Scheme 7, phenyldiazonium salt **1g** was exposed to a catalytic amount of 4-aminomorpholine **3d** in the presence of 2 equiv. of TEMPO. The expected covalent adduct, 2,2,6,6-tetramethyl-1-phenoxy-piperidine **12**, was isolated in 44% yield. Additional experiments (see Supporting Information for details) reveal

that the hydrazine catalyst is present in the reaction mixture (by both LC-MS and NMR analysis) after full conversion has been reached, thus establishing catalytic behaviour. Based on the above observations and previous literature,<sup>3b,10b,15</sup> a general mechanism for the couplings of diazonium salts catalysed by 4-aminomorpholine (typified by the Meerwein arylation) is proposed in Scheme 8.



**Scheme 8.** Proposed catalytic cycle for the hydrazine-catalysed coupling of diazonium salt **13** and heteroarene **17**.

We believe that formation of tetrazene **14**,<sup>15</sup> via reaction of diazonium salt **13** with the hydrazine catalyst **3d**, will generate aryl radical **16** and diazenium radical cation **15** and is driven by the release of nitrogen gas. Radical **16** is subsequently intercepted by a reactive unsaturation typified by **17**, to form radical adduct **18**. Oxidation of **18** via SET of **15**<sup>24</sup> to the corresponding carbocation **19** regenerates the hydrazine catalyst **3d** and aryl radical **16**, enabling turnover.

## Conclusions

In summary, we have shown that simple hydrazines form a new family of low molecular weight reducing agents that allow the catalytic coupling of diazonium salts to electron-rich heteroarenes, enol acetates, alkynes or benzoquinones under simple conditions. Notably, these catalysts' low molecular weight ensures unrivalled practicality when one considers the upscaling of this type of chemistry. That hydrazines can function as catalytic activators for diazonium salts, shown by their presence after full conversion has been achieved, is a new type of reactivity bound to inspire the development of further transformations.

## Acknowledgements

Generous financial support of this research by the Deutsche Forschungsgemeinschaft (Grants MA 4861/1-2 and MA 4861/4-2) and the University of Vienna is acknowledged.

## Notes and references

- 1 a) H. Zollinger, *Acc. Chem. Res.* 1973, **6**, 335–341; b) C. Galli, *Chem. Rev.* 1988, **88**, 765–792; c) F. Minisci, F. Coppa, F. Fontana, G. Pianese and L. Zhao, *J. Org. Chem.* 1992, **57**, 3929–3933.
- 2 a) W. Dedichen, *Chem. Ber.* 1897, **30**, 2657–2658; b) R. Loose, *Chem. Ber.* 1908, **41**, 3906–3907.
- 3 a) H. Meerwein, E. Buchner and K. Van Emsterk, *J. Prakt. Chem.* 1939, **152**, 237; b) M. Heinrich, *Chem. Eur. J.* 2009, **15**, 820–833.
- 4 a) T. Sandmeyer, *Ber. Dtsch. Chem. Ges.* 1884, **17**, 1633, 2650; b) E. Merkushev, *Synthesis* 1988, **12**, 923–937.
- 5 a) R. Pschorr, *Ber. Dtsch. Chem. Ges.* 1896, **29**, 496–501; b) P. Leake, *Chem. Rev.* 1956, **56**, 27–4.
- 6 a) H. Cano-Yelo and A. Deronzier, *J. Chem. Soc., Perkin Trans. 2.* 1984, 1093–1098; b) H. Cano-Yelo and A. Deronzier, *Tetrahedron Lett.* 1984, **25**, 5517–5520.
- 7 For recent reviews on transition metal photocatalysis, see: a) J. Narayanam and C. R. Stephenson, *Chem. Soc. Rev.* 2011, **40**, 102–113; b) C. Prier, D. Rankic and D. W. C. MacMillan, *Chem. Rev.* 2013, **113**, 5322–5363.
- 8 Selected references: a) D. Nagib, M. Scott and D. W. C. MacMillan, *J. Am. Chem. Soc.* 2009, **131**, 10875–10877; b) J. Nguyen, E. D'Amato, J. R. Narayanam and C. Stephenson, *Nature Chem.* 2012, **4**, 854–859; c) B. Sahoo, M. Hopkinson and F. Glorius, *J. Am. Chem. Soc.* 2013, **135**, 5505–5508; d) X. Shu, M. Zhang, Y. He, H. Frei and F. Toste, *J. Am. Chem. Soc.* 2014, **136**, 5844–5847; e) D. Hari, T. Hering and B. König, *Angew. Chem. Int. Ed.* 2014, **53**, 725–728.
- 9 For recent uses of organic dyes as photocatalysts, see: a) S. Pitre, C. Mctiernan, H. Ismaili and J. Scaiano, *J. Am. Chem. Soc.* 2013, **135**, 13286–13289; b) D. Nicewicz and T. Nguyen, *A.C.S. Catal.* 2014, **4**, 355–360; c) D. Hari and B. König, *Chem. Commun.* 2014, **50**, 6688–6699.
- 10 a) A. Penzkofer, A. Beidoun and S. Speiser, *Chem. Phys.* 1993, **170**, 139–148; b) D. Hari, P. Schroll and B. König, *J. Am. Chem. Soc.* 2012, **134**, 2958–2961; c) M. Majeka and A. Wangelin, *Chem. Commun.* 2013, **49**, 5507–5509; d) J. Zhang, L. Wang, Q. Liu, Z. Yang and Y. Huang, *Chem. Commun.* 2013, **49**, 11662–11664.
- 11 a) S. Fukuzumi, H. Kotani, K. Ohkubo, S. Ogo, N. Tkachenko and H. Lemmetyinen, *J. Am. Chem. Soc.* 2004, **126**, 1600–1601; b) D. Hamilton and D. Nicewicz, *J. Am. Chem. Soc.* 2012, **134**, 18577–18580.
- 12 a) F. Würthner, *Chem. Commun.* 2004, **14**, 1564–1579; b) I. Ghosh, T. Ghosh, J. Bardagi and B. König, *Science* 2014, **346**, 725–728.
- 13 Eosin Y Mw: 691.85 g.mol<sup>-1</sup>, acridinium tetrafluoroborate Mw: 388.9 g.mol<sup>-1</sup>, PDI Mw: 710,31 g.mol<sup>-1</sup>. For seminal work on organic superelectron donors, see: J. A. Murphy, *J. Org. Chem.* 2014, **79**, 3731–3746.
- 14 For the use of ascorbic acid as a SET reductant (10 mol%) in the Meerwein arylation, see: F. P. Crisostomo, T. Martin and R. Carrillo, *Angew. Chem. Int. Ed.* 2014, **53**, 2181–2185.
- 15 W. Evanochko and P. Shevlin, *J. Am. Chem. Soc.* 1979, **101**, 4668–4672.
- 16 a) J. Van Haare, L. Groenendaal, H. Peerlings, E. Havinga, J. Vekemans, R. Janssen and E. Meijer, *Chem. Mater.* 1995, **7**, 1984–1989; b) B. Schmidt and D. Geißler, *Eur. J. Org. Chem.* 2011, **25**, 4814–4822.
- 17 a) B. Sreedhar, D. Yada and P. Reddy, *Adv. Synth. Catal.* 2011, **353**, 2823–2836; b) N. Kuhl, M. Hopkinson and F. Glorius, *Angew. Chem. Int. Ed.* 2012, **51**, 8230–8234.
- 18 For homolytic dediazonation induced by solvent, see: G. Pratsch and M. Heinrich, in *Radicals in Synthesis III*, Vol. 320 (Eds.: M. Heinrich, A. Gansäuer), Springer, Berlin, 2012, pp. 33–59;
- 19 It is known that aryl diazonium salts bearing electron-withdrawing substituents give Meerwein products in higher yields: P. Caramella, G. Cellerino, A. C. Coda, A. G. Invernizzi, P. Grunanger, K. N. Houk and F. M. Albini, *J. Org. Chem.* 1976, **41**, 3349–3356.
- 20 a) T. Shen, *Angew. Chem.* 1972, **84**, 512–526; b) V. Aggarwal and B. Olofsson, *Angew. Chem. Int. Ed.* 2005, **44**, 5516–5519.
- 21 a) J. Rutherford, M. Rainka and S. Buchwald, *J. Am. Chem. Soc.* 2002, **124**, 15168–15169; b) T. Hering, D. Hari and B. König, *J. Org. Chem.* 2012, **77**, 10347–10352; c) C. Molinaro, J. Mowat, F. Gosselin, P. O'Shea, J. Marcoux, R. Angelaud and I. Davies, *J. Org. Chem.* 2007, **72**, 1856–1858.
- 22 a) R. Leardini, G. Pedulli, A. Tundo and G. Zanardi, *J. Chem. Soc., Chem. Commun.* 1985, **20**, 1390–1391; b) D. Hari, T. Hering and B. König, *Org. Lett.* 2012, **14**, 5334–5337.
- 23 For arylation of quinones, see: a) Y. Fujiwara, V. Domingo, I. Seiple, R. Gianatassio, M. Del Bel and P. Baran, *J. Am. Chem. Soc.* 2011, **133**, 3292–3295; base-mediated variant: b) M. Lamblin, G. Naturalea, J. Dessolina and F.-X. Felpin, *Synlett.* 2012, **23**, 1621–1624.
- 24 The alternative reoxidation of **18** through SET from **13**, as previously proposed in the literature (cf. refs 10b, 14) cannot be ruled out at this stage.