


Molecular BioSystems


Interfacing chemical biology with the -omic sciences and systems biology

www.rsc.org/molecularbiosystems



Themed issue: Protein Labelling

ISSN 1742-206X



REVIEW ARTICLE

Oliver Seitz, Annette G. Beck-Sickinger *et al.*

Peptide-tags for site-specific protein labelling *in vitro* and *in vivo*

Indexed in
Medline!



Cite this: *Mol. BioSyst.*, 2016,
12, 1731

Received 11th January 2016,
Accepted 25th February 2016

DOI: 10.1039/c6mb00023a

www.rsc.org/molecularbiosystems

Peptide-tags for site-specific protein labelling *in vitro* and *in vivo*

Jonathan Lotze,^{†a} Ulrike Reinhardt,^{†b} Oliver Seitz^{*b} and Annette G. Beck-Sickinger^{*a}

Peptide-tag based labelling can be achieved by (i) enzymes (ii) recognition of metal ions or small molecules and (iii) peptide–peptide interactions and enables site-specific protein visualization to investigate protein localization and trafficking.

Introduction

Visualizing the localization and trafficking of proteins in living systems is the key to understand protein functions in their native environment. Tracking and visualization by fluorescence microscopy require methods that allow the site-specific labelling of a protein of interest (POI). Established almost 20 years ago,¹ the genetic fusion of a POI to an autofluorescent protein such as the green fluorescent protein (GFP) revolutionized protein imaging. The fusion of POIs with engineered, self-modifying enzymes such as SNAP,² CLIP,³ TMP,^{4,5} or Halo-tags⁶ significantly extended the scope to customized labelling by enabling the use of reporter units with unique spectroscopic signatures. Enzyme-based tags provide high specificity and enable fast labelling reactions, which is an advantage when fast biological processes ought to be analysed in time-lapse experiments. Yet, the fusion of enzymes brings along a massive increase in size (18–33 kDa). This can perturb protein trafficking and may impair protein–protein interaction networks that rely on clustering. To avoid potential interference with protein localization and protein function, major efforts are spent to decrease the tag size. The smallest tags currently accessible are provided by the incorporation of unnatural amino acids (UAA) by amber stop codon or related technologies.^{7,8} In addition, the ligand-directed protein labelling or traceless affinity labelling is independent of any tag.^{9,10} However, the establishment and incorporation of UAA in mammalian cells still is very challenging and the traceless affinity labelling requires a POI with a native binding pocket as well as the exact knowledge of the binding site and ligand.

Peptide-based recognition tags offer an attractive alternative for protein labelling. The genetic fusion of peptide-tags is straightforward, the size of 0.6–6 kDa is considerably small and a variety of labelling chemistries allow the introduction of virtually arbitrary reporter groups.

Peptide-tag based labelling methods are multifaceted and can be categorised into three classes: (i) enzyme mediated labelling, (ii) metal ion- or small molecule-dependent labelling and (iii) labelling based on peptide–peptide interactions. We will discuss the characteristics of the different approaches with respect to specificity, speed, stability, tag size, toxicity and versatility. Applications as well as shortcomings will be explained. The review covers the literature until 2015 and particularly emphasizes the most recent and most frequently used developments. Additional information can be taken from other comprehensive reviews.^{11–14}

Peptide-tag based labelling by enzymes – mechanisms and characteristics

Enzymes offer a naturally occurring sequence-specificity for site-selective modifications. Ligases such as biotin ligase, lipoic acid ligase or tubulin tyrosine ligase are capable of transferring molecular probes to a particular amino acid (aa) within a recognition sequence. Similarly, transferases can be used to transfer functional groups from a donor substrate to a specific amino acid within a peptide sequence, as demonstrated for phosphopantetheinyl transferases Sfp and AcpS, transglutaminase or the phosphocholine transferase AnkX. Another possibility is to use transpeptidase activity. Sortase A exchanges a part of the recognition sequence with a separate peptide conjugated to a labelling moiety. The specific modification of an amino acid side chain can also be achieved with formylglycine-generating enzyme,

^a Institut für Biochemie, Universität Leipzig, Brüderstrasse 34, D-04103 Leipzig, Germany. E-mail: abeck-sickinger@uni-leipzig.de

^b Institut für Chemie, Humboldt-Universität zu Berlin, Brook-Taylor-Strasse 2, D-12489 Berlin, Germany. E-mail: oliver.seitz@chemie.hu-berlin.de

[†] Ulrike Reinhardt and Jonathan Lotze contributed equally.



Table 1 Characteristics of enzymatic labelling techniques. One-step labelling achieves the direct modification of a POI with a reporter group (e.g. fluorescent probes). Two-step labelling involves in a first step the transfer of a chemically reactive functional group (e.g. azide, alkyne, biotin, hydrazine conjugates) which is targeted in a second step upon reaction with a suitable modified reporter group (e.g. via SPAAC, biotin–streptavidin binding or hydrazine formation)

Enzyme	Recognition tag	Reported tag positions	Typical/reported concentrations and incubation conditions	Reported applications	One-step or two-step labelling
Biotin ligase (BirA)	GLNDIFEAQKIEWHE (AP-tag)	N-terminal, C-terminal, internal	BirA-coexpression enables biotinylation during protein expression; biotin detection with streptavidin reporter within 15–60 min at RT using fixed cells or at 4 °C using live cells; ^{17,21} Ligation of 1 mM keton probe (biotin analogue) within 10–60 min at 32 °C, subsequent detection with 1 mM hydrazine probe within 10–60 min at 4 °C in live cells ²³	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with fixed or live cells), cell lysate isolated from transgenic mice	Two-step
Lipoic acid ligase (LplA)	DEVLVEIETDKAVLEVPGESEE (LAP-tag 1st generation) GFEIDKVVYDLDA (LAP-tag)	N-terminal, C-terminal, internal	Labelling with 100–350 µM lipoic acid derivative for 15–120 min; probe detection 10–20 min; reaction temperature: 21–37 °C ^{25,28,29}	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with live cell imaging)	One-step and two-step
Tubulin tyrosine ligase (TTL)	VDSVEGEEGEEGEE (Tub-tag)	C-terminal	Labelling with 1 mM tyrosine derivative for 1–3 h; subsequent detection <i>via</i> biocompatible labelling required 4–24 h; reaction temperature: 30 or 37 °C ³³	<i>In vitro</i>	Two-step
Phosphopantetheinyl-transferase (Sfp and AcpS)	DSLEFIASKLA (ybbR-tag) GDSLSWLLRLLN (S6-tag) GDSLDMLEWSLM (A1-tag)	N-terminal, C-terminal, internal	Labelling with 1–10 µM CoA-derivative for 20–40 min; reaction temperature: 37 °C ^{38,39}	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with fixed or live cells)	One-step and two-step
Transglutaminase (TGase)	PNPQLPF (Q1-tag) PKPQQFM (Q2-tag) GQQQLG (Q3-tag) xxQxx (basic Q-tag)	N-terminal, C-terminal, internal	Labelling with 400–500 µM cadaverine probe for 25–30 min or up to 4 h; reaction temperature: 4 or 37 °C ^{41,46}	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with live cell imaging)	One-step and two-step
AnkX/Lem3	TITSSYYR	N-terminal, C-terminal, internal	Labelling with 1 mM CDP-choline probe derivative for ≤1 h; reaction temperature: RT ⁴⁹	<i>In vitro</i>	Two-step
Sortase A (SrtA)	LPXTG	C-terminal	Labelling with 1–5 mM oligoglycine probe for 1–5 h <i>in vitro</i> ^{54,55} /10 µM oligoglycine probe for 15–60 min in living cells; reaction temperature: 37 °C ⁵⁶	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with live cell imaging)	One-step and two-step
Formylglycine-generating enzyme (FGE)	LCxPxR (aldehyde-tag)	N-terminal, C-terminal, internal	Labelling with 30–500 µM hydrazine or aminoxy probes for 2 h or overnight <i>in vitro</i> ^{59,60} /1 mM labelling probe for 1 h in living cells; reaction temperature: RT or 37 °C ⁶¹	<i>In vitro</i> , <i>in vivo</i> (cell surface labelling with live cell imaging)	One-step and two-step

RT – room temperature.


which naturally modifies a cysteine derived thiol group to an aldehyde function (see Table 1).

Already in 1999 a 15 aa long acceptor peptide, termed AP-tag, was reported to serve as a substrate for *E. coli* derived biotin ligase (BirA). This enzyme promotes the conjugation of a biotin to a lysine side chain within the AP-tag (Fig. 1A).¹⁵ BirA-mediated biotinylation was demonstrated for many different AP-tagged proteins in cell lysates including the transcription factors GATA-1 and EKLF.¹⁶ Co-expression of AP-tagged and BirA-tagged proteins allows the so-called proximity biotinylation to study protein–protein interactions.¹⁷ This method is widely used and has enabled investigations of homo- and heterodimerization of chemokine receptors CXCR4, CCR2 and CCR5 in RPE cells¹⁸ or glycoprotein interactions during herpesvirus entry into cells.¹⁹ Of note, the method also enables investigations of proximity between two proteins expressed on different cells. For example, BirA-mediated biotinylation was used to characterize the synaptic connectivity between the presynaptic


adhesion protein NRX-1 and the postsynaptic transmembrane protein NLG-1 *in vivo* in *Caenorhabditis elegans*.²⁰ In a recent study, BirA-mediated biotinylation exposed the anterograde trafficking of calcium activated potassium channels (K_{Ca}3.1) in polarized epithelial cells (including HEK293, MDCK, LLC-PK1 and Caro-2 cells). The AP-tag was introduced into the protein and BirA was fused to an ER-retaining sequence (KDEL) to specifically label AP-tagged K_{Ca}3.1 along the anterograde pathway.²¹ Furthermore the endocytosis of the apical Na⁺/K⁺/2Cl[−] co-transporter (NKCC2) equipped with an AP-tag between the 5th and 6th transmembrane domain was measured in real time by total internal reflection fluorescence (TIRF) microscopy. After BirA-mediated biotinylation, fluorescent staining was performed with fluorescent streptavidin.²² BirA also accepts a biotin derivative containing a ketone group. A comparison of biotin ligases from different organisms revealed that *Saccharomyces cerevisiae* and *Pyrococcus horikoshii* accept alkyne and azide derivatives of biotin.^{23,24} The biotin modifications enable



A) Biotin ligase



B) Lipoic acid ligase



C) Tubulin tyrosine ligase




Fig. 1 Protein labelling by means of (A) biotin ligase (BirA), (B) lipoic acid ligase (LplA) or (C) tubulin tyrosine ligase (TTL). The red star indicates the reporter group.

subsequent labelling of proteins such as epidermal growth factor receptor (EGFR) on live cells by formation of acyl hydrazones or Staudinger ligation.²³

A BirA cross-compatible technique uses *E. coli* derived lipoic acid ligase (LplA), which catalyses the ligation of lipoic acid to a sequence-specific lysine side chain in an ATP-dependent manner (Fig. 1B).^{25,26} LplA specifically recognizes the LAP-tag, which consists of 22 aa, and accepts different lipoic acid derivatives containing alkyne, azide, *trans*-cyclooctene, aryl aldehyde or aryl hydrazine groups as well as fluorophores.^{25,27–29} Strain-promoted alkyne–azide cycloaddition (SPAAC), hydrazone formation or Diels–Alder cycloaddition³⁰ enabled the subsequent chemical modification of the lipoic acid appendage on LAP-tagged proteins such as CFP,²⁵ BFP,²⁸ low-density lipoprotein receptor (LDLR),²⁵ neuroligin-1²⁷ or actin.²⁸ The LplA mutant LplA^{W37V} has been reported to provide higher ligation yields than the wild-type enzyme.²⁷ In contrast to BirA,

LplA accepts fluorescently labelled conjugates and permits one-step labelling of the POI. LplA-mediated protein labelling was demonstrated on HEK293T-, HeLa and COS-7 cells. A minimized LAP-tag with only 13 aa³¹ was used to incorporate a *p*-iodophenyl derivative as new bio-orthogonal handle into proteins, which could be selectively addressed in a second step by palladium-catalysed Sonogashira cross-coupling.³² Wombacher *et al.* reported norbornene-conjugated lipoic acid analogues, which were introduced into LAP-tagged CFP and dihydrofolate reductase from *E. coli* (eDHFR) and fluorescently labelled *via* inverse-electron demand Diels–Alder reaction.³⁰

The enzyme tubulin tyrosine ligase (TTL) allows the C-terminal labelling of proteins with a 14 aa hydrophilic recognition sequence called Tub-tag. TTL is a regulator of microtubule homeostasis and usually promotes the coupling of tyrosine to the α -tubulin C-terminus. Hackenberger and Leonhardt showed that TTL is also able to incorporate tyrosine



derivatives containing azide, formyl, amine or nitril groups (Fig. 1C).³³ This enabled the two step labelling of GFP-specific nanobodies with biotin, fluorescent probes or polyethylene glycol (PEG) chains by means of SPAAC, Staudinger ligation, Staudinger-phosphite reactions and hydrazone or oxime forming reactions. Labelling also succeeded in *E. coli* lysate.


The use of transferases offers interesting alternatives to ligase-mediated labelling. Phosphopantetheinyl transferases (PPTase) such as Sfp and AcpS catalyse the transfer of a phosphopantetheinyl (Ppant) group derived from coenzyme A (CoA) to a serine residue within a specific recognition sequence (Fig. 2A).³⁴ Owing to an impressive substrate promiscuity of Sfp and AcpS, CoA can be modified at its terminal thiol with small molecules, including biotin, sugars, peptides, porphyrin and fluorophores, the latter allowing one-step labelling similar to LplA.^{34–37} Furthermore, a set of three specific tags is available: ybbR-tag (11 aa), S6-tag (12 aa), preferentially recognized by Sfp and A1-tag (12 aa) preferentially recognized by AcpS.³⁸ The ybbR-tag was successfully introduced at the N- and C-terminus, or within the protein sequence as demonstrated for eGFP.³⁷ Orthogonal cell surface labelling with fluorescent CoA conjugates on living HeLa cells was demonstrated for S6-tagged EGFR or A1-tagged transferrin receptor 1 (TfR1).³⁸ The S6/A1 pair of tags was also used to enable labelling of the tropomyosin receptor kinase TrkA and the p75 neurotrophin co-receptor (P75NTR) in SH-SY5Y cells.³⁹ The A1-P75NTR construct was first labelled with biotin-CoA using AcpS and then stained with

streptavidin-QDot525. Subsequently, the S6-TrkA construct was biotinylated by using Sfp followed by labelling with streptavidin-QDot655. To enable imaging of interactions between chemokines and their fluorescent protein (FP)-tagged receptors on live CHO-K1 cells, the chemokines CXCL12, CCL2 and CCL21 were expressed with a S6-tag at the C-terminus and subjected to PPTase-mediated labelling with a fluorophore-CoA conjugate.⁴⁰ Co-localization of signals from the fluorophore and the FP indicated receptor-chemokine interactions.

Transglutaminases (TGase) catalyse the formation of an amide bond between the carboxamide of a glutamine side chain and the ϵ -amino group of a lysine side chain. TGases have been used to attach small molecules to antibodies,⁴¹ modify proteins by PEGylation⁴² or lipidation,⁴³ site-specifically conjugate proteins,⁴⁴ as well as immobilize proteins on solid support.⁴⁵ CFP and EGFR were labelled *in vitro* and on the surface of living cells by means of guinea pig liver TGase (gpTGase), which promotes the conjugation of labelled cadaverine (lysine analogue conjugated to biotin, AlexaFluor586 or photoaffinity moieties) with 6–7 aa long glutamine-rich Q-tags.⁴⁶ Recent studies described the use of tissue TGase (TG2) or microbial TGase (mTGase).^{47,48} The latter provides the advantage of calcium-independent labelling. Transfer of propargylamine with subsequent CuAAC modification was demonstrated for maltose binding protein using mTGase.

Two enzymes from *Legionella pneumophila* were recently investigated by Hedberg *et al.* for the reversible covalent labelling

A) PPTase Sfp



B) AnkX / Lem3




Fig. 2 Protein labelling via (A) phosphopantetheinyl transferase (PPTase) Sfp or (B) phosphocholine transferase AnkX. The hydrolase Lem3 enables removal of the label.




of proteins. The first enzyme AnkX transfers a phosphocholine moiety from a cytidine diphosphate choline (CDP-choline) derivative to the second serine residue within the eight amino acid recognition sequence TITSSYYR (Fig. 2B). The second enzyme Lem3 is able to remove the label upon hydrolytic cleavage of the transferred phosphocholine from the serine side chain. The method tolerates fusion of the recognition sequence to the N- or C-terminus, as well as into internal loops as demonstrated for fluorescein labelling of model proteins such as maltose binding protein, small ubiquitin-like modifier and DrrA enzyme.⁴⁹

Remarkably small tag sizes are found for sortase A and formylglycine-generating enzyme. The transpeptidase sortase A (SrtA) derived from *Staphylococcus aureus* specifically recognizes the short pentapeptide sequence LPxTG. SrtA cleaves the amide bond between threonine and glycine, forms a reactive thioester between the carboxyl group of threonine and an enzyme-derived cysteine and promotes the ligation of this reactive intermediate with an amino group of an oligoglycine substrate by formation of a new amide bond (Fig. 3A). First reports on SrtA as new protein engineering tool described modifications of LPETG-tagged GFP with a set of various oligoglycine substrates conjugated to peptides including unnatural amino acids, proteins or small molecules like folate.⁵⁰ Subsequently, the substrate diversity was extended to PEGylation, peptide-glycosylphosphatidylinositol conjugates or cyclised peptides.^{51–53} SrtA-mediated labelling has been used to label proteins with fluorophores, photocross-linkers, biotin, alkyl chains and cholesterol on intact cells such as HEK293T, CHO and HeLa cells.^{54–56} In a noteworthy

study the bacterial toxin aerolysin was labelled with fluorophores and biotin. The ability to form homo-heptameric pores in membranes of target cells after binding to cell surface receptors was not impaired and SrtA-mediated biotinylation helped to identify aerolysin–receptor interactions, which are responsible for aerolysin uptake.⁵⁷ Recently LPETG-tagged lipopeptides enabled the visualization of liposome trafficking into lung cancer.⁵⁸

The formylglycine-generating enzyme (FGE) naturally co- or post-translationally modifies a cysteine within the conserved recognition motif CxPxR to an aldehyde-bearing formylglycine (Fig. 3B).⁵⁹ The hexapeptide sequence LCxPxR, termed “aldehyde tag”, was included at N-terminal, C-terminal or internal sites as shown for maltose binding protein, human growth hormone, bacterial sulfotransferase or antibodies. Because FGE is endogenously expressed in most prokaryotes and eukaryotes, the aldehyde moiety is formed without additional intervention. For labelling with fluorophores, biotin, peptides or PEG probes the aldehyde group was addressed in hydrazone or oxime forming reactions.^{59,60} The endogenous location of FGE in the endoplasmic reticulum was exploited to specifically trace aldehyde-tagged proteins such as IgG antibody, platelet-derived growth factor receptor (PDGFR) and glycoprotein CD4 along the secretory pathway.⁶¹ Recently, a natural glycosylation site within an Fc fragment was replaced by an aldehyde tag, which allowed the introduction of tailor-made glycoforms by means of oxime-forming reactions.⁶⁰ Recent work with recombinantly expressed FGE suggests that Cu^{II} is required as a cofactor in order to achieve high turnover in formyl generating reactions.⁶²

A) Sortase A



B) Formylglycine-generating enzyme




Fig. 3 Protein labelling via (A) sortase A (SrtA) and (B) formylglycine-generating enzyme (FGE).



Enzymatic labelling techniques provide high specificity, relatively small tag sizes and covalently bound labels. Yet there are disadvantages. To achieve high labelling yields rather long reaction times and/or high concentrations of substrate are required. One of the fastest approaches to stain proteins in living cells was reported for BirA with a total labelling time of 20 min.²³ However, the second step of this two-step labelling method had to be performed at 4 °C to reduce endocytosis of the target protein EGFR. Labelling with SrtA and fluorescent oligoglycine conjugates was reported to be achieved within 15 min on living HEK293T cells. But the concentration of the oligoglycine substrate was considerably high with 10 µM.⁵⁶ At high concentrations of substrate the specificity of labelling can be at stake and excessive washing may be required to remove non-covalently bound substrate. PPTase-mediated protein labelling is even performed at 1–150 µM concentration of biotin-CoA,^{36,37} AlexaFluor488- or TexasRed-CoA³⁸ probes. Most other enzyme-based labelling methods require higher concentrations of substrate. For example, 1 mM ketone-containing biotin derivative is applied in ligations by BirA.²³ Some enzyme-catalysed reactions such as the transpeptidase reaction catalysed by SrtA are reversible which leads to decreases in labelling yields if the substrate is applied at low concentration.^{52,53} Many enzymes depend on cofactors such as ATP (e.g. BirA,^{23,24} LplA,^{25,28,31} TTL³³), Mg²⁺ (e.g. BirA,^{23,24} LplA,^{25,28,31} TTL,³³ PPTase^{38,39}) or Ca²⁺ (e.g. TGase,^{46,47} except mTGase⁴⁸) which may interfere with the biological process under investigation. With regard to the labelling time, the two-step procedures significantly prolong the process. Usual incubation times in the first step take 10–60 min (e.g. LplA,^{25,28} TTL,³³ AnkX,⁴⁹ SrtA,^{54–56} BirA,²³ TGase^{46,47}). The bio-orthogonal labelling reactions in the second

step require up to 3 h (e.g. SPAAC, Staudinger ligation)³³ or even overnight treatment (Cu-promoted alkyne–azide cycloaddition).⁴⁸

However, a rapid labelling at low concentration of labelling agents is desirable in studies of fast biological processes. This feature can be obtained with peptide-tags that have high affinity for reactive metal complexes, small molecules or peptides and thereby enable reactions at high effective molarity, which provides for efficient labelling with reagents at nanomolar concentrations.

Recognition of metal ions and small molecules by peptide-tags – mechanisms and characteristics

The selectivity of labelling can be ensured by the mutual recognition between electron-deficient metal ions and electron-rich ligands. The binding of small metal ion probes, in most cases by a chelating effect, facilitates a one-step labelling of the target protein (see Table 2).

One of the shortest peptide-tags is comprised of tetracysteine or tetraserine motifs. In a pioneering study, Tsien *et al.* showed in 1998 that organobisarsenic acid thioesters form covalent bonds with peptide-tags containing a tetracysteine core CCxxCC, most commonly CCGPCC. Upon binding to the thiol groups of the peptide-tag, the Fluorescein-Arsenical-Helix-Binders (FLAsH) increase their fluorescence approximately 50 000-fold (Fig. 4A). In order to reduce the cytotoxicity and non-specific binding of FLAsH, micromolar concentrations of 1,2-ethandiol (EDT) have to be distributed to the cells simultaneously.^{63,64} Derivatives such as ReAsH,⁶⁴ CHOxAsH,⁶⁴ CrAsH⁶⁵ and Cy3As⁶⁶ expand the scope of available dyes. In many cases,


Table 2 Characteristics of metal ion recognition tags and small molecule binding motifs

Labelling technique	Recognition tag and labelling probes	Reported tag position	Typical/reported concentrations and incubation conditions	Reported applications
Tetracysteine motif	Tag: CCxxCC Probes: FLAsH, ReAsH, CHOxAsH, CrAsH, Cy3As	N-terminal, C-terminal, Internal	Labelling with 1–5 µM FLAsH probe for 30–90 min in living cells; reaction temperature: RT ^{63,64}	Fixed and live cell staining
Tetraserine motif	Tag: SSPGSS Probe: RhoBo	N-terminal, C-terminal, Internal	Labelling with 50 nM probe for 20 min in living cells; reaction temperature: 37 °C ⁶⁸	Live cell staining
His-tag	Tag: His ₆ or His ₁₀ Probes: fluorescent conjugates of Ni ^{II} :bis/tris/tetrakis-NTA; HisZiFit or Zn ^{II} :IDA-Cy5	N-terminal, C-terminal	Labelling with 100–500 nM probe for 1–60 min; reaction temperature: RT ^{72,76,77}	Fixed and live cell staining
Oligo-aspartate-tag	Tags: (D4) ₃ , CA ₆ (D4) ₂ , D3 (DDD or DDDXXDDD), DYKDDDDK, (DDDW) ₃ DDD Probes: fluorescent conjugates of Zn ^{II} :DpaTyr or Ni ^{II} :DpaTyr	N-terminal, C-terminal	Labelling with 1–2 µM for 5–7 min (non-covalent); reaction temperature: 25 °C ⁷⁸ / 20 µM probe for 12 h (covalent); reaction temperature: 4 and 20 °C ⁷⁹	Live cell staining
Lanthanide-binding-tag	Tags: ACADYNKDGWYEELECAA or FIDTNNDGWIEGDELLLEEG Probe: Tb ³⁺	Within target sequence	—	<i>In vitro</i> staining
Fluorett-tag	Tag: CCGGGSKVILFERWTWEPISGAP-GSKVILFEGGPG (ReacTR-tag) Probes: Texas red, Rhodamine red, Oregon green 514, fluorescein	C-terminal	Labelling with 20 µM probe for 1 h; reaction temperature: 37 °C ⁸⁸	Live bacteria staining
dC10α-tag	Tag: LSAAECAAREAAACREAAARAGGK Probes: maleinimide-coumarin compounds	N-terminal, C-terminal	Labelling with 10 µM probe for 30 min; reaction temperature: 37 °C ⁹¹	Live cell staining

RT – room temperature.



A) Tetracysteine - FIAsh tag



B) Tetraserine - RhoBo tag




Fig. 4 Protein labelling by reactions between (A) tetracysteine tag and a bisarsenical profluorescent probe or (B) tetraserine tag and a bisboronate.

flanking sequences have been added to the CCxxCC motif to increase the selectivity.⁶⁷ When administered in 2.5 μM concentration, the bisarsenical dyes permeate the cell membrane and enable intracellular labelling of CCxxCC tagged proteins.⁶⁴ Background signals due to interactions with other thiol-containing proteins often require time-consuming washing procedures. Furthermore, the incorporation of such cysteine-rich sequences can promote incorrect disulphide-bond formation within the POI and disturb the integrity of the protein. Schepartz presented bisboronic acid dyes as alternatives of the bisarsenical dyes (Fig. 4B). A rhodamine-derived bisboronic acid (RhoBo) showed nanomolar affinity for the tetraserine motif SSPGSS and provided increases in fluorescence after binding.⁶⁸ But in contrast to the tetracysteine motif, the target sequence SSPGSS is found in more than 100 human proteins, amongst them the highly abundant myosin heavy chain.

Improved specificity can be achieved by metal-chelating tags, which recognise rather unnatural repetitive sequences such as the His₆-, His₁₀-tag or the oligo-aspartate tag. The hexahistidine tag (His₆-tag), originally developed as an affinity tag for protein purification, is as short as the tetracysteine or tetraserine motif and known to interact tightly with transition-metal complexes, e.g. Ni^{II}:nitrilotriacetic acid (Ni^{II}:NTA).⁶⁹ Since the interactions between metal ion and ligand are non-covalent, high mutual affinities are required to achieve stable labelling. To further increase the stability of the label, Tampé, Piehler and colleagues developed multivalent complexes carrying two to four NTA moieties (called bis-, tris- and tetrakis-NTA, Fig. 5A).⁷⁰ The trisNTA ligands have exceptionally high affinities for His-tags (Ni^{II}:trisNTA-His₁₀; $K_D = 0.1$ nM).⁷¹ The attachment of bright fluorophores allowed the single-molecule super-resolution



A) His₆-tag with Ni^{II}:trisNTAB) CA₆D₄-tag with Zn^{II}:DpaTyr

Fig. 5 Protein labelling via interactions between (A) a histidine tag and a fluorescent Ni^{II}:trisNTA compound or (B) an oligo-aspartate tag (e.g. CA6D4 tag) and a fluorescent Zn^{II}:DpaTyr conjugate.

microscopy of actin and lamin filaments in CHO-K1 cells.⁷² Allbritton designed a trifunctional probe comprised of a Ni^{II}:NTA guiding unit and a photo-reactive arylazide group which enabled the covalent linkage with a His₆-tagged murine dihydrofolate reductase (mDHFR) *in vitro*. The third functionality was either a biotin or a DNA strand for further conjugation or immobilization of the POI *in vitro*.⁷³ Trifunctional probes containing a reporter dye were used by Auer *et al.* for the covalent labelling of a His₆-tagged interleukin-4 receptor on the surface of living cells.⁷⁴ A covalent modification of His-tagged proteins was recently introduced by Tørring and Gothelf. The so-called DNA-templated protein conjugation (DTPC) describes the binding of a trisNTA-oligonucleotide to a His₆-tagged POI. This allows site-selective recognition by a second complementary oligonucleotide carrying a reactive *N*-hydroxysuccinimide (NHS) moiety, which enables subsequent crosslinking to a lysine residue in close proximity within the POI.⁷⁵ This was demonstrated for His₆-tagged serotransferrin and native IgG1 antibodies, which contain a histidine cluster. Tsien *et al.* found that the hexahistidine tag not only binds Ni^{II}:NTA probes, but also a Zn^{II} complex (HisZiFiT-Zn^{II}) with a $K_D \approx 40$ nM.⁷⁶ Since the diamagnetic Zn^{II} is not a fluorescence quencher (other than the paramagnetic Ni^{II}), fluorescein was derivatized in a way to



directly bind two Zn^{2+} ions. Incubation with 100 nM HisZiFiT in buffer with 1 μM free Zn^{2+} and two subsequent washing steps allowed the labelling of cell surface-exposed His₆-STIM1-CFP. Hamachi *et al.* introduced a binuclear Zn^{II} :iminodiacetic acid (IDA) complex to which a Cy5 fluorophore was attached *via* an amide bond. Two His₁₀-tagged G protein-coupled receptors (B2R and m1AChR) could be stained on the surface of HEK293 cells when treated with 0.5 μM Zn^{II} :IDA-Cy5 conjugate for 10 min.⁷⁷

In 2006 Hamachi *et al.* introduced the oligo-aspartate tag (D4 tag) and the corresponding multinuclear Zn^{II} :bis((dipicolylamino)methyl)tyrosine (Zn^{II} :DpaTyr) complexes.⁷⁸ A binuclear Zn^{II} :DpaTyr had a moderate affinity for the D4 tag ($K_{\text{D}} = 1.4 \mu\text{M}$). But by elongation of the tag (D4x3 tag) and utilization of the dimeric Zn^{II} :DpaTyr-FITC complex (harbouring four Zn^{II} ions) at 2 μM concentration, the fluorescence labelling of the muscarinic acetylcholine receptor (m1AChR) was feasible on live cells. However, the interaction is non-covalent and the label will dissociate at prolonged times after labelling. To prevent undesired label losses, a nucleophilic cysteine residue was added to the D4 tag and the binuclear Zn^{II} :DpaTyr complex was equipped with an α -chloroacetyl group (Fig. 5B). After incubation for 12 h with 20 μM of the reactive Zn^{II} :DpaTyr-TAMRA conjugate, the probes were found to enter *E. coli* cells and covalently label intracellular CA₆D₄-tagged maltose binding protein.⁷⁹ The D4/ Zn^{II} :DpaTyr system was reported to be orthogonal to the His₆/Ni^{II}:NTA complex.⁷⁸ Later it was described that Ni^{II}:DpaTyr complexes show remarkably high affinity for the FLAG tag (DYKDDDDK), the D3 tag (DDD) and D3x2 tag (DDDXXDDD), with binding affinities around 100-fold stronger than for the D4/ Zn^{II} :DpaTyr pair.^{80,81} More recently, Komatsu *et al.* designed a Tb³⁺-binding peptide (LBP) *de novo* with aspartic acid as Tb³⁺-binding sites and tryptophan as sensitizing dye. The 15-mer sequence DDDWDDWDDWDDWDD was genetically fused to the C-terminus of glutathione S-transferase (GST) and used for affinity purification on a Tb³⁺-loaded column, as well as for in-gel detection of the fusion protein, rendering the LBP especially useful due to its dual functionality.⁸²

While transition metal ions such as Ni^{II}, Co^{II} and Zn^{II} confer high affinity for recognition tags as described in the labelling techniques above, lanthanide ions add further desirable features; long-lived luminescence in the presence of sensitizing chromophores, excellent X-ray scattering powers and some of them even magnetic properties for NMR studies (*e.g.* Gd^{III}, Dy^{III}, Tb^{III}, Tm^{III}).⁸³ This prompted a search for peptide-tags with high affinity for lanthanides to proteins. Imperiali *et al.* optimized a sequence found in a loop region of the calmodulin protein family, by means of a split-and-pool library and identified an 18-mer lanthanide-binding tag (LBT, ACADYNKDGWYEELECAA). With a K_{D} value of around 220 nM for the binding of Tb³⁺, the LBT tag could be used for concentration determination in cell lysate and for in-gel visualization of a LBT-ubiquitin fusion protein.⁸⁴ Later, Imperiali also found cysteine-free LBTs with nanomolar affinities for Tb³⁺ (YIDTNNDGWYEGDELLA, $K_{\text{D}} = 57 \text{ nM}$;⁸⁵ FIDTNNDGWIEGDELLLEEG, $K_{\text{D}} = 19 \text{ nM}$)⁸⁶.

The search for peptide sequences that enable fluorescence labelling of proteins without added enzymes was extended from




Fig. 6 Covalent labelling with reactive Fluorette (fluorophore binding peptide) tag.

metal binding tags to tags, which have the ability to directly bind small molecule fluorescent probes. Already in 1998, Nolan *et al.* discovered by phage display a panel of Texas red, Rhodamine red, Oregon green 514 and fluorescein binding peptides, named “Fluorettes” (Fig. 6).⁸⁷ Amongst them, the TexasRed-binding peptides (TR) showed the highest, subnanomolar affinities. Jäschke converted the TR peptide into a reactive tag (ReacTR, sequence: CCGGSKVILFEGPAGRWTWEPISEGAPGSKVILFEGGPG) by adding two cysteine residues, which can undergo a proximity-induced nucleophilic substitution reaction with the *N*- α -chloroacetamide-conjugated TexasRed. After incubation with 10 μM reactive TexasRed probe for 1 h, the tagged maltose binding protein (MBP-ReacTR) showed bright fluorescence in the cytosol of transfected *E. coli* cells.⁸⁸

Barbas *et al.* subjected phage-displayed peptide libraries to a reaction-based selection. A 21-mer peptide (rpf1368, sequence: CHNHQKATCRRMRSRETSVKK) was identified which formed an enaminone with 1,3-diketone derivatives. The reaction rates were rather slow with a labelling yield of 90% after 10 h.⁸⁹ Also employing phage display, Weiss *et al.* searched for peptides that react with hydrazides. The resulting hydrazine reactive peptide (HyRe; sequence: HKTNHSCHKREQEHCRTVT) was fused to the T4 lysozyme and underwent selective modification with 1 mM biotin hydrazide as well as with 1 mM rhodamine B hydrazide after 1 h reaction time in a crude cell lysate.⁹⁰

The rational design of a maleimide reactive α -helical peptide led to the dC10 α tag (sequence: LSAAECAAREAACREAAARAGGK), in which two cysteine residues are separated by two turns of an α -helix. The minimalistic probe molecule comprises a coumarin fluorophore and a dimaleimide moiety that quenches the fluorescence until both maleimide groups undergo thiol addition during the labelling reaction. Owing to the small size, the probe was able to enter HEK293T cells after incubation for 30 min at 10 μM concentration and label both, C-terminally dC10 α -tagged histone H2B as well as N-terminally tagged actin.⁹¹

The small molecule-binding peptide-tags provide properties that allow protein labelling in complex mixtures. However, for the achievement of a high affinity tag metal complexes seem to require less amino acid residues than small molecules (*e.g.* 6 aa tag for FLASH vs. 20 aa HyRe tag for rhodamine B hydrazide). High affinity usually provides faster labelling at lower concentrations of labelling agent and, therefore, the reported labelling times for metal-chelating tags in living cells can be as short as 5–7 min in case of oligo-aspartate tagged (D4)₃-m1AChR⁷⁸ or 20 min for tetracysteine-tagged proteins.⁶⁴ Moreover, metal-chelating tags require significant lower amounts of labelling probes with the lowest of 100 nM, as demonstrated for His₁₀/Ni^{II}:trisNTA pair.⁷²



These characteristics demonstrate advantages, also in comparison to enzymatic labelling techniques, however, the labelling of the POI remains non-covalent and thus reversible, hindering unambiguous analyses of low abundance proteins. The reported attempts to establish a covalent linkage between tag and probe employed moderately reactive groups to avoid background labelling. The low reactivity typically resulted in slow rates of covalent labelling, which is disadvantageous for pulse-chase-type experiments.

Peptide and protein based recognition by peptide-tags – mechanisms and characteristics

One way to improve the binding affinity of a tag sequence for the labelling agent is to increase its size. This led to the notion of peptide-binding peptide-tags. Peptides can bind other peptides with high mutual affinity (see Table 3). One example for high affinity peptide–peptide interactions is the coiled-coil motif, in which helical peptides wrap around each other to form a superhelix. In 1996, Hodges *et al.* designed a coiled-coil peptide *de novo*.⁹² The dimerization domain consists of five repeating heptads of the sequences EVSALEK for the negatively charged E-coil and KVSALKE for the positively charged K-coil. Electrostatic interactions drive the high-affinity heterodimer formation ($K_D = 1$ nM). Using the E-coil as genetically encoded tag, the peptides were used for the purification, immobilization


and detection of peptides expressed in *E. coli*. In 2008, Matsuzaki reported the artificial, heterodimeric coiled-coil peptides K3 (KIAALKE)₃, K4 (KIAALKE)₄, E3 (EIAALEK)₃ and E4 (EIAALEK)₄ – originally developed by Hodges *et al.*⁹³ – as tag/probe pair for the live cell imaging of cell surface receptors. The stability of the E3/K3 coiled-coil is in the nanomolar range ($K_D \approx 70$ nM). With a $K_D \approx 6$ nM the E3/K4 coiled-coil provided even higher stability. This allowed the labelling of an E3-tagged prostaglandin E2 receptor EP3 β subtype (EP3 β R) upon incubation with only 20 nM TMR-conjugated K3 and K4 probe peptide within 5 min.⁹⁴ The method was used to investigate the oligomerization state of two membrane receptors by staining the E3-tagged receptors with a mixture of fluorophore-modified K4 peptides.^{95,96} To avoid a potential loss of label during prolonged imaging periods, the coiled-coil motif was repurposed as a scaffold that instructs reactions to form a covalent bond. Ball *et al.* substituted amino acids at the interface of the E3/K3 coiled-coil pair to position a dirhodium catalyst on one coil so that addition of a diazo agent induces the alkylation of a tryptophan residue in the other coil.⁹⁷ The E3_gW-tagged maltose binding protein was selectively biotinylated in cell lysate by treatment with 5 μ M K3_{a,e}Rh₂ catalyst for 16 h in presence of 100 μ M biotin-diazo reagent (Fig. 7A).⁹⁸ Xia *et al.* also harnessed the concept of proximity-induced reactivity by modifying the E3-coil with a nucleophilic thiol provided by a cysteine residue within the E3-coil (CCE probe) and the K3-coil with an electrophilic α -chloroacetyl moiety (CCK probe) (Fig. 7B). Positioning of the two reactive residues at the interface of the coiled-coil

Table 3 Characteristics of tag systems based on peptide–peptide and peptide–protein interactions

Labelling technique	Recognition tag including peptide probes	Reported tag position	Typical/reported concentrations and incubation conditions	Reported applications	Labelling modes
E3/K3 or E3/K4	Tag: (EIAALEK) ₃ Probe: dye-(KIAALKE) ₃ , dye-(KIAALKE) ₄	N-terminal	Labelling with 20–60 nM for 2 min; reaction temperature: 37 °C ⁹⁴	Cell surface labelling, live cell imaging	Non-covalent
CCE/CCK	Tag: GGGE-CAALEKE-VAALEKE-VAALEK (CCE) Probe: GGGK-XAALKEK-VAALKEK-VAALKE (CCK)	N-terminal	Labelling with 0.2–1 μ M probe for 20 min; reaction temperature: RT ⁹⁹	Cell surface labelling, live cell staining, fixed for imaging	Covalent
E3 _g W/ K3 _{a,e} Rh ₂	Tag: EISALEKWISALEQEISALEK (E3 _g W) Catalyst: KISALKQKESALEQKISALEK (K3 _{a,e} Rh ₂ – between E dirhodium probe)	C-terminal	Labelling with 5 μ M K3 catalyst and 100 μ M biotin-diazo probe for 16 h; reaction temperature: 4 °C ⁹⁸	<i>In vitro</i>	Covalent
ER3/R3CL	Tag: M(EIAALER) ₃ -GK (ER3) Probe: dye-(RIAALRE) ₃ -GC (R3CL)	N-terminal	Labelling with 150 μ M probe for 20 min; reaction temperature: 25 °C ¹⁰⁰	Cell surface labelling, live cell imaging	Covalent
Cys-E3/K3 thioester	Tag: C(EIAALEK) ₃ Probe: dye-thioester-(KIAALKE) ₃	N-terminal	Labelling with 100 nM probe for 2–5 min; reaction temperature: RT ^{101,103}	Cell surface labelling, live cell imaging	Covalent
Leucine zipper	ZIP-tag: ALEKELA-EAEKELA-ELEKELA-GGCGG-ALEKKLA-ELEKKA-ELEKKLA Probe: ALKKLA-EzKKLA-ELKKLA	N-terminal	Labelling with 0.1–1 μ M probe for 4–15 min; reaction temperature: 25 or 37 °C ^{104,106}	Cell surface labelling, in combination with CPP intracellular labelling, live cell imaging	Non-covalent and covalent
BTX-SA-tag	BTX-tag: WRYESSLEPYPD SA-tag: MDEKTTGWRGGHVVE-GLAGELEQLRARLEHHPQGQREP	N-terminal	Labelling with 1–2 μ g ml ⁻¹ BTX for 30 min; reaction temperature: 17 or 37 °C ¹⁰⁹	Cell surface labelling, live or fixed cell imaging	Non-covalent
DnaE intein	<i>Npu</i> : N _{int} aa 1–123, C _{int} aa 103–138 <i>Ava</i> : N _{int} aa 1–102, C _{int} aa 103–138	N-terminal, C-terminal	<i>Npu</i> : labelling with 2 μ M C _{int} peptide for 1 h; reaction temperature: 37 °C ¹¹⁴ / <i>Ava</i> : labelling with 2.5 μ M C _{int} for 1 h ¹¹⁶	Intracellular labelling via fusion to CPP	Covalent
AceL/Ter intein	N _{int} aa 1–25, C _{int} aa 26–129	N-terminal	Labelling with 45 μ M N _{int} peptide >1 h; reaction temperature: 8, 25 or 37 °C ¹¹⁷	<i>In vitro</i>	Covalent

(* = temperature was not explicitly specified) RT – room temperature.





The diagram illustrates the crosslinking reaction between a protein of interest (POI) and a cysteine-containing tag (CCE) using a cysteine-containing crosslinker (CCK). The reaction proceeds as follows:

- Reactants:** A protein of interest (POI) fused to a cysteine-containing tag (CCE-SH) and a cysteine-containing crosslinker (CCK-Cl) with a red star indicating its binding site.
- Reaction:** The CCK molecule binds to the CCE tag, forming a crosslink. This process is labeled "crosslinking upon CCE and CCK binding".
- Products:** The final complex shows the CCK molecule bound to the CCE tag, which is still attached to the POI. A chloride ion (Cl^-) is released during the reaction.

With an aim to find a very fast biocompatible reaction for covalent protein labelling that adds only little mass, Seitz *et al.* designed a coiled-coil-promoted acyl transfer reaction, which also bypasses the ligation between tag and probe. For this purpose, an N-terminal cysteine residue was added to the E3 tag and fused to the POI. A thioester-linked fluorophore was conjugated to the N-terminus of the K3 probe.¹⁰¹ The formation

This journal is © The Royal Society of Chemistry 2016





Fig. 8 Covalent protein labelling *via* coiled-coil induced reporter group transfer.

Another small and stable coiled-coil scaffold is the leucine zipper. Based on the crystal structure of a coiled-coil trimer of GCN4, Tamamura *et al.* developed in 2009 artificial leucine zipper peptides with high mutual affinity (ZIP tag/probe pair, $K_D = 18$ nM) for the fluorescence imaging of ZIP-fused membrane proteins. The 23 aa probe helix is equipped with an environmentally sensitive dye (4-nitrobenzo-2-oxa-1,3-diazole, NBD), the other two tag helices are connected *via* a loop sequence and offer a hydrophobic pocket for the dye (A2-tag, 49 aa) (Fig. 9A). The chemokine receptor CXCR4 was genetically fused to the A2-tag and expressed in CHO cells. Incubation of the cells with 1 μ M NBD-probe peptide led to a fluorescence change.¹⁰⁴ To increase the chemical and biological stability, Tamamura *et al.* later incorporated an α -chloroacetyl group to the probe peptide that can crosslink to a cysteine residue in the


loop region of the A2-tag.¹⁰⁵ For intracellular protein targets, an octaarginine sequence was attached to the C-terminus of the probe peptide.¹⁰⁶

Peptide-based recognition of genetically encoded protein tags beyond coiled-coil interactions was introduced by Haganir *et al.* in 2004. The labelling system relied on α -bungarotoxin (BTX, 74 aa); a small protein from snake venom.¹⁰⁷ Peptide sequences that bind to BTX were initially selected by phage display and modified for high affinity binding.¹⁰⁸ The resulting 13 aa BTX-binding site (BBS, sequence: WRYESSLEPYPD, $K_D \approx 60$ nM¹⁰⁹) was placed in the extracellular domain of the AMPA receptor for analysing the expression and membrane trafficking of the tagged receptor. At the same time, Sanes *et al.* not only used the BTX-binding site, but also the streptavidin-binding sequence (SBP, sequence: MDEKTTGWRGG HVVEGLAGELEQLRARLEHHPQGQREP,¹¹⁰ $K_D \approx 25$ nM¹⁰⁹) to stain

A) Leucine Zipper tag



B) Split-intein



C) Cell-penetrating split-intein




Fig. 9 Protein labelling (A) with the leucine zipper tag, which allows covalent labelling and turn-on of fluorescence, (B) by split-intein mediated reactions or (C) by cell-penetrating split-intein mediated reactions.



the C-terminally tagged vesicle-associated protein (VAMP2) on the surface of live cells by incubation with the respective fluorophore-protein (BTX or SA) conjugates.¹⁰⁹ Attempts to use BTX binding site or SBP for covalent labelling have not been reported but seem rather challenging given the size of BTX and SA.

The development of peptide-based tag/probe pairs for protein labelling seems to be characterised by a trade-off between the specificity of a tag and its size. Split intein-mediated protein labelling circumvents this correlation because intein excises itself after ligation of the flanking sequences (Fig. 9B). One intein fragment should be short in order to allow efficient solid-supported peptide synthesis. While such split inteins have been created artificially, they usually lack sufficient splicing performance and solubility.¹¹¹ The naturally occurring *Npu* DnaE intein is one of the fastest inteins reported with a half-life of only 1 min.¹¹² In 2012, Camarero *et al.* equipped the 36 aa C-intein (C_{int}) of *Npu* DnaE with a FRET pair in order to quench the fluorescence in the unspliced state. Using a commercial protein delivery system, the intracellular labelling of transcription factor Yin Yang 1 took place within 2 h using the FRET-quenched protein *trans*-splicing reaction (PTS).¹¹³ In 2015, Kwon engineered a *Npu* DnaE intein and synthesized a C_{int} with a photo-protected ester-linkage for controlled activation, a FRET pair for turn-on fluorescence detection and an octaarginine as cell penetrating peptide (Fig. 9C).¹¹⁴ As a proof of concept, labelling of maltose binding protein *via* PTS was carried out in HeLa cells. Detection of fluorescence was observed within the cells after 30 min incubation time. Beside the *Npu* DnaE intein other fast split inteins were described such as the *Ava* DnaE intein.¹¹⁵ In a recent study, histone H2B was fused to a 102 aa long *Ava* DnaE N-intein (N_{int}) and expressed in HEK293T cells. Live cell imaging was demonstrated after 1 h incubation with *Npu* DnaE C_{int} carrying different reporter groups. *Npu* DnaE C_{int} was used instead of *Ava* DnaE C_{int} due to higher synthesis yields.¹¹⁶ Mootz *et al.* engineered an atypical naturally occurring split intein with an N_{int} fragment consisting of 25 aa only and a C_{int} of 100 aa.¹¹⁷ Recently, Tampé *et al.* significantly improved the affinity of a *Ssp* DnaB M86 split intein composed of an N_{int} with 11 aa and a C_{int} with 143 aa. The N_{int} was C-terminal modified with *tris*NTA moieties and the C_{int} was N-terminal fused to a His₆-tag which increased the affinity up to 40 fold and allowed protein *trans*-splicing at nanomolar concentrations.¹¹⁸

Intein technology is interesting, because the tag is excised upon labelling. Yet, to retain the benefits of small peptide-tags, one short, POI-connected intein fragment has to be combined with a rather large intein fragment that brings in the labelled extein. This construct must be recombinantly expressed and conjugated with a reporter group, which may limit the application.

Conclusion

The ideal protein labelling method should proceed with high tag specificity, require only small tag sizes (to avoid interference with protein function), succeed within short incubation times

(to enable investigations of fast biological processes in pulse-chase-type experiments), confer stability of labelling (preferably by formation of covalent linkages to allow measurements over prolonged time scales), enable versatility as far as the choice of labelling groups is concerned and lay little toxic burden onto the cell or the cellular process under scrutiny. None of the tag-based methods discussed is without shortcomings and there is, at current, no silver bullet available for protein labelling. Enzymatic methods provide high specificity, small tag size and covalent labels, but with the enzymes currently used rather high substrate concentrations and long incubation times are required. The stability and acquisition of the enzymes itself are not always without problems and labelling of intracellular proteins remains challenging. Nevertheless, the growing number of publications in which enzymes such as BirA are used, for example to study protein–protein interactions by proximity biotinylation demonstrate that problems are manageable and that the method allows applications that are difficult to be performed by other means. Notably short incubation times are feasible with metal ion recognition tags. Recent studies demonstrate that the placement of bright dyes such as AlexaFluor 647 in immediate vicinity of the protein *via* the His_{6–10} tags and high affinity Ni^{II}:*tris*NTA complexes facilitates super resolution microscopy of fixed and permeabilized cells. Ideally, the peptide-tag would be targeted by a membrane permeable probe. Though numerous studies demonstrate the feasibility of this approach, the available peptide-tags require rather long sizes to confer affinities for fluorescent dyes in a range that require staining with $\geq 10 \mu\text{M}$ concentration; which may lead to high background. Higher tag specificities are achievable with peptide based affinity reagents. For example, the *de novo* designed E3/K3 coiled-coil offers high affinity and selectivity at comparably small tag size (21 aa). This enables rapid labelling at low concentrations of probe. A recently developed peptide-templated acyl transfer technique combines the speed of complex formation *via* coiled-coil interactions with the advantages of covalent labelling. The method involves a highly reactive thioester-linked fluorophore K3 conjugate, which rapidly reacts with a cysteine terminated E3-coil. The covalent labelling is extremely fast (2–5 min) and only little cargo is added to the protein of interest. Yet at current, the method is restricted to N-terminal labelling of cell surface proteins.

It is obvious that the peptide-tag based labelling methods available as of today still do not offer the robustness and wide applicability known from autofluorescent proteins and enzyme based tags (*e.g.* SNAP, CLIP, Halo). Yet, given the scientific need for less perturbing labelling methods, considering the pace of the current development of the field and taking into account that an ever increasing number of chemists joins the race, the authors are convinced that peptide based tags will become game changing enablers in a not too distant future.

References

- 1 R. Rizzuto, M. Brini, P. Pizzo, M. Murgia and T. Pozzan, *Curr. Biol.*, 1995, **5**, 635–642.
- 2 A. Keppler, S. Gendreizig, T. Gronemeyer, H. Pick, H. Vogel and K. Johnsson, *Nat. Biotechnol.*, 2003, **21**, 86–89.



- 3 A. Gautier, A. Juillerat, C. Heinis, I. R. Corrêa Jr., M. Kindermann, F. Beaufls and K. Johnsson, *Chem. Biol.*, 2008, **15**, 128–136.
- 4 C. Jing and V. W. Cornish, *ACS Chem. Biol.*, 2013, **8**, 1704–1712.
- 5 L. W. Miller, Y. Cai, M. P. Sheetz and V. W. Cornish, *Nat. Methods*, 2005, **2**, 255–257.
- 6 G. V. Los, L. P. Encell, M. G. McDougall, D. D. Hartzell, N. Karassina, C. Zimprich, M. G. Wood, R. Learish, R. F. Ohana, M. Urh, D. Simpson, J. Mendez, K. Zimmerman, P. Otto, G. Vidugiris, J. Zhu, A. Darzins, D. H. Klaubert, R. F. Bulleit and K. V. Wood, *ACS Chem. Biol.*, 2008, **3**, 373–382.
- 7 E. A. Rodriguez, H. A. Lester and D. A. Dougherty, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 8650–8655.
- 8 H. Neumann, *FEBS Lett.*, 2012, **586**, 2057–2064.
- 9 S. Tsukiji, M. Miyagawa, Y. Takaoka, T. Tamura and I. Hamachi, *Nat. Chem. Biol.*, 2009, **5**, 341–343.
- 10 T. Hayashi and I. Hamachi, *Acc. Chem. Res.*, 2012, **45**, 1460–1469.
- 11 M. Rashidian, J. K. Dozier and M. D. Distefano, *Bioconjugate Chem.*, 2013, **24**, 1277–1294.
- 12 Z. Ma, L. Du and M. Li, *J. Med. Chem.*, 2014, **57**, 8187–8203.
- 13 D. Jung, K. Min, J. Jung, W. Jang and Y. Kwon, *Mol. Biosyst.*, 2013, **9**, 862–872.
- 14 Y. Takaoka, A. Ojida and I. Hamachi, *Angew. Chem., Int. Ed.*, 2013, **52**, 4088–4106.
- 15 D. Beckett, E. Kovaleva and P. J. Schatz, *Protein Sci. Publ. Protein Soc.*, 1999, **8**, 921–929.
- 16 E. de Boer, P. Rodriguez, E. Bonte, J. Krijgsveld, E. Katsantoni, A. Heck, F. Grosveld and J. Strouboulis, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 7480–7485.
- 17 M. Fernández-Suárez, T. S. Chen and A. Y. Ting, *J. Am. Chem. Soc.*, 2008, **130**, 9251–9253.
- 18 E. Steel, V. L. Murray and A. P. Liu, *PLoS One*, 2014, **9**, e93646.
- 19 M. Lajko, A. F. Haddad, C. A. Robinson and S. A. Connolly, *J. Virol. Methods*, 2015, **221**, 81–89.
- 20 M. Desbois, S. J. Cook, S. W. Emmons and H. E. Bülow, *Genetics*, 2015, **200**, 697–705.
- 21 C. A. Bertuccio, S.-L. Lee, G. Wu, M. B. Butterworth, K. L. Hamilton and D. C. Devor, *PLoS One*, 2014, **9**, e92013.
- 22 A. B. Jaykumar, P. S. Caceres, I. Sablaban, B. A. Tannous and P. A. Ortiz, *Am. J. Physiol. Renal Physiol.*, 2016, **310**, F183–F191.
- 23 I. Chen, M. Howarth, W. Lin and A. Y. Ting, *Nat. Methods*, 2005, **2**, 99–104.
- 24 S. A. Slavoff, I. Chen and A. Y. Ting, *J. Am. Chem. Soc.*, 2008, **130**, 1160–1162.
- 25 M. Fernández-Suárez, H. Baruah, L. Martínez-Hernández, K. T. Xie, J. M. Baskin, C. R. Bertozzi and A. Y. Ting, *Nat. Biotechnol.*, 2007, **25**, 1483–1487.
- 26 D. E. Green, T. W. Morris, J. Green, J. E. Cronan and J. R. Guest, *Biochem. J.*, 1995, **309**(pt 3), 853–862.
- 27 J. D. Cohen, P. Zou and A. Y. Ting, *ChemBioChem*, 2012, **13**, 888–894.
- 28 D. S. Liu, A. Tangpeerachaikul, R. Selvaraj, M. T. Taylor, J. M. Fox and A. Y. Ting, *J. Am. Chem. Soc.*, 2012, **134**, 792–795.
- 29 J. Z. Yao, C. Uttamapinant, A. Poloukhine, J. M. Baskin, J. A. Codelli, E. M. Sletten, C. R. Bertozzi, V. V. Popik and A. Y. Ting, *J. Am. Chem. Soc.*, 2012, **134**, 3720–3728.
- 30 M. Best, A. Degen, M. Baalman, T. T. Schmidt and R. Wombacher, *ChemBioChem*, 2015, **16**, 1158–1162.
- 31 S. Puthenveetil, D. S. Liu, K. A. White, S. Thompson and A. Y. Ting, *J. Am. Chem. Soc.*, 2009, **131**, 16430–16438.
- 32 S. Hauke, M. Best, T. T. Schmidt, M. Baalman, A. Krause and R. Wombacher, *Bioconjugate Chem.*, 2014, **25**, 1632–1637.
- 33 D. Schumacher, J. Helma, F. A. Mann, G. Pichler, F. Natale, E. Krause, M. C. Cardoso, C. P. R. Hackenberger and H. Leonhardt, *Angew. Chem., Int. Ed.*, 2015, **54**, 13787–13791.
- 34 J. Yin, F. Liu, X. Li and C. T. Walsh, *J. Am. Chem. Soc.*, 2004, **126**, 7754–7755.
- 35 J. Yin, F. Liu, M. Schinke, C. Daly and C. T. Walsh, *J. Am. Chem. Soc.*, 2004, **126**, 13570–13571.
- 36 J. Yin, A. J. Lin, P. D. Buckett, M. Wessling-Resnick, D. E. Golan and C. T. Walsh, *Chem. Biol.*, 2005, **12**, 999–1006.
- 37 J. Yin, P. D. Straight, S. M. McLoughlin, Z. Zhou, A. J. Lin, D. E. Golan, N. L. Kelleher, R. Kolter and C. T. Walsh, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 15815–15820.
- 38 Z. Zhou, P. Cironi, A. J. Lin, Y. Xu, S. Hrvatin, D. E. Golan, P. A. Silver, C. T. Walsh and J. Yin, *ACS Chem. Biol.*, 2007, **2**, 337–346.
- 39 L. Marchetti, T. De Nadai, F. Bonsignore, M. Calvello, G. Signore, A. Viegi, F. Beltram, S. Luin and A. Cattaneo, *PLoS One*, 2014, **9**, e113708.
- 40 T. Kawamura, B. Stephens, L. Qin, X. Yin, M. R. Does, T. H. Smith, N. Grimsey, R. Abagyan, J. Trejo, I. Kufareva, M. M. Fuster, C. L. Salanga and T. M. Handel, *PLoS One*, 2014, **9**, e81454.
- 41 S. Jeger, K. Zimmermann, A. Blanc, J. Grünberg, M. Honer, P. Hunziker, H. Struthers and R. Schibli, *Angew. Chem., Int. Ed.*, 2010, **49**, 9995–9997.
- 42 A. Mero, B. Spolaore, F. M. Veronese and A. Fontana, *Bioconjugate Chem.*, 2009, **20**, 384–389.
- 43 H. Abe, M. Goto and N. Kamiya, *Chem. – Eur. J.*, 2011, **17**, 14004–14008.
- 44 T. Tanaka, N. Kamiya and T. Nagamune, *FEBS Lett.*, 2005, **579**, 2092–2096.
- 45 J. Tominaga, N. Kamiya, S. Doi, H. Ichinose, T. Maruyama and M. Goto, *Biomacromolecules*, 2005, **6**, 2299–2304.
- 46 C.-W. Lin and A. Y. Ting, *J. Am. Chem. Soc.*, 2006, **128**, 4542–4543.
- 47 C. Gnaccarini, W. Ben-Tahar, A. Mulani, I. Roy, W. D. Lubell, J. N. Pelletier and J. W. Keillor, *Org. Biomol. Chem.*, 2012, **10**, 5258–5265.
- 48 S. K. Oteng-Pabi, C. Pardin, M. Stoica and J. W. Keillor, *Chem. Commun.*, 2014, **50**, 6604–6606.
- 49 K. Heller, P. Ochtrup, M. F. Albers, F. B. Zauner, A. Itzen and C. Hedberg, *Angew. Chem., Int. Ed.*, 2015, **54**, 10327–10330.



- 50 H. Mao, S. A. Hart, A. Schink and B. A. Pollok, *J. Am. Chem. Soc.*, 2004, **126**, 2670–2671.
- 51 X. Guo, Q. Wang, B. M. Swarts and Z. Guo, *J. Am. Chem. Soc.*, 2009, **131**, 9878–9879.
- 52 R. Parthasarathy, S. Subramanian and E. T. Boder, *Bioconjugate Chem.*, 2007, **18**, 469–476.
- 53 Z. Wu, X. Guo and Z. Guo, *Chem. Commun.*, 2011, **47**, 9218–9220.
- 54 J. M. Antos, G. M. Miller, G. M. Grotenbreg and H. L. Ploegh, *J. Am. Chem. Soc.*, 2008, **130**, 16338–16343.
- 55 M. W. Popp, J. M. Antos, G. M. Grotenbreg, E. Spooner and H. L. Ploegh, *Nat. Chem. Biol.*, 2007, **3**, 707–708.
- 56 T. Tanaka, T. Yamamoto, S. Tsukiji and T. Nagamune, *ChemBioChem*, 2008, **9**, 802–807.
- 57 I. Wuethrich, J. G. C. Peeters, A. E. M. Blom, C. S. Theile, Z. Li, E. Spooner, H. L. Ploegh and C. P. Guimaraes, *PLoS One*, 2014, **9**, e109883.
- 58 A. Tabata, Y. Ohkubo, N. Anyoji, K. Hojo, T. Tomoyasu, Y. Tatematsu, K. Ohkura and H. Nagamune, *Anticancer Res.*, 2015, **35**, 4411–4417.
- 59 I. S. Carrico, B. L. Carlson and C. R. Bertozzi, *Nat. Chem. Biol.*, 2007, **3**, 321–322.
- 60 E. L. Smith, J. P. Giddens, A. T. Iavarone, K. Godula, L.-X. Wang and C. R. Bertozzi, *Bioconjugate Chem.*, 2014, **25**, 788–795.
- 61 P. Wu, W. Shui, B. L. Carlson, N. Hu, D. Rabuka, J. Lee and C. R. Bertozzi, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 3000–3005.
- 62 P. G. Holder, L. C. Jones, P. M. Drake, R. M. Barfield, S. Bañas, G. W. de Hart, J. Baker and D. Rabuka, *J. Biol. Chem.*, 2015, **290**, 15730–15745.
- 63 B. A. Griffin, S. R. Adams and R. Y. Tsien, *Science*, 1998, **281**, 269–272.
- 64 S. R. Adams, R. E. Campbell, L. A. Gross, B. R. Martin, G. K. Walkup, Y. Yao, J. Llopis and R. Y. Tsien, *J. Am. Chem. Soc.*, 2002, **124**, 6063–6076.
- 65 H. Cao, B. Chen, T. C. Squier and M. U. Mayer, *Chem. Commun.*, 2006, 2601–2603.
- 66 H. Cao, Y. Xiong, T. Wang, B. Chen, T. C. Squier and M. U. Mayer, *J. Am. Chem. Soc.*, 2007, **129**, 8672–8673.
- 67 C. Hoffmann, G. Gaietta, A. Zürn, S. R. Adams, S. Terrillon, M. H. Ellisman, R. Y. Tsien and M. J. Lohse, *Nat. Protoc.*, 2010, **5**, 1666–1677.
- 68 T. L. Halo, J. Appelbaum, E. M. Hobert, D. M. Balkin and A. Schepartz, *J. Am. Chem. Soc.*, 2009, **131**, 438–439.
- 69 E. Hochuli, W. Bannwarth, H. Döbeli, R. Gentz and D. Stüber, *Nat. Biotechnol.*, 1988, **6**, 1321–1325.
- 70 A. Tinazli, J. Tang, R. Valiokas, S. Picuric, S. Lata, J. Piehler, B. Liedberg and R. Tampé, *Chem. – Eur. J.*, 2005, **11**, 5249–5259.
- 71 S. Lata, A. Reichel, R. Brock, R. Tampé and J. Piehler, *J. Am. Chem. Soc.*, 2005, **127**, 10205–10215.
- 72 R. Wieneke, A. Raulf, A. Kollmannsperger, M. Heilemann and R. Tampé, *Angew. Chem., Int. Ed.*, 2015, **54**, 10216–10219.
- 73 G. D. Meredith, H. Y. Wu and N. L. Allbritton, *Bioconjugate Chem.*, 2004, **15**, 969–982.
- 74 M. Hintersteiner, T. Weidemann, T. Kimmerlin, N. Filiz, C. Buehler and M. Auer, *ChemBioChem*, 2008, **9**, 1391–1395.
- 75 C. B. Rosen, A. L. B. Kodal, J. S. Nielsen, D. H. Schaffert, C. Scavenius, A. H. Okholm, N. V. Voigt, J. J. Enghild, J. Kjems, T. Tørring and K. V. Gothelf, *Nat. Chem.*, 2014, **6**, 804–809.
- 76 C. T. Hauser and R. Y. Tsien, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 3693–3697.
- 77 S. Fujishima, H. Nonaka, S. Uchinomiya, Y. A. Kawase, A. Ojida and I. Hamachi, *Chem. Commun.*, 2011, **48**, 594–596.
- 78 A. Ojida, K. Honda, D. Shinmi, S. Kiyonaka, Y. Mori and I. Hamachi, *J. Am. Chem. Soc.*, 2006, **128**, 10452–10459.
- 79 H. Nonaka, S. Tsukiji, A. Ojida and I. Hamachi, *J. Am. Chem. Soc.*, 2007, **129**, 15777–15779.
- 80 H. Nonaka, S. Fujishima, S. Uchinomiya, A. Ojida and I. Hamachi, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 6696–6699.
- 81 A. Ojida, S.-H. Fujishima, K. Honda, H. Nonaka, S.-H. Uchinomiya and I. Hamachi, *Chem. – Asian J.*, 2010, **5**, 877–886.
- 82 S. Sueda, S. Tanaka, S. Inoue and H. Komatsu, *Anal. Biochem.*, 2012, **422**, 52–54.
- 83 S. Aime, M. Botta, M. Fasano and E. Terreno, *Chem. Soc. Rev.*, 1998, **27**, 19–29.
- 84 K. J. Franz, M. Nitz and B. Imperiali, *ChemBioChem*, 2003, **4**, 265–271.
- 85 M. Nitz, K. J. Franz, R. L. Maglathlin and B. Imperiali, *ChemBioChem*, 2003, **4**, 272–276.
- 86 L. J. Martin, B. R. Sculimbrene, M. Nitz and B. Imperiali, *QSAR Comb. Sci.*, 2005, **24**, 1149–1157.
- 87 M. N. Rozinov and G. P. Nolan, *Chem. Biol.*, 1998, **5**, 713–728.
- 88 M. Sunbul, L. Nacheva and A. Jäschke, *Bioconjugate Chem.*, 2015, **26**, 1466–1469.
- 89 F. Tanaka, R. Fuller, L. Asawapornmongkol, A. Warsinke, S. Gobuty and C. F. Barbas, *Bioconjugate Chem.*, 2007, **18**, 1318–1324.
- 90 G. M. Eldridge and G. A. Weiss, *Bioconjugate Chem.*, 2011, **22**, 2143–2153.
- 91 Y. Chen, C. M. Clouthier, K. Tsao, M. Strmiskova, H. Lachance and J. W. Keillor, *Angew. Chem., Int. Ed.*, 2014, **53**, 13785–13788.
- 92 B. Tripet, L. Yu, D. L. Bautista, W. Y. Wong, R. T. Irvin and R. S. Hodges, *Protein Eng.*, 1996, **9**, 1029–1042.
- 93 J. R. Litowski and R. S. Hodges, *J. Biol. Chem.*, 2002, **277**, 37272–37279.
- 94 Y. Yano, A. Yano, S. Oishi, Y. Sugimoto, G. Tsujimoto, N. Fujii and K. Matsuzaki, *ACS Chem. Biol.*, 2008, **3**, 341–345.
- 95 K. Kawano, Y. Yano, K. Omae, S. Matsuzaki and K. Matsuzaki, *Anal. Chem.*, 2013, **85**, 3454–3461.
- 96 H. Yamashita, Y. Yano, K. Kawano and K. Matsuzaki, *Biochim. Biophys. Acta, Biomembr.*, 2015, **1848**, 1359–1366.
- 97 B. V. Popp and Z. T. Ball, *J. Am. Chem. Soc.*, 2010, **132**, 6660–6662.



- 98 Z. Chen, B. V. Popp, C. L. Bovet and Z. T. Ball, *ACS Chem. Biol.*, 2011, **6**, 920–925.
- 99 J. Wang, Y. Yu and J. Xia, *Bioconjugate Chem.*, 2014, **25**, 178–187.
- 100 Y. Yano, N. Furukawa, S. Ono, Y. Takeda and K. Matsuzaki, *Pept. Sci.*, 2015, DOI: 10.1002/bip.22715.
- 101 U. Reinhardt, J. Lotze, S. Zernia, K. Mörl, A. G. Beck-Sickinger and O. Seitz, *Angew. Chem., Int. Ed.*, 2014, **53**, 10237–10241.
- 102 T. N. Grossmann and O. Seitz, *J. Am. Chem. Soc.*, 2006, **128**, 15596–15597.
- 103 U. Reinhardt, J. Lotze, K. Mörl, A. G. Beck-Sickinger and O. Seitz, *Bioconjugate Chem.*, 2015, **26**, 2106–2117.
- 104 H. Tsutsumi, W. Nomura, S. Abe, T. Mino, A. Masuda, N. Ohashi, T. Tanaka, K. Ohba, N. Yamamoto, K. Akiyoshi and H. Tamamura, *Angew. Chem., Int. Ed.*, 2009, **48**, 9164–9166.
- 105 W. Nomura, T. Mino, T. Narumi, N. Ohashi, A. Masuda, C. Hashimoto, H. Tsutsumi and H. Tamamura, *Pept. Sci.*, 2010, **94**, 843–852.
- 106 W. Nomura, N. Ohashi, A. Mori and H. Tamamura, *Bioconjugate Chem.*, 2015, **26**, 1080–1085.
- 107 Y. Sekine-Aizawa and R. L. Haganir, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 17114–17119.
- 108 M. Harel, R. Kasher, A. Nicolas, J. M. Guss, M. Balass, M. Fridkin, A. B. Smit, K. Brejc, T. K. Sixma, E. Katchalski-Katzir, J. L. Sussman and S. Fuchs, *Neuron*, 2001, **32**, 265–275.
- 109 C. M. McCann, F. M. Bareyre, J. W. Lichtman and J. R. Sanes, *BioTechniques*, 2005, **38**, 945–952.
- 110 A. D. Keefe, D. S. Wilson, B. Seelig and J. W. Szostak, *Protein Expression Purif.*, 2001, **23**, 440–446.
- 111 H. D. Mootz, *ChemBioChem*, 2009, **10**, 2579–2589.
- 112 J. Zettler, V. Schütz and H. D. Mootz, *FEBS Lett.*, 2009, **583**, 909–914.
- 113 R. Borra, D. Dong, A. Y. Elnagar, G. A. Woldemariam and J. A. Camarero, *J. Am. Chem. Soc.*, 2012, **134**, 6344–6353.
- 114 D. Jung, K. Sato, K. Min, A. Shigenaga, J. Jung, A. Otaka and Y. Kwon, *Chem. Commun.*, 2015, **51**, 9670–9673.
- 115 N. H. Shah, G. P. Dann, M. Vila-Perelló, Z. Liu and T. W. Muir, *J. Am. Chem. Soc.*, 2012, **134**, 11338–11341.
- 116 Y. David, M. Vila-Perelló, S. Verma and T. W. Muir, *Nat. Chem.*, 2015, **7**, 394–402.
- 117 I. V. Thiel, G. Volkmann, S. Pietrokovski and H. D. Mootz, *Angew. Chem., Int. Ed.*, 2014, **53**, 1306–1310.
- 118 M. Braner, A. Kollmannsperger, R. Wieneke and R. Tampé, *Chem. Sci.*, 2016, DOI: 10.1039/C5SC02936H.

