

INORGANIC CHEMISTRY

FRONTIERS







INORGANIC CHEMISTRY









View Article Online View Journal | View Issue

RESEARCH ARTICLE



Cite this: Inorg. Chem. Front., 2016,

Enhancing the photoluminescence quantum yields of blue-emitting cationic iridium(III) complexes bearing bisphosphine ligands†

Diego Rota Martir, a Ashu K. Bansal, b Vincent Di Mascio, a David B. Cordes, a Adam F. Henwood, Alexandra M. Z. Slawin, Paul C. J. Kamer, A Laura Martínez-Sarti, Antonio Pertegás, Henk J. Bolink, Ifor D. W. Samuel and Eli Zysman-Colman*a

Herein we present a structure-property relationship study of thirteen cationic iridium(III) complexes of the form of [Ir(C^N)₂(P^P)]PF₆ in both solution and the solid state through systematic evaluation of six bisphosphine (P^P) ligands (xantphos, dpephos, dppe, Dppe, nixantphos and isopropxantphos). All of the complexes are sky-blue emissive, but their photoluminescence quantum yields (Φ_{Pl}) are generally low. However, strong and long-lived blue luminescence ($\lambda_{\rm em}$ = 471 nm; $\phi_{\rm PL}$ = 52%; $\tau_{\rm e}$ = 13.5 μs) can be obtained by combining the reduced bite angle of the 1,2-bis-diphenylphosphinoethene (dppe) chelate with the bulky 2-(4,6-difluorophenyl)-4-mesitylpyridinato (dFmesppy) cyclometalating ligand. To the best of our knowledge this is the highest Φ_{PL} and the longest au_e reported for cyclometalated iridium(IIII) complexes bearing bisphosphine ligands. Light-emitting electrochemical cells (LEECs) were fabricated using lead complexes from this study, however due in part to the irreversible electrochemistry, no functional LEEC was achieved. Organic light-emitting diodes were successfully fabricated but only attained maximum external quantum efficiencies of 0.25%.

Received 12th September 2015, Accepted 28th October 2015 DOI: 10.1039/c5qi00177c

rsc.li/frontiers-inorganic

Introduction

The expansion of low-cost and more efficient lighting sources based on phosphorescent materials is one of the greatest challenges of our century. In particular, iridium complexes have attracted much attention due to their relatively short-lived triplet excited states (τ_e) and high photoluminescence quantum yields (Φ_{PL}), which are crucial properties for emitters employed in organic light emitting diodes (OLEDs) or lightemitting electrochemical cells (LEECs).2 In addition, due to

the large variety of ligand families that are readily synthetically accessible, the emission color of iridium(III) complexes can be easily tuned across the entirety of the visible spectrum.³ In the last decade, despite the reports of many phosphorescent green and red emitters⁴ having been synthesised, the design of highperformance blue-emitting materials remains a challenge.5 Recently, examples of sky-blue and deep-blue emitting cationic iridium complexes bearing biimidazole,6 bis(NHC),7 substituted triazole or tetrazole,8 or pyrazolyl-pyridine9 as ancillary ligands have been explored, but challenges still remain regarding efficiencies and stabilities of these emitters in devices. 10 Thus, there is still a demand for blue-emitting phosphors as emitters in lighting devices.

The vast majority of cationic iridium complexes possess the motif $[Ir(C^N)_2(N^N)]^+$, where C^N is a cyclometalating ligand with 2-phenylpyridinato (ppy) as the archetype and N^N is an ancillary ligand with 2,2'-bipyridine (bpy) as the archetype. 11 Despite their prominence as ligands in homogeneous catalysis, 12 phosphorus(III)-based ligands have remained underexplored in their complexation to iridium(III). Indeed, only a few examples have been reported implicating ligands with a P-Ir bond for optoelectronic applications; these include monodentate phosphines, 13 phosphinites, phosphonites, 14 phosphites, 15 bidendate benzyldiphenylphosphine P^C chelators, 16

^aOrganic Semiconductor Centre, EaStCHEM School of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST, UK.

E-mail: eli.zysman-colman@st-andrews.ac.uk; http://www.zysman-colman.com; Fax: +44 (0)1334 463808; Tel: +44 (0)1334 463826

^bOrganic Semiconductor Centre, SUPA, School of Physics and Astronomy, University of St. Andrews, St. Andrews, Fife KY16 9SS, UK. E-mail: idws@st-andrews.ac.uk ^cInstituto de Ciencia Molecular, Universidad de Valencia, C/ Catedrático J. Beltrán, ES-46980 Paterna Valencia Snain E-mail: henk holink@w es

[†] Electronic supplementary information (ESI) available: NMR spectra for all C^N ligands and complexes. MS spectra for all complexes. Supplementary crystal structure data of complexes 2a, 4c and 1e. Supplementary optoelectronic data for complexes. Supplementary LEEC data for 1b, 4a and 4b. CCDC 1423251-1423259 for 1a, 1b, 1c, 1e, 2a, 3c, 4a, 4b, 4c. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5qi00177c

other P^C ligands, 17 P^N18 and P^P chelators. 19 Given the paucity of photoluminescent iridium complexes bearing phosphine ligands in general and P^P chelators in particular, we have sought to systematically explore the effect of differing the bisphosphine ancillary ligand on the properties of [Ir(C^N)₂(P^P)]PF₆ complexes. Bisphosphines potentially offer several advantages: (1) they confer chemical stability both in the solid state and in organic solvents due in part to their steric bulk; 16c,19a (2) the electronic properties of the complexes remain highly tunable; 13a (3) due to their strong σ -donating character, they inhibit d-d transitions through destabilization of the metal-centered (MC) antibonding orbitals, making radiative decay more favourable. 15,16d,19a Improved organic light emitting diode (OLED) device performance has been achieved as a consequence of the use of phosphine-based ligands. 15,16c,17a

We^{2b,6a,8c,20} and others, ^{9,17c,19b,21} have been recently targeting the development of deep blue-emitting cationic iridium complexes, which are particularly relevant for light emitting electrochemical cells (LEECs), as an alternative solid-state lighting device architecture to OLEDs. In the context of [Ir(C^N)₂(P^P)]PF₆ complexes, Hor et al. recently reported the synthesis and the photophysical investigation of iridium(III) complexes bearing dppb (1,4-bis(diphenylphosphino)butane), dpephos (bis(2-diphenylphosphinophenyl) ether), nixantphos (2,4-bis(diphenylphosphino)phenoxazine) and xantphos (4,5bis-diphenylphosphino-9,9-dimethylxanthene) P^P ancillary



Eli Zysman-Colman

Eli Zysman-Colman obtained his Ph.D. from McGill University in 2003 under the supervision of Prof. David N. Harpp as an FCAR scholar where he conducting research in physical organic sulfur chemistry. He then completed two postdoctoral fellowships, one in supramolecular chemistry with Jay Siegel at the Organic Chemistry Institute, University of Zurich as an FQRNT fellow and the other in inorganic materials chemistry with Stefan

Bernhard at Princeton University as a PCCM fellow. He joined the department of chemistry at the Université de Sherbrooke in Quebec, Canada as an assistant professor in 2007. In 2013, he moved to the University of St Andrews where he is an EaStCHEM Research Fellow/Lecturer (tenured). His research program focuses on the rational design of: (I) luminophores for energy-efficient visual displays and flat panel lighting based on organic light emitting diode (OLED) and light-emitting electrochemical cell (LEEC) device architectures; (II) light harvesting dyes for dye-sensitized solar cells (DSSCs) and organic photovoltaics; (III) sensing materials employed in electrochemiluminescence; and (IV) photoredox catalysts for organic reactions.

ligands twinned with either ppy, 2-(4,6-difluorophenyl)pyridinato (dFppy) or 2'-phenyl-9-O-benzyl-10,11-dihydrocinchonine (pcn) C^N ligands. ^{19a} The bluest emission in their study (λ_{em} : 472 nm) was achieved with the complex [Ir(dFppy)₂(dppb)]PF₆, which exhibited a $\Phi_{\rm PL}$ of 30% and bi-exponential emission decay kinetics (τ_e : 0.22, 0.02 µs) at room temperature in dichloromethane. Similar sky-blue emission (λ_{em} : 476 nm) was observed for the complex [Ir(dFppy)₂(xantphos)]PF₆, but in this case the Φ_{PL} was very low at 0.02 with very short bi-exponential lifetimes (τ_e : 0.004, 0.06 µs). Other instances of blue emission in MeCN solution within the same class of complexes include $[Ir(ppy)_2(dppe)]^+$ (λ_{em} : 459 nm, Φ_{PL} : 0.44%, τ_e : 0.29 µs), τ_e $[Ir(ppy)_2(dppb)]^+$ (λ_{em} : 459 nm, Φ_{PL} : 0.68%, τ_e : 0.52 µs), 19d and $[Ir(Fmppy)_2(dppe)]^+$ (λ_{em} : 490 nm, Φ_{PL} : 18%, τ_e : 8.25 μ s)^{19b} (Fmppy = 2-[(4-fluorophenyl)-5-methyl]pyridinato); there is significant inconsistency in the reported properties of [(ppy)2Ir-(dppe)]PF₆ with Lowry et al., quoting an emission maximum of 490 nm, a $\Phi_{\rm PL}$ of 7.3% and a $\tau_{\rm e}$ of 3.9 $\mu {\rm s.}^{19b}$ None of this class of cationic complexes to date have been explored as emitters in solid-state lighting devices.

In the present work, we systematically study the properties of thirteen complexes of the form [Ir(C^N)2(P^P)]PF6 (Chart 1), with the bisphosphine ligand as one of six commonly used in the literature: xantphos, nixantphos, dpephos, dppe, Dppe and isopropxantphos, Scheme 1). In a bid to push the emission further into the blue and to enhance the photoluminescence quantum yield, a series of four C^N ligands were used: ppy, mesppy, dFppy and dFmesppy (mesppy = 2-phenyl-4-mesitylpyridinato; dFmesppy = 2-(4,6-difluorophenyl)-4-mesitylpyridinato). The bulky mesppy and dFmesppy ligands were incorporated into the study as the latter had been previously shown by Bryce²² and co-workers in [Ir(dFmesppy)₂(pic)] (pic = 2-piconilate) to (1) improve the solubility of the complex in organic solvents, which impacts positively homogeneous film deposition;23 (2) promote reduced intermolecular interaction due to the bulky mesityl group and therefore reduced quenching phenomena, giving rise to increased Φ_{PL} in the solid state; (3) negligibly impact the emission energy, as the mesityl moiety is disposed in an nearly orthogonal and locked conformation with respect to the plane of the pyridine ring. Of all the bisphosphines, complexes with either dppe or Dppe as the ancillary ligand exhibited the highest photoluminescence quantum yield and longest emission lifetimes.

Results and discussion

Synthesis

The syntheses of the C^N ligands and the heteroleptic iridium complexes are shown in Scheme 1. Each of dFppy, mesppy and dFmesppy were prepared in high yield via Suzuki-Miyaura²⁴ palladium-catalysed cross-coupling reactions. The C^N ligands were complexed with IrCl3·3H2O and the resulting the μ-dichloro-bridged iridium dimers [Ir- $(ppy)_2Cl]_2$, **D1**; $[Ir(dFppy)_2Cl]_2$, **D2**; $[Ir(mesppy)_2Cl]_2$, **D3**; and

[Ir(dFmesppy)₂Cl]₂, **D4** were formed in good yield under standard conditions.²⁵ The iridium complexes 1-4 (Chart 1) were isolated in high yield through cleavage of D1-D4 with the corresponding bisphosphine ligands: 4,5-bis-diphenylphosphino-9,9-dimethylxanthene (xantphos), La; bis(2-diphenylphosphinophenyl) ether (dpephos), 1,2-bis-(diphenylphosphino)ethene (dppe), Lc; 1,2-bis(diphenylphosphino)ethane (Dppe), L_d; 4,5-bis(diphenylphosphino)-9-isopropylenxanthene (isopropxantphos), Le; and 2,4-bis-(diphenylphosphino)phenoxazine (nixantphos), Lf. All complexes were purified by column chromatography and isolated as the PF₆⁻ salt following an anion metathesis reaction using NH₄PF₆. The purity of the complexes was confirmed by ¹H, ¹³C, ¹⁹F and ³¹P NMR spectroscopy, HRMS and melting point

analyses (1H, 13C, 31P, 19F-NMR and HRMS spectra are reported in the ESI†). HRMS analysis showed in each case the diagnostic molecular ion peak. The 31P NMR spectra showed one singlet between -15 and -25 ppm, corresponding to the two magnetically equivalent phosphorous atoms in the P^P ligand and one septet at around -150 ppm, characteristic of the PF₆ anion. The 19F NMR spectra for 4a, 4b, 3c, 4c showed two singlets between -105 and -110 ppm corresponding to the two fluorine atoms present on each C^N ligand and a doublet at around -70, -80 ppm, characteristic to the six magnetically equivalent fluorine atoms coupled to the phosphorous nucleus of the PF₆ anion. In addition, crystal structures of 1a, 2a, 4a, 1b, 4b, 1c, 3c, 4c, 1e, were obtained through single crystal X-ray analysis.

Chart 1 Chemical structure of complexes 1-4

70% - 95% 1) P^P (Ln*) IrCl₃ 3H₂O 2) NH₄PF₆ d 45% - 90% R1,R2: H, ppy R1: H. R2: mesityl mesppy R1: F, R2:H, dFppy R1: F,R2: mesityl dFmesppy R1.R2: H. D1 R1: H, R2: mesityl D2 R1: F, R2: H, D3 R1: F,R2: mesityl D4 B(OH)₂ dpe phos. Lb dppe, L, isopropxantphos, L

Scheme 1 Synthesis of C^N ligands and [Ir(C^N)₂(P^P)]PF₆ complexes 1-4. Reagents and conditions. a2.0 equiv. K₂CO₃, 5 mol% Pd(PPh₃)₄, N₂, 1,4 $dioxane/H_{2}O~(4:1~v/v),~105~^{\circ}C,~56~h;~^{b}2.0~equiv.~K_{2}CO_{3},~5~mol\%~Pd(PPh_{3})_{4},~N_{2},~1,2-dimethoxyethane/H_{2}O~(4:1~v/v),~130~^{\circ}C,~19~h;~^{c}2-EtOC_{2}H_{4}OH/mol_{2}OH/mol_{$ H_2O (4 : 1 v/v), 110 °C, N_2 , 19 h. di. $CH_2Cl_2/MeOH$ (2 : 1 v/v), 55 °C, 19 h, N_2 ; ii. Excess solid NH_4PF_6 .

Crystal structures

Crystal structures of complexes 1a, 2a, 4a, 1b, 4b, 1c, 3c, 4c and 1e were obtained through X-ray diffraction studies. The structures of the Ir(III) complexes 1a, 1b and 1c are shown in Fig. 1, those of complexes 4a, 4b and 4c are shown in Fig. 2, whereas the remaining structures are reported in the ESI.†

In all the structures, the iridium centre adopts a distortedoctahedral geometry, as expected, with the two nitrogen atoms of the C^N ligands mutually trans to each other, and the two phosphorous atoms of the P^P ligand disposed trans with respect to the carbon atoms of the C^N ligands. Both the Ir-C and Ir-N bond lengths fall in the range expected for [Ir-(C^N)₂(L^L)] complexes (where L^L is any bidentate ligand),

and the Ir-P bond lengths also fall into the range expected for [Ir(C^N)₂(P^P)] complexes with two phosphorus donors (CSD version 5.36).²⁶ Those complexes with a mesityl-functionalized C^N ligand (2a, 4a, 2b and 4c), all display the desired orthogonal orientation of the mesityl with respect to the pyridinato fragment The structures of 1b and 1e are closely related to those of the known structures of these complexes, ^{19a,c} including the arrangement of the complexes themselves. The differences primarily arise from changes in packing and solvation.

The P-Ir-P angles can be seen to vary in a systematic manner. For complexes 1a, 2a and 4a (with xantphos), 1b and 4b (with dpephos) and 1e (with nixantphos), the P-Ir-P bond angles are distorted to greater than 90° [angles varying between $98.74(7)^{\circ}$ and $102.53(7)^{\circ}$; whereas for complexes 1c,




Fig. 1 Molecular structure of 1a (a), 1b (b) and 1c (c). Hydrogen atoms, PF₆⁻ counterions, solvent molecules, minor components of disordered molecules and additional independent molecules are omitted for clarity. Selected bond lengths (Å) and angles (°): (1a) Ir_1-N_1 2.079(6), Ir_1-N_{13} $2.084(6),\ |r_1-C_8|\ 2.042(7),\ |r_1-C_{20}|\ 2.041(8),\ |r_1-P_1|\ 2.482(2),\ |r_1-P_2|\ 2.523(2),\ |r_1-N_{71}|\ 2.074(6),\ |r_1-N_{83}|\ 2.077(6),\ |r_1-C_{78}|\ 2.051(7),\ |r_1-C_{90}|\ 2.063(7),\ |r_1-C_{90}|\ 2.0$ $|r_{71}-P_{71}| 2.483(2), |r_{71}-P_{72}| 2.537(2), |P_{1}-|r_{1}-P_{2}| 102.53(7), |P_{1}-|r_{71}-P_{72}| 102.19(7); \\ \text{(1b)} |r_{1}-N_{1}| 2.092(10), |r_{1}-N_{13}| 2.057(10), |r_{1}-C_{8}| 2.022(13), |r_{1}-C_{10}| 2.092(13), |r_{1}-C_{10}| 2.092$ $2.039(12),\; |r_1-P_1|\; 2.448(4),\; |r_1-P_2|\; 2.537(4),\; |P_1-P_1-P_2|\; 99.53(12);\; \textbf{(1c)}\; |r_1-N_1|\; 2.076(3),\; |r_1-N_1|\; 2.072(3),\; |r_1-C_8|\; 2.058(4),\; |r_1-C_2|\; 2.059(4),\; |r_1-P_1|\; 2.076(3),\; |r_1-P_1|\; 2.076$ $2.3680(11),\ |r_{1}-P_{2}|2.3593(11),\ |r_{51}-N_{51}|2.072(7),\ |r_{51}-N_{51}|2.080(11),\ |r_{51}-C_{58}|2.060(5),\ |r_{51}-P_{51}|2.3558(13),\ P_{1}-|r_{1}-P_{2}|84.49(4),\ P_{51}-|r_{51}-P_{51}(A)|84.1(3).$ [Symmetry code: (A) -x + 1/2, y, 1 - z].




Fig. 2 Molecular structure of 4a (a), 4b (b) and 4c (c). Hydrogen atoms, PF₆⁻ counterions, solvent molecules, minor components of disordered molecules and additional independent molecules are omitted for clarity. Selected bond lengths (Å) and angles (°): (4a) Ir₁-N₁ 2.064(7), Ir₁-N₂₂ $2.068(7), \; |r_1 - C_8| \; 2.037(8), \; |r_1 - C_2| \; 2.026(9), \; |r_1 - P_1| \; 2.476(2), \; |r_1 - P_2| \; 2.531(2), \; |P_1 - |r_1 - P_2| \; 100.81(8); \; (4b) \; |r_1 - N_1| \; 2.084(6), \; |r_1 - N_2| \; 2.068(6), \; |r_1 - C_8| \; (1.00016)$ $2.061(8), Ir_1 - C_{29} \ 2.052(8), Ir_1 - P_1 \ 2.461(2), Ir_1 - P_2 \ 2.473(2), P_1 - Ir_1 - P_2 \ 98.74(7); \\ (4c) Ir_1 - N_1 \ 2.063(2), Ir_1 - N_{22} \ 2.067(3), Ir_1 - C_8 \ 2.053(3), Ir_1 - C_{29} \ 2.064(3), Ir_1 - C_{29} \ 2.061(3), Ir_2 - C_{29} \ 2.061(3), Ir_3 - C$ Ir_1-P_1 2.3642(8), Ir_1-P_2 2.3653(8), $P_1-Ir_1-P_2$: 83.06(3).

3c and 4c (with dppe) the P-Ir-P bond angles are reduced to less than 90° [angles varying between 83.06(3)° and 84.49(4)°]. This appears to arise from the size of the chelate formed by the P^P ligand, and its flexibility. In each of 1a, 2a, 4a, 1b, 4b and 1e an eight-membered chelate exists whereas in 1c, 3c and 4c a more rigid five membered chelate is present. Additionally, the complexes with an eight-membered chelate break down into two sub-groups depending on the flexibility of the backbone of the P^P ligand, those complexes of xantphos and nixantphos showing a range of P-Ir-P bond-angles [100.81(8)-102.53(7)°] slightly above those seen in the more conformationally flexible dpephos complexes [98.74(7)-99.53(12)°]. To accommodate the varying sizes of chelate rings, the Ir-P bonds are found to vary in consequence, the complexes with smaller chelate rings showing shorter Ir-P bonds [2.3642(8)-2.3687(5) Å]

than those in the complexes with larger chelate rings [2.448(4)-2.537(2) Å].

Photophysical properties

The optoelectronic properties of complexes 1-4 have been investigated both in MeCN solution and in the solid state and are summarized in Table 1. The absorption spectra are shown in Fig. 3. The absorption spectra of all the complexes are characterized by two intense bands between 260 nm and 320 nm and a broad lower intensity band at around 360 nm. Similar to many other cationic iridium complexes of the form $[Ir(C^N)_2(P^P)]^+$ found in the literature, 16c,d,18,19 the two higher energetic bands can be attributed to the spin-allowed ${}^{1}\pi \to \pi^{*}$ ligand centered (1LC) transitions localized on the C^N ligands, while the broad bands at wavelengths longer than 350 nm can

Table 1 Relevant photophysical data for 1-4^a

	$\lambda_{\mathrm{em}}^{a,b}$ (nm)			b)	$\tau_{\mathrm{e}}^{\;\;a}$		
	MeCN	Film ^d	MeCN	Film ^d	$MeCN^e$ (ns)	Film ^{d,e} (ns)	
1a	484 [0.9], 510 [1]	507 [1], 657 [0.18]	0.3	0.8	6 (34.9), 20 (65.1)	5 (38.7), 40 (61.3)	
2a	485 [0.90], 509 [1]	490 [0.90], 509 [1], 652 [0.18]	0.6	1.5	4 (18.7), 28 (81.3)	2 (23.0), 17 (22.8), 186 (54.2)	
4a	461 [0.80], 489 [1]	474 [0.87], 500 [1], 630 [0.18]	0.7	5.7	17 (12), 36 (88)	50 (3.6), 306 (39.2), 882 (59.2)	
1b	465 [0.80], 498 [1], 530 [0.60]	472 [0.65], 514 [1]	0.6	1.4	3 (5.9), 27 (94.0)	2 (16.3), 21 (17.2), 196 (66.5)	
2b	466 [0.8], 501 [1], 539 [0.5]	472 [0.84], 504 [1], 538 [0.6]	0.9	1.7	4 (8.5), 28 (91.4)	3 (4.3), 44 (20.0), 303 (75.7)	
4b	459 [0.82], 484 [1]	462 [0.82]; 486 [1]; 515 [0.70]	0.9	7.2	4 (9.5), 89 (90.4)	8 (10.6), 143 (19.1), 1070 (70.2)	
1c	457 [0.95], 488 [1], 510 [0.58]	460 [0.56], 499 [1], 512 [0.92]	4.2	6.0	123 (2.7), 1105 (97.3)	6 (4.7), 150 (38.4), 1036 (56.9)	
2c	456 [0.94], 486 [1], 516 [0.58]	462 [0.84], 492 [1], 517 [0.67]	4.5	7.8	5 (2.5), 1570 (97.5)	5 (2.9), 254 (14.2), 1183 (83.0)	
4c	444 [0.78], 471 [1], 504 [0.63]	448 [0.77]; 470 [1]; 504 [0.66]	52	37.2	179 (2.1), 13 468 (97.9)	3 (32.7), 245 (4.95), 3946 (62.38)	
3c	445 [0.76], 471 [1], 503 [0.67]	446 [0.77]; 472 [1], 500 [0.78]	18	20.7	25 (1.6), 9211 (98.7)	2 (30.2), 21 (5.6), 2598 (64.2)	
1d	457 [0.95], 488 [1], 516 [0.59]	465 [0.37], 514 [1]	1.6	4.2	31 (1.7), 497 (98.2)	5 (6.1), 168 (30.0), 836 (64.0)	
1f	485 [0.9], 509 [1]	478 [0.90], 499 [1], 598 [0.36]	0.6	0.7	6 (31.0), 26 (69.0)	1 (42.5), 12 (57.5)	

^a Measurements in degassed MeCN at 298 K. ^b Principal emission peaks listed with values in brackets indicating relative intensity. ^c Quinine sulfate employed as the external reference (Φ_{PL} = 54.6% in 0.5 M H₂SO₄ at 298 K). ²⁹ ^d Films formed by dip-coating deposition on pristine quartz substrates. PLQY measurements were carried out under nitrogen. Values in parentheses are pre-exponential weighting factor, in relative % intensity, of the emission decay kinetics.




Fig. 3 UV-Vis spectra of complexes 1a, 2a, 4a, 1b, 1d, 1e, 1f (a) and 2b, 4b, 1c, 2c, 4c, 3c (b) recorded in MeCN at 298 K.

be assigned to a mixture of spin-allowed and spin-forbidden metal-to-ligand charge transfer transitions (¹MLCT/³MLCT). Indeed, as previously reported and predicted by TD-DFT calculations, 4a,27 spin-forbidden transitions directly to the triplet state are accessible in iridium(III) complexes due to the large spin-orbit coupling exhibited by the heavy iridium atom. The presence of the mesityl moiety in 2a, 4a, 2b, 4b, 2c and 4c leads to enhanced molar absorptivities in the UV region of the spectrum while also introducing a strongly absorbing shoulder at around 260-270 nm, the transition of which is assigned as ¹LC in nature.

Fig. 4 illustrates the normalized room temperate emission spectra of 1-4 upon photoexcitation into the $^1\pi \to \pi^*$ band (at 360 nm). Emission is blue to blue-green with emission maxima bunched between 460 nm and 510 nm. There are two distinct families of complexes based on their emission profiles. A structured vibronic emission profile is observed for b, c and d complexes (Fig. 4b) bearing one of dpephos, dppe and Dppe P^P ligands, which is indicative of a 3LC excited state. 3c,11,19b,28 The emission is less structured for 1a, 2a, 4a, 1f (Fig. 4a) bearing xantphos-type ancillary ligands, which implicates greater ³MLCT character. ^{19a,27a,e} No emission was detected for complex 1e.

Incorporation of electron-withdrawing fluorine atoms on the C^N ligand promoted the expected stabilization of the frontier molecular orbitals and the blue-shift in the emission observed for complexes 4a, 4b, 4c and 3c compared with their fluorine-free congeners. 3b,20b

Despite only minute variations in the emission energy, remarkable modulation of both $\Phi_{\rm PL}$ and $au_{\rm e}$ could be obtained through choice of C^N and P^P ligands. Firstly, of the ppy-containing complexes the highest photoluminescence quantum yields arise from complexes using dppe as the ancillary ligand (cf. 1c), where Φ_{PL} at 4.2% is an order of magnitude higher than the other complexes in the series, save 1d. Bi-exponential decay kinetics were observed for each of these complexes, similar to other $[Ir(C^N)_2(P^P)]^+$ complexes ^{19a} however, only in 1c was there an observed microsecond component. Complex 1d, bearing the Dppe P^P ligand, showed a modestly reduced $\Phi_{\rm PL}$ of 1.2% and shorter bi-exponential decay (31 ns and 497 ns). Thus, the stronger bonding of the dppe (and Dppe) ligand to the iridium center has two effects: (1) there is a stron-




Fig. 4 Normalized photoluminescence spectra of complexes 1a, 2a, 4a, 1f (a) and 1b, 2b, 4b, 1c, 2c, 4c, 3c, 1d (b) recorded in degassed MeCN at 298 K (λ_{exc} : 360 nm).

ger destabilization of the non-emissive MC states leading to an enhanced photoluminescence quantum yields and (2) there is a blue-shifted emission due to the stronger ligand field. ^{14–16,30}

In an effort to improve the Φ_{PL} of these complexes, we evaluated the family of mesppy-containing complexes (2a, 2b, 2c). Unfortunately only a slight improvement in the solution-state photophysical properties was observed. For instance, the Φ_{PL} for complex 2c, [Ir(ppymes)₂(dppe)]PF₆, was 4.5% ν s. 4.2% obtained for 1c, [Ir(ppy)₂(dppe)]PF₆. Consistent with the energy gap law, along with the expected blue-shift in the solution-state emission (*e.g.*, 3c: λ_{em} = 471 nm; 1c: λ_{em} = 488 nm) upon fluorine incorporation onto the C^N ligands, there is an observed enhanced photoluminescence quantum yield and longer emission lifetimes (*e.g.*, 3c: 18%; 1c: 4.2%). ^{19a,b}

Considering the improved photophysical properties observed for 3c, we decided to evaluate the family of dFmesppy cyclometalated complexes bearing xantphos, dpephos and dppe as ancillary ligands, with a view to reducing the intramolecular interactions and therefore the quenching phenomena following introduction of the bulky mesityl substituent. Disappointingly, the observed Φ_{PL} and τ_e values for 4a and 4b remained essentially unaltered compared to analogs 2a and 2b. We were however delighted to observe the much-enhanced Φ_{PL} for 4c at 52% and emission dominated by a component of lifetime of 13.5 µs compared to analog 3c.

To discern if the effects observed in MeCN solution are transposable to the solid state, dip-coated neat films were analyzed. Generally, the trends observed in solution are mirrored in the films (Table 1), with slight enhancement in $\Phi_{\rm PL}$. It is worth noting that a remarkable improvement of the $\Phi_{\rm PL}$ in the solid state was observed for 4a and 4b, which contain the dFmesppy C^N ligands. These two complexes are nearly nonemissive in MeCN ($\Phi_{\rm PL}$: 0.7% and 0.9%, respectively) with very short emission lifetimes ($\tau_{\rm e}$: ca. 36 ns and 89 ns, respectively) but significantly brighter in the solid state ($\Phi_{\rm PL}$: 5.7% and 7.4%, respectively and $\tau_{\rm e}$: ca. 882 ns and 1070 ns, respectively). Thus, the non-radiative pathways in solution that result from the flexibility of the eight-membered chelate of the P^P ligand are significantly mitigated as the complexes are rigidified in the solid state.

For 1b, 1c and 1d bearing, respectively, dpephos, dppe and Dppe P^P ligands, a red shift in the emission energy (by up to 4.5×10^2 cm⁻¹) and broader emission are observed going from solution to the solid state (Fig. S77†). This is in agreement with many reports on aggregation phenomena of iridium(III) complexes in the solid state, in which either or both a red-shift in the emission and broadness in the emission bands is observed.31 In order to overcome such effects, the ligands are often modified to incorporate bulky groups. 21i,31,32 Our results show that for 2b, 4b, 2c and 4c the bulky mesityl substituent suppresses aggregation in the solid state, with their thin film emission profiles mirroring those observed in MeCN solution. A similar, but less pronounced effect, is observed for complexes 1a and 1f, where the red-shifted emission in the solid state is noticeably reduced due to the bulkiness of the xantphos and isopropxantphos P^P ligands, respectively.

Electrochemical properties

The electrochemical properties of the complexes have been investigated by cyclic voltammetry in degassed MeCN and the oxidation potentials, $E_{\rm pa}$ (V), reported with respect to SCE (Fc/Fc⁺ = 0.38 V in MeCN)³³ are compiled in Table S1.† For complexes bearing ppy and mesppy C^N ligands a single irreversible oxidation wave around 1.4 V is observed. When the complexes possess dFppy or dFmesppy C^N ligands, the oxidation potentials are more positive ($E_{\rm pa}$: ca. 1.7 V) due to stabilization of the HOMO promoted by the fluorine electron-withdrawing substituents. Similar to other cationic iridium(III) complexes, these oxidation processes are assigned to the Ir^{III}/Ir^{IV} redox couple with significant involvement of the C^N ligands.^{11,14} The absence of an observable reduction process within the solvent window points to a large electrochemical gap promoted by the strongly σ-donating P^P ligands.

Electroluminescent devices: light-emitting electrochemical cells (LEECs)

In view of the promising thin film photoluminescence properties of some of the ionic transition metal complexes (iTMC) in this study, light-emitting electrochemical cells (LEECs) based on compounds 1b, 4a, 4b and 4c were prepared, according to the procedure described in the Experimental section. LEECs consisted of a 100 nm thick emitting layer, sandwiched between an ITO/PEDOT:PSS (80 nm) anode (ITO = indium tin oxide; PEDOT:PSS = poly(3,4-ethylenedioxythiophene) polystyrene sulfonate) and a thermally evaporated aluminum cathode (70 nm) (Fig. 5c). The PEDOT:PSS layer was used to smoothen the ITO surface and hence to increase the yield and reproducibility of working devices. The device lifetime was measured by applying a pulsed current and by monitoring the voltage and luminance versus time, with a True Colour Sensor MAZeT (MTCSiCT Sensor) and a Botest OLT OLED Lifetime-Test System. The LEECs were driven using a pulsed current of 765 A m⁻², which is a high current density for this type of devices that leads to rapid degradation. The high current density, however, was necessary to increase the light emission to a level sufficient to be detectable by our equipment. All four LEECs showed a fast voltage drop at the initial operation stage, a consequence of fast ionic motion towards the electrode interfaces, which decreases the injection barrier for electrons and holes (see Fig. S99-102†).34 Following the electrical double layers formation at the electrode interface, the injected charges create p- and n-doped regions in the active layer close to the correspondent electrode. 35 In spite of the typical electronic behavior of the LEECs, the luminance detected was rather low for all four devices. Assuming that electrons and holes are effectively injected into the active layer, the poor performance for the LEECs is attributed to the irreversibility of the phosphine ligand oxidation and to the high driving current density.

In a first attempt to exploit the potential of these new complexes, in view of their high photoluminescence efficiency, a host-guest LEEC was prepared using complex **4c** in the configuration as previously described, ³⁶ which consisted on an




Fig. 5 (a) Luminance (solid blue line) and average voltage (open red squares) for ITO/PEDOT:PSS/active layer/Al (active layer = 1:1 by mass of NMS25: SPPO13 with 10 wt% of 4c) with a pulsed driving regime (block-wave pulsed current; 1000 Hz; 50% duty cycle; average density 100 A m⁻²). (b) Electroluminescence spectrum with inset photo of LEEC. (c) Schematic representation of the LEEC architecture.

ionic carbazole (NMS25) and a spirobifluorene (SPPO13) as hole and electron transport materials, respectively. The hostguest layer was prepared using a NMS25: SPPO13 mixture (1:1 mass ratio) doped with 10 wt% of 4c. The host-guest device showed electroluminescence at a lower current density compared to the devices employing solely the iTMC (Fig. 5a). This is an indication of a better electronic transport through the host materials compared to the pure iTMC. The absolute electroluminescence of the host-guest system is somewhat low, which could be most likely due to an inadequate alignment of the lowest unoccupied molecular orbital (LUMO) of the iTMC with respect to the LUMO of the SPPO13 (-2.9 eV), ³⁷ resulting in inefficient energy transfer from the host (SPPO13) to the guest (4c). The reduction wave in 4c was found to lie outside the electrochemical window of the solvent (see electrochemistry section), which means that the LUMO energy of this material should be less than -2.8 eV (Fc/Fc⁺ reference). The

host-guest LEEC showed a sky-blue electroluminescence, where the maximum emission corresponds to 479 nm (Fig. 5b).

We are currently working on the preparation of more suitable polar electron transport materials to attempt to improve these host guest devices.

Electroluminescent devices: organic light-emitting diodes (OLEDs)

OLEDs employing **3c** and **4c** were also fabricated and tested in light of their promising thin film photophysical properties. The device architecture consisted of the following structure: ITO/PEDOT:PSS (40 nm)/PVK (35 nm)/(mCP + Active layer + OXD7) (30 nm)]/B3PyMPM (60 nm)/Ca (20 nm)/Al (200 nm) structure, where **3c** or **4c** act as the active layer. PEDOT:PSS, PVK, mCP, OXD7 and B3PyMPM denote poly(3,4-ethylenedioxythiophene):poly(styrenesulfonate), poly(*N*-vinylcarbazole),

1,3-bis(N-carbazolyl)benzene, 2,2'-(1,3-phenylene)bis[5-(4-tertbutylphenyl)-1,3,4-oxadiazole] and bis-4,6-(3,5-di-3-pyridylphenyl)-2-methylpyrimidine, respectively (Fig. 6d). PVK facilitates the injection of holes and the electron transport layer B3PyMPM blocks the holes from penetration into cathode due to its deep lying HOMO and reduces the leakage current. Except for B3PyMPM and the contacts, all the layers were deposited by solution-processing methods. Device performance is summarized in Table 2.

The electroluminescence (EL) spectra are very similar to the PL spectra of the neat films as shown in Fig. 6a, an indication that in both cases the complexes are emitting from the same excited state. Fig. 6b shows the current-voltage-luminance characteristics of two devices. The turn-on voltages for both devices are similar. Fig. 6c shows the external quantum efficiency of the devices together with the power efficiency. Both the devices exhibit similar CIE coordinates (0.20, 0.29). Despite 4c exhibiting a higher thin film Φ_{PL} compared to 3cboth OLED devices show similar poor efficiencies, which suggest that the performance of device 1 can be optimized further by improving the charge balance.

Conclusions

Research Article

summary, thirteen blue to blue-green emitting $(\lambda_{\rm em} 477-510 \text{ nm})$ cationic heteroleptic iridium(III) complexes bearing bisphosphine ancillary ligands have been synthesized and their optoelectronic properties investigated. Through optimization of the bite angle of the P^P chelate and introduction of bulky and fluorinated C^N ligands, we have rationally designed the brightest reported blue-emitting (λ_{em} : 471 nm) complex of this class in 4c (Φ_{PL} : 52%). However, these complexes exhibit irreversible electrochemistry, which perhaps accounts for the poor LEEC performances observed using complexes 1b, 4a, 4b and 4c. The EL study shows that further refinement in charge balance is necessary in order to improve the performance of the OLED devices.

Experimental section

General synthetic procedures

Commercial chemicals were used as supplied. All reactions were performed using standard Schlenk techniques under inert (N2) atmosphere with reagent grade solvents. Flash column chromatography was performed using silica gel (Silia-P from Silicycle, 60 Å, 40-63 µm). Analytical thin layer chromatography (TLC) was performed with silica plates with aluminum backings (250 µm with indicator F-254). Compounds were visualized under UV light. 1H, 13C, 31P NMR 19F NMR spectra were recorded on a Bruker Avance spectrometer at 500 MHz, 126 MHz and 471 MHz respectively. The following abbreviations have been used for multiplicity assignments: "s" for singlet, "d" for doublet, "t" for triplet, "m" for multiplet and "br" for broad. Deuterated chloroform (CDCl₃) and deuterated

dichloromethane (CD2Cl2) were used as the solvents of record. Melting points (Mp's) were recorded using open-ended capillaries on an Electrothermal melting point apparatus and are uncorrected. High-resolution mass spectra were recorded at the EPSRC UK National Mass Spectrometry Facility at Swansea University on a quadrupole time-of-flight (ESI-Q-TOF), model ABSciex 5600 Triple TOF in positive electrospray ionization mode and spectra were recorded using sodium formate solution as the calibrant.

Ligand syntheses

2-Chloro-4-(2,4,6-trimethylphenyl)pyridine (A). The thesis of this ligand is by a modified method to a previously method.22 2-Chloro-4-Iodopyridine 16.74 mmol), 2,4,6-trimethylphenyl boronic acid (4.0 g, 25.11 mmol), potassium carbonate (50 mmol) were added to a round bottomed flask containing 50 mL of a mixture of 1,4dioxane and distilled water (4:1 v/v). The reaction mixture was degassed by multiple vacuum and N2 purging cycles, and Pd(PPh₃)₄ (0.693 g, 0.60 mmol) was added to the flask under positive nitrogen pressure. The mixture was refluxed under nitrogen atmosphere for 48 h and then cooled to room temperature. The mixture was poured onto distilled water and extracted multiple times with dichloromethane. The organic fractions were combined, washed with a portion of brine and dried over magnesium sulfate. Filtration and evaporation under reduced pressure gave the crude product (3.80 g). The crude product was purified by flash column chromatography (5% ethyl acetate/hexane on silica) to give 3.30 g of pure compound as a colourless oil. Yield: 85%. Rf: 0.40 (20% EtOAc/ hexanes on silica). 1 H NMR (500 MHz, CDCl₃) δ (ppm): 8.46 (d, J = 5.1 Hz, 1H), 7.18 (s, 1H), 7.06 (dd, J = 5.1, 1.4 Hz, 1H),6.99 (s, 2H), 2.36 (s, 3H), 2.04 (s, 6H). The characterisation matches that reported.²²

2-Phenyl-4-(2,4,6-trimethylphenyl)pyridine (mesppy). 2-Chloro-4-(2,4,6-trimethylphenyl)pyridine (A) (3.3 g, 14.24 mmol), phenyl boronic acid (2.78 g, 22.78 mmol), potassium carbonate (40 mmol) were added to a round bottomed flask containing 50 mL of a mixture of 1,2-dimethoxyethane and distilled water (4:1 v/v). The reaction mixture was degassed by multiple vacuum and N_2 purging cycles, and $Pd(PPh_3)_4$ (0.548 g, 0.48 mmol) was added to the flask under positive nitrogen pressure. The mixture was refluxed under nitrogen atmosphere for 19 h and then cooled to room temperature. The mixture was poured onto distilled water and extracted multiple times with dichloromethane. The organic fractions were combined, washed with a portion of brine and dried over magnesium sulfate. Filtration and evaporation under reduced pressure gave the crude product (3.90 g). The crude product was purified by flash column chromatography (silica, hexane/ethyl acetate gradient 100:0 to 80:20) to give 3.50 g of pure compound as a colourless oil. Yield: 87%. Rf: 0.38 (10% EtOAc/ hexanes on silica). ¹H NMR (500 MHz, CDCl₃) δ (ppm): 8.77 (dd, J = 5.0, 0.8 Hz, 1H), 8.05 (m, 2H), 7.60 (q, J = 0.8 Hz, 1H),7.54–7.42 (m, 3H), 7.09 (dd, J = 5.3, 1.3 Hz, 1H), 7.01 (s, 2H), 2.4 (s, 3H), 2.08 (s, 3H). 13 C NMR (126 MHz, CD₂Cl₂) δ (ppm):




Fig. 6 (a) Normalized electroluminescent emission spectra of 4c (black) and 3c (red) in thin films. Inset shows the photograph of the working OLED with 4c. (b) Current density and luminance versus applied voltage for OLEDs made with 4c (black) and 3c (red). (c) External quantum efficiency and Power efficiency versus applied voltage for OLEDs made with 4c (black) and 3c (red). (d) Schematic representation of the OLED architectures.

Table 2 OLED performance data^a

Device	Complex	Turn on voltage (V@ 1 cd m ⁻²)		Luminance (cd m ⁻²)		
1	4c	8.6	22.3	95.6	0.24	0.10
2	3c	8.0	21.2	79.9	0.21	0.09

 $[^]aJ=$ current density; EQE = external quantum efficiency; PE = power efficiency.

157.7, 150.3, 149.9, 139.4, 137.6, 136.5, 135.2, 129.1, 128.8, 128.6, 128.4, 127.0, 123.2, 121.5, 21.1, 20.7. FT-MS: Calculated: $(C_{20}H_{19}N)$ 273.1585, found: 274.1585.

2-(2,4-Difluorophenyl)pyridine (dFppy). The synthesis of this ligand is by a modified method to a previously reported method.6a 2,4-Difluorophenylboronic acid 17.70 mmol), 2-bromopyridine (2.00 g, 12.60 mmol), sodium carbonate (4.37 g, 31.60 mmol) were added to a round bottom flask containing a mixture of 1,4-dioxane and distilled water (4:1 v/v) to obtain a concentration of 0.15 to 0.20 M. The reaction mixture was degassed via bubbling nitrogen for 20 minutes. Pd(PPh₃)₄ (0.731 g, 0.633 mmol) was quickly added to the round bottom flask. The mixture was degassed again by evacuating and then backfilling with nitrogen three times, before heating to reflux. The mixture was refluxed under nitrogen atmosphere for 19 h, before cooling to room temperature and pouring onto distilled water. Extraction of the product with multiple portions of ethyl acetate was followed by combining the organic fractions, washing with water and then saturated sodium bicarbonate to remove residual boronic acid. Evaporation under reduced pressure gave the crude product. Purification by flash column chromatography (silica, hexane/ ethyl acetate 95:5) gave 0.554 g of pure compound as a colourless oil. Yield: 70%. R_f: 0.48 (20% EtOAc/hexanes on silica). ¹H NMR (500 MHz, CDCl₃) δ (ppm): 8.72 (dt, J = 4.5 Hz, 1.0 Hz, 1H), 8.03-7.99 (m, 1H), 7.75 (d, J = 4.0 Hz, 2H), 7.28-7.23 (m, 1H), 7.03-6.99 (m, 1H), 6.93-6.90 (m, 1H). ¹³C NMR (126 MHz, $CDCl_3$) δ (ppm): 164.6, 162.1, 159.5, 152.7, 149.9, 136.6, 132.3, 124.4, 122.6, 112.1, 104.5. ¹⁹F { ¹H } NMR (471 MHz, CDCl₃) δ (ppm): -109.3 (d, J = 6.8 Hz, 1F), -113.0 (d, J = 6.8 Hz, 1F). GCMS: (13.6 min) [M]⁺: 191. The characterisation matches that reported.6a

2-(2,4-Difluorophenyl)-4-(2,4,6-trimethylphenyl)pyridine (dFmesppy). The synthesis of this ligand is by a modified method to that previously reported. ²² 2-Chloro-4-(2,4,6-trimethylphenyl)pyridine (A) (2.28 g, 9.88 mmol), 2,4-Difluorphenylphenyl boronic acid (2.18 g, 13.82 mmol), potassium carbonate (3.41 g, 24.51 mmol) were added to a Schlenk tube containing a mixture of 1,2-dimethoxyethane and distilled water (4:1 v/v). The reaction mixture was degassed by multiple vacuum and N₂ purging cycles, and Pd(PPh₃)₄ (0.571 g, 0.494 mmol) was added to the tube under positive nitrogen pressure. The mixture was refluxed under nitrogen atmosphere

for 19 h and then cooled to room temperature. The mixture was poured onto distilled water and extracted multiple times with dichloromethane. The organic fractions were combined, washed with a portion of brine and dried over magnesium sulfate. Filtration and evaporation under reduced pressure gave the crude product (3.59 g). The crude product was purified by flash column chromatography (silica, hexane/ethyl acetate 9:1) to give 2.88 g of pure compound as a colourless oil. Yield: 94%. Rf: 0.43 (hexane/ethyl acetate, 6:1 on silica). ¹H NMR (500 MHz, CDCl₃) δ (ppm): 8.75 (dd, J = 7.5 Hz, 1.5 Hz, 1H), 8.08 (dt, J = 8.5, 10 Hz, 1H), 7.59 (m, 1H), 7.09 (dd, J = 5.0 Hz, 1.5 Hz, 1H, 7.02 (tdd, J = 10.0, 3.3, 1.0 Hz, 1H) 6.97(d, J = 1.0 Hz, 2H), 6.90 (dt, J = 3.0, 12.5 Hz, 1H), 2.34 (s, 3H),2.05 (s, 6H). 13 C NMR (126 MHz, CDCl₃) δ (ppm): 164.67, 161.97, 159.45, 152.85, 122.83, 150.16, 150.05, 137.78, 136.39, 135.37, 132.40, 132.36, 132.30, 132.26, 128.53, 125.48, 125.38, 123.67, 112.15, 112.12, 111.94, 111.91, 104.80, 104.54, 104.28, 21.20, 20.76. ¹⁹F { ¹H} NMR (471 MHz, CDCl₃) δ (ppm): -109.3 $(d, J = 9.4 \text{ Hz}, 1\text{F}), -112.7 (d, J = 9.4 \text{ Hz}, 1\text{F}). \text{ HR-MS (APCI}^+)$: $[M + H]^+$ Calculated: $(C_{20}H_{17}F_2NH)$ 310.1402; Found: 310.1402. The characterisation matches that reported. 22

General procedure for the synthesis of [(C^N)2Ir]2Cl2 dimers

The iridium(III) dimers, $[(ppy)_2Ir(\mu\text{-Cl})]_2$, $[(mesppy)_2Ir(\mu\text{-Cl})]_2$, $[(dFppy)_2Ir(\mu\text{-Cl})]_2$, $[(dFppy)_2Ir(\mu\text{-Cl})]_2$ were prepared according to the procedure described by Nonoyama. Briefly, to a round bottomed flask containing $IrCl_3\cdot 3H_2O$ (1.0 equiv.) and C^N ligand (2.2 equiv.) was added 2-ethoxyethanol to give a concentration of 0.02 M. The reaction mixture was degassed by multiple vacuum and N_2 purging cycles and the mixture was heated to reflux at 120 °C for 19 h. The solution was cooled to room temperature and the yellow precipitate was filtered, washed with water, hexane and ether and finally dried under vacuum to afford the pure material.

Tetrakis[2-(4',6'-difluorophenyl)-4-(2,4,6-trimethylphenyl)-pyridinato-N, C^2 ']-bis(μ-chloro)diiridium(m) [Ir(mesppy)₂-(μ-Cl)]₂ (D2). Yield: 70%. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 9.69 (d, J = 6.4 Hz, 2H), 7.77 (s, 4H), 7.55 (d, J = 9.0 Hz, 4H), 7.05 (d, J = 10.9 Hz, 8H), 6.86 (m, 7H), 6.72 (dt, J = 8.1 Hz, 4H), 5.95 (d, J = 8.1 Hz, 4H), 2.42 (s, 12H), 2.16 (s, 12H), 2.15 (s, 12H).

Tetrakis[2-(4',6'-difluorophenyl)-pyridinato-N, C^2]-bis(μ-chlorodiiridium(ш), [Ir(dFppy)₂(μ-Cl)]₂ (D3). Yield: 46%. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 9.12 (d, J = 5.0 Hz, 4H), 8.33 (d, J = 8.5 Hz, 4H), 7.88 (t, J = 7.0 Hz, 4H), 6.87 (td, J = 5.8, 1.0 Hz, 4H), 6.38 (td, J = 11.0, 2.0 Hz, 4H) 5.31 (dd, J = 9.3 Hz, 2.0 Hz, 4H). ¹⁹F {¹H} NMR (471 MHz, CD₂Cl₂) δ (ppm): -108.4 (d, J = 9.4 Hz, 4F), -110.6 (d, J = 9.4 Hz, 4F).

Tetrakis[2-(4',6'-difluorophenyl)-4-(2,4,6-trimethylphenyl)-pyridinato-N, C^2]-bis(μ -chloro)diiridium(m), [Ir(dFmesppy)₂-(μ -Cl)]₂ (D4). Yield: 81%. ¹H {¹⁹F} NMR (400 MHz, CD₂Cl₂) δ (ppm): 9.57 (d, J = 6.0 Hz, 4H), 8.13 (s, 4H), 7.02 (d, J = 10.4 Hz, 8H), 6.89 (dd, J = 6.0, 2.0 Hz, 4H), 5.29 (m, 4H) 2.38 (s, 12H), 2.12 (s, 24H). ¹⁹F {¹H} NMR (371 MHz, CD₂Cl₂) δ (ppm): -108.1 (d, J = 11.1 Hz, 4F), -110.2 (d, J = 10.8 Hz, 4F).

General procedure for the synthesis of [Ir(C^N)₂(P^P)]PF₆ complexes

To a Schlenk tube containing [Ir(ppy)₂Cl]₂, (D1), [Ir(mesppy)₂Cl]₂ (D2), ([Ir(dFppy)₂Cl]₂, (D3) or [Ir(dFmesppy)₂Cl]₂, (D4) (1.0 equiv.) and P^P ligand (2.5 equiv.) were added DCM and MeOH (2:1 v/v) to give a concentration of 0.03 M. The reaction mixture was degassed by multiple vacuum and N2 purging cycles and the mixture and the mixture was heated to 55 °C for 19 h under nitrogen atmosphere. The solution was cooled to room temperature and solid NH₄PF₆ (10.0 equiv.) was added and the reaction mixture was left to stir for a further 1 h. The resulting suspension was evaporated to dryness, with the residue then copiously washed with Et2O and distilled water. This crude product was purified by flash column chromatography (silica, DCM/MeOH gradient 100:0 to 90:10). Fractions containing the desired complex were combined and solid NH₄PF₆ (10 equiv.) was added. The suspension was stirred at room temperature for 0.5 h. This mixture was then evaporated to dryness, washed vigorously with distilled water and dried to afford the pure material.

Iridium(III)bis[2-phenylpyridinato]-4,5-bis(diphenylphosphino)-9,9-dimethylxanthenehexafluorophosphate, [Ir(ppy)₂(xantphos)]-(PF₆) (1a). Yield: 73%. R_f : 0.24 (DCM + 3% MeOH). Mp: 292–302 °C (degraded). 1 H NMR (500 MHz, CDCl₃) δ (ppm): 8.61 (d, J = 5.7 Hz, 2H), 7.88 (d, J = 4.1 Hz, 4H), 8.5 (dd, J = 1.3, 7.6 Hz, 2H), 7.32-7.31 (m, 4H), 7.12-7.07 (m, 7H), 6.84 (m, 8H), 6.78 (m, 2H), 6.71 (m, 2H), 6.64 (t, J = 7.4 Hz, 2H), 6.42 (t, J = 8.4 Hz, 4H), 6.28 (t, J = 7.8 Hz, 2H), 5.20 (d, J = 7.9 Hz, 2H), 1.15 (s, 3H), 1.13 (s, 3H). 13 C NMR (126 MHz, CDCl₃) δ (ppm): 154.44, 141.66, 138.79, 134.11, 134.03, 132.36, 131.12, 130.94, 129.87, 129.61, 129.52, 128.53, 128.02, 125.12, 125.10, 124.15, 123.58, 122.90, 120.32, 109.99, 30.97, 29.59, 22.88 ppm. ³¹P NMR (162 MHz, CDCl₃) δ (ppm): -20.2, -145.0 (hept). HR $NSI^{+}MS: [M - PF_{6}]^{+}$ Calculated: $(C_{61}H_{48}IrN_{2}OP_{2}): 1079.2871;$ Found: 1079.2856. The characterization matches that reported. 19a

Iridium(III)bis[2-phenyl-4-(2,4,6-trimethylphenyl)pyridinato]-4,5-bis(diphenylphosphino)-9,9-dimethylxanthene hexafluorophosphate, [Ir(mesppy)₂(xantphos)](PF₆) (2a). Yield: 89%. $R_{\rm f}$: 0.28 (DCM + 3% MeOH). Mp: 234–237 °C. ¹H NMR (500 MHz, CD_2Cl_2) δ (ppm): 8.76 (d, J = 5.9 Hz, 2H), 7.70 (s, 2H), 7.60 (d, J = 7.0 Hz, 2H), 7.31 (t, J = 7.6 Hz, 2H), 7.30 (d, J =8.0 Hz, 2H), 7.15 (m, 9H), 6.95 (m, 14H), 6.70 (t, J = 6.8 Hz, 2H), 6.63 (t, J = 6.8 Hz, 4H), 6.58 (dd, J = 1.5, 6.1 Hz, 2H), 6.30 (t, J = 6.9 Hz, 2H), 5.16 (d, J = 7.9 Hz, 2H), 2.36 (s, 6H), 2.01 (s, 6H)6H), 1.49 (s, 6H). 13 C NMR (126 MHz, CD₂Cl₂) δ (ppm): 154.17, 138.66, 134.08, 132.63, 132.59, 131.22, 131.16, 129.72, 129.65, 128.71, 128.51, 128.38, 128.19, 128.01, 127.11, 125.10, 124.93, 124.16, 122.86, 121.89, 29.68, 28.57, 20.88, 20.81, 20.11. ³¹P NMR (162 MHz, CD_2Cl_2) δ (ppm): –20.13, –144.41 (hept). HR NSI^+ MS: $[M - PF_6]^+$ Calculated: $(C_{79}H_{68}IrN_2OP_2)$: 1315.4436; Found: 1315.4419.

Ir(III)bis[2-(2,4-difluoro)-phenyl-4-(2,4,6-trimethylphenyl)pyridinato]-4,5-bis(diphenylphosphino)-9,9-dimethylxanthene hexafluorophosphate, [Ir(dFmesppy)₂(xantphos)](PF₆) (4a). Yield:

89%. R_f : 0.25 (DCM + 5% MeOH). Mp: 234–238 °C. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm) 8.85 (d, J = 5.5 Hz, 2H), 8.15 (s, 2H), 7.69 (d, J = 6.8 Hz, 2H), 7.48 (t, J = 7.9 Hz, 2H), 7.28 (t, J = 7.1 Hz, 6H), 7.22 (t, J = 7.1 Hz, 3H), 7.05 (m, 17H), 6.75 (t, J = 7.3, 4H), 6.68 (dd, J = 5.6, 1.9 Hz, 2H), 6.22 (dt, J = 7.4, 1.2 Hz, 2H), 2.39 (s, 6H), 2.15 (s, 6H), 2.03 (s, 6H), 1.57 (s, 6H), 1.53 (s, 6H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 164.80, 154.15, 153.91, 153.09, 138.82, 134.71, 134.23, 134.09, 133.85, 132.69, 132.51, 131.26, 130.29, 128.95, 128.82, 128.61, 128.20, 125.82, 125.64, 125.39, 116.86, 112.57, 112.42, 99.16, 28.77, 20.82, 20.11. ¹⁹F NMR (162 MHz, CD₂Cl₂) δ (ppm): -72.8, -74.4, -104.5, -108.5. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm) -20.13, -145.23 (hept). FT NSI⁺ MS: [M - PF₆]⁺ Calculated ($C_{79}H_{64}F_4$ IrN₂OP₂): 1387.4059; Found: 1357.4041.

Iridium(m)bis[2-phenylpyridinato]-bis[(2-diphenylphosphino)-phenyl]methane hexafluorophosphate, [Ir(ppy)₂(dpephos)](PF₆) (1b). Yield: 48%. R_f : 0.21 (DCM + 3% MeOH). Mp: 291–299 °C (degraded). 1 H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.76 (d, J = 6.4 Hz 2H), 7.70 (t, J = 8.0 Hz, 2H), 7.57 (d, J = 8.0 Hz, 2H), 7.16 (t, J = 8.5, 2H), 7.30–7.11 (m, 20H), 7.06 (t, J = 7.1 Hz, 2H), 6.92 (t, J = 7.2 Hz, 4H), 6.80–6.72 (m, 4H), 6.66 (m, 2H), 6.48–6.39 (m, 6H), 5.35 (dd, J = 8.4 Hz, 2H). 13 C NMR (126 MHz, CD₂Cl₂) δ (ppm): 166.60, 152.87, 140.85, 136.46, 133.22, 130.09, 128.48, 127.15, 126.03, 123.25, 123.33, 121.33, 120.84, 118.11, 117.09. 31 P NMR (162 MHz, CD₂Cl₂) δ (ppm): δ −15.98, −145.23 (hept). HR NSI $^+$ MS: [M − PF $_6$] $^+$ Calculated: (C $_{58}$ H $_{44}$ IrN $_2$ OP $_2$): 1039.2558; Found: 1039.2543. The characterization matches that reported. 19c

Iridium(III)bis[2-phenyl-4-(2,4,6-trimethylphenyl)pyridinato]bis[(2-diphenylphosphino)phenyl]methane hexafluorophosphate, [Ir(mesppy)₂(dpephos)](PF₆) (2b). Yield: 83%. $R_{\rm f}$: 0.27 (DCM + 3% MeOH). Mp 234–236 °C. ¹H NMR (500 MHz, CD_2Cl_2) δ (ppm): 8.86 (d, J = 5.2 Hz, 2H), 7.48 (s, 2H), 7.38-7.32 (m, 6H), 7.27 (q, J = 8.4 Hz, 4H), 7.20 (t, J =7.5 Hz, 8H), 7.12 (t, J = 9.3 Hz, 2H), 7.02–6.97 (m, 8H), 6.82 (t, J = 6.8 Hz, 2H, 6.70 (d, J = 10.1 Hz, 2H), 6.59-6.51 (m, 8H),5.39 (d, J = 6.2 Hz, 2H), 2.34 (s, 6H), 2.03 (s, 12H), 1.97 (s, 6H). 13 C NMR (126 MHz, CD₂Cl₂) δ (ppm): 154.58, 135.32, 135.24, 132.86, 132.72, 132.23, 130.56, 130.29, 129.94, 129.23, 128.67, 128.43, 128.36, 127.82, 125.16, 124.33, 124.25, 123.23, 121.78, 118.97, 20.78, 20.13, 0.75. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): -16.09, -145.23 (hept). HR NSI⁺ MS: $[M - PF_6]^+$ Calculated: $(C_{76}H_{64}IrN_2OP_2)$: 1275.4123; Found: 1275.4121.

Iridium(m)bis[2-(2,4-difluorophenyl)-4-(2,4,6-trimethylphenyl)-pyridinato]-bis[(2-diphenylphosphino)phenyl]methane hexafluorophosphate, [(dFmesppy)₂Ir(dpephos)](PF₆) (4b). Yield: 85%. R_f : 0.22 (DCM + 5% MeOH). Mp: 303–306. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm) 8.92 (d, J = 6.4 Hz, 2H), 7.90 (s, 2H), 7.45–7.13 (m, 20H), 7.10 (t, J = 6.8, 4H), 7.00 (d, J = 12.8 Hz, 4H), 6.74 (d, J = 5.5 Hz, 2H), 6.67 (t, J = 8.6 Hz, 3H), 6.61 (dd, J = 6.1, 1.1 Hz, 2H), 6.38 (dt, J = 9.6 Hz, 1.5 Hz, 2H), 2.34 (s, 6H), 2.03 (s, 6H), 1.96 (s, 6H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm) 164.79, 156.63, 154.62, 153.44, 138.68, 135.07, 134.76, 134.16, 133.38, 132.78, 132.09, 130.97, 129.35, 128.68, 128.02, 127.41, 125.50, 125.29, 124.74, 119.07, 112.91, 99.72, 65.67, 20.79, 20.15. ¹⁹F NMR (162 MHz, CD₂Cl₂) δ (ppm):

 $\begin{array}{l} -72.8,\ -74.4,\ -106.4,\ -108.4.\ ^{31}P\ \ NMR\ \left(162\ \ MHz,\ CD_{2}Cl_{2}\right)\\ \delta\left(ppm\right):\ -17.66,\ -145.23\ \left(hept\right).\ FT\ NSI^{^{+}}\ MS:\ \left[M-PF_{6}\right]^{^{+}}\ Calculated:\ \left(C_{70}H_{60}F_{4}IrN_{2}OP_{2}\right):\ 1347.3746;\ Found:\ 1347.3731. \end{array}$

Iridium(m)bis[2-phenylpyridine]-Bis[1,2-bis(diphenylphosphino)ethenato] hexafluorophosphate, [Ir(ppy)₂(dppe)](PF₆) (1c). Yield: 78%. R_f : 0.23 (DCM + 3% MeOH). Mp: 331–335 °C. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.42 (dd, J = 52 Hz, 2H), 7.77 (t, J = 8 Hz, 4H), 7.54 (d, J = 4.2 Hz, 2H), 7.51 (t, J = 4.2, 2H), 7.50 (t, J = 8 Hz, 4H), 7.45 (t, J = 8.3 Hz, 6H), 7.12 (t, J = 4.1 Hz, 2H), 7.07 (t, J = 5.1 Hz, 2H), 7.02 (t, J = 4.4 Hz, 2H), 6.88 (t, J = 8.5 Hz, 4H), 6.51 (t, J = 8 Hz, 4H), 6.32 (m, 4H), 1.27 (s, 2H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 167.49, 152.99, 152.96, 148.88, 148.62, 148.34, 143.26, 137.65, 132.69, 132.66, 132.62, 131.82, 130.36, 129.73, 129.63, 129.60, 129.56, 129.46, 129.42, 128.42, 128.34, 125.12, 123.64, 122.90, 120.18. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): −20.86, −145.23 (hept). HR NSI MS: [M − PF₆] Calculated: (C₄₈H₃₈IrN₂P₂): 897.2139; Found: 897.2121. The characterization matches that reported. ^{19d}

Iridium(m)bis[2-phenyl-4-(2,4,6-trimethylphenyl)pyridinato]bis[1,2-bis(diphenylphosphino)ethene] hexafluorophosphate, [Ir(mesppy)₂(dppe)](PF₆) (2c). Yield: 88%. $R_{\rm f}$: 0.28 (DCM + 3% MeOH). Mp 230–235 °C. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.59 (dd, J = 50 Hz, 2H), 7.84 (t, J = 7.7 Hz, 4H), 7.52 (d, J = 9.8 Hz, 2H), 7.50 (d, J = 10.2, 2H), 7.48 (m, 6H), 7.43 (d, J = 2.3 Hz, 2H), 7.11 (t, J = 6.1 Hz, 2H), 7.05 (q, J = 15.6, 4H), 7.00 (t, J = 9.8 Hz, 4H), 6.97 (m, 4H), 6.72 (t, J = 10.25 Hz, 4H), 6.24 (dd, J = 5.4 Hz, 4H), 6.19 (dd, J = 4.8 Hz, 2H), 1.97 (s, 6H), 1.85 (s, 6H), 1.27 (s, 6H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 173.80, 132.51, 131.81, 131.58, 130.42, 130.28, 129.72, 129.69, 129.53, 128.39, 125.18, 122.14, 123.77, 121.64, 83.41, 80.11, 20.74, 20.05. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): −19.71, −145.23 (hept). HR NSI⁺ MS: [M − PF₆]⁺ Calculated (C₆₆H₅₈IrN₂P₂): 1133.3704; Found: 1133.3695.

Iridium(III)bis[2-(2,4-difluorophenyl)-4-(2,4,6-trimethylphenyl)pyridinato]-bis[1,2-bis(diphenylphosphino)ethene] hexafluorophosphate, $[Ir(dFmesppy)_2(dppe)](PF_6)$ (4c). Yield: 88%. R_f : 0.30 (DCM + 5% MeOH). Mp: 369-372 °C. ¹H NMR (500 MHz, CD_2Cl_2) δ (ppm) 8.68-8.50 (m, 2H), 7.84-8.76 (m, 6H), 7.60–7.46 (m, 8H), 7.12 (t, J = 7.7 2H), 7.01 (dt, J = 7.7, 1.8 Hz, 4H), 6.94 (s, 4H), 6.79 (t, J = 8.9 Hz, 4H), 6.63 (dt, J = 8.9, 1.2 Hz, 2H), 6.22 (dd, J = 6.4, 1.7 Hz, 1H), 5.59 (m, 2H), 2.30 (s, 6H), 1.94 (s, 6H), 1.87 (s, 6H), 1.53 (s, 2H). ¹³C NMR (126 MHz, CD_2Cl_2) δ (ppm): 164.16, 153.02, 152.92, 148.84, 148.30, 138.57, 134.60, 134.38, 134.15, 132.47, 132.14, 130.63, 129.73, 128.66, 128.58, 128.42, 127.89, 127.98, 125.50, 125.33, 124.46, 113.84, 113.70, 100.00, 20.75, 20.04. ¹⁹F NMR (162 MHz, CD_2Cl_2) δ (ppm): -72.8, -74.4, -106.4, -108.0 ³¹P NMR (162 MHz, CD_2Cl_2) δ (ppm): -20.2, -145.23 (hept). FT NSI^+ MS: $[M - PF_6]^+$ Calculated $(C_{66}H_{54}F_4IrN_2P_2)$: 1205.3328; Found: 1205.3302.

Iridium(III)bis[2-(2,4-difluorophenyl)pyridinato]-bis[1,2-bis-(diphenylphosphino)ethene] hexafluorophosphate, [Ir(dFppy)₂-(dppe)](PF₆) (3c). Yield: 90%. $R_{\rm f}$: 0.20 (DCM + 5% MeOH). Mp: 346–348 °C. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.52–8.35 (m, 2H), 8.87 (dd, J = 7.9, 2.8 Hz, 2H), 7.74 (t, J = 8.5 Hz, 4H), 7.57 (t, J = 8.0 Hz, 2H), 7.55–7.42 (m, 6H), 7.11 (t, J = 7.4 Hz,

2H), 6.92 (t, J = 7.7, 4H), 6.63 (dd, J = 9.2, 2.4 Hz, 2H), 6.56 (t, J = 9.5 Hz, 4H), 6.35 (t, J = 6.3 Hz, 2H), 5.79 (m, 2H), 2.12 (s, 2H). ¹³C NMR (126 MHz, CD_2Cl_2) δ (ppm): 164.05, 160.64, 153.16, 148.59, 148.32, 148.05, 138.70, 132.62, 132.56, 132.19, 130.17, 129.92, 129.86, 129.50, 129.46, 129.43, 128.68, 128.62, 128.50, 128.14, 127.44, 126.78, 126.39, 124.14, 123.97, 123.35, 113.93, 113.78, 100.19, 99.98, 99.76. ¹⁹F NMR (162 MHz, CD_2Cl_2) δ (ppm): -72.8, -74.4, -106.3, -108.0. ³¹P NMR (162 MHz, CD_2Cl_2) δ (ppm): -21.56, -145.23 (hept). FT NSI MS: $[M - PF_6]^+$ Calculated ($C_{48}H_{34}F_4IrN_2P_2$): 969.1763; Found: 969.1733.

Iridium(m)bis[2-phenylpyridinato]-bis[1,2-bis(diphenylphosphino)ethane] hexafluorophosphate, [Ir(ppy)₂(sdppe)](PF₆) (1d). Yield: 82%. R_f : 0.23 (DCM + 3% MeOH). Mp: 156–164 °C. ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 7.63 (q, J = 8.6 Hz, 6H), 9.03 (t, J = 7.9 Hz, 4H), 7.37 (t, J = 8.4 Hz, 2H), 7.30 (t, J = 8.2 Hz, 2H), 7.23 (dd, J = 6.2 Hz, 4H), 6.98 (t, J = 7.3 Hz, 2H), 6.89 (q, J = 14.6 Hz, 4H), 6.81 (t, J = 6.0 Hz, 4H), 6.58 (t, J = 8.92 Hz, 4H), 6.25 (m, 4H), 3.84 (dd, J = 10.0 Hz, 2H), 2.74 (d, J = 6.08 Hz, 2H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 151.51, 141.23, 135.31, 131, 130.21, 129.29, 128.30, 127.20, 127.16, 126.62, 126.58, 122.83, 121.47, 112.04, 118.11, 66.42, 20.96. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): -10.79, -145.23 (hept). HR NSI⁺ MS: [M - PF₆]⁺ Calculated: (C₄₈H₄₀IrN₂P₂): 899.2296; Found: 899.2297.

Iridium(m)bis[2-phenylpyridinato]-4,5-bis(diphenylphosphino)-9-isopropylkanthene hexafluorophosphate, [Ir(ppy)₂(isopropxantphos)](PF₆) (1f). Yield: 55%. $R_{\rm f}$: 0.25 (DCM + 3% MeOH). Mp: 285–292 °C (degraded). ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.43 (s, 2H), 7.87 (t, J = 7.2 Hz, 4H), 7.60 (d, J = 7.6 Hz, 2H), 7.17 (t, J = 7.6 Hz, 2H), 7.09 (m, 6H), 6.87 (t, J = 6,6 Hz, 4H), 6.79 (t, J = 9.4 Hz, 4H), 6.64 (t, J = 7.4 Hz, 4H), 6.38 (t, J = 8.3 Hz, 4H), 6.30 (t, J = 7.6 Hz, 2H), 2.14 (s, 6H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 168.11, 141.86, 138.80, 135.41, 134.48, 132.44, 132.37, 130.96, 129.67, 128.55, 128.49, 128.15, 128.08, 124.24, 123.65, 122.99, 121.33, 120.34, 68.33, 30.97, 23.66, 22.86. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): −19.09, −145.23 (hept). HR NSI⁺ MS: [M − PF₆]⁺ Calculated (C₆₂H₄₈IrN₂OP₂): 1091.2871; Found: 1091.2863.

Iridium(m)bis[2-phenylpyridinato]-4,6-bis(diphenylphosphino)-phenoxazine hexafluorophosphate, [Ir(ppy)₂(nixantphos)](PF₆) (1e). Yield: 62%. $R_{\rm f}$: 0.24 (DCM + 3% MeOH). Mp: 297–305 °C (degraded). ¹H NMR (500 MHz, CD₂Cl₂) δ (ppm): 8.92 (d, J = 5.2 Hz, 2H), 7.90 (m, 4H), 7.30 (t, J = 8.0 Hz, 4H), 7.17 (t, J = 7.8 Hz, 2H), 7.07 (t, J = 8.0 Hz, 4H), 6.88 (m, 14H), 6.65 (t, J = 8.1 Hz, 2H), 6.46 (t, J = 7.9 Hz, 4H), 6.29 (t, J = 8.2 Hz, 4H), 5.27 (d, J = 8.0 Hz, 2H). ¹³C NMR (126 MHz, CD₂Cl₂) δ (ppm): 154.82, 138.50, 134.42, 134.38, 132.30, 132.27, 130.85, 130.00, 129.68, 129.39, 128.46, 128.06, 128.03, 127.99, 125.48, 124.52, 124.10, 123.69, 122.79, 120.33, 116.81. ³¹P NMR (162 MHz, CD₂Cl₂) δ (ppm): -21.51, -145.23 (hept). HR NSI⁺ MS: [M - PF₆]⁺ Calculated (C₅₈H₄₃IrN₃OP₂): 1052.25; Found: 1052.2487.

X-ray crystallography

Single crystals were grown by vapour diffusion of ether into concentrated CH₂Cl₂ solution (2a, 3c, 4b and 4c), and by slow

evaporation of mixed solutions of CH₂Cl₂/hexane (1a, 1b, 1c and 1e) or CH₂Cl₂/heptane (4a). Data were collected at either 173 K (1a, 1b, 1c, 1e, 3c, 4b and 4c) or 93 K (4a) on a Rigaku FR-X Ultrahigh brilliance Microfocus RA generator/confocal optics and Rigaku XtaLAB P200 system, with Mo Kα radiation $(\lambda = 0.71075 \text{ Å})$, or (2a) at 173 K on a Rigaku MM-007HF High brilliance RA generator/confocal optics and Rigaku XtaLAB P100 system, with Cu K α radiation ($\lambda = 1.54187$ Å). Intensity data were collected using ω steps (or ω and φ steps for 2a) accumulating area detector images spanning at least a hemisphere of reciprocal space. All data were corrected for Lorentz polarization effects, and a multiscan absorption correction was applied by using CrystalClear.³⁸ Structures were solved by Patterson methods (PATTY)³⁹ and refined by full-matrix leastsquares against F² (SHELXL-2013). 40 Non-hydrogen atoms were refined anisotropically, and hydrogen atoms were refined using a riding model, except for the NH hydrogen in 1e, which was located from the difference Fourier map and refined subject to a distance restraint. All calculations were performed using the CrystalStructure interface.41

Photophysical measurements

All samples were prepared in HPLC grade acetonitrile with varying concentrations in the order of µM. Absorption spectra were recorded at room temperature using a Shimadzu UV-1800 double beam spectrophotometer. Molar absorptivity determination was verified by linear least-squares fit of values obtained from at least four independent solutions at varying concentrations with absorbance ranging from 6.05×10^{-5} to $2.07 \times 10^{-5} \text{ M}.$

The sample solutions for the emission spectra were prepared in HPLC grade MeCN and degassed via three freezepump-thaw cycles using an in-house designed quartz cuvette. Steady state emission and excitation spectra and time-resolved emission spectra were recorded at 298 K using an Edinburgh Instruments F980. All samples for steady state measurements were excited at 360 nm while samples for time-resolved measurements were excited at 378 nm using a PDL 800-D pulsed diode laser. Emission quantum yields were determined using the optically dilute method.³² A stock solution with absorbance of ca. 0.5 was prepared and then four dilutions were prepared with dilution factors between 2 and 20 to obtain solutions with absorbances of ca. 0.095, 0.065, 0.05 and 0.018, respectively. The Beer-Lambert law was found to be linear at the concentrations of the solutions. The emission spectra were then measured after the solutions were rigorously degassed via three freeze-pump-thaw cycles prior to spectrum acquisition. For each sample, linearity between absorption and emission intensity was verified through linear regression analysis and additional measurements were acquired until the Pearson regression factor (R^2) for the linear fit of the data set surpassed 0.9. Individual relative quantum yield values were calculated for each solution and the values reported represent the slope value. The equation $\Phi_s = \Phi_r(A_r/A_s)(I_s/I_r)(n_s/n_r)^2$ was used to calculate the relative quantum yield of each of the sample, where $\Phi_{\rm r}$ is the absolute quantum yield of the reference, n is the refractive index of the solvent, A is the absorbance at the excitation wavelength, and I is the integrated area under the corrected emission curve. The subscripts s and r refer to the sample and reference, respectively. A solution of quinine sulfate in 0.5 M H_2SO_4 ($\Phi_r = 54.6\%$) was used as the external reference. 29,42

Electrochemistry measurements

Cyclic voltammetry (CV) measurements were performed on an Electrochemical Analyzer potentiostat model 600D from CH Instruments. Solutions for cyclic voltammetry were prepared in MeCN and degassed with MeCN-saturated nitrogen by bubbling for about 10 min prior to scanning. Tetra(n-butyl)ammoniumhexafluorophosphate (TBAPF6; ca. 0.1 M in MeCN) was used as the supporting electrolyte. An Ag/Ag⁺ electrode (silver wire in a solution of 0.1 M KCl in H₂O) was used as the pseudoreference electrode; a Pt electrode was used for the working electrode and a Pt electrode was used as the counter electrode. The redox potentials are reported relative to a saturated calomel electrode (SCE) electrode with a ferrocene/ferrocenium (Fc/Fc⁺) redox couple as an internal reference (0.38 V vs. SCE).33

Electroluminescent devices

All commercial materials were used as received: aqueous dispersion of poly(3,4-ethylenedioxythiophene):poly(styrenesulfonate) (PEDOT:PSS, CLEVIOSTM P VP Al 4083; Heraeus); ionic liquid (IL) 1-butyl-3-methyl-imidazolium tetrafluoroborate [Bmim][PF₆] (Aldrich) and SPPO13 (Luminescence Technology Corp.). NS25 was synthetized according to the previously reported procedure.36 The LEEC devices were made as follows. Indium tin oxide ITO-coated glass plates were patterned by conventional photolithography (Naranjo Substrates). The substrates were cleaned ultrasonically in water-soap, water and 2-propanol baths. After drying, the substrates were placed in a UV-ozone cleaner (Jelight 42-220) for 20 min. An 80 nm layer of PEDOT:PSS was spin-coated on the ITO-glass substrate. To prepare the emitting layer, the devices were spin-coated at 1000 rpm for 30 s from a solution containing either the ionic transition metal complex (iTMC) and the IL at a 4:1 molar ratio (iTMC:IL) or a 1:1 mass ratio mixture of NMS25: SPPO13 (with 10 wt% of 4c). After deposition of the emitting layer, the devices were transferred into an inert atmosphere glovebox, where the aluminium electrode was thermally evaporated using a shadow mask. The size of the LEEC was 6.5 mm^2 .

The thickness of the films was determined with an Ambios XP-1 profilometer. Time dependence of luminance and voltage was measured by applying pulsed current and by monitoring the voltage and the luminance simultaneously by a True Colour Sensor MAZeT (MTCSICT Sensor) using a Lifetime Test System designed by BoTEST (Botest OLT OLED Lifetime-Test System). Electroluminescence spectra were recorded with an Avantes fiber-optics photo-spectrometer.

For OLED devices, ITO-coated soda lime glass substrates were cleaned by ultrasound in acetone and 2-propanol, followed by an oxygen plasma treatment. A 40 nm thick poly(3,4ethylenedioxythiophene):poly(styrenesulfonate) (PEDOT:PSS) was spin-coated on the ITO and baked on a hot plate at 120 °C for 10 min. A hole-transport layer of 35 nm-thick poly(N-vinylcarbazole) (PVK) was spin-coated on the PEDOT:PSS layer and baked at 80 °C for 2 hours in a nitrogen glove box. An emissive layer of complexes blended with 1,3-bis(N-carbazolyl)benzene (mCP) and 4-diphenylphosphoryl dibenzofuran (o-DBFPPO) were spin-coated on top of the PVK layer. An electron-transport layer of 60 nm-thick bis-4,6-(3,5-di-3-pyridylphenyl)-2-methylpyrimidine (B3PyMPM) was deposited through a shadow mask. A cathode of Ca/Al (20 nm/200 nm) was then deposited on the B3PyMPM layer in the same vacuum system. After the evaporation, the devices were encapsulated with optical curing adhesive (Norland NOA68) and glass coverslips in the glove box. The device has an active area of 2×1.5 mm². The current-voltage-light output characteristics were measured using a Keithley source measure unit with a calibrated silicon photodiode. The EL spectra were measured using a charge coupled device spectrograph.

Acknowledgements

EZ-C acknowledges the University of St Andrews for financial support. We thank Johnson Matthey and Umicore AG for the gift of materials and Cihang Yu for the preparation of isoprop-xantphos. We thank Dr Nail Shavaleev for the synthesis of NMS25. IDWS and AKB acknowledge support from EPSRC (EP/J01771X). We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University for analytical services. This work has been supported by the Spanish Ministry of Economy and Competitiveness (MINECO) MAT2014–55200.

References

- 1 H. Sasabe and J. Kido, Development of high performance OLEDs for general lighting, *J. Mater. Chem. C*, 2013, 1, 1699.
- 2 (a) F. Kessler, Y. Watanabe, H. Sasabe, H. Katagiri, M. K. Nazeeruddin, M. Grätzel and J. Kido, High-performance pure blue phosphorescent OLED using a novel bisheteroleptic iridium(iii) complex with fluorinated bipyridyl ligands, J. Mater. Chem. C, 2013, 1, 1070; (b) S. Evariste, M. Sandroni, T. W. Rees, C. Roldan-Carmona, L. Gil-Escrig, H. J. Bolink, E. Baranoff and E. Zysman-Colman, Fluorine-free blue-green emitters for light-emitting electrochemical cells, J. Mater. Chem. C, 2014, 2, 5793; (c) J. Zhuang, W. Li, W. Su, Y. Liu, Q. Shen, L. Liao and M. Zhou, Highly efficient phosphorescent organic light-emitting diodes using a homoleptic iridium(III) complex as a sky-blue dopant, Org. Electron., 2013, 14, 2596.
- 3 (a) For a perspective on the correlation between structure and quantum yields in neutral iridium(III) complexes see: Y. You and S. Y. Park, Phosphorescent iridium(iii) com-

- plexes: toward high phosphorescence quantum efficiency through ligand control, Dalton Trans., 2009, 1267; (b) M. S. Lowry and S. Bernhard, Synthetically Tailored Excited States: Phosphorescent, Cyclometalated Iridium (III) Complexes and Their Applications, Chem. - Eur. J., 2006, **12**, 7970; (c) S. Lamansky, P. Djurovich, D. Murphy, F. Abdel-Razzaq, H.-E. Lee, C. Adachi, P. E. Burrows, S. R. Forrest and M. E. Thompson, Highly Phosphorescent Bis-Cyclometalated Iridium Complexes: Synthesis, Photophysical Characterization, and Use in Organic Light Emitting Diodes, J. Am. Chem. Soc., 2001, 123, 4304; (d) S. Lamansky, P. Djurovich, D. Murphy, F. Abdel-Razzaq, R. Kwong, I. Tsyba, M. Bortz, B. Mui, R. Bau and M. E. Thompson, Synthesis and Characterization of Phosphorescent Cyclometalated Iridium Complexes, Inorg. Chem., 2001, 40, 1704.
- 4 (a) C. Ulbricht, B. Beyer, C. Friebe, A. Winter and U. S. Schubert, Recent Developments in the Application of Phosphorescent Iridium(III) Complex Systems, *Adv. Mater.*, 2009, 21, 4418; (b) L. Wang, N. Wang, Y. Zhang, H. He and J. Zhang, Color tuning from red to green of bis-cyclometalated iridium(III) emitters based on benzoimidazole ligands in OLEDs: A DFT and TD-DFT investigation, *Synth. Met.*, 2014, 194, 160.
- 5 (a) J.-S. Lu, H.-F. Chen, J.-C. Kuo, R. Sun, C.-Y. Cheng, Y.-S. Yeh, H.-C. Su and K.-T. Wong, Efficient solid-state white light-emitting electrochemical cells employing embedded red color conversion layers, *J. Mater. Chem. C*, 2015, 3, 2802; (b) J. Qiao, L. Duan, L. Tang, L. He, L. Wang and Y. Qiu, High-efficiency orange to near-infrared emissions from bis-cyclometalated iridium complexes with phenyl-benzoquinoline isomers as ligands, *J. Mater. Chem.*, 2009, 19, 6573.
- 6 (a) A. F. Henwood, S. Evariste, A. M. Z. Slawin and E. Zysman-Colman, Rigid biimidazole ancillary ligands as an avenue to bright deep blue cationic iridium(iii) complexes, *Faraday Discuss.*, 2014, 174, 165; (b) S.-J. Yun, H.-J. Seo, M. Song, S.-H. Jin and Y. I. Kim, Blue Emitting Cationic Iridium Complexes Containing Two Substituted 2-Phenylpyridine and One 2,2'-Biimidazole for Solution-Processed Organic Light-Emitting Diodes (OLEDs), *Bull. Korean Chem. Soc.*, 2012, 33, 3645.
- 7 S. B. Meier, W. Sarfert, J. M. Junquera-Hernández, M. Delgado, D. Tordera, E. Ortí, H. J. Bolink, F. Kessler, R. Scopelliti, M. Grätzel, M. K. Nazeeruddin and E. Baranoff, A deep-blue emitting charged bis-cyclometallated iridium(iii) complex for light-emitting electrochemical cells, J. Mater. Chem. C, 2013, 1, 58.
- 8 (a) E. Orselli, G. S. Kottas, A. E. Konradsson, P. Coppo, R. Fröhlich, L. De Cola, A. van Dijken, M. Büchel and H. Börner, Blue-Emitting Iridium Complexes with Substituted 1,2,4-Triazole Ligands, Äâ Synthesis, Photophysics, and Devices, *Inorg. Chem.*, 2007, 46, 11082; (b) C.-H. Yang, Y.-M. Cheng, Y. Chi, C.-J. Hsu, F.-C. Fang, K.-T. Wong, P.-T. Chou, C.-H. Chang, M.-H. Tsai and C.-C. Wu, Blue-Emitting Heteroleptic Iridium(III) Complexes Suitable for

- High-Efficiency Phosphorescent OLEDs, Angew. Chem., Int. Ed., 2007, 46, 2418; (c) J. M. Fernandez-Hernandez, S. Ladouceur, Y. Shen, A. Iordache, X. Wang, L. Donato, S. Gallagher-Duval, M. de Anda Villa, J. D. Slinker, L. De Cola and E. Zysman-Colman, Blue light emitting electrochemical cells incorporating triazole-based luminophores, J. Mater. Chem. C, 2013, 1, 7440.
- 9 L. He, L. Duan, J. Qiao, R. Wang, P. Wei, L. Wang and Y. Qiu, Blue-Emitting Cationic Iridium Complexes with 2-(1H-Pyrazol-1-yl)pyridine as the Ancillary Ligand for Efficient Light-Emitting Electrochemical Cells, Adv. Funct. Mater., 2008, 18, 2123.
- 10 T. Hu, L. He, L. Duan and Y. Qiu, Solid-state light-emitting electrochemical cells based on ionic iridium(iii) complexes, J. Mater. Chem., 2012, 22, 4206.
- 11 S. Ladouceur and E. Zysman-Colman, A Comprehensive Survey of Cationic Iridium(III) Complexes Bearing Nontraditional Ligand Chelation Motifs, Eur. J. Inorg. Chem., 2013, 2013, 2985.
- 12 J. A. Gillespie, E. Zuidema, P. W. N. M. van Leeuwen and P. C. J. Kamer, in Phosphorus(III) Ligands in Homogeneous Catalysis: Design and Synthesis, John Wiley & Sons, Ltd, 2012, p. 1.
- 13 (a) P. Alam, P. Das, C. Climent, M. Karanam, D. Casanova, A. R. Choudhury, P. Alemany, N. R. Jana and I. R. Laskar, Facile tuning of the aggregation-induced emission wavelength in a common framework of a cyclometalated iridium(iii) complex: micellar encapsulated probe in cellular imaging, J. Mater. Chem. C, 2014, 2, 5615; (b) C. S. Chin, M.-S. Eum, S. y. Kim, C. Kim and S. K. Kang, New Type of Photoluminescent Iridium Complex: Novel Synthetic Route for Cationictrans-Bis(2-phenylpyridinato)iridium(III) Complex, Eur. J. Inorg. Chem., 2006, 2006, 4979; (c) Y. Koga, N. Yoshida and K. Matsubara, PL and EL behavior of nearred luminescent metallopolymer easily prepared from phosphine-containing copolymer and chloro{bis(1-phenylisoquinolinato-N,C2')}iridium(III), and its monomeric analog, J. Polym. Sci., Part A: Polym. Chem., 2009, 47, 4366; (d) X. Shen, H. Yang, X.-H. Hu, Y. Xu, F.-L. Wang, S. Chen and D.-R. Zhu, Synthesis, crystal structure and photophysical properties of a series of new neutral iridium(III) complexes with 2-phenylpyridine, Inorg. Chem. Commun., 2009, 12, 785; (e) P. Alam, G. Kaur, C. Climent, S. Pasha, D. Casanova, P. Alemany, A. Roy Choudhury and I. R. Laskar, New 'aggregation induced emission (AIE)' active cyclometalated iridium(iii) based phosphorescent sensors: high sensitivity for mercury(ii) ions, Dalton Trans., (f) P. Alam, 16431; M. D. Bandyopadhyay, A. R. Choudhury and I. R. Laskar, Aggregation-Induced Emission Activity in Iridium(III) Diimine Complexes: Investigations of Their Vapochromic Properties, Eur. J. Inorg. Chem., 2014, 2014, 3710.
- 14 Y.-Y. Lyu, Y. Byun, O. Kwon, E. Han, W. S. Jeon, R. R. Das and K. Char, Substituent Effect on the Luminescent Properties of a Series of Deep Blue Emitting Mixed Ligand Ir(III) Complexes, J. Phys. Chem. B, 2006, 110, 10303.

- 15 C. H. Lin, Y. Y. Chang, J. Y. Hung, C. Y. Lin, Y. Chi, M. W. Chung, C. L. Lin, P. T. Chou, G. H. Lee, C. H. Chang and W. C. Lin, Iridium(III) complexes of a dicyclometalated phosphite tripod ligand: strategy to achieve blue phosphorescence without fluorine substituents and fabrication of OLEDs, Angew. Chem., Int. Ed., 2011, 50, 3182.
- 16 (a) Y. C. Chiu, C. H. Lin, J. Y. Hung, Y. Chi, Y. M. Cheng, K. W. Wang, M. W. Chung, G. H. Lee and P. T. Chou, Authentic-blue phosphorescent iridium(III) complexes bearing both hydride and benzyl diphenylphosphine; control of the emission efficiency by ligand coordination geometry, Inorg. Chem., 2009, 48, 8164; (b) C. H. Lin, Y. Chi, M. W. Chung, Y. J. Chen, K. W. Wang, G. H. Lee, P. T. Chou, W. Y. Hung and H. C. Chiu, Heteroleptic Ir(III) complexes containing both azolate chromophoric chelate and diphenylphosphinoaryl cyclometalates; reactivities, electronic properties and applications, Dalton Trans., 2011, 40, 1132; (c) Y.-C. Chiu, J.-Y. Hung, Y. Chi, C.-C. Chen, C.-H. Chang, C.-C. Wu, Y.-M. Cheng, Y.-C. Yu, G.-H. Lee and P.-T. Chou, En Route to High External Quantum Efficiency (~12%), Organic True-Blue-Light-Emitting Diodes Employing Novel Design of Iridium (III) Phosphors, Adv. Mater., 2009, 21, 2221; (d) J.-Y. Hung, C.-H. Lin, Y. Chi, M.-W. Chung, Y.-J. Chen, G.-H. Lee, P.-T. Chou, C.-C. Chen and C.-C. Wu, Phosphorescent Ir(iii) complexes bearing double benzyldiphenylphosphine cyclometalates; strategic synthesis, fundamental and integration for white OLED fabrication, J. Mater. Chem., 2010, 20, 7682.
- 17 (a) B.-S. Du, C.-H. Lin, Y. Chi, J.-Y. Hung, M.-W. Chung, T.-Y. Lin, G.-H. Lee, K.-T. Wong, P.-T. Chou, W.-Y. Hung and H.-C. Chiu, Diphenyl(1-naphthyl)phosphine Ancillary for Assembling of Red and Orange-Emitting Ir(III) Based Phosphors; Strategic Synthesis, Photophysics, and Organic Light-Emitting Diode Fabrication, Inorg. Chem., 2010, 49, 8713; (b) X. Shang, D. Han, S. Guan, X. Wang and G. Zhang, Theoretical investigation on the electronic structures and phosphorescent properties of a series of Ir(III) complexes with the diphenyl(1-naphthyl)phosphine ancillary ligand, Chem. Phys. Lett., 2014, 614, 110; (c) F. Zhang, D. Ma, L. Duan, J. Qiao, G. Dong, L. Wang and Y. Qiu, Synthesis, characterization, and photophysical and electroproperties luminescent of blue-emitting cationic iridium(III) complexes bearing nonconjugated ligands, Inorg. Chem., 2014, 53, 6596.
- 18 A.-F. Ma, H.-J. Seo, S.-H. Jin, U.-C. Yoon, M.-H. Hyun, S.-K. Kang and Y.-I. Kim, Novel Cationic 2-Phenylpyridinebased Iridium(III) Complexes Bearing an Ancillary Phosphine Ligand: Synthesis, Photophysics and Crystal Structure, Bull. Korean Chem. Soc., 2009, 30, 2754.
- (a) S.-X. Luo, L. Wei, X.-H. Zhang, M. H. Lim, K. X. V. Lin, M. H. V. Yeo, W.-H. Zhang, Z.-P. Liu, D. J. Young and T. S. A. Hor, Enhanced Emission and Analyte Sensing by Iridium(III) Cyclometalated Bearing Bent Diphosphine Chelators, Organometallics, 2013, 32, 2908; (b) M. S. Lowry, W. R. Hudson, R. A. Pascal Jr. and S. Bernhard, Accelerated Luminophore Discovery

through Combinatorial Synthesis, J. Am. Chem. Soc., 2004, 126, 14129; (c) Y. Jian, S. Peng, X. Li, X. Wen, J. He, L. Jiang and Y. Dang, Novel solid state emitting iridium complexes with reduced phosphorescence-self-quenching triggered by steric blocking phosphine ligands, Inorg. Chim. Acta, 2011, 368, 37; (d) E. C. Constable, C. E. Housecroft, E. Schönhofer, J. Schönle and J. A. Zampese, Softening the donor set for light-emitting electrochemical cells: [Ir(ppv)2 (N^N) +, $[Ir(ppy)2(P^P)]$ + and $[Ir(ppy)2(P^S)]$ + salts, Polyhedron, 2012, 35, 154.

- 20 (a) L. Donato, P. Abel and E. Zysman-Colman, Cationic iridium(iii) complexes bearing a bis(triazole) ancillary ligand, Dalton Trans., 2013, 42, 8402; (b) S. Ladouceur, D. Fortin and E. Zysman-Colman, Enhanced Luminescent Iridium(III) Complexes Bearing Aryltriazole Cyclometal-50, lated Ligands, Inorg. Chem., 2011, 11514; (c) S. Ladouceur, K. N. Swanick, S. Gallagher-Duval, Z. Ding and E. Zysman-Colman, Strongly Blue Luminescent Cationic Iridium(III) Complexes with an Electron-Rich Ancillary Ligand: Evaluation of Their Optoelectronic and Electrochemiluminescence Properties, Eur. J. Inorg. Chem., 2013, 2013, 5329.
- 21 (a) J. M. Fernández-Hernández, C.-H. Yang, J. I. Beltrán, V. Lemaur, F. Polo, R. Frőhlich, J. Cornil and L. De Cola, Control of the Mutual Arrangement of Cyclometalated Ligands in Cationic Iridium(III) Complexes. Synthesis, Spectroscopy, and Electroluminescence of the Different Isomers, J. Am. Chem. Soc., 2011, 133, 10543; (b) M. K. Nazeeruddin, R. T. Wegh, Z. Zhou, C. Klein, Q. Wang, F. DeAngelis, S. Fantacci and M. Grätzel, Efficient Green-Blue-Light-Emitting Cationic Iridium Complex for Light-Emitting Electrochemical Cells, Inorg. Chem., 2006, 45, 9245; (c) F. De Angelis, S. Fantacci, N. Evans, C. Klein, S. M. Zakeeruddin, J.-E. Moser, K. Kalyanasundaram, H. J. Bolink, M. Gratzel and M. K. Nazeeruddin, Controlling Phosphorescence Color and Quantum Yields in Cationic Iridium Complexes: and A Combined Experimental and Theoretical Study, Inorg. Chem., 2007, 46, 5989; (d) C. Shik Chin, M.-S. Eum, S. yi Kim, C. Kim and S. Kwon Kang, Blue-Light-Emitting Complexes: Cationic (2-Phenylpyridinato)iridium(III) Complexes with Strong-Field Ancillary Ligands, Eur. J. Inorg. Chem., 2007, 2007, 372; (e) L. He, J. Qiao, L. Duan, G. Dong, D. Zhang, L. Wang and Y. Qiu, Toward Highly Efficient Solid-State White Light-Emitting Electrochemical Cells: Blue-Green to Red Emitting Cationic Iridium Complexes with Imidazole-Type Ancillary Ligands, Adv. Funct. Mater., 2009, 19, 2950; (f) M. Mydlak, C. Bizzarri, D. Hartmann, W. Sarfert, G. Schmid and L. De Cola, Positively Charged Iridium(III) Triazole Derivatives as Blue Emitters for Light-Emitting Electrochemical Cells, Adv. Funct. Mater., 2010, 20, 1812; (g) L. He, L. Duan, J. Qiao, D. Zhang, L. Wang and Y. Qiu, Enhanced stability of blue-green light-emitting electrochemical cells based on a cationic iridium complex with 2-(1-phenyl-1H-pyrazol-3yl)pyridine as the ancillary ligand, Chem. Commun., 2011, 47, 6467; (h) M. Felici, P. Contreras-Carballada, Y. Vida,

J. M. M. Smits, R. J. M. Nolte, L. De Cola, R. M. Williams and M. C. Feiters, IrIII and RuII Complexes Containing Triazole-Pyridine Ligands: Luminescence Enhancement upon Substitution with b-Cyclodextrin, Chem. - Eur. J., 2009, 15, 13124; (i) L. He, L. Duan, J. Qiao, G. Dong, L. Wang and Y. Oiu, Highly Efficient Blue-Green and White Light-Emitting Electrochemical Cells Based on a Cationic Iridium Complex with a Bulky Side Group, Chem. Mater., 2010, 22, 3535; (j) A. B. Tamayo, S. Garon, T. Sajoto, P. I. Djurovich, I. M. Tsyba, R. Bau and M. E. Thompson, Cationic Biscyclometalated Iridium(III) Diimine Complexes and Their Use in Efficient Blue, Green, and Red Electroluminescent Devices, Inorg. Chem., 2005, 44, 8723; (k) C.-H. Yang, J. Beltran, V. Lemaur, J. Cornil, D. Hartmann, W. Sarfert, R. Fröhlich, C. Bizzarri and L. De Cola, Iridium Metal Complexes Containing N-Heterocyclic Carbene Ligands for Blue-Light-Emitting Electrochemical Cells, Inorg. Chem., 2010, 49, 9891; (l) H.-F. Chen, W.-Y. Hung, S.-W. Chen, T.-C. Wang, S.-W. Lin, S.-H. Chou, C.-T. Liao, H.-C. Su, H.-A. Pan, P.-T. Chou, Y.-H. Liu and K.-T. Wong, Cationic Iridium Complexes with Intramolecular π - π Interaction and Enhanced Steric Hindrance for Solid-State Light-Emitting Electrochemical Cells, Inorg. Chem., 2012, 51, 12114; (m) S. B. Meier, W. Sarfert, J. M. Junquera-Hernandez, M. Delgado, D. Tordera, E. Orti, H. J. Bolink, F. Kessler, R. Scopelliti, M. Gratzel, M. K. Nazeeruddin and E. Baranoff, A deep-blue emitting charged bis-cyclometallated iridium(iii) complex for light-emitting electrochemical cells, J. Mater. Chem. C, 2013, 1, 58; (n) S. Soman, J. C. Manton, J. L. Inglis, Y. Halpin, B. Twamley, E. Otten, W. R. Browne, L. De Cola, J. G. Vos and M. T. Pryce, New synthetic pathways to the preparation of near-blue emitting heteroleptic Ir(iii)N6 coordinated compounds with microsecond lifetimes, Chem. Commun., 2014, 50, 6461; (o) N. Darmawan, C. H. Yang, M. Mauro, M. Raynal, S. Heun, J. Pan, H. Buchholz, P. Braunstein and L. De Cola, Efficient near-UV emitters based on cationic bis-pincer iridium(III) carbene complexes, Inorg. Chem., 2013, 52, 10756; (p) W.-T. Chen, Y.-J. Chen, C.-S. Wu, J.-J. Lin, W.-L. Su, S.-H. Chen and S.-P. Wang, Two new bluephosphorescent Ir(III) cyclometalated complexes demonstrating the pushing-up effects of amino on levels of Pi-type molecular orbitals, *Inorg. Chim. Acta*, 2013, 408,

22 V. N. Kozhevnikov, Y. Zheng, M. Clough, H. A. Al-Attar, G. C. Griffiths, K. Abdullah, S. Raisys, V. Jankus, M. R. Bryce and A. P. Monkman, Cyclometalated Ir(III) Complexes for High-Efficiency Solution-Processable Blue PhOLEDs, Chem. Mater., 2013, 25, 2352.

225.

- 23 C. Zhong, C. Duan, F. Huang, H. Wu and Y. Cao, Materials and Devices toward Fully Solution Processable Organic Light-Emitting Diodes, Chem. Mater., 2011, 23, 326.
- 24 (a) N. Miyaura, T. Yanagi and A. Suzuki, The Palladium-Catalyzed Cross-Coupling Reaction of Phenylboronic Acid with Haloarenes in the Presence of Bases, Synth. Commun., 1981, 11, 513; (b) N. Miyaura and A. Suzuki, Palladium-

- Catalyzed Cross-Coupling Reactions of Organoboron Compounds, Chem. Rev., 1995, 95, 2457.
- 25 M. Nonoyama, Benzo[h]quinolin-10-yl-N Iridium(III) Complexes, Bull. Chem. Soc. Jpn., 1974, 47, 767.
- 26 F. H. Allen, The Cambridge Structural Database: a quarter of a million crystal structures and rising, Acta Crystallogr., Sect. B: Struct. Sci., 2002, 58, 380.
- 27 (a) S. Ladouceur, D. Fortin and E. Zysman-Colman, The role of substitution on the photophysical properties of 5,5'diaryl-2,2'-bipyridine (bpy*) in [Ir(ppy)2(bpy*)]PF6 complexes: A combined experimental and theoretical study, Inorg. Chem., 2010, 49, 5625; (b) K. K.-W. Lo and J. S.-Y. Lau, Cyclometalated Iridium(III) Diimine Bis(biotin) Complexes as the First Luminescent Biotin-Based Cross-Linkers for Avidin, Inorg. Chem., 2007, 46, 700; (c) J. B. Waern, C. Desmarets, L.-M. Chamoreau, H. Amouri, A. Barbieri, C. Sabatini, B. Ventura and F. Barigelletti, Luminescent Cyclometalated RhIII, IrIII, and (DIP)2RuII Complexes with Carboxylated Bipyridyl Ligands: Synthesis, X-ray Molecular Structure, and Photophysical Properties, Inorg. Chem., 2008, 47, 3340; (d) H. J. Bolink, E. Coronado, R. n. D. Costa, N. Lardiés and E. Ortì, Near-Quantitative Internal Quantum Efficiency in a Light-Emitting Electrochemical Cell, Inorg. Chem., 2008, 47, 9149; (e) A. B. Tamayo, B. D. Alleyne, P. I. Djurovich, S. Lamansky, I. Tsyba, N. N. Ho, R. Bau and M. E. Thompson, Synthesis and Characterization of Facial and Meridional Tris-cyclometalated Iridium(III) Complexes, J. Am. Chem. Soc., 2003, 125, 7377.
- 28 For recent reviews see: L. Flamigni, A. Barbieri, C. Sabatini, B. Ventura and F. Barigelletti, Photochemistry and Photophysics of Coordination Compounds: Iridium, Top. Curr. Chem., 2007, 281, 143.
- 29 W. H. Melhuish, Quantum Efficiences Of Fluorescence Of Organic Substances: Effect Of Solvent And Concentration Of The Fluorescent Solute 1, J. Phys. Chem., 1961, **65**, 229.
- 30 J. Wang, F.-Q. Bai, B.-H. Xia and H.-X. Zhang, Efficient Blue-Emitting Ir(III) Complexes with Phosphine Carbanion-Based Ancillary Ligand: A DFT Study, J. Phys. Chem. A, 2011, 115, 11689.
- 31 N. M. Shavaleev, F. Monti, R. D. Costa, R. Scopelliti, H. J. Bolink, E. Ortí, G. Accorsi, N. Armaroli, E. Baranoff, M. Grätzel and M. K. Nazeeruddin, Bright Blue Phosphor-

- escence from Cationic Bis-Cyclometalated Iridium(III) Isocyanide Complexes, Inorg. Chem., 2012, 51, 2263.
- 32 (a) H.-C. Su, F.-C. Fang, T.-Y. Hwu, H.-H. Hsieh, H.-F. Chen, G.-H. Lee, S.-M. Peng, K.-T. Wong and C.-C. Wu, Highly Efficient Orange and Green Solid-State Light-Emitting Electrochemical Cells Based on Cationic IrIII Complexes with Enhanced Steric Hindrance, Adv. Funct. Mater., 2007, 17, 1019; (b) C. Rothe, C.-J. Chiang, V. Jankus, K. Abdullah, X. Zeng, R. Jitchati, A. S. Batsanov, M. R. Bryce and A. P. Monkman, Ionic Iridium(III) Complexes with Bulky Side Groups for Use in Light Emitting Cells: Reduction of Concentration Quenching, Adv. Funct. Mater., 2009, 19, 2038.
- 33 V. V. Pavlishchuk and A. W. Addison, Conversion constants for redox potentials measured versus different reference electrodes in acetonitrile solutions at 25 °C, Inorg. Chim. Acta, 2000, 298, 97.
- 34 S. van Reenen, P. Matyba, A. Dzwilewski, R. A. J. Janssen, L. Edman and M. Kemerink, Salt Concentration Effects in Planar Light-Emitting Electrochemical Cells, Adv. Funct. Mater., 2011, 21, 1795.
- 35 S. van Reenen, R. A. J. Janssen and M. Kemerink, Doping dynamics in light-emitting electrochemical cells, Org. Electron., 2011, 12, 1746.
- Pertegás, N. M. Shavaleev, D. Tordera, E. Ortí, M. K. Nazeeruddin and H. J. Bolink, Host-guest blue lightemitting electrochemical cells, J. Mater. Chem. C, 2014, 2, 1605.
- 37 (a) S. E. Jang, K. S. Yook and J. Y. Lee, High power efficiency in simplified two layer blue phosphorescent organic light-emitting diodes, Org. Electron., 2010, 11, 1154; (b) J. Wang, X. Xu, Y. Tian, C. Yao, R. Liu and L. Li, A novel blue fluorescent polymer for solution-processed fluorescent-phosphorescent hybrid WOLEDs, J. Mater. Chem. C, 2015, 3, 2856.
- 38 CrystalClear-SM Expert v. 2.1, The Woodlands, Texas, USA, 2010-2014.
- 39 P. T. Beurskens, G. Beurskens, R. de Gelder, S. Garcia-Granda, R. O. Gould, R. Israel and J. M. M. Smits, DIRDIF-99. DIRDIF-99, University of Nijmegen, The Netherlands, 1999.
- 40 G. Sheldrick, Crystal structure refinement with SHELXL, Acta Crystallogr., Sect. C: Cryst. Struct. Commun., 2015, 71, 3.
- 41 CrystalStructure v 4.1, The Woodlands, Texas, USA, 2014.
- 42 A. M. Brouwer, Standards for photoluminescence quantum yield measurements in solution (IUPAC Technical Report)*, Pure Appl. Chem., 2011, 83, 2213.