

CrossMark
click for updatesCite this: *Chem. Sci.*, 2016, 7, 1276

Boronic acid–DMAPO cooperative catalysis for dehydrative condensation between carboxylic acids and amines†

Kazuaki Ishihara*^{ab} and Yanhui Lu^a

Arylboronic acid and 4-(*N,N*-dimethylamino)pyridine *N*-oxide (DMAPO) cooperatively catalyse the dehydrative condensation reaction between carboxylic acids and amines to give the corresponding amides under azeotropic reflux conditions. This cooperative use is much more effective than their individual use as catalysts, and chemoselectively promotes the amide condensation of (poly)conjugated carboxylic acids. The present method is practical and scalable, and has been applied to the synthesis of sitagliptin and a drug candidate.

Received 4th October 2015
Accepted 2nd November 2015

DOI: 10.1039/c5sc03761a

www.rsc.org/chemicalscience


Introduction

The catalytic dehydrative condensation reaction between carboxylic acids and amines is one of the most ideal methods for synthesizing the corresponding amides.¹ In 1996, Yamamoto *et al.* reported the first example of the dehydrative amide condensation reaction catalysed by *meta*- or *para*-electron-deficient group-substituted phenylboronic acids such as 3,4,5-trifluorophenylboronic acid (**1**) ($pK_a = 6.8$)^{2a} and 3,5-bis(trifluoromethyl)phenylboronic acid (**2**) ($pK_a = 7.2$)^{2b} under azeotropic reflux conditions (Scheme 1).³ These boronic acids are more acidic than phenylboronic acid ($pK_a = 8.8, 8.9$).²

In 2006, Whiting *et al.* reported that *ortho*-Brønsted base-substituted phenylboronic acids such as 2-(*N,N*-diisopropylaminomethyl)phenylboronic acid (**3**) were effective catalysts for the amide condensation of aromatic carboxylic acids under the same conditions as above.^{1a,4} In 2008 and 2012, Hall *et al.* reported that 2-iodophenylboronic acid (**4a**) and 2-iodo-5-methoxyphenylboronic acid (**4b**) were also effective catalysts for the amide condensation in the presence of drying agents (activated 4 Å molecular sieves) at lower temperature.⁵ The *o*-iodo group of **4a** and **4b** assists the catalysis of amide condensation as a weak base.⁶ In addition to these boronic acids, boric acid,^{7a,c} benzo[1,3,2]dioxaborol-2-ol,^{7b} methylboronic acid,^{7d} and some *o*-Brønsted base-substituted boronic acids⁸ have been reported to be useful as amidation catalysts. However, the substrate scope is still quite limited. For example,

harsh conditions (higher temperature, prolonged reaction time, excess amounts of substrates, increased amounts of catalysts, *etc.*) are required for sterically hindered α -branched carboxylic acids and conjugated carboxylic acids. In 2013, Whiting *et al.* discovered an interesting synergistic catalytic effect between *o*-tolylboronic acid (50 mol%) and *o*-nitrophenylboronic acid (50 mol%) in dipeptide synthesis.^{3f} To the best of our knowledge, this was the first example of two cooperative promoters for direct amidation.^{3f,9}

In the process catalysed by arylboronic acid, a mixed anhydride intermediate **5** is generated from the carboxylic acid and arylboronic acid under azeotropic reflux conditions or in the presence of drying agents in the first stage (Schemes 1 and 2). This is the first activation of the carboxylic acid. If a nucleophilic additive (Nu) reacts with **5** to generate a more active cationic intermediate **7** (ref. 10) (second activation) *via*




Scheme 1 Dehydrative condensation of carboxylic acids with amines catalysed by arylboronic acids, and representative examples of catalysts.

^aGraduate School of Engineering, Nagoya University, B2-3(611), Furo-cho, Chikusa, Nagoya 464-8603, Japan. E-mail: ishihara@cc.nagoya-u.ac.jp

^bJST, CREST, B2-3(611), Furo-cho, Chikusa, Nagoya 464-8603, Japan

† Electronic supplementary information (ESI) available: Experimental procedure and characterization data of new compounds are provided. CCDC 1429213. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc03761a





Scheme 2 Our proposal: the second activation of a mixed anhydride **5** with a nucleophilic additive (Nu) to generate a more active cationic intermediate **7**.

a tetrahedral intermediate **6**, the amide condensation may proceed more rapidly. However, if Nu preferentially coordinates as a Lewis base to the boron atom of **5**, a less active species **8** is generated and the amide condensation may be suppressed. Here we report that arylboronic acids and *N,N*-dimethylaminopyridine *N*-oxide (DMAPO) cooperatively promote the dehydrative condensation between various carboxylic acids and amines.

Results and discussion

First, the amide condensation reaction between 2-phenylbutyric acid and benzylamine was examined in the presence of 5 mol% each of boronic acid **2** and a nucleophilic additive under azeotropic reflux conditions in fluorobenzene (bp. 85 °C)^{3f} for 17 h (Table 1). Boronic acid **2** did not promote the reaction in the absence of additive under these conditions (entry 1). Tertiary amines such as *N,N*-diisopropylethylamine and 4-(*N,N*-dimethylamino)pyridine (DMAP)¹¹ were not effective as additives (entries 2 and 3). 4-Methoxypyridine *N*-oxide (MPO) was also

Table 1 Effect of additives on the dehydrative condensation between 2-phenylbutyric acid and benzylamine^a

Entry	Additive	Yield ^b (%)	Entry	Additive	Yield ^b (%)
1	None	<5	4	MPO	<5
2	<i>i</i> -Pr ₂ EtN	<5	5	DMAPO	99
3	DMAP	<5	6	PPYO	27

^a A solution of 2-phenylbutyric acid (0.5 mmol) and benzylamine (0.5 mmol) in fluorobenzene was heated in the presence of **2** (5 mol%) and additive (0 or 5 mol%) under azeotropic reflux conditions. ^b Isolated yield.

less active (entry 4). In contrast, a more nucleophilic but weak base, DMAPO,¹² was quite effective for the amide condensation (entry 5). However, a more nucleophilic additive such as 4-(pyrrolidin-1-yl)pyridine *N*-oxide (PPYO) was less effective than DMAPO (entry 6), perhaps because the strong nucleophilicity of PPYO might reduce the activity of **7**.

Next, the cooperative effects of several boronic acids (5 mol%) were compared in the condensation reaction between 2-phenylbutyric acid or benzoic acid and benzylamine in the presence of DMAPO (5 mol%) (Table 2). These less reactive carboxylic acids were not activated by the individual use of boronic acids under the same conditions. As expected, 2-DMAPO and **4b**-DMAPO efficiently activated 2-phenylbutyric acid (entries 1 and 3). Phenylboronic acid and **3** were almost inert, even in the presence of DMAPO (entries 2 and 4). Interestingly, 2-DMAPO was more effective than **4b**-DMAPO for the amide condensation of benzoic acid (entries 1 and 3). While Whiting's catalyst **3** was quite effective for the amide condensation of benzoic acid, the catalytic activity of **3** was suppressed in the presence of DMAPO (entry 4).¹³

To explore the substrate scope using the cooperative catalysts, 2-DMAPO, the amide condensation reactions of several less reactive α -branched carboxylic acids and arenecarboxylic acids were examined under azeotropic reflux conditions in fluorobenzene (bp. 85 °C) or toluene (bp. 110 °C). As shown in Table 3, in each example, the cooperative catalysts were much more effective than **2** alone, the results for which are shown in brackets. Notably, not only aliphatic primary amines but also sterically hindered aliphatic secondary amines, less nucleophilic anilines and alkoxyamines reacted with these carboxylic acids. In particular, 2-DMAPO was effective in the amidation of arenecarboxylic acids with sterically hindered amines, in comparison with **3** and **4b** (entries 9–14). This cooperative method is scalable to practical volumes: the catalytic loading of 2-DMAPO could be reduced to 2.5 mol% for the dehydrative condensation on an 80 mmol scale (entry 4).

Table 2 Cooperative effects of boronic acid–DMAPO on the dehydrative condensation reaction^a

Entry	ArB(OH) ₂	Yield ^{b,c} (%) of PhEtCHCONHBn	Yield ^{b,d} (%) of PhCONHBn
1	2	99 [<5]	97 [<5]
2	PhB(OH) ₂	<5 [<5]	<5 [<5]
3	4b	92 [<5]	20 [8]
4	3	7 [15]	80 [95]

^a 0.5 mmol of carboxylic acid and 0.5 mmol of benzylamine were used in the presence of 5 mol% of ArB(OH)₂ and 0 or 5 mol% of DMAPO. ^b The results when both catalysts were used are shown. For comparison, the results without DMAPO are shown in brackets. ^c Conditions: fluorobenzene (bp. 85 °C), 17 h. ^d Conditions: toluene (bp. 110 °C), 4 h.




Fig. 1 X-ray structure of **9z**.


the undesired complex **8** would be generated in an equilibrium mixture. Complex **8** might be converted to the more stable complex **9**, which is inert to the amide condensation. In fact, the generation of inert complex **9** was ascertained by ^{11}B and ^1H NMR analysis in the amidation of less-hindered carboxylic acids.¹⁵ Also, the chemical structure of the cyclic complex prepared from **2**, phthalic acid, and DMAPO was determined to be that of **9z** by X-ray diffraction analysis (Fig. 1).¹⁶

For sterically hindered carboxylic acids such as arene-carboxylic acids and α -branched carboxylic acids, the desired intermediate **7** was preferentially generated. Thus, *o*-non-substituted and *m*- or *p*-electron-deficient group-substituted phenylboronic acids such as **1** and **2** were more suitable. In contrast, for less sterically hindered α -nonbranched carboxylic acids, the undesired complex **9** was generated more easily. In addition, the strong Lewis acidity of **2** helped to stabilize **9** by the tight coordination of DMAPO to the boron centre. This is why **4b** and phenylboronic acid were slightly more effective than **2** for the condensation of α -nonbranched carboxylic acids. Not only Lewis acidity, but also the bulkiness of the *o*-substituent of the boronic acid might suppress the generation and stability of **9**. It is noted that the effect of DMAPO was not striking at ambient temperature. Heating was required to accelerate the equilibrium between **6** and **8**.

The utility of the cooperative catalysts was also demonstrated for the selective amide condensation of β -substituted acrylic acids to give the corresponding amides **10** (Table 6). The production of Michael adducts **11** was fairly minimal. In contrast, when boronic acids were used in the absence of DMAPO, the yield and selectivity of the reaction for **10** were moderate. Control experiments ascertained that **10** ($n = 1$) was selectively obtained from **13**,¹⁷ and **11** ($n = 1$) was not generated from **10** ($n = 1$) but **14**. Amide **10c** is known to be a potential antimitotic agent, especially for brain cancers (entry 6).¹⁸ The cooperative catalysts were effective for the selective amide condensation of not only β -substituted acrylic acids, but also polyconjugated carboxylic acids and but-2-ynoic acid (entries 12–17).

Table 6 Cooperative effects of boronic acid–DMAPO on the dehydrative condensation of conjugated carboxylic acids^a

Entry	ArB(OH) ₂	Time (h)	Product 10 or 12	
			Yield ^b (%)	Selectivity ^b (%)
1	2	12	10a , 87 [42]	93 [74] ^c
2	4b	10	10a , 82 [79]	94 [91] ^c
3	PhB(OH) ₂	12	10a , 15 [4]	95 [78] ^c
4	2	15	10b , 77 [17]	89 [30] ^c
5	4b	16	10b , 65 [29]	89 [47] ^c
6 ^{d,e}	2	15	10c , 96 [49]	96 [64] ^c
7	2	38	10d , 90 [78]	>95 [>95] ^c
8	2	8	10e , 68 [19]	82 [37] ^c
9	4b	7	10e , 72 [62]	84 [78] ^c
10 ^d	PhB(OH) ₂	19	10e , 81 [56]	92 [76] ^c
11 ^d	PhB(OH) ₂	22	10f , 73 [32]	96 [55] ^c
12 ^d	2	16	10g , 98 [45]	>99 [45] ^f
13 ^d	2	16	10h , 97 [69]	>99 [79] ^f
14 ^d	2	16	10i , 73 [<5]	97 [<5] ^f
15 ^d	2	23	10j , 92 [–]	>99 [50] ^f
16 ^d	2	24	10k , 99 [25]	>99 [56] ^f
17 ^d	2	14	12 , 98 [46]	>99 [82] ^f



^a Unless noted otherwise, 0.5 mmol of carboxylic acid and 0.5 mmol of amine were used in the presence of 5 mol% of ArB(OH)₂ and 0 or 5 mol% of DMAPO. ^b The results when both catalysts were used are shown. For comparison, the results without DMAPO are shown in brackets. ^c β -Aminoamide **11** ($n = 1$) was obtained as the sole minor product. ^d 10 mol% of each of the catalysts was used. ^e Toluene was used as a solvent. ^f Several minor products including **11** were obtained.

Conclusions

In conclusion, this new cooperative catalytic system is quite effective for the amidation reaction of less reactive carboxylic acids, such as sterically hindered α -branched carboxylic acids and arene-carboxylic acids, and the chemoselective amidation reaction of conjugated carboxylic acids. Based on the NMR



spectra and X-ray diffraction analysis of inert species **9**, a preliminary mechanism was proposed. Further mechanistic studies are in progress. We believe that these findings will trigger the further development of high-performance amidation catalysts.

Acknowledgements

Financial support for this project was partially provided by the Program for Leading Graduate Schools: IGER Program in Green Natural Sciences (MEXT), JSPS KAKENHI Grant Numbers 26620079, 15K13692, and 15H05755.

Notes and references

- For selected reviews, see: (a) I. Georgiou, G. Ilyashenko and A. Whiting, *Acc. Chem. Res.*, 2009, **42**, 756; (b) K. Ishihara, *Tetrahedron*, 2009, **65**, 1085; (c) H. Charville, D. Jackson, G. Hodges and A. Whiting, *Chem. Commun.*, 2010, **46**, 1813; (d) C. Grosjean, J. Parker, C. Thirsk and A. R. Wright, *Org. Process Res. Dev.*, 2012, **16**, 781; (e) E. Dimitrijević and M. S. Taylor, *ACS Catal.*, 2013, **3**, 945; (f) H. Zheng and D. G. Hall, *Aldrichimica Acta*, 2014, **47**, 41; (g) H. Lundberg, F. Tinnis, N. Selander and H. Adolfsson, *Chem. Soc. Rev.*, 2014, **43**, 2714; (h) K. Ishihara, in *Synthesis and Application of Organoboron Compounds, Topics in Organometallic Chemistry*, ed. E. Fernández and A. Whiting, Springer, Heidelberg, 2015, vol. 49, pp. 243–270.
- (a) J. Yan, G. Springsteen, S. Deeter and B. Wang, *Tetrahedron*, 2004, **60**, 11205; (b) S. Kheirjou, A. Abedin and A. Fattahi, *Comput. Theor. Chem.*, 2012, **1000**, 1.
- For the electron-deficient arylboronic acid-catalysed condensation of carboxylic acids, see: (a) K. Ishihara, S. Ohara and H. Yamamoto, *J. Org. Chem.*, 1996, **61**, 4196; (b) K. Ishihara, S. Ohara and H. Yamamoto, *Macromolecules*, 2000, **33**, 3511; (c) K. Ishihara, S. Kondo and H. Yamamoto, *Synlett*, 2001, 1371; (d) K. Ishihara, S. Ohara and H. Yamamoto, *Org. Synth.*, 2002, **79**, 176; (e) T. Maki, K. Ishihara and H. Yamamoto, *Synlett*, 2004, 1355; (f) S. Liu, Y. Yang, X. Liu, F. K. Ferdousi, A. S. Batsanov and A. Whiting, *Eur. J. Org. Chem.*, 2013, 5692.
- (a) K. Arnold, A. S. Batsanov, B. Davies and A. Whiting, *Green Chem.*, 2008, **10**, 124; (b) I. Georgiou, G. Ilyashenko and A. Whiting, *Acc. Chem. Res.*, 2009, **42**, 756.
- (a) R. M. Al-Zoubi, O. Marion and D. G. Hall, *Angew. Chem., Int. Ed.*, 2008, **47**, 2876; (b) R. M. Al-Zoubi and D. G. Hall, *Org. Lett.*, 2010, **12**, 2480; (c) N. Gernigon, R. M. Al-Zoubi and D. G. Hall, *J. Org. Chem.*, 2012, **77**, 8386; (d) N. Gernigon, H. Zheng and D. G. Hall, *Tetrahedron Lett.*, 2013, **54**, 4475.
- (a) T. Marcelli, *Angew. Chem., Int. Ed.*, 2010, **49**, 6840; (b) C. Wang, H.-Z. Yu, Y. Fu and Q.-X. Guo, *Org. Biomol. Chem.*, 2013, **11**, 2140.
- (a) P. Tang, *Org. Synth.*, 2005, **81**, 262; (b) T. Maki, K. Ishihara and H. Yamamoto, *Org. Lett.*, 2006, **8**, 1431; (c) R. K. Mylavarapu, K. Gcm, N. Kolla, R. Veeramalla, P. Koilkonda, A. Bhattacharya and R. Bandichhor, *Org. Process Res. Dev.*, 2007, **11**, 1065; (d) R. Yamashita, A. Sakakura and K. Ishihara, *Org. Lett.*, 2013, **15**, 3654.
- For other examples of the *o*-Brønsted base-substituted phenylboronic acid-catalysed condensation of carboxylic acids, see: (a) A. Sakakura, T. Ohkubo, R. Yamashita, M. Akakura and K. Ishihara, *Org. Lett.*, 2011, **13**, 892; (b) A. Sakakura, R. Yamashita, M. Akakura and K. Ishihara, *Aust. J. Chem.*, 2011, **64**, 1458; (c) T. M. El Dine, W. Erb, Y. Berhault, J. Rouden and J. Blanchet, *J. Org. Chem.*, 2015, **80**, 4532; (d) E. K. W. Tam, Rita, L. Y. Liu and A. Chen, *Eur. J. Org. Chem.*, 2015, 1100.
- For recent examples of boron catalysts for reactions involving carboxylic acids, see: (a) T. Azuma, A. Murata, Y. Kobayashi, T. Inokuma and Y. Takemoto, *Org. Lett.*, 2014, **16**, 4256; (b) Y. Morita, T. Yamamoto, H. Nagai, Y. Shimizu and M. Kanai, *J. Am. Chem. Soc.*, 2015, **137**, 7075.
- The unstable counter anion of **7**, ArB(OH)O^- , may be immediately converted to $[\text{ArBL}_3]^-$ in the presence of carboxylic acids and amines.
- (a) Review: N. de Rycke, F. Couty and O. R. P. David, *Chem.–Eur. J.*, 2011, **17**, 12852; (b) A. Sakakura, K. Kawajiri, T. Ohkubo and K. Ishihara, *J. Am. Chem. Soc.*, 2007, **129**, 14775; (c) R. Tandon, T. Unzer, T. A. Nigst, N. de Rycke, P. Mayer, B. Wendt, O. P. R. David and H. Zipse, *Chem.–Eur. J.*, 2013, **19**, 6435.
- I. Shina, H. Ushiyama, Y. Yamada, Y. Kawakita and K. Nakata, *Chem.–Asian J.*, 2008, **3**, 454.
- The bulkiness of the *o*-substituent of **3** may suppress the second activation from **6** to **7** with DMAPO. Alternatively, DMAPO may suppress the Brønsted base role of the *o*-substituent of **3**.
- D. Kim, L. Wang, M. Beconi, G. J. Eiermann, M. H. Fisher, H. He, G. J. Hickey, J. E. Kowalchick, B. Leiting, K. Lyons, F. Marsilio, M. E. McCann, R. A. Patel, A. Petrov, G. Scapin, S. B. Patel, R. S. Roy, J. K. Wu, M. J. Wyvrat, B. B. Zhang, L. Zhu, N. A. Thornberry and A. E. Weber, *J. Med. Chem.*, 2005, **48**, 141.
- The reactivity and stability of **9** were investigated. See the ESI for details.†
- 9z** was produced with *N*-benzylphthalimide through the dehydrative condensation reaction between phthalic acid and benzylamine in the presence of **2** (5 mol%) and DMAPO (5 mol%) in fluorobenzene under azeotropic reflux conditions. In this reaction, the cooperative use of **2** and DMAPO was less effective than the single use of **2**. See the ESI (Table S2†) for details.
- Intermediate **13** was prepared from DMAPO and the corresponding acryl chloride. The reaction of **13** with amine gave **10** as the sole product. See the ESI for details.†
- J. B. Leslie, R. C. Holaday, T. Nguyen and P. J. Hergenrother, *J. Med. Chem.*, 2010, **53**, 3964.

