


 Cite this: *RSC Adv.*, 2025, 15, 33816

# Structural modifications of toxyloxanthone C and macluraxanthone isolated from *Maclura cochinchinensis*: cytotoxicity, antibacterial activity, and *in silico* studies

 Chaiwat Linphosan,<sup>a</sup> Waranya Klangswad,<sup>b</sup> Jantana Yahuafai,<sup>c</sup> Jinda Jandaruang,<sup>d</sup> Trinop Promgool,<sup>e</sup> Siripit Pitchuanom,<sup>ib</sup> Jakkapat Paluka,<sup>f</sup> Sophon Boonlue,<sup>g</sup> Kitisak Poopasit,<sup>b</sup> Kwanjai Kanokmedhakul<sup>ib</sup> and Oue-artorn Limtragool<sup>ib</sup>\*<sup>a</sup>

Toxyloxanthone C and macluraxanthone, isolated from the roots of *Maclura cochinchinensis*, have been reported to exhibit promising cytotoxic and antibacterial activities. Accordingly, thirteen xanthone derivatives were synthesized from these two parent xanthones by simple acylation, alkylation, sulfonylation, and bromination reactions. All derivatives were evaluated for their cytotoxicity against three cancer cell lines, HelaS3, A549, and HepG2, and their antibacterial activity against four Gram-positive bacterial strains, namely methicillin-resistant *Staphylococcus aureus*, *S. aureus*, *Bacillus subtilis*, and *Bacillus cereus*. 5,6-Diacetoxytoxyloxanthone C (**1a**) displayed cytotoxicity against three cancer cells with IC<sub>50</sub> values ranging from 12.20 to 22.61 μM. Additionally, **1a** demonstrated potent cytotoxicity against A549 cells with IC<sub>50</sub> = 12.20 μM, without showing cytotoxicity toward Vero cells. Moreover, 1,5,6-tripentanoyloxytoxyloxanthone C (**1b**) and 4-bromotoxyloxanthone C (**1f**) exhibited significant cytotoxicity against A549 cells with IC<sub>50</sub> values of 5.77 and 7.52 μM, respectively. In addition, **1a** and **1f** showed potent antibacterial activity against all bacteria tested with the same MIC value of 4 μg mL<sup>-1</sup>, which was stronger than the parent xanthone **1**. Molecular docking studies revealed that **1a**, **1b**, and **1f** interacted with CDK2 through a competitive inhibition mechanism. Additionally, the binding conformations of **1a** and **1f** within the active sites of key enzymes involved in bacterial cell wall synthesis were similar to that of the tetracycline drug. The *in silico* physicochemical investigation indicated that **1a** and **1f** exhibited a favorable drug-likeness. Based on this finding, **1a** represents a promising lead candidate for further study as an anticancer and an antibacterial agent.

 Received 7th August 2025  
 Accepted 29th August 2025

DOI: 10.1039/d5ra05758b

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## 1 Introduction

*Maclura cochinchinensis* (Lour.) Corner is a shrub belonging to the Moraceae family and is widely distributed across various regions of Asia, including China, Japan, Korea, Taiwan, India,

Vietnam, Laos, and Thailand. Traditionally, the heartwood of this plant has been utilized in the treatment of chronic fever, diarrhea, and skin infections.<sup>1,2</sup> Additionally, its leaves have been used for wound healing, while the roots have been employed in traditional medicine to treat bruising, boils, scabies, rheumatism, blood stasis, dysmenorrhea, and contusions.<sup>2,3</sup> Xanthones and flavonoids, the major bioactive constituents isolated from *M. cochinchinensis*,<sup>2-7</sup> have demonstrated a variety of biological activities, including antioxidant,<sup>1</sup> antibacterial,<sup>4,7</sup> anticancer,<sup>1,3,4</sup> and anti-inflammatory.<sup>2,6</sup>

Several xanthone derivatives have been reported for structural modifications to enhance their biological activity. Three xanthones, α-mangostin, β-mangostin, and γ-mangostin, were partially modified under acidic conditions and subsequently evaluated for their antibacterial activities. Several of their analogues demonstrated improved pharmacokinetic properties. However, all modified derivatives showed weaker antibacterial activities against methicillin-resistant *Staphylococcus aureus* (MRSA), *Bacillus subtilis*, and *Pseudomonas aeruginosa*

<sup>a</sup>Multidisciplinary Research Unit of Pure and Applied Chemistry, Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Mahasarakham University, Maha Sarakham, Thailand. E-mail: oueartorn.l@msu.ac.th; Fax: +66-43 754246; Tel: +66-43 754246

<sup>b</sup>Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Khon Kaen University, Khon Kaen, Thailand

<sup>c</sup>Clinical Research Section, Division of Research and Academic Support, National Cancer Institute, Bangkok, Thailand

<sup>d</sup>Innovation in Chemistry for Community Research Unit, Chemistry Program, Faculty of Science and Technology, Sakon Nakhon Rajabhat University, Sakon Nakhon, Thailand

<sup>e</sup>Department of Industrial Chemistry, Faculty of Applied Science, King Mongkut's University of Technology North Bangkok, Bangkok, Thailand

<sup>f</sup>Research Administration Division, Khon Kaen University, Thailand

<sup>g</sup>Department of Microbiology, Faculty of Science, Khon Kaen University, Khon Kaen, Thailand



compared to their parent compounds.<sup>8</sup> Furthermore,  $\alpha$ -mangostin was partially synthesized by cationic modification of the free 3-OH and 6-OH groups with amine moieties, followed by antibacterial evaluation. Some of its derivatives exhibited enhanced antibacterial potency against MRSA and *S. aureus*.<sup>9</sup> Ananixanthone isolated from *Calophyllum teysmannii* stem bark, underwent structural modifications through acetylation, methylation, and benzylation at the 1-OH and 5-OH positions. The parent compound exhibited more potent cytotoxicity against SNU-1 (stomach cancer) and K562 (leukemia) cell lines. Among its derivatives, only 5-methoxyananixanthone exhibited superior cytotoxicity against LS174T (colon cancer) cells.<sup>10</sup>

Besides *M. cochinchinensis*, toxyloxanthone C (**1**) has been isolated from various plants such as *Cudrania*,<sup>11–15</sup> *Cratoxylum*,<sup>16</sup> and *Rheedia*<sup>17</sup> species. It has been reported to exhibit antifungal,<sup>12,18</sup> antibacterial,<sup>14,19</sup> and cytotoxicity against HCT-116, SMMC-7721, SGC-7901, and BGC-823 cancer cell lines.<sup>13,18</sup> Whereas, macluraxanthone (**2**) has been reported in the *Garcinia*,<sup>20–22</sup> *Mesua*,<sup>23</sup> and *Calophyllum*<sup>24</sup> species. It has been shown cytotoxicity against several cancer cell lines such as HeLa, A549, PC-3, HT-29, WPMY-1, Hep G2, NCI-H23, and KB.<sup>20,21,24,25</sup> Additionally, **2** exhibited *anti*-HIV,<sup>20</sup> antibacterial,<sup>4</sup> and antimalarial.<sup>26</sup> From our previous report,<sup>4</sup> xanthenes **1** and **2** were isolated from *M. cochinchinensis*, they showed potent cytotoxicity toward three cancer cell lines, HeLaS3, A549, and HepG2 and also exhibited antibacterial activity against four Gram-positive bacteria MRSA, *S. aureus*, *B. subtilis*, and *Bacillus cereus*. Herein, further study of these bioactive xanthenes is reported. The chemical structural modifications of parent xanthenes **1** and **2** using simple organic reactions yielded thirteen xanthone derivatives. All compounds were evaluated for their cytotoxicity and antibacterial activity. Additionally, molecular docking simulations and drug-like property are also presented.

## 2 Result and discussion


### 2.1 Chemistry

From our previous reported,<sup>4</sup> toxyloxanthone C (**1**) and macluraxanthone (**2**) were the main compounds isolated from the roots of *M. cochinchinensis*. Xanthone **1** exhibited cytotoxicity against the HeLaS3, A549, HepG2, and Vero cell lines with IC<sub>50</sub> values of 13.55 ± 1.11, 21.78 ± 4.89, 20.81 ± 2.57, and 8.52 ± 0.64  $\mu$ M, respectively. While xanthone **2** showed cytotoxicity against the HeLaS3, A549, HepG2, and Vero cell lines with IC<sub>50</sub> values of 1.59 ± 0.12, 6.46 ± 0.98, 5.26 ± 0.41, and 4.29 ± 0.60, respectively. In addition, **1** displayed antibacterial activity against MRSA, *S. aureus*, *B. subtilis*, and *B. cereus* with MIC values of 128  $\mu$ g mL<sup>-1</sup> for all strains, and **2** exhibited activities toward MRSA, *B. subtilis*, and *B. cereus* with MIC values of 16, 64, and 64  $\mu$ g mL<sup>-1</sup>, respectively.<sup>4</sup> To improve their bioactivities, structural modifications of **1** and **2** were performed. Thirteen xanthone derivatives were successfully synthesized using simple organic reactions, including acylation, alkylation, sulfonation, and bromination, as shown in Schemes 1 and 2.


According to literature,<sup>27</sup> acetyl substitution at 6-OH of  $\alpha$ -mangostin exhibited significant antibacterial activity against

MRSA, which was better than the standard drug penicillin. Furthermore, the acetyl substitutions at 1-OH and 6-OH of  $\alpha$ -mangostin, when combined with penicillin, exhibited greater antibacterial efficacy against MRSA compared to penicillin alone. Therefore, acylation reactions were applied to the parent xanthenes **1** and **2** to investigate the antibacterial activity. Acetylation of **1** and **2** using acetic anhydride in pyridine afforded derivatives **1a** (41%) and **2a** (65%). The <sup>1</sup>H NMR data of **1a** showed two singlet signals of two methyl groups at  $\delta_{\text{H}}$  2.43 and 2.35, while the <sup>13</sup>C NMR data of **1a** showed carbonyl signals at  $\delta_{\text{C}}$  167.7 and 167.4, corresponding to acetoxy groups. Similarly, the <sup>1</sup>H NMR data signals of **2a** at  $\delta_{\text{H}}$  2.39 and 2.34, together with its <sup>13</sup>C NMR signals at  $\delta_{\text{C}}$  167.7 and 167.6, confirmed the presence of two acetoxy groups. Additionally, the <sup>1</sup>H NMR data of **1a** and **2a** exhibited intramolecular H-bonding signals at  $\delta_{\text{H}}$  12.84 and 13.40, respectively. Treatments of **1** and **2** with pentanoyl chloride in the presence of pyridine provided xanthone derivatives **1b** (11%) and **2b** (19%), respectively. Notably, the <sup>1</sup>H NMR spectra of **1b** and **2b** displayed no intramolecular H-bonding signal around  $\delta_{\text{H}}$  12.8 and 13.4, respectively. In addition, their <sup>13</sup>C NMR spectra revealed three carbonyl signals at  $\delta_{\text{C}}$  172.6, 170.6, and 170.1 for **1b** and at  $\delta_{\text{C}}$  172.1, 170.6, and 170.5 for **2b**. In the acylation reactions, the use of a more electrophilic acid chloride resulted in acyl substitution at the 1-OH, 5-OH, and 6-OH positions.<sup>27</sup> Structural modifications were further carried out by treated **1** and **2** with methanesulfonyl chloride in pyridine, afforded mesylated derivatives **1c** (77%) and **2c** (88%). The <sup>1</sup>H NMR spectra of **1c** and **2c** showed singlet signals of two methyl groups at  $\delta_{\text{H}}$  3.49 and 3.38 for **1c**, as well as 3.43 and 3.33 for **2c**, corresponding to mesyl groups. According to a previous report,<sup>10</sup> methylation of 5-OH group in ananixanthone resulted in enhanced cytotoxicity against the LS174T cell line compared to its parent compound. Therefore, alkylation reactions were carried out on xanthenes **1** and **2** to further investigate their cytotoxic potential. Methylation of **1** using methyl iodide in the presence of sodium carbonate produced the dimethylated product **1d** (94%), whereas methylation of **2** yielded both a dimethylated product **2d** (45%) and a monomethylated product **2e** (14%). The <sup>1</sup>H NMR data of **1d** showed two methoxy signals at  $\delta_{\text{H}}$  4.00 and 3.99, whereas **2d** displayed two methoxy signals at  $\delta_{\text{H}}$  4.00 and 3.92. The <sup>1</sup>H NMR data of **2e** exhibited a singlet signal of a methoxy group at  $\delta_{\text{H}}$  4.01, while the HMBC spectrum revealed a correlation between these methoxy protons to C-5, confirming the substitution at the C-5 position. Ethylation of **1** using ethyl iodide in the presence of sodium carbonate provided diethylated product **1e** (86%), while **2** produced diethylated **2f** (70%) and monoethylated **2g** (27%) derivatives. In the alkylation reactions, **2** reacted with alkyl halide to yield both mono- and dialkylated products, which might due to the steric hindrance of the isoprenyl group at the C-4 position. Lastly, brominated xanthenes was reported to demonstrate notable antibacterial activities against *S. aureus* and MRSA.<sup>28</sup> Consequently, the bromination reaction was applied to the parent xanthenes **1** and **2** to analyze their antibacterial efficacy. Bromination of **1** using *N*-bromosuccinimide (NBS) in aqueous acetonitrile at room temperature afforded 4-bromoxanthone **1f** (36%). The <sup>1</sup>H





Scheme 1 Derivatives of toxyloxanthone C (1).



Scheme 2 Derivatives of macluraxanthone (2).



NMR data of **1f** showed the absence of the aromatic singlet signal around  $\delta_{\text{H}}$  6.4 corresponding to H-4, while the  $^{13}\text{C}$  NMR data of **1f** showed a characteristic signal for the C-4 brominated carbon at  $\delta_{\text{C}}$  81.9, supporting the bromination position. In contrast, the reaction of **2** under the same bromination conditions did not yield any brominated product, due to the less nucleophilicity of the aromatic ring B.

## 2.2 Biological activity

Thirteen xanthone derivatives **1a–f** and **2a–g** were evaluated for their cytotoxicity against human cervical cancer cells (HelaS3), human lung cancer cells (A549), human liver cancer cells (HepG2), and African green monkey kidney cells (Vero), and the result is summarized in Table 1. The acetyl derivative **1a** displayed cytotoxicity against HelaS3, A549, and HepG2 cell lines, with  $\text{IC}_{50}$  values of  $22.61 \pm 2.02$ ,  $12.20 \pm 1.08$ , and  $21.10 \pm 1.18$   $\mu\text{M}$ , respectively. Notably, **1a** exhibited twice cytotoxic potency against A549 cells compared to its parent xanthone **1**. Moreover, **1a** showed no cytotoxicity against normal cells ( $\text{IC}_{50} > 100$   $\mu\text{M}$ ). Compound **1b** displayed strong toward A549 with  $\text{IC}_{50}$  values of  $5.77 \pm 2.14$   $\mu\text{M}$ , approximately fourfold higher than its parent **1**. However, **1b** also exhibits strong cytotoxicity to normal cells. In addition, **1b** demonstrated cytotoxicity against HepG2 with  $\text{IC}_{50}$  values of  $17.15 \pm 4.50$   $\mu\text{M}$  comparable to **1**. The acetyl derivative **2a** exhibited moderate cytotoxicity against three cancer cell lines with  $\text{IC}_{50}$  values in the range of  $15.66 \pm 1.17$  to  $25.55 \pm 4.77$   $\mu\text{M}$ . The pentanoyl derivative **2b** showed weak cytotoxicity against HelaS3 ( $\text{IC}_{50} = 55.31 \pm 2.18$   $\mu\text{M}$ ) and A549 ( $\text{IC}_{50} = 41.84 \pm 1.91$   $\mu\text{M}$ ) cancer cells. This result indicated that the acyl derivatives of **2** decrease the cytotoxicity against the cancer cell lines tested. The mesylated derivatives, **1c** and **2c**, displayed weak cytotoxicity against HelaS3 cell line, with  $\text{IC}_{50}$  values of

$65.33 \pm 3.92$  and  $84.26 \pm 7.86$   $\mu\text{M}$ , respectively. In contrast, the dimethylated products **1d** and **2d**, as well as the ethylated derivatives **2f** and **2g**, showed no cytotoxicity for all cancer cells tested. This suggested that the alkylation derivatives at the 5-OH and 6-OH positions of **1** and **2** reduce their cytotoxic potency. The bromoxanthone derivative **1f** demonstrated moderate cytotoxicity to HelaS3 and HepG2 cell lines, with  $\text{IC}_{50}$  values of  $17.43 \pm 1.75$  and  $19.42 \pm 1.52$   $\mu\text{M}$ , respectively. In addition, **1f** exhibited strong cytotoxicity against A549 ( $\text{IC}_{50} = 7.52 \pm 0.36$   $\mu\text{M}$ ), which is threefold higher than that of its parent **1**. However, **1f** showed cytotoxic toward normal cells. It was found that others derivative **2** in this study eliminated the cytotoxicity of all cancer cell lines tested. In conclusion, acylation and bromination of **1** enhance the cytotoxicity toward A549 cell line, but only acetyl **1a** has the potential to be developed as an anticancer drug targeting A549 cells, as it has anticancer activity without causing toxicity to normal cells ( $\text{IC}_{50} > 100$   $\mu\text{M}$ ).

All xanthone derivatives **1a–f** and **2a–g** were further evaluated for antibacterial activity against four Gram-positive bacteria, including MRSA, *S. aureus*, *B. subtilis*, and *B. cereus*, using the microdilution method (Table 2). Compounds **1a** and **1f** exhibited significant antibacterial efficacy against all bacterial strains with the same MIC value at  $4$   $\mu\text{g mL}^{-1}$ , representing 32-fold improvement in potency compared to their parent **1**. Similarly, **1b** also showed notable activity with MIC values ranging from 8 to  $16$   $\mu\text{g mL}^{-1}$ , indicated their increasing potency activity compared to its parent **1**. This suggested that acylation and bromination of **1** enhanced antibacterial activity. Compounds **1c**, **1d**, and **1e** showed no antibacterial activity against all bacterial strains. It should be noted that sulfonylation and alkylation at the 5-OH and 6-OH positions decreased the antibacterial activity. In case of derivatives of **2**, only the acetyl derivative **2a** exhibited antibacterial activity against all tested

Table 1 *In vitro* cytotoxicity of toxylloxanthone C and macluraxanthone derivatives<sup>a</sup>

Compound	Cytotoxicity ( $\text{IC}_{50}$ , $\mu\text{M}$ )			
	HelaS3	A549	HepG2	Vero
<b>1a</b>	$22.61 \pm 2.02$	$12.20 \pm 1.08$	$21.10 \pm 1.18$	>100
<b>1b</b>	$20.46 \pm 4.92$	$5.77 \pm 2.14$	$17.15 \pm 4.50$	$5.59 \pm 0.36$
<b>1c</b>	$65.33 \pm 3.92$	>100	>100	>100
<b>1d</b>	>100	>100	>100	>100
<b>1e</b>	$50.39 \pm 8.45$	>100	>100	>100
<b>1f</b>	$17.43 \pm 1.75$	$7.52 \pm 0.36$	$19.42 \pm 1.52$	$9.52 \pm 1.02$
<b>2a</b>	$25.55 \pm 4.77$	$15.66 \pm 1.17$	$24.54 \pm 2.98$	$10.10 \pm 1.67$
<b>2b</b>	$55.31 \pm 2.18$	$41.84 \pm 1.91$	>100	$35.95 \pm 2.32$
<b>2c</b>	$84.26 \pm 7.86$	$48.83 \pm 4.73$	>100	>100
<b>2d</b>	>100	>100	>100	>100
<b>2e</b>	>100	$90.46 \pm 6.23$	>100	>100
<b>2f</b>	>100	>100	>100	>100
<b>2g</b>	>100	>100	>100	>100
<b>1</b>	$13.55 \pm 1.11$	$21.78 \pm 4.89$	$20.81 \pm 2.57$	$8.52 \pm 0.64$
<b>2</b>	$1.59 \pm 0.12$	$6.46 \pm 0.98$	$5.26 \pm 0.41$	$4.29 \pm 0.60$
Doxorubicin	$0.29 \pm 0.02$	$0.48 \pm 0.036$	$0.37 \pm 0.02$	$1.95 \pm 0.13$

<sup>a</sup> HelaS3 = human cervical carcinoma, A549 = human lung carcinoma, HepG2 = human hepatocellular carcinoma, Vero = African green monkey kidney.

Table 2 Antibacterial activity of toxylloxanthone C and macluraxanthone derivatives

Compound	MIC ( $\mu\text{g mL}^{-1}$ )			
	MRSA <sup>a</sup>	<i>S. aureus</i> <sup>b</sup>	<i>B. subtilis</i> <sup>c</sup>	<i>B. cereus</i> <sup>d</sup>
<b>1a</b>	4	4	4	4
<b>1b</b>	8	8	8	16
<b>1c</b>	>128	>128	>128	>128
<b>1d</b>	>128	>128	>128	>128
<b>1e</b>	>128	>128	>128	>128
<b>1f</b>	4	4	4	4
<b>2a</b>	8	16	32	16
<b>2b</b>	>128	>128	>128	>128
<b>2c</b>	>128	>128	>128	>128
<b>2d</b>	>128	>128	>128	>128
<b>2e</b>	>128	>128	>128	>128
<b>2f</b>	>128	>128	>128	>128
<b>2g</b>	>128	>128	>128	>128
<b>1</b>	128	128	128	128
<b>2</b>	16	>128	64	64
Vancomycin	2	2	2	1

<sup>a</sup> Methicillin resistant *Staphylococcus aureus* <sup>b</sup> *S. aureus* ATCC 25923.


<sup>c</sup> *Bacillus subtilis* ATCC 6633. <sup>d</sup> *Bacillus cereus* ATCC 11778.





**Table 3** Summary of the binding interactions and energies of active xanthone with CDK2, PBP2a of MRSA, and SCWP *O*-acetyltransferase of *B. cereus*

Compound	Binding energy (kcal mol <sup>-1</sup> )	Protein-ligand interactions		
		Hydrogen bond	Hydrophobic	Electrostatic
<b>Interactions of xanthone with CDK2</b>				
<b>1</b>	-8.62	Glu81, Leu83, Asp86	Ile10, Ala31, Phe82, Lys89, Leu134, Ala144	—
<b>1a</b>	-9.43	Lue83, Asp86, Asp145	Ile10, Ala31, His84, Lys89, Leu134, Ala144	—
<b>1b</b>	-9.78	Asp86, Lys89	Ile10, Val18, Ala31, Phe80, Phe82, Leu134, Ala144	—
<b>1f</b>	-8.52	Leu83, His84, Gln131	Ile10, Val18, Ala31, Phe80, Phe82, Ala144, Lue134	—
RC-3-89	-10.51	Glu81, Leu83, Asp86, Lys89	Ile10, Val18, Ala31, Phe82, Leu134, Ala144, Asp145	—
<b>Interactions of xanthone with PBP2a of MRSA</b>				
<b>1a</b>	-8.02	Ser403, Glu602, Asn464, Met641	Tyr446, His583, Ala642	—
<b>1f</b>	-7.29	Ser403, Ser462, Ser598	Asn464, Tyr519, Met641	—
Tetracycline	-8.35	Ser403, Tyr444, Try446, Ser462, Asn464, Ser598	Gln521, Ala646, Met641	—
<b>Interactions of xanthone with <i>B. subtilis</i> LCP enzyme</b>				
<b>1a</b>	-7.55	Thr197, Arg210, Gln214	Asp75, Asp85, Arg106, Arg210	Asp75, Asp85
<b>1f</b>	-7.32	Gln211	Pro105, Arg106, Asp207, Phe208, Leu273, Tyr286	Asp107, Asp207
Tetracycline	-5.86	Asp75, Arg210, Gln214	Asp75, Asp85, Arg198, Arg210	Arg198
<b>Interactions of xanthone with SCWP <i>O</i>-acetyltransferase of <i>B. cereus</i></b>				
<b>1a</b>	-7.18	Lys103, Asn104, Asn251, Arg359	His201, His202, Ser249, Phe250	His201
<b>1f</b>	-6.97	Lys103, Asn251, Ser385	His201	Asp92, His201, Arg359
Tetracycline	-5.75	Lys103, Asn104, Asn251, Ser385	His201, His202, Asn386	Arg359

**Fig. 2** (a) Docking poses of **1** (green), **1a** (purple), **1b** (yellow), and **1f** (pink) onto the active site of CDK2 and (b) superimposed conformation of co-crystallized ligand (green) and **1a** (purple) onto the active site of CDK2.

bacterial cell wall, resulting in the enhanced structural integrity of cell wall.<sup>37</sup> Therefore, the molecular docking of these key enzymes involved in cell wall synthesis of bacterial pathogens was performed to elucidate their potential for treating bacterial infections. Tetracycline was investigated as a control for all proteins and the results are summarized in Table 3. The active site of the PBP2a of MRSA consists of the amino acid residues Ser403, Lys406, Tyr446, Ser462, Asn464, Ser598, Gly599, Thr600, and Met641.<sup>34</sup> Tetracycline exhibited the lowest binding energy of  $-8.35$  kcal mol<sup>-1</sup> and formed hydrogen bonds with key amino acid residues Ser403, Tyr446, Ser462, Asn464, and Ser598 within the PBP2a active site of the transpeptidase domain (Fig. 3). Compounds **1a** and **1f** demonstrated the binding energy

of  $-8.02$  and  $-7.29$  kcal mol<sup>-1</sup>, respectively, within the PBP2a active site. Compound **1a** established hydrogen bonds with Ser403, Asn464, and Met641 and formed pi-pi stacking with Tyr446. Significantly, the carbonyl of acetyl groups formed hydrogen bonds with Glu602 and Asn464, indicating that the acetyl groups may play an important role in antibacterial against MRSA. Compound **1f** established hydrogen bonds with Ser403, Ser462, and Ser598. Moreover, the superimposed conformation of **1f** in the PBP2a active site was similar to those of tetracycline, as shown in Fig. 6a. According to the docking results of *B. subtilis*, the amino acids Asp75, Arg83, Asp85, Arg106, Asp107, Lys120, Phe156, Arg198, Arg200, Asp207, and Arg210 are considered as the active site of *B. subtilis* LCP






Fig. 3 The binding interactions of tetracycline, **1a**, and **1f** with the PBP2a of MRSA.




Fig. 4 The binding interactions of tetracycline, **1a**, and **1f** with *B. subtilis* LCP enzyme.

enzyme.<sup>35</sup> Tetracycline showed a binding energy of  $-5.86$  kcal mol<sup>-1</sup> and formed hydrogen bonds with Asp75, Asp207, Arg210, and Gln214 (Fig. 4). Compound **1a** demonstrated the lowest binding energy of  $-7.55$  kcal mol<sup>-1</sup>. Its xanthone core structure formed electrostatic interactions with key amino residues Asp75, Asp85, and Arg210. The carbonyl group at C-5 formed three hydrogen bonds with Thr197, Arg210, and Gln214 amino acids. The docking poses of **1a** exhibited similar orientation to that of tetracycline in the active site (Fig. 6b). Compound **1f** established a hydrogen bond with Gln211, while its bromine atom interacted with Pro105 and Phe208 through hydrophobic interactions. For the SCWP *O*-acetyltransferase molecular docking result of *B. cereus*, the amino acid residues Lys186, His201, His202, Ser337, Ser354, Arg359, and Ser364 are considered as the active site of SCWP *O*-acetyltransferase of *B.*

*cereus*.<sup>37</sup> The binding energy of tetracycline was calculated to be  $-5.75$  kcal mol<sup>-1</sup> (Table 3). Docking analysis revealed that tetracycline formed hydrogen bonds with Lys103, Asn104, Asn251, and Ser385. Additionally, it exhibited a pi-cation interaction with Arg359 and pi-alkyl interactions with His201 and His202 (Fig. 5). Compound **1a** displayed a hydrogen bond and a pi cation interaction with the key amino acids Arg359 and His201, respectively. Furthermore, the acetyl groups of **1a** established hydrogen bonds with the Lys103 and Asn104 residues, indicating that the acetyl groups could improve antibacterial toward *B. cereus*. Compound **1f** formed three hydrogen bonds with Lys103, Asn251, and Ser385 and pi-cation interactions with the key amino acids His201 and Arg359. The bromine atom established a halogen bond with Asp92, suggesting its role in stabilizing the protein-ligand interaction. Moreover, the






Fig. 5 The binding interactions of tetracycline, **1a**, and **1f** with SCWP *O*-acetyltransferase of *B. cereus*.




Fig. 6 The superimposed conformations of (a) tetracycline (purple) and **1f** (green) with the PBP2a of MRSA, (b) tetracycline (purple) and **1a** (yellow) with *B. subtilis* LCP enzyme, and (c) tetracycline (purple), **1a** (yellow), and **1f** (green) with SCWP *O*-acetyltransferase of *B. cereus*.

binding energy of **1a** and **1f** was lower than that of tetracycline. Structural superimpositions of **1a**, **1f**, and tetracycline further confirmed their similar binding interaction within the active site (Fig. 6c).

#### 2.4 Physicochemical properties

Lipinski's and Veber's rules were used to evaluate the drug-likeness of all compounds, and the results are summarized in Table 4. Lipinski's rule considers many criteria, including molecular weight ( $MW \leq 500$ ), hydrogen bond acceptors ( $HBA \leq 10$ ), hydrogen bond donors ( $HBD \leq 5$ ), and  $\log P$  (lipophilicity index  $< 5$ ).<sup>39</sup> A compound that violates more than one of these criteria is less likely to be an orally active drug. All compounds, except for **1b** and **2b**, meet these criteria, suggesting their potential for good oral bioavailability. These indicated that the structural modification of both parent compounds with pentanoyl chloride tends to compromise their drug-like properties. While Veber's rule considers the number of rotatable bonds ( $NORTB \leq 10$ ) and topological polar surface area ( $TPSA \leq 140$ ).<sup>40</sup> Most compounds, with the exception of **1b**, **2b**, **1c**, and **2c**, comply with both criteria, supporting the favorable oral absorption. The results of Lipinski's and Veber's rules indicated that compounds **1a** and **1f** could be developed to be the orally active drug.

## 3 Experimental

### 3.1 General experimental procedures

Melting points were determined on a SANYO Gallenkamp (Leicester, UK) melting point apparatus and are uncorrected. IR spectra were taken on a PerkinElmer Spectrum One spectrophotometer (Agilent Technologies, U.S.A.). NMR spectra were recorded in  $CDCl_3$  and  $CD_3OD$  on a Bucker Ascend 400 MHz (Bruker, Germany). Mass spectra were performed on an Agilent 1260 Infinity series high-performance liquid chromatography system (Agilent, Waldbronn, Germany), coupled with a 6540 ultra-high-definition accurate mass spectrometer (Agilent Technologies, Singapore). Column chromatography was carried out on SiliCycle silica gel (40–63  $\mu m$ , SiliCycle, Inc., Canada), TLC was performed with precoated Merck silica gel 60 PF254 on an aluminum (Merck, Darmstadt, Germany).

### 3.2 Plant material

Roots of *Maclura cochinchinensis* were collected from Pattani province, Thailand.<sup>4</sup> The specimens were identified by Prof. Pranom Chantaranothai, Department of Biology, Khon Kaen University, Thailand, where a voucher specimen (SK17) was deposited.



Table 4 Physicochemical properties of modified and parent compounds by *in silico* analysis

Compound	MW <sup>a</sup> (g mol <sup>-1</sup> )	LogP <sup>b</sup>	HBA <sup>c</sup>	HBD <sup>d</sup>	NORTB <sup>e</sup>	TPSA <sup>f</sup> (Å <sup>2</sup> )
1a	412.39	3.35	8	1	4	112.27
1b	580.67	6.64	9	0	15	118.34
1c	484.50	2.93	10	1	4	163.17
1d	356.37	3.49	6	1	2	78.13
1e	384.42	4.14	6	1	4	78.13
1f	407.21	3.32	6	3	0	100.13
2a	478.49	4.59	8	1	6	112.27
2b	646.77	7.69	9	0	17	118.34
2c	550.60	4.09	10	1	6	163.17
2d	422.47	4.68	6	1	4	78.13
2e	408.44	4.37	6	2	3	89.13
2f	450.52	5.33	6	1	6	78.13
2g	422.47	4.64	6	2	4	89.13
1	328.32	2.74	6	3	0	100.13
2	394.42	3.95	6	3	2	100.13

<sup>a</sup> Molecular weight. <sup>b</sup> Octanol/water partition coefficient. <sup>c</sup> Number of hydrogen bond acceptors. <sup>d</sup> Number of hydrogen bond donors. <sup>e</sup> Number of rotatable bonds. <sup>f</sup> Topological polar surface area.

### 3.3 Preparation of xanthone derivatives

**3.3.1 Isolation of toxylloxanthone C (1) and macluraxanthone (2).** The starting xanthone, macluraxanthone (2), was obtained from the crude hexane and EtOAc extracts of the root of *M. cochinchinensis*. Meanwhile, toxylloxanthone C (1) was recrystallized from the solid of subfraction E7.1 using MeOH as a solvent, following the procedure described in a previous publication.<sup>4</sup>

Toxylloxanthone C (1); yellow solid;  $[\alpha]_D^{24} +17.7$  (*c* 0.36, acetone);<sup>41</sup> <sup>1</sup>H NMR (400 MHz, (CH<sub>3</sub>)<sub>2</sub>CO)  $\delta$  13.45 (1H s, 1-OH), 7.60 (1H, d, *J* = 8.4 Hz, H-8), 6.96 (1H, d, *J* = 8.4 Hz, H-7), 6.30 (1H, s, H-4), 4.50 (1H, q, *J* = 6.8 Hz, H-12), 1.46 (s, 3H, H-14), 1.36 (3H, d, *J* = 6.4 Hz, H-13), 1.22 (3H, s, H-15); <sup>13</sup>C NMR (100 MHz, (CH<sub>3</sub>)<sub>2</sub>CO)  $\delta$  181.2 (C-9), 166.2 (C-1), 159.1 (C-3), 158.2 (C-4a), 151.6 (C-6), 146.5 (C-10a), 132.7 (C-5), 117.0 (C-8), 116.9 (C-8a), 114.2 (C-2), 113.3 (C-7), 103.6 (C-9a), 91.3 (C-12), 89.6 (C-4), 43.6 (C-11), 25.0 (C-14), 20.5 (C-15), 14.2 (C-13); HRESI-MS *m/z* 329.1025 [M + H]<sup>+</sup> (Calc. for 329.1025, C<sub>18</sub>H<sub>17</sub>O<sub>6</sub>).

Macluraxanthone (2); yellow crystals; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  7.56 (1H, d, *J* = 8.8 Hz, H-8), 6.88 (1H, d, *J* = 8.4 Hz, H-7), 6.69 (1H, d, *J* = 10.0 Hz, H-11), 6.44 (1H, dd, *J* = 17.6, 10.4 Hz, H-19), 5.64 (1H, d, *J* = 10.0 Hz, H-12), 4.99 (1H, dd, *J* = 17.6, 1.2 Hz, H-20a), 4.84 (1H, dd, *J* = 10.4, 1.2 Hz, H-20b), 1.73 (6H, s, H-17, H-18), 1.47 (6H, s, H-14, H-15); <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD)  $\delta$  182.5 (C-9), 160.1 (C-1), 157.3 (C-3), 156.6 (C-4a), 152.8 (C-19), 152.7 (C-6), 147.5 (C-10a), 134.2 (C-5), 128.4 (C-12), 117.1 (C-11), 116.8 (C-8), 114.9 (C-8a), 114.6 (C-7), 113.7 (C-4), 107.7 (C-20), 106.1 (C-2), 103.9 (C-9a), 79.3 (C-13), 42.2 (C-16), 30.2 (C-17, C-18), 28.1 (C-14, C-15); HRESI-MS: *m/z* 395.1496 [M + H]<sup>+</sup> (Calc. for C<sub>23</sub>H<sub>23</sub>O<sub>6</sub> 395.1495).

**3.3.2 General procedure to prepare 1a-c and 2a-c.**<sup>42,43</sup> Toxylloxanthone C (1) (24.3 mg, 0.074 mmol) or macluraxanthone (2) (27.9 mg, 0.071 mmol) was dissolved in pyridine (2 mL), followed by the dropwise addition of an excess of acetic anhydride (Ac<sub>2</sub>O) at 0 °C. The reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into

cold water (20 mL) and extracted with EtOAc (2 × 20 mL). The organic layer was combined, washed with water and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure to obtain a crude solid. Recrystallization of a crude solid from hexane yielded **1a** (12.8 mg, 41%) or **2a** (22 mg, 65%).

The reactions of **1** or **2** with pentanoyl chloride or methanesulfonyl chloride were performed following the same procedure described above, affording **1b** (10.2 mg, 11%), **1c** (27.3 mg, 77%), **2b** (19.9 mg, 19%), and **2c** (34.3 mg, 88%), respectively.

**3.3.2.1 General procedure to prepare 1d-e and 2d-g.**<sup>42</sup> To a solution of toxylloxanthone C (1) (30.0 mg, 0.089 mmol) or macluraxanthone (2) (59.2 mg, 0.15 mmol) in dry acetone (2 mL), anhydrous sodium carbonate (160 mg), and methyl iodide (0.12 mL) were added. The reaction mixture was refluxed overnight. The reaction was quenched by the addition of cold water and extracted with EtOAc (2 × 20 mL). The organic combined layer was washed with water, brine, and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent gave a crude oil. The crude product was purified by silica gel column chromatography using 20% EtOAc/hexane as an eluent, yielding **1d** (30.2 mg, 94%) or **2d** (28.6 mg, 45%) and **2e** (8.6 mg, 14%).

The reaction of **1** (49.0 mg, 0.149 mmol) or **2** with ethyl iodide (EtI) (59.1 mg, 0.149 mmol) was performed in the same procedure as described above, yielding **1e** (42.3 mg, 86%), or **2f** (41.7 mg, 70%) and **2g** (13.4 mg, 27%).

**3.3.3 Bromination using N-bromosuccinimide (NBS).**<sup>43</sup> Toxylloxanthone C (1) (30.8 mg, 0.094 mmol) was dissolved in EtOAc (2 mL), and a solution of NBS (0.155 mmol) in MeCN:H<sub>2</sub>O (1:1, 2 mL) was added dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 1 hour, followed by stirring at room temperature for 1 hour. The reaction mixture was poured into cold water (20 mL) and extracted with EtOAc (2 × 20 mL). The organic layer was combined, washed with water and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure to afford a crude solid. Recrystallization from hexane



yielded **1f** (24.2 mg, 36%). Bromination of macluraxanthone (**2**) (37.3 mg, 0.094 mmol) using the same procedure resulted in no product formation.

### 3.4 Spectroscopic data

5,6-Diacetoxyxyloxyloxanthone C (**1a**); yellow solid; mp 164.1–165.3 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 2965, 2915, 1775, 1648, 1579, 1428, 1370, 1260; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  12.84 (1H, s, 1-OH), 8.14 (1H, d, *J* = 8.8, 0.8 Hz, H-8), 7.20 (1H, d, *J* = 8.8, 0.8 Hz, H-7), 6.29 (1H, s, *J* = 0.8 Hz, H-4), 4.52 (1H, q, *J* = 6.8 Hz, H-12), 2.43 (3H, s, H-2''), 2.35 (3H, s, H-2'), 1.50 (3H, s, H-14), 1.39 (3H, d, *J* = 6.8 Hz, H-13), 1.26 (3H, s, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  179.9 (C-9), 167.7 (C-1''), 167.4 (C-1'), 166.5 (C-1), 158.8 (C-3), 157.6 (C-4a), 149.2 (C-6), 147.4 (C-10a), 130.1 (C-5), 123.5 (C-8), 119.6 (C-8a), 118.6 (C-2), 117.6 (C-7), 104.2 (C-9a), 91.4 (C-12), 90.1 (C-4), 43.4 (C-11), 25.3 (C-2''), 20.8 (C-2'), 20.7 (C-14), 20.4 (C-15), 14.5 (C-13); HRESI-MS *m/z* 413.1245 [M + H]<sup>+</sup> (Calc. for 413.1236, C<sub>22</sub>H<sub>21</sub>O<sub>8</sub>).

1,5,6-Tripentanoyloxyxyloxyloxanthone C (**1b**); white solid; mp 85.2–87.9 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 2958, 2935, 2873, 1765, 1658, 1456, 1418, 1257; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.13 (1H, d, *J* = 8.8 Hz, H-8), 7.15 (1H, d, *J* = 8.8 Hz, H-7), 6.47 (1H, s, H-4), 4.59 (1H, q, *J* = 6.4 Hz, H-12), 2.77 (2H, t, *J* = 7.6 Hz, H-2''), 2.67 (2H, t, *J* = 7.6 Hz, H-2'), 2.59 (2H, t, *J* = 7.6 Hz, H-2'), 1.97–1.67 (6H, m, H-3'', H-3' and H-3'), 1.54 (3H, s, H-14), 1.48 (6H, m, H-4'', H-4' and H-4'), 1.43 (3H, d, *J* = 6.4 Hz, H-13), 1.43 (3H, s, H-15), 1.10–0.92 (9H, m, H-5'', H-5', H-5'); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  173.8 (C-9), 172.6 (C-12), 170.6 (C-1''), 170.1 (C-1'), 164.1 (C-1), 154.0 (C-3), 152.2 (C-4a), 148.5 (C-6), 147.3 (C-10a), 130.8 (C-5), 124.4 (C-8), 121.2 (C-8a), 120.1 (C-2), 118.7 (C-7), 109.5 (C-9a), 103.6 (C-12), 91.4 (C-4), 44.5 (C-11), 34.2 (C-2''), 33.9 (C-2'), 33.6 (C-2'), 27.0 (C-3''), 26.9 (C-3''), 26.7 (C-3'), 25.8 (C-14), 22.42 (C-4''), 22.38 (C-4''), 22.38 (C-4'), 21.3 (C-15), 14.4 (C-13), 14.0 (C-5''), 13.9 (C-5''), 13.8 (C-5'); HRESI-MS *m/z* 581.2718 [M + H]<sup>+</sup> (Calc. for 581.2751, C<sub>33</sub>H<sub>41</sub>O<sub>9</sub>).

5,6-Dimesyloxyxyloxyloxanthone C (**1c**); yellow solid; mp 191.1–192.3 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 3020, 2940, 1648, 1597, 1448, 1355, 1274; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  12.65 (1H, s, 1-OH), 8.20 (1H, d, *J* = 8.8 Hz, 1H, H-8), 7.48 (1H, d, *J* = 8.8 Hz, H-7), 6.30 (1H, s, H-4), 4.55 (1H, q, *J* = 6.8 Hz, H-12), 3.49 (3H, s, 6-OSO<sub>2</sub>CH<sub>3</sub>), 3.38 (3H, s, 5-OSO<sub>2</sub>CH<sub>3</sub>), 1.50 (3H, s, H-14), 1.40 (d, *J* = 6.8 Hz, 3H, H-13), 1.26 (s, 3H, H-15); <sup>13</sup>C NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  180.0 (C-9), 166.9 (C-1), 158.9 (C-3), 157.3 (C-4a), 149.5 (C-6), 146.6 (C-10a), 130.1 (C-5), 125.1 (C-8), 120.6 (C-8a), 118.6 (C-2), 118.3 (C-7), 104.0 (C-9a), 91.7 (C-12), 89.9 (C-4), 43.5 (C-11), 40.3 (6-OSO<sub>2</sub>CH<sub>3</sub>), 39.5 (5-OSO<sub>2</sub>CH<sub>3</sub>), 25.3 (C-14), 20.7 (C-15), 14.5 (C-13); HRESI-MS *m/z* 485.0579 [M + H]<sup>+</sup> (Calc. for 485.0576, C<sub>20</sub>H<sub>21</sub>O<sub>10</sub>S<sub>2</sub>).

5,6-Dimethoxyxyloxyloxanthone C (**1d**); yellow solid; mp 142.7–145.9 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 3069, 2958, 2938, 2880, 2844, 1648, 1590, 1428, 1282, 1212; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  13.10 (1H, s, 1-OH), 7.97 (1H, d, *J* = 8.8 Hz, H-8), 7.95 (1H, d, *J* = 8.8 Hz, H-7), 6.40 (1H, s, H-4), 4.50 (1H, q, *J* = 6.8 Hz, H-12), 4.00 (3H, d, 6-OCH<sub>3</sub>), 3.99 (3H, d, 5-OCH<sub>3</sub>), 1.50 (s, 3H, H-14), 1.40 (3H, d, *J* = 6.4 Hz, H-13), 1.25 (3H, s, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  180.7 (C-9), 166.1 (C-1), 158.8 (C-3), 158.0 (C-4a), 157.6 (C-6),

150.3 (C-10a), 136.2 (C-5), 121.5 (C-8), 117.0 (C-8a), 115.5 (C-2), 108.6 (C-7), 103.8 (C-9a), 91.2 (C-12), 89.9 (C-4), 61.7 (6-OCH<sub>3</sub>), 56.6 (5-OCH<sub>3</sub>), 43.4 (C-11), 25.3 (C-14), 20.7 (C-15), 14.4 (C-13); HRESI-MS *m/z* 357.1346 [M + H]<sup>+</sup> (Calc. for 357.1338, C<sub>20</sub>H<sub>21</sub>O<sub>6</sub>).

5,6-Diethoxyxyloxyloxanthone C (**1e**); yellow solid; mp 132.3–132.5 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 2977, 1656, 1604 1569, 1440, 1284, 1146; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  13.13 (1H, s, 1-OH), 7.92 (1H, d, *J* = 8.8 Hz, H-8), 6.94 (1H, d, *J* = 8.8 Hz, H-7), 6.38 (1H, s, H-4), 4.50 (1H, q, *J* = 6.4 Hz, H-12), 4.19 (4H, q, *J* = 6.8 Hz, H-1'', 1'), 1.50 (6H, m, H-14, H-2'') 1.41 (6H, m, H-13, H-2'), 1.25 (s, 3H, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  180.7 (C-9), 165.9 (C-1), 158.8 (C-3), 158.1 (C-4a), 157.3 (C-6), 150.8 (C-10a), 135.3 (C-5), 121.2 (C-8), 116.9 (C-8a), 115.2 (C-2), 109.5 (C-7), 103.8 (C-9a), 91.1 (C-12), 89.9 (C-4), 69.8 (C-1''), 64.9 (C-1'), 43.4 (C-11), 25.3 (C-15), 20.7 (C-14), 15.7 (C-2''), 14.9 (C-2') 14.4 (C-13); HRESI-MS *m/z* 385.1651 [M + H]<sup>+</sup> (Calc. for C<sub>22</sub>H<sub>25</sub>O<sub>6</sub>, 385.1651).

4-Bromotoxyloxyloxanthone C (**1f**); brown solid; mp 162.4–162.7 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 3595, 3165, 2959, 1646, 1607, 1579, 1425, 1309 1247, 1203, 698; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  7.59 (1H, d, *J* = 8.4 Hz, H-8), 6.92 (1H, d, *J* = 8.4 Hz, H-7), 4.61 (1H, q, *J* = 6.4 Hz, H-12), 1.51 (3H, s, H-14), 1.44 (3H, d, *J* = 6.4 Hz, H-13), 1.25 (3H, s, H-15); <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD)  $\delta$  182.1 (C-9), 164.2 (C-1), 158.7 (C-3), 155.1 (C-4a), 153.7 (C-6), 147.8 (C-10a), 133.9 (C-5), 118.3 (C-8), 117.7 (C-8a), 114.6 (C-2), 114.1 (C-7), 105.3 (C-9a), 92.8 (C-12), 81.9 (C-4), 45.8 (C-11), 25.6 (C-14), 20.9 (C-15), 14.5 (C-13); HRESI-MS *m/z* 407.1025 [M + H]<sup>+</sup> (Calc. for C<sub>18</sub>H<sub>16</sub>BrO<sub>6</sub>, 407.1030), *m/z* 409.0111 [M+2 + H]<sup>+</sup> (Calc. for C<sub>18</sub>H<sub>16</sub>BrO<sub>6</sub> + 2, 409.0110).

5,6-Diacetoxymacluraxanthone (**2a**); yellow solid; mp 165.9–168.2 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 3728, 2927, 1625, 1453, 1374; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  13.4 (1H, s, 1-OH), 8.13 (1H, d, *J* = 8.8 Hz, H-8), 7.22 (1H, d, *J* = 8.8 Hz, H-7), 6.73 (1H, d, *J* = 10.0 Hz, H-11), 6.24 (1H, dd, *J* = 17.6, 10.4 Hz, H-19), 5.58 (1H, d, *J* = 10.0 Hz, H-12), 4.88 (1H, dd, *J* = 17.6, 1.2 Hz, H-20a) 4.84 (1H, dd, *J* = 10.4, 1.2 Hz, H-20b), 2.39 (3H, s, H-2'') 2.34 (3H, s, H-2') 1.65 (6H, s, H-17, H-18) 1.46 (6H, s, H-14, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  180.3 (C-9), 167.7 (C-1'), 167.6 (C-1''), 160.1 (C-3), 156.7 (C-1), 154.9 (C-4a), 150.2 (C-19), 149.2 (C-6), 147.7 (C-10a), 130.8 (C-5), 127.7 (C-12), 123.5 (C-11), 119.1 (C-8), 118.8 (C-8a), 115.8 (C-7), 113.5 (C-4), 108.5 (C-20), 105.8 (C-2), 103.7 (C-9a), 78.7 (C-13), 41.2 (C-16), 30.1 (C-17, C-18), 28.1 (C-14, C-15), 20.8 (C-2''), 20.6 (C-2'); HRESI-MS *m/z* 479.1706 [M + H]<sup>+</sup> (Calc. for C<sub>27</sub>H<sub>27</sub>O<sub>8</sub>, 479.1706).

1,5,6-Tripentanoyloxymacluraxanthone (**2b**); yellow solid; mp 99.0–100.2 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>); 2958, 2933, 1766, 1652, 1595, 1460, 1130; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.09 (2H, d, *J* = 8.8 Hz, H-8), 7.15 (2H, d, *J* = 8.8 Hz, H-7), 6.44 (2H, d, *J* = 10.0 Hz, H-11), 6.28 (2H, dd, *J* = 17.6, 10.4 Hz, H-19), 5.72 (2H, d, *J* = 10.0 Hz, H-12), 4.86 (1H, dd, *J* = 17.6, 1.2 Hz, H-20a), 4.84 (1H, dd, *J* = 10.4, 1.2 Hz, H-20b), 2.82 (2H, t, *J* = 7.6 Hz, H-2''), 2.66 (2H, t, *J* = 7.6 Hz, H-2'), 2.57 (2H, t, *J* = 7.6 Hz, H-2'), 1.78 (6H, m, H-3', H-3'', H-3'''), 1.67 (6H, s, H-17, H-18), 1.48 (6H, m, H-4', H-4'', H-4'''), 1.46 (6H, s, H-14, H-15), 1.00 (9H, m, H-5', H-5'', H-5'''); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  174.5 (C-9), 172.1 (C-1''), 170.6 (C-1'), 170.5 (C-1'), 157.6 (C-1), 156.4 (C-3), 150.0 (C-4a), 148.5 (C-19), 147.3 (C-6), 144.4 (C-10a), 131.5 (C-5), 130.7 (C-12), 123.9 (C-11), 120.6 (C-8), 120.5 (C-8a), 118.7 (C-7), 115.7 (C-4), 112.9 (C-



20), 109.4 (C-2), 108.5 (C-9a), 78.4 (C-13), 41.7 (C-16), 34.2 (C-2'''), 33.9 (C-2''), 33.8 (C-2'), 30.0 (C-17, C-18), 28.0 (C-14, C-15), 26.9 (C-3''', C-3''), 26.8 (C-3'), 22.6 (C-4'''), 22.4 (C-4''), 22.3 (C-4'), 14.0 (C-5'''), 13.9 (C-5''), 13.9 (C-5'); HRESI-MS:  $m/z$  647.3219 [M + H]<sup>+</sup> (Calc. for C<sub>38</sub>H<sub>47</sub>O<sub>9</sub>, 647.3220).

5,6-Dimesyloxymacluraxanthone (**2c**); yellow solid; mp 179.8–181.9 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>): 3011 2978 1652 1600 1576 1455 1408 1352 1291 1175; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  13.26 (1H, s, 1-OH), 8.24 (1H, d,  $J$  = 8.8 Hz, H-8), 7.48 (2H, d,  $J$  = 8.8 Hz, H-7), 6.73 (1H, d,  $J$  = 10.0 Hz, H-11), 6.34 (1H, dd,  $J$  = 17.6, 10.4 Hz, H-19), 5.61 (1H, d,  $J$  = 10.0 Hz, H-12), 4.91 (1H, dd,  $J$  = 17.6, 1.2 Hz, H-20a), 4.84 (1H, dd,  $J$  = 10.4, 1.2 Hz, H-20b), 3.43 (3H, s, 6-OSO<sub>2</sub>CH<sub>3</sub>), 3.33 (3H, s, 5-OSO<sub>2</sub>CH<sub>3</sub>), 1.72 (6H, s, H-17, H-18), 1.47 (6H, s, H-14, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  179.5 (C-9), 160.7 (C-1), 156.6 (C-3), 154.9 (C-4a), 150.6 (C-19), 149.9 (C-6), 146.5 (C-10a), 129.7 (C-5), 128.0 (C-12), 125.3 (C-11), 120.4 (C-8), 117.6 (C-8a), 115.6 (C-7), 114.6 (C-4), 108.2 (C-20), 106.1 (C-2), 103.7 (C-9a), 79.0 (C-13), 41.2 (C-16), 40.3 (C-6-OSO<sub>2</sub>CH<sub>3</sub>), 39.0 (C-5-OSO<sub>2</sub>CH<sub>3</sub>), 30.0 (C-17, C-18), 28.1 (C-14, C-15); HRESI-MS:  $m/z$  551.1012 [M + H]<sup>+</sup> (Calc. for C<sub>25</sub>H<sub>27</sub>O<sub>10</sub>S<sub>2</sub>, 551.1046).

5,6-Dimethoxymacluraxanthone (**2d**); yellow solid; mp 138.1–141.2 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>): 2966 2917 1648 1596 1458 1428 1286; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  13.63 (1H, s, 1-OH), 7.96 (2H, d,  $J$  = 8.8 Hz, H-8), 6.98 (1H, d,  $J$  = 8.8 Hz, H-7), 6.73 (2H, d,  $J$  = 10.0 Hz, H-11), 6.37 (1H, dd,  $J$  = 17.6, 10.4 Hz, H-19), 5.58 (2H, d,  $J$  = 10 Hz, H-12), 4.92 (1H, dd,  $J$  = 17.6, 1.2 Hz, H-20a), 4.84 (1H, dd,  $J$  = 10.4, 1.2 Hz, H-20b), 4.00 (3H, s, 6-OCH<sub>3</sub>), 3.92 (3H, s, 5-OCH<sub>3</sub>), 1.727 (6H, s, H-17, H-18), 1.47 (6H, s, H-14, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.1 (C-9), 159.4 (C-1), 158.1 (C-3), 156.6 (C-4a), 155.4 (C-19), 150.8 (C-6), 150.1 (C-10a), 136.5 (C-5), 127.3 (C-12), 121.4 (C-11), 116.1 (C-8), 114.9 (C-8a), 113.8 (C-7), 108.7 (C-4), 107.9 (C-20), 105.3 (C-2), 103.4 (C-9a), 78.3 (C-13), 61.6 (6-OCH<sub>3</sub>), 56.5 (5-OCH<sub>3</sub>), 41.2 (C-16), 30.0 (C-17, C-18), 27.9 (C-14, C-15); HRESI-MS:  $m/z$  423.1795 [M + H]<sup>+</sup> (Calc. for C<sub>25</sub>H<sub>27</sub>O<sub>6</sub>, 423.1808).

5-Methoxymacluraxanthone (**2e**); yellow crystals; mp 151.4–158.8 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>): 3279 2933 2840 1619 1567 1412 1282; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  13.54 (1H, s, 1-OH), 7.73 (1H, d,  $J$  = 8.8 Hz, H-8), 6.95 (1H, d,  $J$  = 8.8 Hz, H-7), 6.75 (1H, d,  $J$  = 10.0 Hz, H-11), 6.65 (1H, dd,  $J$  = 17.6, 10.4 Hz, H-19), 6.22 (1H, s, 6-OH), 5.60 (1H, d,  $J$  = 10.0 Hz, H-12), 5.18 (1H, dd,  $J$  = 17.6, 1.2 Hz, H-20a), 5.04 (1H, dd,  $J$  = 10.4, 1.2 Hz, H-20b), 4.01 (3H, s, 5-OCH<sub>3</sub>), 1.65 (6H, s, H-17, H-18), 1.50 (6H, s, H-14, H-15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.1 (C-9), 159.2 (C-1), 156.8 (C-3), 155.1 (C-4a), 154.6 (C-19), 151.6 (C-6), 144.5 (C-10a), 133.6 (C-5), 127.3 (C-12), 116.9 (C-11), 116.1 (C-8), 114.4 (C-8a), 113.5 (C-7), 108.5 (C-4), 105.5 (C-20), 104.8 (C-2), 103.2 (C-9a), 78.4 (C-13), 56.7 (5-OCH<sub>3</sub>), 41.4 (C-16), 28.6 (C-17, C-18), 28.0 (C-14, C-15); HRESI-MS:  $m/z$  409.1657 [M + H]<sup>+</sup> (Calc. for C<sub>24</sub>H<sub>25</sub>O<sub>6</sub>, 409.1651).

5,6-Diethoxymacluraxanthone (**2f**); yellow solid; mp 118.6–120.3 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>): 3599 2979 1644 1594 1571 1446 1287; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  13.68 (1H, s, 1-OH), 7.91 (1H, d,  $J$  = 8.8 Hz, H-8), 6.94 (1H, d,  $J$  = 8.8 Hz, H-7), 6.74 (1H, d,  $J$  = 10.0 Hz, H-11), 6.38 (1H, dd,  $J$  = 17.6, 10.4 Hz, H-19), 5.57 (1H, d,  $J$  = 10.0 Hz, H-12), 4.90 (1H, dd,  $J$  = 17.6, 1.2 Hz, H-20a), 4.83 (1H, dd,  $J$  = 10.4, 1.2 Hz, H-20b), 4.20 (2H, m, H-1''), 4.15 (2H, m,

H-1'), 1.73 (6H, s, H-18, 17), 1.47 (12H, m, 2'', 2', 14, 15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.1 (C-9), 159.3 (C-1), 157.5 (C-3), 156.6 (C-4a), 155.5 (C-19), 150.9 (C-6), 150.3 (C-10a), 135.5 (C-5), 127.3 (C-12), 121.1 (C-11), 116.1 (C-8), 114.7 (C-8a), 113.7 (C-7), 109.5 (C-4), 107.8 (C-20), 105.2 (C-2), 103.4 (C-9a), 78.2 (C-13), 69.5 (C-1''), 64.9 (C-1'), 41.2 (C-16), 30.0 (C-17, C-18), 27.9 (C-14, C-15), 15.6 (C-2''), 14.9 (C-2'); HRESI-MS:  $m/z$  451.2121 [M + H]<sup>+</sup> (Calc. for C<sub>27</sub>H<sub>31</sub>O<sub>6</sub>, 451.2121).

5-Ethoxymacluraxanthone (**2g**); yellow solid; mp 126.3–128.6 °C; IR  $\nu_{\max}$  (cm<sup>-1</sup>): 3272 2964 1647 1619 1594 1568 1460 1417 1139; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  13.57 (1H, s, 1-OH), 7.71 (1H, d,  $J$  = 8.8 Hz, H-8), 6.94 (1H, d,  $J$  = 8.8 Hz, H-7), 6.76 (1H, d,  $J$  = 10.0 Hz, H-11), 6.63 (1H, dd,  $J$  = 17.6, 10.4 Hz, H-19), 6.18 (1H, s, 6-OH), 5.60 (1H, d,  $J$  = 10.0 Hz, H-12), 5.16 (1H, dd,  $J$  = 17.6, 1.2 Hz, H-20a), 5.04 (1H, dd,  $J$  = 10.4, 1.2 Hz, H-20b), 4.25 (2H, m, H-1''), 1.66 (6H, s, H-17, 18), 1.52 (3H, d, H-2''), 1.50 (6H, s, 14, 15); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.1 (C-9), 159.2 (C-1), 156.8 (C-3), 154.7 (C-4a), 154.6 (C-19), 151.0 (C-6), 144.6 (C-10a), 133.7 (C-5), 127.2 (C-12), 116.8 (C-11), 116.2 (C-8), 114.3 (C-8a), 113.5 (C-7), 109.4 (C-4), 105.5 (C-20), 105.1 (C-2), 103.3 (C-9a), 78.3 (C-13), 65.2 (C-1''), 41.4 (C-16), 28.7 (C-17, C-18), 28.0 (C-14, C-15), 14.9 (C-2''); HRESI-MS:  $m/z$  423.1808 [M + H]<sup>+</sup> (Calc. for C<sub>25</sub>H<sub>27</sub>O<sub>6</sub>, 423.1808).

### 3.5 Cytotoxicity and cell proliferation assays

All types of cancer and non-tumorigenic cells were purchased from ATCC. Cytotoxicity assays against human cervical carcinoma (HelaS3; ATCC CCL2-2), human hepatocellular carcinoma (HepG2; ATCC HB-8065), human lung carcinoma (A549; ATCC CCL-185), and African green monkey kidney (Vero; ATCC CCL-81) cell lines were performed by terminal deoxynucleotidyl transferase (TdT) dUTP Nick-End Labeling (TUNEL) assay.<sup>44</sup> Cells proliferation was performed using the MTT colorimetric assay,<sup>44</sup> as in our previous publication.<sup>4</sup>

### 3.6 Antibacterial assay

Four bacterial cultures *Bacillus cereus* ATCC 11778, *Bacillus subtilis* ATCC 6633, *Staphylococcus aureus* ATCC 25923, and Methicillin resistant *Staphylococcus aureus* (MRSA) were employed as the test organisms. Minimum inhibitory concentration (MIC) values were determined using the dilution method, as described in the M07-A9,<sup>45</sup> as described in our previous paper.<sup>4,46</sup>

### 3.7 Molecular docking simulation

All the calculations were performed on Intel® Core™ i7-12700F CPU 2.11 GHz with 32 GB DDR4 RAM. The crystallographic structures of cyclin-dependent kinase 2 (CDK2) in complex with the inhibitor RC-3-89 (PDB code: 4GCJ), methicillin acylpenicillin binding protein 2a from methicillin resistant *Staphylococcus aureus* (PDB code: 1MWU), and *Bacillus cereus* PatB1 (PDB code: 5V8E) were obtained from the Protein Data Bank. The existing ligands and water molecules were removed from the crystal structure. Molecular docking studies were performed using the AutoDock 4.2.6 program.<sup>47</sup> The 3D structures of compounds **1**, **1a**, **1b**, and **1f** were minimized at the B3LYP/6-



31G(d) levels using the Gaussian 03 program. All hydrogens were added and Gasteiger charges were assigned by using AutoDockTools.<sup>47</sup> A grid box of 40 × 40 × 40 points was centered at the coordinates of the RC-3-89 inhibitor ( $x = -51.563$ ,  $y = 98.287$ ,  $z = -62.755$ ) to define the binding site of the CDK2 protein. Similarly, the grid box was generated at the center of the coordinate's ligand with dimensions of 40 × 40 × 40 points, centered at  $x = 28.221$ ,  $y = 28.839$ , and  $z = 87.515$  for PBP2a from MRSA, and centered at  $x = -67.474$ ,  $y = -1.104$ , and  $z = -15.158$  for *Bacillus cereus* PatB1. For *B. subtilis* TagU, a grid box was generated in the absence of a co-crystallized ligand to cover all active site residues<sup>35</sup> with dimensions of 70 × 70 × 70 points, centered at  $x = 8.698$ ,  $y = 24.542$ , and  $z = 24.418$ . All grid boxes were constructed with a grid spacing of 0.375 Å. The optimized ligands were docked onto protein templates using the Lamarckian genetic algorithm. The docking protocol was set to default, with 150 independent docking runs. The protein templates of CDK2, PBP2a from MRSA, and SCWP *O*-acetyltransferase of *B. cereus* exhibited the best results of redocking experiments with the RMSD of 0.68, 1.67, and 1.42 Å, respectively (Table S1). Finally, the docking results were then analyzed to identify the best cluster of each compound with the lowest free binding energy. Protein–ligand interactions were visualized by using the Discovery Studio 2021 Client program (Accelrys, Inc., San Diego, CA, USA).

### 3.8 *In silico* physicochemical properties

The physicochemical properties of all compounds were evaluated to assess their potential as drug candidates using the SwissADME<sup>48</sup> web server. Drug-likeness was verified using the Lipinski's and Veber's rules.

## 4 Conclusions

Two main compounds, toxylloxanthone C (**1**) and macluraxanthone (**2**) from *M. cochinchinensis* roots, have been structurally modified through simple organic reactions including acylation, alkylation, sulfonylation, and bromination yielding thirteen derivatives **1a–f** and **2a–g**. All compounds were evaluated for their cytotoxic and antibacterial activities. Compounds **1a**, **1b**, and **1f** exhibited greater cytotoxic potency against A549 cells compared to their parent compound. Molecular docking studies revealed that **1a**, **1b**, and **1f** bound to CDK2 through hydrogen bonding and hydrophobic interactions with key amino acid residues. Furthermore, **1a** and **1b** demonstrated lower binding energies than **1**. Notably, **1a** exhibited potential as a lead anticancer candidate for further development due to its strong cytotoxicity with minimal toxicity toward normal cells. Additionally, **1a**, **1b**, **1f**, and **2a** displayed stronger antibacterial activity against four bacterial strains than their parent compounds. The docking results indicated that **1a** and **1f** interacted with key amino acids in the PBP2a active site of MRSA. In addition, the binding conformation of **1a** in the active site of LCP enzyme from *B. subtilis* aligned with that of tetracycline. While **1a**, **1f**, and tetracycline exhibited similar binding orientations with the key amino

acids within the active site of *B. cereus*. The physicochemical properties of **1a** and **1f** satisfied Lipinski's and Veber's rules, supporting their potential as drug-like candidates. Most derivatives of **2** exhibited reduced anticancer and antibacterial activities compared to the parent xanthone. This decrease in activity is likely due to steric hindrance between the prenyl group at the C-4 position and the additional substituents introduced in the derivatives, a structural effect consistent with previous reports on modified prenylated xanthones.<sup>10,27</sup> Overall, **1a** and **1f** exhibited significant dual bioactivities, showing cytotoxicity against A549 cells and antibacterial activity against four bacterial cultures; however, **1f** also displayed cytotoxicity toward Vero cells. Therefore, 5,6-diacetoxytoxylloxanthone C (**1a**) is interesting candidate for further study as anticancer and antibacterial agents.

## Author contributions

C. Linphosan performed the experiments, analyzed the data, evaluated antibacterial activity, and drafted the initial manuscript. W. Klangawad isolated parent compounds. J. Yahuafai performed cytotoxicity assay. J. Jandaruang and T. Promgool evaluated antibacterial activity. S. Pitchuanom supervised the molecular docking studies. J. Paluka and S. Boonlue provided antibacterial testing facilities. K. Poopasit carried out the NMR experiment. K. Kanokmedhakul to consultant and manuscript editing. O. Limtragool, the principal investigator, designed the experiments, analyzed the data, conducted *in silico* studies, and prepared the manuscript.

## Conflicts of interest

The authors declare no conflict of interest.

## Data availability

The datasets supporting this article have been uploaded as part of the SI. See DOI: <https://doi.org/10.1039/d5ra05758b>.

## Acknowledgements

This research project was financially supported by Thailand Science Research and Innovation (TSRI). Linphosan C. thanks the Center of Excellence for Innovation in Chemistry (PERCH-CIC). We acknowledge the Office of the Permanent Secretary, Ministry of Higher Education, Science, Research and Innovation (OP S MHESI, Grant No. RGNS 64–140) for partial support. We thank Prof. Chavi Yenjai for providing chemical support for this research.

## References

- 1 L. Rueankham, P. Panyajai, A. Saiai, M. Rungrojsakul, S. Tima, S. Chiampanichayakul, K. Yeerong, S. Somwongin, W. Chaiyana, P. Dejkriengkraikul, S. Okonogi, T. Katekunlaphan and S. Anuchapreeda, *BMC Complement. Med. Ther.*, 2023, **23**, 1–20.



- 2 C. Jansakun, W. Chulrik, J. Hata, T. Utaipan, W. Pabuprapap, N. Supaweera, O. Mueangson, A. Suksamrarn and W. Chunglok, *Inflammopharmacology*, 2023, **31**, 529–541.
- 3 T. Van Chien, N. T. Anh, N. T. Thanh, T. T. P. Thao, T. Van Loc and T. Van Sung, *Nat. Prod. Res.*, 2019, **33**, 212–218.
- 4 W. Lakornwong, K. Kanokmedhakul, J. Masranoi, S. Tontapha, J. Yahuafai, S. Laphookhieo, V. Suthiphasilp and S. Kanokmedhakul, *Nat. Prod. Res.*, 2022, **36**, 6021–6030.
- 5 K. I. Nakashima, T. Tanaka, H. Murata, K. Kaburagi and M. Inoue, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 1998–2001.
- 6 K. ichi Nakashima, T. Ogiwara, T. Hirai, T. Tanaka, H. Murata, K. Kaburagi, Y. Fujii-Kuriyama, H. Hayashi and M. Inoue, *Bioorg. Med. Chem.*, 2017, **25**, 4253–4258.
- 7 I. Polbuppha, V. Suthiphasilp, T. Maneerat, R. Charoensup, T. Limtharakul, S. Cheenpracha, S. G. Pyne and S. Laphookhieo, *Phytochemistry*, 2021, **187**, 112773.
- 8 N. Boonnak, S. Chantrapromma, K. Sathirakul and C. Kaewpiboon, *Bioorg. Med. Chem. Lett.*, 2020, **30**, 127494.
- 9 H. Zou, J. J. Koh, J. Li, S. Qiu, T. T. Aung, H. Lin, R. Lakshminarayanan, X. Dai, C. Tang, F. H. Lim, L. Zhou, A. L. Tan, C. Verma, D. T. H. Tan, H. S. O. Chan, P. Saraswathi, D. Cao, S. Liu and R. W. Beuerman, *J. Med. Chem.*, 2013, **56**, 2359–2373.
- 10 L. Kar Wei, N. H. Zamakshshari, G. C. L. Ee, S. H. Mah and S. M. Mohd Nor, *Nat. Prod. Res.*, 2018, **32**, 2147–2151.
- 11 A. J. Hou, T. Fukai, M. Shimazaki, H. Sakagami, H. D. Sun and T. Nomura, *J. Nat. Prod.*, 2001, **64**, 65–70.
- 12 T. Fukai, M. Yonekawa, A. J. Hou, T. Nomura, H. D. Sun and J. Uno, *J. Nat. Prod.*, 2003, **66**, 1118–1120.
- 13 Y. S. Zou, A. J. Hou, G. F. Zhu, Y. F. Chen, H. D. Sun and Q. S. Zhao, *Bioorg. Med. Chem.*, 2004, **12**, 1947–1953.
- 14 T. Fukai, Y. Oku, A. J. Hou, M. Yonekawa and S. Terada, *Chem. Biodivers.*, 2004, **1**, 1385–1390.
- 15 C. Lei, C. C. Liu, E. H. Pi and A. J. Hou, *Helv. Chim. Acta*, 2014, **97**, 1683–1688.
- 16 Y. H. Duan, Y. Dai, G. H. Wang, L. Y. Chen, H. F. Chen, D. Q. Zeng, Y. L. Li and X. S. Yao, *J. Asian Nat. Prod. Res.*, 2015, **17**, 519–531.
- 17 G. D. Monache, F. D. Monache, G. B. M. Bettolo and R. A. de Lima, *J. Nat. Prod.*, 1983, **46**, 655–659.
- 18 Y. H. Wang, A. J. Hou, G. F. Zhu, D. F. Chen and H. D. Sun, *Planta Med.*, 2005, **71**, 273–274.
- 19 T. Fukai, Y. Oku, A. J. Hou, M. Yonekawa and S. Terada, *Phytomedicine*, 2005, **12**, 510–513.
- 20 P. Pailee, C. Kuhakarn, C. Sangsuwan, S. Hongthong, P. Piyachaturawat, K. Suksen, S. Jariyawat, R. Akkarawongsapat, J. Limthongkul, C. Napaswad, P. Kongsaree, S. Prabpai, T. Jaipetch, M. Pohmakotr, P. Tuchinda and V. Reutrakul, *Phytochemistry*, 2018, **147**, 68–79.
- 21 B. J. Zhang, W. W. Fu, R. Wu, J. L. Yang, C. Y. Yao, B. X. Yan, H. S. Tan, C. W. Zheng, Z. J. Song and H. X. Xu, *Planta Med.*, 2019, **85**, 444–452.
- 22 E. R. Sukandar, S. Kaennakam, K. Rassamee, T. Ersam, P. Siripong and S. Tip-Pyang, *J. Nat. Prod.*, 2019, **82**, 1312–1318.
- 23 I. M. J. Ng, S. H. Mah and C. L. L. Chua, *Nat. Prod. Res.*, 2021, **35**, 5409–5414.
- 24 S. N. Haerani, A. Raksat and K. Pudhom, *J. Nat. Med.*, 2021, **75**, 670–674.
- 25 S. Kaennakam, K. Mudsing, K. Rassamee, P. Siripong and S. Tip-pyang, *J. Nat. Med.*, 2019, **73**, 257–261.
- 26 S. Laphookhieo, W. Maneerat and S. Koysomboon, *Molecules*, 2009, **14**, 1389–1395.
- 27 R. Ge, H. Zhao, Q. Tang, K. Chandarajoti, H. Bai, X. Wang, K. Zhang, W. Ye, X. Han, C. Wang and W. Zhou, *Microbiol. Spectr.*, 2024, **12**, e01631–24.
- 28 D. I. S. P. Resende, P. Pereira-Terra, J. Moreira, J. Freitas-Silva, A. Lemos, L. Gales, E. Pinto, M. E. de Sousa, P. M. da Costa and M. M. M. Pinto, *Mol.*, 2020, **25**, 2405.
- 29 E. Schonbrunn, S. Betzi, R. Alam, M. P. Martin, A. Becker, H. Han, R. Francis, R. Chakrasali, S. Jakkaraj, A. Kazi, S. M. Sebti, C. L. Cubitt, A. W. Gebhard, L. A. Hazlehurst, J. S. Tash and G. I. Georg, *J. Med. Chem.*, 2013, **56**, 3768–3782.
- 30 A. Al Sheikh Ali, D. Khan, A. Naqvi, F. F. Al-Blewi, N. Rezki, M. R. Aouad and M. Hagar, *ACS Omega*, 2021, **6**, 301–316.
- 31 N. Khamto, L. Chaichuang, P. Rithchumpon, W. Phupong, P. Bhoopong, S. Tateing, W. Pompimon, N. Semakul, N. O. Chomsri and P. Meepowpan, *RSC Adv.*, 2021, **11**, 31433–31447.
- 32 S. Bondock, N. Alabbad, A. Hossan, M. M. Abdou, A. A. Shati, M. Y. Alfaifi, S. E. I. Elbehairi and N. M. Mohamed, *RSC Adv.*, 2024, **14**, 18838–18855.
- 33 Y. Li, J. Zhang, W. Gao, L. Zhang, Y. Pan, S. Zhang and Y. Wang, *Int. J. Mol. Sci.*, 2015, (16), 9314–9340.
- 34 D. Lim and N. C. J. Strynadka, *Nat. Struct. Biol.*, 2002, **9**, 870–876.
- 35 F. K. K. Li, F. I. Rosell, R. T. Gale, J. P. Simorre, E. D. Brown and N. C. J. Strynadka, *J. Biol. Chem.*, 2020, **295**, 2629–2639.
- 36 M. R. Guda, G. V. Zyryanov, A. Dubey, V. S. Munagapati and J. C. Wen, *Pharmaceuticals*, 2023, **16**, 1159.
- 37 D. Sychantha, D. J. Little, R. N. Chapman, G. J. Boons, H. Robinson, P. L. Howell and A. J. Clarke, *Nat. Chem. Biol.*, 2017, **14**, 79–85.
- 38 M. Masumi, F. Noormohammadi, F. Kianisaba, F. Nouri, M. Taheri and A. Taherkhani, *Int. J. Microbiol.*, 2022, **2022**, 9130700.
- 39 C. A. Lipinski, F. Lombardo, B. W. Dominy and P. J. Feeney, *Adv. Drug Deliv. Rev.*, 1997, **23**, 3–25.
- 40 D. F. Veber, S. R. Johnson, H. Y. Cheng, B. R. Smith, K. W. Ward and K. D. Kopple, *J. Med. Chem.*, 2002, **45**, 2615–2623.
- 41 Y. H. Wang, A. J. Hou, G. F. Zhu, D. F. Chen and H. D. Sun, *Planta Med.*, 2005, **71**, 273–274.
- 42 S. Arthan, P. Posri, S. Walunchapruk, T. Senawong and C. Yenjai, *RSC Adv.*, 2022, **12**, 17837–17845.



- 43 J. Suthiwong, J. Wandee, S. Pitchuanchom, P. Sojikul, V. Kukongviriyapan and C. Yenjai, *Med. Chem. Res.*, 2018, **27**, 2042–2049.
- 44 P. Siripong, J. Yahuafai, K. Shimizu, K. Ichikawa, S. Yonezawa, T. Asai, K. Kanokmedakul, S. Ruchirawat and N. Oku, *Biol. Pharm. Bull.*, 2006, **29**, 2070–2076.
- 45 M. Balouiri, M. Sadiki and S. K. Ibnsouda, *J. Pharm. Anal.*, 2016, **6**, 71–79.
- 46 O. artorn Rajachan, W. Lakornwong, S. Pitchuanchom, N. P. Suchaichit, J. Boonmak, S. Youngme, K. Kanokmedhakul and S. Kanokmedhakul, *Fitoterapia*, 2021, **152**, 104912.
- 47 G. M. Morris, H. Ruth, W. Lindstrom, M. F. Sanner, R. K. Belew, D. S. Goodsell and A. J. Olson, *J. Comput. Chem.*, 2009, **30**, 2785–2791.
- 48 A. Daina, O. Michielin and V. Zoete, *Sci. Rep.*, 2017, **7**, 1–13.

