



Cite this: *Environ. Sci.: Processes Impacts*, 2020, **22**, 663

Insights into origins and function of the unexplored majority of the reductive dehalogenase gene family as a result of genome assembly and ortholog group classification[†]

Olivia Molenda, ^a Luz A. Puentes Jácome, ^a Xuan Cao, ^b Camilla L. Nesbø, ^{ab} Shuiquan Tang, ^a Nadia Morson, ^c Jonas Patron, ^b Line Lomheim, ^a David S. Wishart^{bd} and Elizabeth A. Edwards ^{*ac}

Organohalide respiring bacteria (OHRB) express reductive dehalogenases for energy conservation and growth. Some of these enzymes catalyze the reductive dehalogenation of chlorinated and brominated pollutants in anaerobic subsurface environments, providing a valuable ecosystem service. *Dehalococcoides mccartyi* strains have been most extensively studied owing to their ability to dechlorinate all chlorinated ethenes – most notably carcinogenic vinyl chloride – to ethene. The genomes of OHRB, particularly obligate OHRB, often harbour multiple putative reductive dehalogenase genes (*rdhA*), most of which have yet to be characterized. We recently sequenced and closed the genomes of eight new strains, increasing the number of available *D. mccartyi* genomes in NCBI from 16 to 24. From all available OHRB genomes, we classified predicted translations of reductive dehalogenase genes using a previously established 90% amino acid pairwise identity cut-off to identify Ortholog Groups (OGs). Interestingly, the majority of *D. mccartyi* dehalogenase gene sequences, once classified into OGs, exhibited a remarkable degree of synteny (gene order) in all genomes sequenced to date. This organization was not apparent without the classification. A high degree of synteny indicates that differences arose from *rdhA* gene loss rather than recombination. Phylogenetic analysis suggests that most *rdhA* genes have a long evolutionary history in the Dehalococcoidia with origin prior to speciation of *Dehalococcoides* and *Dehalogenimonas*. We also looked for evidence of synteny in the genomes of other species of OHRB. Unfortunately, there are too few closed *Dehalogenimonas* genomes to compare at this time. There is some partial evidence for synteny in the *Dehalobacter restrictus* genomes, but here too more closed genomes are needed for confirmation. Interestingly, we found that the *rdhA* genes that encode enzymes that catalyze dehalogenation of industrial pollutants are the only *rdhA* genes with strong evidence of recent lateral transfer – at least in the genomes examined herein. Given the utility of the *RdhA* sequence classification to comparative analyses, we are building a public web server (<http://RDaseDB.biozone.utoronto.ca>) for the community to use, which allows users to add and classify new sequences, and download the entire curated database of reductive dehalogenases.

Received 23rd December 2019
Accepted 20th February 2020

DOI: 10.1039/c9em00605b

rsc.li/espi

Environmental significance

Organohalide-respiring bacteria (OHRB) including *Dehalococcoides mccartyi*, *Dehalobacter restrictus* and *Dehalogenimonas* are important biocatalysts of a wide variety of dehalogenation reactions. An OHRB genome can harbour over 30 different copies of reductive dehalogenase-encoding genes and public databases currently contain thousands of representatives, yet fewer than 1% of all sequences have a characterized function. We used an amino acid sequence similarity-based method to classify all publicly available reductive dehalogenase sequences from closed genomes or metagenome-assembled genomes. Comparative phylogenetics combined with classification gave us new insights into origins and function of the unexplored majority of the reductive dehalogenase gene family. Given the utility of the approach, we created a public web server (<http://RDaseDB.biozone.utoronto.ca>) with a curated database of reductive dehalogenases and a classification tool to easily situate newly identified sequences.

^aDepartment of Chemical Engineering and Applied Chemistry, University of Toronto, Ontario, Canada. E-mail: elizabeth.edwards@utoronto.ca

^bDepartment of Biological Sciences, Faculty of Science, University of Alberta, Edmonton, Canada

^cDepartment of Cell and Systems Biology, University of Toronto, Ontario, Canada

^dDepartment of Computing Science, University of Alberta, Edmonton, Canada

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/c9em00605b

Introduction

Exposure to toxic or carcinogenic halogenated organic compounds, particularly chlorinated aliphatic hydrocarbons (CAHs), *via* groundwater and soil is of public concern.¹ While biological transformation processes proceed to some extent



naturally, these processes can be very effectively accelerated.² Natural attenuation, biostimulation, and bioaugmentation using mixed microbial consortia capable of reductive dechlorination are now commonly used to remediate sites contaminated with CAHs. Biologically-mediated reductive dechlorination of chlorinated pesticides was reported as early as 1967.³ Dechlorination of CAHs such as perchloroethene (PCE) to less chlorinated ethenes under methanogenic conditions was first reported in 1983.⁴ The idea to deploy specific bacteria to degrade CAHs became more viable after 1987, upon the discovery of anaerobic mixed cultures that could dechlorinate the last and most toxic daughter product, vinyl chloride, from PCE stepwise dechlorination⁵ and the subsequent isolation of *Dehalococcoides* as the responsible organism.^{6,7} At present, the efficacy of *in situ* bioaugmentation to transform contaminants such as PCE and trichloroethene (TCE), and daughter products *cis*-dichloroethene (cDCE) and vinyl chloride (VC) to ethene is well established.^{8,9}

Organohalide respiring bacteria (OHRB) conserve energy for growth from cleavage of carbon–halogen bonds. These bacteria use halogenated organic compounds as terminal electron acceptors in their metabolism. DCB-1,¹⁰ later named *Desulfovimonile tiedjei*¹¹ was the first characterized OHRB (~1986) capable of growing with 3-chlorobenzoate as an electron acceptor. OHRB come from many different phyla including Deltaproteobacteria (*Geobacter*, *Desulfuromonas*, *Anaeromyxobacter*, *Desulfoluna*, *Desulfomonile* and *Desulfovibrio*), Epsilonproteobacteria (*Sulfurospirillum*), Betaproteobacteria (*Shewanella* and *Comamonas*), Firmicutes (*Dehalobacter* and *Desulfitobacterium*) and Chloroflexi (*Dehalococcoides*, *Dehalogenimonas* and *Dehalobium*)¹² – see recent reviews.^{13–16} The success of bioremediation is largely due to the discovery of highly substrate-specific OHRB that link dehalogenation with growth, and thus whose populations increase as dehalogenation proceeds. These OHRB express respiratory reductive dehalogenases that can be categorized as part of a quinone-dependent or quinone-independent electron transport chain.¹⁴ The Dehalococcoidia, including *Dehalococcoides* and *Dehalogenimonas*, are quinone-independent, obligate organohalide respiring bacteria for which no other growth supporting electron acceptors are known. Characterized members of the Dehalococcoidia are capable of using more difficult-to-dechlorinate CAHs that only have one or two chlorine substituents. This ability of the Dehalococcoidia is key to complete detoxification. The enrichment culture studied herein (called KB-1) is routinely used for bioaugmentation and contains multiple organohalide-respiring bacteria (OHRB), including *Dehalococcoides mccartyi*, *Dehalobacter restrictus* and *Geobacter lovleyi*, which, in concert, are capable of complete dechlorination of many chlorinated ethenes, ethanes and benzenes.^{8,17} From the lens of geologic time, OHRB are phylogenetically diverse microbes of ancient origin that participate not only in detoxification of man-made contaminants, but also in the global halogen cycle, recycling a multitude of naturally-occurring organohalides.^{12,18,19}

Many OHRB genomes contain multiple distinct genes predicted to be reductive dehalogenases, termed *rdhA* genes¹⁶ sometimes accounting for a significant proportion of their

genomes. For example, *D. mccartyi* have remarkably small genomes (~1.4 Mbp) yet some strains contain up to 37 *rdhA* genes. *D. mccartyi* contain a core syntenic region encoding for “housekeeping” genes such as biosynthesis of amino acids, cell components, transcription/translation, nutrient transport and energy conservation.^{20,21} The differences between strains in this region are no more than single nucleotide polymorphisms (SNPs) with average nucleotide identity (ANI) of 85–99%.²² This stable core genome is interrupted by two variable regions flanking the origin of replication commonly referred to as High Plasticity Regions (HPRs, individually HPR1 and HPR2).²¹ HPRs show signs of both lateral gene transfer and homologous recombination including repeats, duplication events, insertion sequences, genomic islands, and phage-related sequences. The majority of reductive dehalogenases in *D. mccartyi* are located on these HPRs. In addition, several characterized dehalogenases are within genomic islands with evidence of site-specific integration, such as the VC reductase genes *vcra* and *bvcA*,²³ and TCE reductase gene *tceA*.²⁴ It is generally thought that recombination events have allowed *D. mccartyi* to adapt to naturally-occurring and anthropogenic halogenated compounds.^{21,23}

Reductive dehalogenase genes occur in an operon containing the *rdhA*, *rdhB*, and sometimes, additional genes. The *rdhA* gene encodes the catalytically active enzyme (RdhA or RDase if characterized) and the *B* gene encodes a membrane anchor protein containing two or sometimes three transmembrane helices.^{13–15,25,26} *D. mccartyi* RdhA and RdhB subunits are part of a unique protein complex which is currently considered to be a fully functional stand-alone respiratory chain without quinone or cytochrome involvement.²⁷ Other than RdhAB, the protein complex includes the organohalide respiration molybdoenzyme (OmeA), formally called the complex iron–sulfur cluster molybdoenzyme (CISM),²⁷ found to have hydrogenase activity,²⁸ its putative membrane anchor (OmeB), hydrogen uptake hydrogenase (HupX) with a [NiFe] large subunit (HupL) and iron–sulfur containing small subunit (HupS).

RdhAs are identified based on the presence of two motifs: a twin-arginine TAT membrane export sequence (RRXXFXK), and an eight-iron ferredoxin cluster binding motif (CXXCXXCXXXCP). RDases can have one or two iron–sulfur binding motifs. In certain RDases, such as *Dehalogenimonas lykanthroporepellens* BL-DC-9 DepA, a cobalamin binding motif can be identified (DXHXXGSXLGG).^{29–31} A corrinoid molecule is a critical co-factor for folding and function of RdhA proteins.^{32–34} However, outside of motif regions, RdhAs can have as little as 30% full-length pairwise identity (PID) even when found in the same genome. While many putative dehalogenases sequences can be found in any particular genome, only a select few have been found expressed in response to different CAHs.¹⁴ As a result, without a universal functioning heterologous expression system, only a few dehalogenases have been characterized out of the thousands of sequences in the National Center for Biotechnology Information (NCBI) database. To help classify newly found putative dehalogenases with unknown functions, Hug *et al.* developed a classification system in 2013, grouping RdhAs with upwards of 90% amino acid PID into sets



of orthologous sequences (*i.e.* Ortholog Groups, or OGs) which most likely share similar function.^{19,35} Since 2013, two crystal structures of RdhA proteins have been solved,^{36,37} confirming previous features and providing more resolution to the critical substrate-binding domain.

The first objective of this study was to identify and sequence the different strains of OHRB in a set of related KB-1 subcultures amended with different chlorinated electron acceptors. We had previously noted that different electron acceptors enriched for different strains of *D. mccartyi*.³⁸ From three cultures, 8 new *D. mccartyi* genomes were closed, increasing the total number of publicly available *D. mccartyi* genomes to 24. As well, one new *Dehalobacter restrictus* genome was closed increasing the number publicly available to 4 (as of December 2019). The second objective was to compare these genomes to identify key trends in genome evolution, homologous recombination, and lateral gene transfer. We first specifically investigated the evolution of reductive dehalogenase genes in *D. mccartyi* since most genomes were available for comparison. We then tried to extend the analysis to *Dehalogenimonas*, a close relative of *Dehalococcoides*, and finally to *Dehalobacter* – a distinct genus in the Firmicutes. The Ortholog Group classification method was extremely useful in this analysis, which led us to the development of a public web server to make it easier for the community to compare and classify new sequences (<http://RDaseDB.biozone.utoronto.ca>). The user can also download part of or the entire manually curated reductive dehalogenase sequence database for further analyses.

Methods

Enrichment cultures analysed

The KB-1 cultures were originally enriched from aquifer materials at a TCE contaminated site in southern Ontario as previously described.^{8,9,39} A commercial version of the culture is maintained at SiREM (Guelph, ON) for bioaugmentation. The parent enrichment culture maintained at the University of Toronto, referred to as KB-1/TCE-MeOH, has been maintained since 1998 in a 2 L glass bottle (~1.3 L culture with 0.7 L headspace) in anaerobic mineral medium amended with 100 mg L⁻¹ TCE as electron acceptor and methanol (MeOH) as electron donor, added at 5× the electron equivalents (eeq.) required for complete dechlorination, as previously described.^{39–41} TCE is completely dechlorinated to ethene prior to refeeding, approximately every 2–4 weeks. Acetogens in the mixed culture ferment added methanol to hydrogen and acetate required by *D. mccartyi*. In 2001, a sub-culture was created with a 2% transfer into pre-reduced anaerobic mineral medium and maintained on VC and hydrogen as described in³⁹ and is referred to as KB-1/VC-H₂. This 200 mL VC enrichment was maintained in a 250 mL glass bottle sealed using a Teflon mininert cap and was amended with 55 mg L⁻¹ VC (supplied as 5 mL of pure VC gas) and 5× eeq. hydrogen gas (supplied as 5 mL 80% H₂ : 20% CO₂, Praxair). The KB-1/VC-H₂ enrichment bottle was also amended with 0.5 mM sodium acetate as a carbon source every ten feedings. In 2003, a 1,2-dichloroethane (DCA)-fed enrichment culture was created (KB-1/1,2-

DCA-MeOH culture) fed with 250 mg L⁻¹ 1,2-DCA and 5× eeq. MeOH, in a 2 L glass Pyrex bottle containing 1.3 L culture with 0.7 L headspace. Approximately every 6 months, half of each culture was removed and substituted with fresh pre-reduced anaerobic mineral medium to replenish vitamins and buffer. In all enrichment cultures, chlorinated ethenes/ethanes, methane and ethene concentrations were monitored using gas chromatography with an FID detector (Agilent 7890A GC system, G1888 auto-sampler, helium used as carrier gas, packed inlet, Agilent GSQ-Plot column 0.53 mm × 30 mm), calibrated with external standards.

Metagenomic sequencing and genome assembly

DNA for metagenome sequencing was extracted from larger samples (40–615 mL) taken from the three functionally-stable enrichment cultures described above: KB-1/VC-H₂ (40 mL sample), KB-1/TCE-MeOH (500 mL sample), KB-1/cDCE-MeOH (300 mL sample) and KB-1/1,2-DCA-MeOH (615 mL sample). Extractions were conducted between February and May 2013. Cultures were filtered using Sterivex™ filters (Millipore 0.2 µm) and the DNA was extracted using the CTAB method (JGI bacterial genomic DNA isolation using CTAB protocol v.3). DNA was sequenced at the Genome Quebec Innovation Sequencing Centre using Illumina HiSeq 2500 technology. Paired-end sequencing with an insert size of ~400 bp and read length of ~150 bp provided roughly 50 million reads per culture. Additional mate-pair sequencing with insert size of ~8000 bp and read length of ~100 bp was conducted for the KB-1/TCE-MeOH and KB-1/1,2-DCA-MeOH cultures where we had more DNA (>10 µg each). For metagenomic sequencing using short-read Next Generation Sequencing (NGS), we have demonstrated the utility of integrating long-insert mate-pair data to resolve challenges in the metagenomic assembly, especially those related to repeat elements and strain variation.⁴² Raw sequences were trimmed with Trimmomatic⁴³ to remove bases of low quality and to remove adapters.

The *D. mccartyi* genomes were assembled in six steps as described in supplemental methods. Quantitative PCR to quantify the abundance of specific *rhdA* genes and 16S rRNA amplicon sequencing for relative abundances of different organisms (methods provided in ESI†) were also performed to guide the metagenomic assembly (illustrated in Fig. S1†). In all cases, multiple genomes could be closed (assembled into a single circular scaffold without gaps) from a single enrichment culture because the different populations of *D. mccartyi* were present at different abundances (as inferred from read depth) at the time of sampling. Two complete genomes, each containing a vinyl chloride reductase gene (*vcrA*), were closed from the KB-1/VC-H₂. The naming convention developed identifies culture (KB-1) electron acceptor (in this case vinyl chloride or VC) and relative abundance (number 1 for highest abundance, 2 for next and so on). Thus, the *D. mccartyi* strains from culture KB-1/VC-H₂ are named strains KBVC1 and KBVC2. Three *D. mccartyi* genomes containing the vinyl chloride reductase *bvcA* were closed from the KB-1/1,2-DCA enrichment culture referred to as strains KBDCA1, KBDCA2 and KBDCA3



and one *Dehalobacter restrictus* genome named strain 12DCA. A *D. mccartyi* complete genome containing a *vcrA* gene was assembled from KB-1/TCE-MeOH culture, named strain KBTCE1. Additionally, two genomes containing trichloroethene reductase *tceA* genes were assembled from the KB-1/TCE-MeOH culture, strains KBTCE2 and KBTCE3. The genomes were annotated using the RAST⁴⁴ and BASyS⁴⁵ servers; results were manually inspected and corrected where required and automatically annotated after submission to NCBI by PGAP.⁴⁶ Additional searches for conserved domains were conducted using NCBI conserved domain search (*E*-value threshold of 0.01). The origin of replication was identified using OriLoc in R.⁴⁷

Alignments and phylogenetic trees

A concatenated gene alignment was created by aligning a set of 109 homologous genes found in genomes of *Dehalococcoides mccartyi* (22 strains), *Dehalogenimonas lykanthroporepellens*, *D. alkenigignens*, *Dehalogenimonas* sp. WBC-2, and a Chloroflexi out-group *Sphaerobacter thermophilus*. Outgroups were used to better visualize branching and to improve the estimate of the number of mutations along each branch. Outgroups were selected to be as closely related to in-group as reasonably possible without being part of the in-group. Homologous genes were identified using reciprocal BLASTp and OrthoMCL followed by manual inspection using the Get Homologues package.⁴⁸ Each gene was aligned using MUSCLE v3.8.3.1⁴⁹ with default settings. The alignments were concatenated to create one long alignment (138 334 bp long alignment, 26 sequences, 83% average pairwise identity, gaps included). A maximum likelihood (ML) tree was built using the RAxML⁵⁰ plugin in Geneious 8.1.8⁵¹ with the GTR substitution model and 100 bootstrap replicates. The best scoring ML tree was chosen as the final tree.

Five-hundred and fifty-one reductive dehalogenase sequences were selected from *Dehalococcoidia* to create an

amino acid and a separate nucleotide phylogenetic tree using Geneious 8.1.8. These included all *rdhA* which have been assigned an Ortholog Group (OG) number from the RDase database, all *rdhA* from three species (*D. lykanthroporepellens*, *D. alkenigignens*, and *Dehalogenimonas* sp. WBC-2) and a reductive dehalogenase from *Desulfoluna spongiiphila* as the out-group. *D. spongiiphila* is a reductive dehalogenating, anaerobic, sulfate-reducing bacterium isolated from a marine sponge.⁵² RDases which cannot be classified into a group at this time have still been assigned a number in the database along with the letter "S" to indicate that this sequence is a singleton. A second tree was created using the same technique from all *Dehalobacter* reductive dehalogenase gene and protein sequences, and a third tree with a combination of previously functionally characterized RDases from *Dehalococcoides*, *Dehalobacter* and *Dehalogenimonas*. Alignments and tree-building were conducted using MUSCLE and RAxML as described above. FigTree 1.4.2 was used to visualize and further edit the tree to generate figures in this study (<http://tree.bio.ed.ac.uk/software/figtree/>).

Time since divergence analysis and detection of positive selection

The time since divergence was calculated using the number of mutations identified from phylogenetic concatenated gene alignments and OG *rdhA* gene alignments to date (in substitutions per site). Substitutions per site identified from *rdhA* OG alignments were compared to see whether *rdhA* could have originated before or after (1) the divergence of *Dehalococcoidia* into *Dehalococcoides* and *Dehalogenimonas* and (2) the divergence of *Dehalococcoides* into three clades: Pinellas, Cornell and Victoria.

We compared the ratio of non-synonymous (K_a) to synonymous substitutions rates (K_s) in *rdhA* OG alignments because the K_a/K_s ratio can identify positive selection. If $K_a/K_s < 1$, then non-synonymous mutations are either neutral or deleterious, whereas if $K_a/K_s > 1$, then positive selection occurred.⁵³ K_a/K_s

Table 1 General features of *Dehalococcoides mccartyi* and *Dehalobacter restrictus* genomes closed from KB-1 trichloroethene (TCE), 1,2-dichloroethane (1,2-DCA) and vinyl chloride (VC) enrichment cultures

	<i>Dehalococcoides</i>								<i>Dehalobacter</i>
	KBVC1	KBVC2	KBTCE1	KBTCE2	KBTCE3	KBDCA1	KBDCA2	KBDCA3	12DCA
Genome size (mbp)	1.39	1.35	1.39	1.33	1.27	1.43	1.39	1.34	3.05
G + C content (%)	47.3	47.2	47.3	49.1	49.3	47.4	47.5	47.6	44.5
Protein coding genes	1468	1432	1451	1381	1319	1496	1462	1404	2906
Hypothetical genes (%)	31.1	30.2	30.3	29.1	26.8	32.8	31.9	29.1	19.1
tRNA	47	48	47	45	45	47	46	46	52
CRISPR-Cas genes	7	0	0	0	0	0	0	6	0
Nitrogen fixation genes	0	0	0	9	9	0	0	0	0
Sub-group/Clade	Pinellas	Pinellas	Pinellas	Cornell	Cornell	Pinellas	Pinellas	Pinellas	n/a
Electron acceptor provided to culture	VC	VC	TCE	TCE	TCE	1,2-DCA	1,2-DCA	1,2-DCA	1,2-DCA
<i>rdhA</i> genes	22	16	16	5	5	7	7	9	22
Identifiable <i>rdhA</i> ^a	<i>vcrA</i> , <i>pceA</i>	<i>vcrA</i> , <i>pceA</i>	<i>vcrA</i> , <i>pceA</i>	<i>tceA</i>	<i>tceA</i>	<i>bvcA</i>	<i>bvcA</i>	<i>bvcA</i>	Unknown
NCBI accession number	CP019968	CP019969	CP019999	CP019865	CP019866	CP019867	CP019868	CP019946	CP046996

^a Identifiable *rdhA* indicates the presence of a *rdhA* gene whose protein product was characterized in a different study. It is not known whether these *rdhA* are expressed by these specific strains.



ratios were calculated using the PAML package.^{54,55} Only OGs with five or more *rdhA* were used for the analysis since using fewer than five sequences is not recommended.⁵⁶ Two models were run: one which described neutral or nearly neutral evolution and one which additionally accounted for adaptive evolution. The χ^2 test was used to determine whether the K_a/K_s gene ratios were significantly greater than 1, suggesting positive selection.

Nucleotide sequence accession numbers

The KB-1 sequencing bioproject can be accessed in the National Center for Biotechnology Information (NCBI) database using bioproject no. PRJNA376155. *Dehalococcoides mccartyi* KB-1 closed genome nucleotide accession numbers in NCBI are: strain KBDCA1 CP019867, strain KBDCA2 CP019868, strain KBDCA3 CP019946, strain KBVC1 CP019968, strain KBVC2 CP019969, strain KBTCE1 CP01999, strain KBTCE2 CP019865,




Fig. 1 Overview of reductive dehalogenase genes found in eight *D. mccartyi* genomes originating from different geographical locations. Top left shows the general architecture of *D. mccartyi* genomes. Genomes are identified by strain name and country of origin indicated beneath. Reductive dehalogenase homologous genes (*rdhA*) are marked on each genome, coloured by Ortholog Group (OG). Genomes are aligned (dotted lines indicating gaps) to visualize OGs. Genome high plasticity region (HPR), individually HPR1 and HPR2.



and strain KBCTCE3 CP019866. KB-1 *Dehalobacter restrictus* 12DCA closed genome: CP046996. All metagenomic raw reads are available in NCBI short-read archive (SRA) accession no. SRR11085567-71. 16S rRNA amplicon sequences have been deposited in NCBI SRA accession no. SRP144609.

Results & discussion

General features of *Dehalococcoides* and *Dehalobacter* genomes from KB-1 enrichment cultures

Eight complete genomes of *D. mccartyi* strains and one *Dehalobacter restrictus* strain were assembled (Fig. S1† for assembly methodology) and annotated from three different enrichment cultures (Table 1) using Illumina mate-pair and paired-end metagenomic sequencing in combination with 16S rRNA amplicon sequencing and qPCR of functional *rdhA* to guide assembly (Fig. S2†). In all KB-1 enrichments, we found lower abundance, fragmented *Dehalococcoides* and *Dehalobacter* contigs which could not be assembled into complete genomes indicating that each of the three KB-1 sub-cultures must have at least one additional strain which could not be assembled totalling at least eleven strains of *Dehalococcoides mccartyi* and at least two strains of *Dehalobacter restrictus*. The strains whose genomes were assembled and closed have been named based on contaminant/electron acceptor amended and a number to indicate rank abundance with respect to other strains found in that same enrichment culture (Table 1). *D. mccartyi* strains, overall, share at least 98% nucleotide sequence identity of the 16S rRNA gene⁵⁷ (Fig. S3†) and fall into three clades known as the Pinellas, Victoria and Cornell groupings.⁵⁸ Most KB-1 strains (KBVC1, KBVC2, KBDCA1, KBDCA2, KBDCA3 and KBTCE1) fall into the Pinellas clade (which also contains strains CBDB1,²⁰ BTF08,⁵⁹ DCMB5,⁵⁹ 11a5,⁶⁰ WBC-2,³⁵ GT,⁶¹ IBARAKI⁶² and BAV1), while only two KB-1 strains (KBTCE2 and KBTCE3) fall into the Cornell clade (containing strain 195, MB⁶³ and CG4⁶⁴) (ESI Fig. S3†). All genomes have a clear GC skew with one origin of replication, and the majority of reductive dehalogenase genes are found flanking the origin of replication, primarily coded on the leading strands.

Strains KBVC1 and KBDCA3 contain complete CRISPR-Cas systems⁶⁵ that are similar to those in published strains 11a,⁶⁶ CBDB1,²⁰ DCMB5⁵⁹ and GT.⁶¹ Cornell-clade strains KBTCE2 and KBTCE3 contain nitrogen fixation genes similar to type strain 195.⁶⁷ The number of putative reductive dehalogenase genes (*rdhA*) among all strains varies from five to twenty-two (Table 1). Strain KBTCE3 has the smallest genome of all known *D. mccartyi* (1.27 Mbp), with the fewest number of *rdhA* (only five in both KBTCE3 and KBTCE2) and has highest GC content (49.3% in KBTCE3) (Table 1). We consider all eight KB-1 *D. mccartyi* genomes to be representative of independent strains because the number of accumulated SNPs and genomic rearrangements has clearly resulted in functional changes and differential growth patterns as inferred from the observed different abundances in the same KB-1 sub-culture.

The KB-1 1,2-DCA enrichment culture contains a population of *Dehalobacter*, as determined by 16S rRNA amplicon analysis (Fig. S2†). We closed a genome from this population whose 16S rRNA gene sequence shares 99.9% nucleotide PID with

Dehalobacter restrictus PER-K23⁶⁸ and 99.8% average nucleotide identity (ANI). Several rearrangements and insertions have occurred which makes the whole genome alignment of the two strains 76.9% nucleotide PID. This KB-1 *Dehalobacter* genome will be further referred to as *Dehalobacter restrictus* strain 12DCA.

Reductive dehalogenase genes in *Dehalococcoides mccartyi* and conserved synteny

Hug *et al.* (2013)¹⁹ developed a classification system for reductive dehalogenases where sequences were assigned to Ortholog Groups (OGs) in an attempt to cluster *rdhA* sequences into groups with similar activity on specific halogenated electron acceptors.³⁵ Previous studies identified 32 distinct *rdhA* genes in the KB-1/TCE-MeOH culture,^{38,69,70} with two being identical to genes found previously in *D. mccartyi* CBDB1. Although 104 *RdhA* were found in this study, only three new OGs could be described, and the remaining *RdhA* sequences fell into previously identified groupings (Table S2†). As more *D. mccartyi* genomes are closed, fewer new dehalogenases are discovered that do not belong to an existing OG (Fig. 2); as of December 2019, a total of 84 OGs from *D. mccartyi* have been identified. We further applied this classification to 144 *Dehalogenimonas* and 356 *Dehalobacter* reductive dehalogenases. A similar trend occurred with *Dehalobacter restrictus* *RdhAs*, but not with *Dehalogenimonas*, because too few *Dehalogenimonas* genomes are available, and those that are available, are not from the same species (Fig. 2). With the strict 90% amino acid full-length alignment identity cut off, resulting ortholog groups were found to be genus-specific for *Dehalococcoides* and *Dehalogenimonas*.

Whole genome alignments of *D. mccartyi* generally result in strong core-genome region alignments (~90% nucleotide pairwise identity), with poor alignments of two regions flanking the origin, deemed the high plasticity regions (HPR) (<30% pairwise nucleotide identity) suggesting that HPRs are unique per strain. Analysing twenty-four closed genomes has revealed that most of the *rdhA* in HPRs are actually syntenous, and that poor alignments are a result of gene loss of specific *rdhA* genes (Fig. 1 and




Fig. 2 The number of new Ortholog Groups (OGs) of reductive dehalogenases (RDases) found with each new *Dehalococcoides mccartyi*, *Dehalobacter restrictus*, or *Dehalogenimonas* (various species) genome or Metagenome Assembled Genome (MAG) available from NCBI or IMG.






Fig. 3 Synteny in *Dehalococcoides mccartyi* genomes. (a) Order of *rdhA* found in high plasticity region one (HPR1) in 22 *D. mccartyi* genomes labeled by strain name. *rdhA* are labeled by Ortholog Group (OG) number. *rdhA* of the same OG share >90% amino acid pairwise identity. Unpaired *rdhA* (i.e. singletons) are not included due to space limitations (see Table S2† for all genomes and *rdhA*). Green (leading strand) and red (lagging strand) accents indicate DNA strand clockwise from the *oriC*. The majority of *rdhA* are on the leading strand. * found in all *D. mccartyi* strains. (b) Order of *rdhA* found in high plasticity region two (HPR2) in 22 *D. mccartyi* genomes labeled by strain name. *rdhA* are labeled as in panel (a) except colors reflect different leading strands. Green (lagging strand) and red (leading strand) accents indicate DNA strand anticlockwise from the *oriC*. The majority of *rdhA* are on the leading strand. HPR2 starts at tRNA-Leu/Arg/Val at approximately 1.2 Mbp from the origin. OG 35 is only in strain 195 after tRNA-Ala, 1.3 Mbp from the origin. +OG known to be expressed during starvation.

3). Therefore, most strain HPRs are much more similar than previously thought.²¹ A conserved sequential order of *rdhA* exists in all *D. mccartyi* strains (Fig. 3a HPR1, Fig. 3b HPR2, with accession numbers Table S2†). For example, at the end of HPR2, the most common order of RdhA OGs is 5'-40:30:34:11:10:17-3'. Some strains have all of these OGs (CG5, BTF08, CBDB1, KBVC1, GT, KBVC2, KBTCE1, 11a5) while certain strains only have a select few, such as IBARAKI 5'-30:11-3', CG4 5'-10:17-3', or GY50, that have 5'-40:30:10:17-3' (Fig. 3B). The visualization of this synteny would not have been possible without grouping of *rdhA* sequences, nor without the numerous complete genomes currently publicly available in NCBI.

Trends in *Dehalococcoides* *rdhA*: acquisition, loss and evolution

There are two evolutionary time points which can be identified from the concatenated homologous gene alignment in the *Dehalococcoides* and *Dehalogenimonas* phylogenetic tree (Fig. 4). First, an estimate of the point of divergence of *Dehalococcoides* and *Dehalogenimonas* (Dehalococcoidia speciation) and secondly the separation of *Dehalococcoides* *mccartyi* into three distinct clades: the Pinellas, Cornell and Victoria (clade speciation). The clade separation in *D. mccartyi* is evident not only at the 16S rRNA level (Fig. S3†) but also in each homologous *rdhA* gene alignment

(Fig. 5). *D. mccartyi* strains that are in the same clade typically have the same SNPs. These divergence time points could also be seen in *rdhA* alignments within OG groups. Each OG *rdhA* alignment includes sequences with varying numbers of nucleotide substitutions. These substitutions are sometimes clade-specific, meaning that the OG was present prior to clade speciation, and the divergences occurred after the clade speciation event. Other OGs include sequences with very few substitutions, which are not clade-specific, suggesting more recent acquisition (Fig. 4 and 5 with specific examples; all OG analyzed in Fig. S4 and S5†).

Reductive dehalogenase sequences belonging to an OG with a long history with *Dehalococcoides* (before the clade speciation event, or even before the Dehalococcoidia speciation event, Tables S3 and S4†) occur syntenically and are typically preceded by MarR-type regulators, such as MarR-type regulator *Rdh2R* (cbdb1456), which was found to suppress downstream *rdhA* expression in CBDB1.⁷¹ Syntenic OG alignments have small (~ 0) K_a/K_s ratios meaning that syntenic OG *rdhAs* are not currently under selective pressure (Fig. 5). The first and most common trend observed from sequence classification and analysis is a consensus order of *rdhA* genes which are ancestral and not found expressed or synthesized at this time¹⁵ (summarized in Table 2). *Dehalococcoides* OGs whose level of divergence predates the Dehalococcoidia speciation event, such as OG 15 and OG 34 have homologous representatives in




Fig. 4 Phylogenetic tree created from an alignment of 109 concatenated core genes from *Dehalococcoides* *mccartyi* closed genomes and *Dehalogenimonas* closed genomes with chloroflexi *Sphaerobacter thermophilus* as out-group. Best scoring maximum likelihood tree is displayed. Bottom scale indicates timing of key events including clade separation in *D. mccartyi* and the Most Recent Common Ancestor (MRCA) of several dehalogenases listed by name or by Ortholog Group (OG) if uncharacterized. *D. mccartyi* clades are highlighted in common colour. Scale indicates number of substitutions per site. Double cross-hatching indicates this branch was reduced in length by half for visualization purposes.






Fig. 5 Phylogenetic tree of reductive dehalogenase genes which belong to Ortholog Group (OG) 5, 13, 71, 15 and 34. Best scoring maximum likelihood tree displayed. Scale shows number of substitutions per site. A trichloroethene dehalogenase from *Desulfitobacterium* (Dhaf_0696) used as out-group. The *rdhA* in the tree are coloured by clade: blue (Pinellas), green (Victoria), red (Cornell) and identified with strain name followed by locus tag of *rdhA* in parentheses. *Dehalogenimonas* *rdhA* similar to these OG groups are shown in black.

Dehalogenimonas, including *Dehalogenimonas* sp. strains WBC-2, GP, *D. alkenigignens* and *D. formicexedens*.⁷²

For some OGs (Table 2), there is evidence for positive selection on *RdhA* according to K_a/K_s ratios. For example, we found that OGs 13 and 71 occur in the exactly the same position in different genomes, and that genomes contain either one or the other, but not both. OG 13 (Fig. 3) is found in both Pinellas and Victoria clades and is highly similar to OG 71 which is only in Cornell strains (Fig. 3). OG 13 dehalogenases have small K_a/K_s ratios while *RdhAs* in OG 71 have high ratios, suggesting positive selection. The conditions experienced by Cornell strains likely led to the use and specialization of this dehalogenase. Therefore, the second trend in dehalogenases in *D. mccartyi* is the evolution of new dehalogenases from existing ones (Summarized in Table 2).

Some *rdhA* are thought to have been acquired laterally by *D. mccartyi*, including *tceA* (OG 5, Fig. 4),²⁴ *bvcA* (OG 28),²¹ and *vcrA*

(OG 8)^{21,23} due to their location on genomic islands, and very high nucleotide and amino acid sequence conservation among strains from distant geographical locations and sometimes different clades. Additionally, *cbrA* (OG 53), *mbrA* (OG 52) *pcba* (OG 35) and *pteA* (OG 16) currently also appear to have very few mutations and occur in the vicinity of mobile elements. McMurdie *et al.* (2011)²³ established that *vcrA* nucleotide polymorphisms indicate that it was acquired by *Dehalococcoides* approximately ~1000 years ago, after the *Dehalococcoides* clade speciation event, meaning *vcrA* looks identical across all clades. All *rdhA* predicted to occur on mobile elements also have few mutations within OGs like *vcrA*, suggesting that they were acquired in a similarly recent time frame (Table S3†). Interestingly, all of these mobile OGs have members which have been biochemically characterized due to their association with dehalogenation of industrial pollutants and growth on these substrates in a laboratory setting.



Table 2 Summary of common features differentiating Ortholog Groups (OGs) of reductive dehalogenase sequences from *D. mccartyi*^a

Type	OG #s	General features	Protein information
1	10, 11, 15, 17, 19, 21, 22, 23, 30, 48, 55, 60, 66	Present in more than one clade Found in <i>D. mccartyi</i> before clades separated. Synteny within genomes conserved. OG 15 illustrated in Fig. 5	OG 15 known to be expressed by <i>D. mccartyi</i> during starvation ^{101,102} OG 23 found in all <i>D. mccartyi</i> genomes at time of analysis
2	13/71, 18/40, 32/69, 33/38	Pairs of OGs which have recently diverged from one another. Location of <i>rdhA</i> is syntetic across all genomes. OG 13 and 71 illustrated in Fig. 5	No characterized members at time of study
3	5, 8, 16, 28, 52, 53, 35	Highly similar PID in different strains, regardless of clades. Often associated with mobile genes or on a genomic island. OG 5 illustrated in Fig. 5	5 – <i>TceA</i> ¹⁰³ 8 – <i>VcrA</i> ¹⁰⁰ 16 – <i>PteA</i> ⁶⁰ 28 – <i>BvcA</i> ¹⁰⁴ 52 – <i>Mbra</i> ⁶³ 53 – <i>CbrA</i> ¹⁰⁵ 35 – <i>PcbA</i> ⁶⁴
4	26, 34, 57	Duplicated within the same genome. OG 34 illustrated in Fig. 5	No characterized members at time of study

^a PID – pairwise identity.

Some OGs display movement within a genome, rather than between genomes. Victoria clade strains VS, CG1 and GY50 contain examples of an OG occurring twice in the same genome suggesting that a duplication event occurred. Duplicated OGs occur in different HPRs, and not in tandem (Fig. 3). In one case, *pceA* (OG 30) appears to have moved within a genome and without duplication. OG 30 is syntetic in HPR2 in 14 strains, with the exception of strain 195 where it occurs in HPR1 (Fig. 3). In strain 195, *pceA* is located near a serine recombinase which possibly mediated its movement. Additionally, the strain 195 *pceA* is present without its usual upstream transcriptional regulator, which was thought to be responsible for *rdhA* regulation.^{71,73} Transcriptomic studies show that strain 195 will continue to produce high transcript levels of *pceA* regardless of starvation or TCE amendment.⁷⁴ Furthermore, strain 195 is the only strain to produce *pceA* in the presence of PCE, where other strains, such as CG5, transcribe *pceA* in the presence of multiple PCB congeners⁶⁴ and CBDB1 was found to transcribe *pceA* in the presence of 2,3-dichlorophenol (DCP).^{75,76} Thus, the third trend we observed in dehalogenase evolution is mobility, either within or between strains (summarized as types 3 (lateral transfer) and 4 (within genome duplication or rearrangement) in Table 2).

Gene loss and genomic streamlining in *Dehalococcoides mccartyi*

D. mccartyi genomes are unique in that they are among the smallest genomes found in free-living bacteria (avg. 1.4 Mbp and 1451 protein-coding genes). A common theme among all small free-living prokaryotes is their high niche specialization and low-nutrient level environments.⁷⁷ *D. mccartyi* genomes probably required an extensive period of time to become as specialized and as small as they currently are. Wolf *et al.* (2013) proposed a theory that gene loss is equally or even more

important than lateral gene transfer in shaping genomes.⁷⁸ The high level of synteny and number of mutations found in Ortholog Groups of dehalogenases supports this theory in *D. mccartyi*. Given that all dehalogenases capable of dehalogenating industrial contaminants also appear to be mobile, it is possible that anthropogenic releases of organohalides have caused *D. mccartyi* genomes to enter a period of complexification involving sharing select *rdhA*. The exchange of key reductive dehalogenases among *D. mccartyi* can be akin to the recent dissemination of antibiotic resistance genes in the natural environment.^{79–81} Transfer mechanisms may even be related considering the homology observed between the integration module of the *vcrA* containing genomic island and the Staphylococcal cassette chromosome that carries the broad spectrum beta-lactam resistance gene.^{23,82}

Reductive dehalogenases in other obligate dechlorinating organisms: *Dehalobacter* and *Dehalogenimonas*

We analyzed the reductive dehalogenases from other OHRB including *Dehalobacter* and *Dehalogenimonas* to extend the OG classification system beyond *Dehalococcoides*. In both cases, there are very few publicly available closed genomes, which limited our analysis. Members of the genus *Dehalobacter* are also organohalide-respiring bacteria that use hydrogen or formate as electron donor, acetate as carbon source, and organohalides as electron acceptors.^{83,84} *Dehalobacter* spp. have been shown to respire and/or be implicated in the reductive dechlorination of chlorinated methanes,⁸⁵ ethanes,^{86–88} ethenes,^{68,89} cyclohexanes,^{90,91} benzenes,^{92,93} phenols,⁹⁴ biphenyls,⁹⁵ and phthalides.⁹⁵ As of November of 2019, ten *Dehalobacter* genome assemblies, corresponding to strains PER-K23 (DSM 9455), CF, DCA, UNSWDHB, TeCB1, MCB1, 12DCB1, 14DCB1, E1, and FTH1 were publicly available NCBI, but only



three of these genomes are closed (strains PER-K23, CF, DCA) (Table S6† for all genomes analyzed in this study). Four additional metagenome-assembled-genomes of *Dehalobacter* from KB-1 subcultures are found in the Integrated Microbial Genomes & Microbiomes (IMG/M) system, corresponding to strains TCB1 (2775506732), TCB2 (2786546105), TCB3 (2821530269), and HCH1 (2823892816). Like *Dehalococcoides*, many *rdhA* genes (~10 to 30) can be found in each *Dehalobacter* genome, but only a subset of genes has been characterized (see Table S5†). As part of this study, a new closed *Dehalobacter restrictus* genome was submitted to NCBI, strain 12DCA (CP046996).

The *rdhA* genes of the 15 publicly available *Dehalobacter restrictus* genome assemblies were translated and classified into OGs. Like *Dehalococcoides mccartyi*, fewer new *Dehalobacter restrictus* dehalogenases were classified into new OGs with an increasing number of distinct genome assemblies (Fig. 2). A total of 80 OGs resulted from the classification of 356 *RdhA* protein sequences from 15 genomes (or metagenome assembled genomes/binned genomes), approaching saturation, as with

Dehalococcoides mccartyi sequences (Fig. 2). A full *Dehalobacter* RDase tree annotated by OG can be found in Fig. S6.† An initial examination of the distribution of OGs in closed *Dehalobacter* genomes as well as in some of the contigs from other *Dehalobacter* spp., (Fig. S7†) suggests that some *Dehalobacter* *RdhA* OGs also display synteny across genomes. As additional *Dehalobacter* complete and closed genomes become available, trends such as synteny and gene evolution will become easier to see.

With respect to *Dehalogenimonas*, to-date only six genomes have been closed or binned (*D. lykanthroporepellans*,⁹⁶ *Dehalogenimonas* sp. strain WBC-2,³⁵ *D. alkenigignens* strain IP3-3,⁹⁷ *D. alkenigignens* strain BRE15M, *D. formicexedens*⁹⁸ and *Dehalogenimonas* sp. strain GP⁹⁹) contributing 144 reductive dehalogenase sequences. These dehalogenases formed only 18 OGs, where 16 of the 18 come from two similar strains of *Dehalogenimonas alkenigignens*. Unlike *Dehalococcoides mccartyi* and *Dehalobacter restrictus*, *Dehalogenimonas* is a more divergent group at this time (Fig. 4 and 2) with multiple species identified. Much more sequencing is required to more clearly define the extent and organization of OGs in *Dehalogenimonas*.




Fig. 6 Phylogenetic amino acid tree of reductive dehalogenases which have been at least partially characterized from *Dehalococcoides* (Dhc, in blue), *Dehalogenimonas* (Dhg, in grey) and *Dehalobacter* (Dhb, in yellow). Best scoring maximum likelihood tree displayed. Scale indicates number of substitutions per site. Ortholog Groups (OGs) of dehalogenases with upwards of 90% amino acid identity are highlighted and identified by number. OGs containing a functionally characterized representative are annotated by dehalogenase name. *rdhA* sequences are coloured by genome they originated from. *rdhA* are named by NCBI locus tag.



We next considered only the reductive dehalogenases from *Dehalococcoides*, *Dehalobacter* and *Dehalogenimonas* that have been characterized, either biochemically or genetically, to reveal relationships between function and OG classification (Fig. 6). From *Dehalococcoides* dehalogenases, all characterized representatives in the same OG have activity on the same substrates. For example VcrA from different *D. mccartyi* (strain VS,¹⁰⁰ WBC-2³⁵ or from the KB-1 consortium¹⁰¹) are active on the same tested substrates (see Table S5† for full list of characterized dehalogenases). We found the same to be true with BvcA (*D. mccartyi* BAV1 and KB-1) and TceA (*D. mccartyi* 195 and KB-1). It is important to note that the number of substrates tested was a limited list in all cases. From *Dehalogenimonas* there are no OGs with more than one characterized representative. From *Dehalobacter*, OG 97 can be described as a 1,1,1-trichloroethane/chloroform/1,1-dichloroethane group of dehalogenases with several overlapping functions, and OG 96 includes a perchloroethene dehalogenase and a tetrachlorobenzene/trichlorobenzene dehalogenase. At present, the members of these OGs have not been extensively tested on different substrates. Future work and new protein structures will refine the definition of OGs to better associate dehalogenase function to sequence.

New community resource: the reductive dehalogenase database web server

A database of reductive dehalogenase gene sequences and protein sequences was previously located in a publicly accessible Google Doc.¹⁹ We have transformed this resource into a web-based database where the data can be interactively viewed and downloaded. Additionally, users can check and add new sequences, classify RdhA into Ortholog Groups and visualize sequences in a tree with other RdhA sequences. The webserver can be accessed at: <http://RDaseDB.biozone.utoronto.ca>.

Conclusions

Sequencing and completely closing multiple genomes of *D. mccartyi* has provided new insights into the evolution of *rdhA* genes in this genus. The majority of *rdhA* found in *D. mccartyi* display a much higher degree of synteny between genomes than previously appreciated. Evolutionary analysis revealed that most *rdhA* sequences are old, and have likely been in *D. mccartyi* genomes since the time of a *Dehalococcoidia* common ancestor (Fig. 4 and Table S4-4†). It is possible that the relatively recent anthropogenic release of specific industrial chemicals has enabled the dissemination of specific reductive dehalogenases capable of solvent dehalogenation, initiating a period of adaptation and complexification of the *D. mccartyi* genome. These genome comparisons have provided specific examples of how the complement of *D. mccartyi* *rdhA* has been shaped, revealing (1) adaptation of existing *rdhA* to new substrates, (2) lateral gene transfer of *rdhA* from the environment and (3) duplication or movement within genomes. While too few genomes are currently available from other OHRB, it is anticipated that these trends will still prevail. The classification of reductive

dehalogenase sequences into Ortholog Groups has enabled a better understanding of the evolution and function of *rdhA* sequences, clearly showing that they are of pre-industrial origin. The OG classification prevents over-reporting of "novel" sequences, but rather allows for rapid situation of newly sequenced members in the greater context. To assist future studies of organohalide-respiring bacteria and their reductive dehalogenases, we have begun a public resource to assist the community in the classification of reductive dehalogenase sequences by building a high-quality curated database of all reductive dehalogenases, and maintaining a simple classification system and nomenclature.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Support was provided by the Government of Canada through Genome Canada and the Ontario Genomics Institute (no. 2009-OGI-ABC-1405 to E. A. E.), the Natural Sciences and Engineering Research Council of Canada (NSERC PGS-D to O. M.), the NSERC CREATE RENEW program [180804567], and the Government of Ontario through the Ontario Graduate Scholarship program (OGS to L. A. P. J.). Support was also provided by the Government of Ontario through the ORF-GL2 program and the United States Department of Defense through the strategic Environmental Research and Development Program (SERDP) under contract (W912HQ-07-C-0036 project ER-1586). Special thanks to Laura Hug for initially coming up with the OG classification and her input on the reductive dehalogenase web server. Enormous thanks to Herbert Molenda for programming the website and database. Also special thanks to anonymous reviewers for their thorough comments and suggestions.

References

- 1 H. F. Stroo, A. Leeson and C. H. Ward, *Bioaugmentation for Groundwater Remediation*, Springer, New York, 2012.
- 2 S. S. Suthersan, M. Schnobrich, J. Martin, J. F. Horst and E. Gates, Three Decades of Solvent Bioremediation: The Evolution from Innovation to Conventional Practice, *Groundwater Monit. Rem.*, 2017, **37**, 14–23.
- 3 D. W. Hill and P. L. McCarty, Anaerobic degradation of selected chlorinated hydrocarbon pesticides, *J. - Water Pollut. Control Fed.*, 1967, **39**, 1259–1277.
- 4 E. J. Bouwer and P. L. McCarty, Transformations of 1- and 2-carbon halogenated aliphatic organic compounds under methanogenic conditions, *Appl. Environ. Microbiol.*, 1983, **45**, 1286–1294.
- 5 D. L. Freedman and J. M. Gossett, Biological reductive dechlorination of tetrachloroethylene and trichloroethylene to ethylene under methanogenic conditions, *Appl. Environ. Microbiol.*, 1989, **55**, 2144–2151.
- 6 R. Seshardi, L. Adrian, D. E. Fouts, J. A. Eisen, A. M. Phillippe, B. A. Methe, N. L. Ward, W. C. Nelson,



R. J. Dobson, S. C. Daugherty, L. M. Brinkac, S. A. Sullivan, R. Madupu, K. E. Nelson, K. H. Kang, M. Impraim, K. Tran, J. M. Robinson, H. A. Forberger, C. M. Fraser, S. H. Zinder and J. F. Heidelberg, Genome sequence of the PCE-dechlorinating bacterium *Dehalococcoides ethenogenes*, *Science*, 2005, **307**, 105–108.

7 X. Maymo-Gatell, Y. Chien, J. M. Gossett and S. H. Zinder, Isolation of a bacterium that reductively dechlorinates tetrachloroethene to ethene, *Science*, 1997, **276**, 1568–1571.

8 D. J. Major, M. L. McMaster, E. E. Cox, E. A. Edwards, S. M. Dworatzek, E. R. Hendrickson, M. G. Starr, J. A. Payne and L. W. Buonamici, Field demonstration of successful bioaugmentation to achieve dechlorination of tetrachloroethene to ethene, *Environ. Sci. Technol.*, 2000, **36**, 5106–5116.

9 M. Duhamel, S. D. Wehr, L. Yu, H. Rizvi, D. Seepersad, S. Dworatzek, E. E. Cox and E. A. Edwards, Comparison of anaerobic dechlorinating enrichment cultures maintained on tetrachloroethene, trichloroethene, *cis*-dichloroethene and vinyl chloride, *Water Res.*, 2002, **36**, 4193–4202.

10 J. Dolffing and J. M. Tiedje, Hydrogen cycling in a three-tiered food web growing on the methanogenic conversion of 3-chlorobenzoate, *FEMS Microbiol. Ecol.*, 1986, **2**, 293–298.

11 K. A. DeWeerd, L. Mandelco, R. S. Tanner, C. R. Woese and J. M. Suflita, *Desulfomonile tiedjei* gen. nov. and sp. nov., a novel anaerobic, dehalogenating, sulfate-reducing bacterium, *Arch. Microbiol.*, 1990, **154**, 23–30.

12 F. E. L. Lorenz Adrian, *Organohalide-Respiring Bacteria*, ed. F. L. Lorenz Adrian, Heidelberg, Berlin, 2006.

13 T. Schubert, L. Adrian, R. G. Sawers and G. Diekert, Organohalide respiratory chains: composition, topology and key enzymes, *FEMS Microbiol. Ecol.*, 2018, 94.

14 M. Fincker and A. M. Spormann, Biochemistry of Catabolic Reductive Dehalogenation, *Annu. Rev. Biochem.*, 2017, **86**, 357–386.

15 D. Türkowsky, N. Jehmlich, G. Diekert, L. Adrian, M. von Bergen and T. Goris, An integrative overview of genomic, transcriptomic and proteomic analyses in organohalide respiration research, *FEMS Microbiol. Ecol.*, 2018, 94.

16 S. Atashgahi, Y. Lu and H. Smidt, *Organohalide-Respiring Bacteria*, ed. L. Adrian and F. E. Löffler, Springer Berlin Heidelberg, Berlin, Heidelberg, 2016, pp. 63–105, DOI: 10.1007/978-3-662-49875-0_5.

17 L. A. Puentes Jácome and E. A. Edwards, A switch of chlorinated substrate causes emergence of a previously undetected native *Dehalobacter* population in an established *Dehalococcoides*-dominated chloroethene-dechlorinating enrichment culture, *FEMS Microbiol. Ecol.*, 2017, 93.

18 M. J. Krzmarzick, B. B. Crary, J. J. Harding, O. O. Oyerinde, A. C. Leri, S. C. B. Myneni and P. J. Novak, Natural Niche for Organohalide-Respiring Chloroflexi, *Appl. Environ. Microbiol.*, 2012, **78**, 393–401.

19 L. A. Hug, F. Maphosa, D. Leys, F. E. Löffler, H. Smidt, E. A. Edwards and L. Adrian, Overview of organohalide-respiring bacteria and a proposal for a classification system for reductive dehalogenases, *Philos. Trans. R. Soc., B*, 2013, **368**, 1–10.

20 M. Kube, A. Beck, S. H. Zinder, H. Kuhl, R. Reinhardt and L. Adrian, Genome sequence of the chlorinated compound-respiring bacterium *Dehalococcoides* species strain CBDB1, *Nat. Biotechnol.*, 2005, **23**, 1269–1273.

21 P. J. McMurdie, S. F. Behrens, J. A. Müller, J. Goke, K. M. Ritalahti, R. Wagner, E. Goltsman, A. Lapidus, S. Holmes, F. E. Löffler and A. M. Spormann, Localized plasticity in the streamlined genomes of vinyl chloride respiring *Dehalococcoides*, *PLoS Genet.*, 2009, **5**, e1000714.

22 C. Ding, M. J. Rogers, K.-L. Yang and J. He, Loss of the *ssrA* genome island led to partial debromination in the PBDE respiring *Dehalococcoides mccartyi* strain GY50, *Environ. Microbiol.*, 2017, **19**, 2906–2915.

23 P. J. McMurdie, L. A. Hug, E. A. Edwards, S. Holmes and A. A. Spormann, Site-specific mobilization of vinyl-chloride respiration islands by a mechanism common in *Dehalococcoides*, *BMC Genomics*, 2011, **12**, 287–302.

24 R. Krajmalnik-Brown, Y. Sung, K. M. Ritalahti, F. M. Saunders and F. E. Löffler, Environmental distribution of the trichloroethene reductive dehalogenase gene (*tceA*) suggests lateral gene transfer among *Dehalococcoides*, *FEMS Microbiol. Ecol.*, 2007, **59**, 206–214.

25 H. Smidt, M. Van Leest, J. van der Oost and W. M. De Vos, Transcriptional regulation of the *cpr* gene cluster in orthochlorophenol respiring *Desulfitobacterium dehalogenans*, *J. Bacteriol.*, 2000, **182**, 5683–5691.

26 K. Chen, L. Huang, C. Xu, X. Liu, J. He, S. H. Zinder, S. Li and J. Jiang, Molecular characterization of the enzymes involved in degradation of a brominated aromatic herbicide, *Mol. Microbiol.*, 2013, **89**, 1121–1139.

27 A. Kublik, D. Deobald, S. Hartwig, C. L. Schiffmann, A. Andrades, M. von Bergen, R. G. Sawers and L. Adrian, Identification of a multi-protein reductive dehalogenase complex in *Dehalococcoides mccartyi* strain CBDB1 suggests a protein-dependent respiratory electron transport chain obviating quinone involvement, *Environ. Microbiol.*, 2016, **18**, 3044–3056.

28 S. Hartwig, N. Dragomirova, A. Kublik, D. Türkowsky, M. von Bergen, U. Lechner, L. Adrian and R. G. Sawers, A H₂-oxidizing, 1,2,3-trichlorobenzene-reducing multienzyme complex isolated from the obligately organohalide-respiring bacterium *Dehalococcoides mccartyi* strain CBDB1, *Environ. Microbiol. Rep.*, 2017, **9**, 618–625.

29 R. Banerjee and S. W. Ragsdale, The many faces of vitamin B12: catalysis by cobalamin-dependent enzymes, *Annu. Rev. Biochem.*, 2003, **72**, 209–247.

30 Y. T. Marcy, R. S. Ishoey, T. B. Laksen, B. P. Stockwell, A. L. Walenz, K. Y. Halpern, K. Y. Beeson, S. M. Goldberg and S. R. Quake, Nanoliter reactors improve multiple displacement amplification of genomes from single cells, *PLoS Genet.*, 2007, **3**, 1702–1708.

31 Y. Morita, T. Futagami, M. Goto and K. Furukawa, Functional characterization of the trigger factor protein PceT of a tetrachloroethene-dechlorinating *Desulfitobacterium hafniense* Y51, *Appl. Microbiol. Biotechnol.*, 2009, **83**, 775–781.





32 A. Sunyama, M. Yamashita, S. Yoshino and K. Furukawa, Molecular characterization of the PceA dehalogenase of *Desulfitobacterium* sp. strain Y51, *Bacteriol.*, 2002, **184**, 3419–3425.

33 J. Yan, K. M. Ritalahti, D. D. Wagner and F. L. Löffler, Unexpected specificity of interspecies cobamide transfer from *Geobacter* spp. to organohalide-respiring *Dehalococcoides mccartyi* strains, *Appl. Environ. Microbiol.*, 2012, **78**, 6630–6636.

34 A. Parthasarathy, T. A. Stich, S. T. Lohner, A. Lesnefsky, R. D. Britt and A. Spormann, Heterologously expressed vinyl chloride reductive dehalogenase (VcrA) from *Dehalococcoides mccartyi* strain VS, *J. Am. Chem. Soc.*, 2015, **137**, 3525–3532.

35 O. Molenda, A. T. Quaile and E. A. Edwards, *Dehalogenimonas* sp. strain WBC-2 genome and identification of its *trans*-dichloroethene reductive dehalogenase, TdrA, *Appl. Environ. Microbiol.*, 2016, **82**, 40–50.

36 K. A. Payne, C. P. Quezada, K. Fisher, M. S. Dunstan, F. A. Collins, H. Sjuts, C. Levy, S. Hay, S. E. Rigby and D. Leys, Reductive dehalogenase structure suggests a mechanism for B12-dependent dehalogenation, *Nature*, 2015, **517**, 513–516.

37 M. Bommer, C. Kunze, J. Fesseler, T. Schubert, G. Diekert and H. Dobbek, Structural basis for organohalide respiration, *Science*, 2014, **346**, 455–458.

38 A. Pérez-de-Mora, A. Lacourt, M. L. McMaster, X. Liang, S. M. Dworatzek and E. A. Edwards, Chlorinated electron acceptor abundance drives selection of *Dehalococcoides mccartyi* (*D. mccartyi*) strains in dechlorinating enrichment cultures and groundwater environments, *Front. Microbiol.*, 2018, **9**.

39 M. Duhamel, K. Mo and E. A. Edwards, Characterization of a highly enriched *Dehalococcoides*-containing culture that grows on vinyl chloride and trichloroethene, *Appl. Environ. Microbiol.*, 2004, **70**, 5538–5545.

40 M. Duhamel and E. A. Edwards, Growth and yields of dechlorinators, acetogens, and methanogens during reductive dechlorination of chlorinated ethenes and dihaloelimination of 1,2-dichloroethane, *Environ. Sci. Technol.*, 2007, **41**, 2303–2310.

41 M. Duhamel, S. D. Wehr, L. Yu, H. Rizvi, D. Seepersad, S. Dworatzek, E. E. Cox and E. A. Edwards, Comparison of anaerobic dechlorinating enrichment cultures maintained on tetrachloroethene, trichloroethene, *cis*-dichloroethene and vinyl chloride, *Water Res.*, 2002, **36**, 4193–4202.

42 S. Tang, Y. Gong and E. A. Edwards, Semi-automatic *in silico* gap closure enabled *de novo* assembly of two *Dehalobacter* genomes from metagenomic data, *PLoS One*, 2012, **7**.

43 A. M. Bolger, M. Lohse and B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data, *Bioinformatics*, 2014, **30**, 2114–2120.

44 R. Aziz, D. Bartels, A. D. Best, M. DeJongh, T. Disz, R. Edwards, K. Forsma, S. Gerdes, E. M. Glass, M. Kubal, F. Meyer, G. Olsen, R. Olson, A. Ostermn, R. Overbeek, L. McNeil, D. Paarmann, T. Paczian, B. Parello, G. Push, C. Reich, R. Stevens, O. Vassieva, V. Vonstein, A. Wilke and O. Zagnitko, The RAST server: rapid annotations using subsystems technology, *BMC Genomics*, 2008, **9**.

45 G. H. Van Domselaar, P. Stothard, S. Shrivastava, J. A. Cruz, A. Guo, X. Dong, P. Lu, D. Szafron, R. Greiner and D. S. Wishart, BASys: a web server for automated bacterial genome annotation, *Nucleic Acids Res.*, 2005, **33**, W455–W459.

46 T. Tatusova, M. DiCuccio, A. Badretdin, V. Chetvernin, E. P. Nawrocki, L. Zaslavsky, A. Lomsadze, K. D. Pruitt, M. Borodovsky and J. Ostell, NCBI prokaryotic genome annotation pipeline, *Nucleic Acids Res.*, 2016, **44**, 6614–6624.

47 A. Frank and J. Lobry, Oriloc: prediction of replication boundaries in unannotated bacterial chromosomes, *Bioinformatics*, 2000, **16**, 566–567.

48 B. Contreras-Moreira and P. Vinuesa, GET_HOMOLOGUES, a versatile software package for scalable and robust microbial pangenome analysis, *Appl. Environ. Microbiol.*, 2013, **79**, 7696–7701.

49 R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput, *Nucleic Acids Res.*, 2004, **32**, 1792–1797.

50 A. Stamatakis, RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models, *Bioinformatics*, 2006, **22**, 2688–2690.

51 M. Kearse, R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes and A. Drummond, Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data, *Bioinformatics*, 2012, **28**, 1647–1649.

52 Y. B. Ahn, L. J. Kerkhof and M. M. Haggblom, *Desulfoluna spongiiphila* sp. nov., a dehalogenating bacterium in the Desulfobacteraceae from the marine sponge *Aplysina aerophoba*, *Int. J. Syst. Evol. Microbiol.*, 2009, **59**, 2133–2139.

53 A. Nekrutenko, K. D. Makova and W.-H. Li, The K(A)/K(S) Ratio Test for Assessing the Protein-Coding Potential of Genomic Regions: An Empirical and Simulation Study, *Genome Res.*, 2002, **12**, 198–202.

54 B. Xu and Z. Yang, pamlX: A Graphical User Interface for PAML, *Mol. Biol. Evol.*, 2013, **30**, 2723–2724.

55 Z. Yang, PAML: a program package for phylogenetic analysis by maximum likelihood, *CABIOS, Comput. Appl. Biosci.*, 1997, **13**, 555–556.

56 Z. Yang and J. P. Bielawski, Statistical methods for detecting molecular adaptation, *Trends Ecol. Evol.*, 2000, **15**, 496–503.

57 F. E. Löffler, J. Yan, K. M. Ritalahti, L. Adrian, E. A. Edwards, K. T. Konstantinidis, J. A. Müller, H. Fullerton, S. H. Zinder and A. M. Spormann, *Dehalococcoides mccartyi* gen. nov., sp. nov., obligately organohalide-respiring anaerobic bacteria relevant to halogen cycling and bioremediation, belong to a novel bacterial class, Dehalococcoidia class nov., order *Dehalococcoidales* ord. nov. and family *Dehalococcoidaceae* fam. nov., within the phylum Chloroflexi, *Int. J. Syst. Evol. Microbiol.*, 2013, **63**, 625–635.

58 E. R. Hendrickson, J. A. Payne, R. M. Young, M. G. Starr, M. P. Perry, S. Fahnestock, D. E. Ellis and R. C. Ebersole,

Molecular Analysis of *Dehalococcoides* 16S Ribosomal DNA from Chloroethene-Contaminated Sites throughout North America and Europe, *Appl. Environ. Microbiol.*, 2002, **68**, 485–495.

59 M. Pörz, T. Goris, T. Wubet, M. T. Tarkka, F. Buscot, I. Nijenhuis, U. Lechner and L. Adrian, Genome sequences of two dehalogenation specialists – *Dehalococcoides mccartyi* strains BTF08 and DCMB5 enriched from the highly polluted Bitterfeld region, *FEMS Microbiol. Lett.*, 2013, **343**, 101–104.

60 S. Zhao, C. Ding and J. He, Genomic characterization of *Dehalococcoides mccartyi* strain 11a5 reveals a circular extrachromosomal genetic element and a new tetrachloroethene reductive dehalogenase gene, *FEMS Microbiol. Ecol.*, 2017, **93**, fiw235.

61 Y. Sung, K. M. Ritalahti, R. P. Apkarian and F. E. Löffler, Quantitative PCR confirms purity of strain GT, a novel trichloroethene-to-ethene-respiring *Dehalococcoides* isolate, *Appl. Environ. Microbiol.*, 2006, **72**, 1980–1987.

62 M. Yohda, O. Yagi, A. Takechi, M. Kitajima, H. Matsuda, N. Miyamura, T. Aizawa, M. Nakajima, M. Sunairi, A. Daiba, T. Miyajima, M. Teruya, K. Teruya, A. Shiroma, M. Shimoji, H. Tamotsu, A. Juan, K. Nakano, M. Aoyama, Y. Terabayashi, K. Satou and T. Hirano, Genome sequence determination and metagenomic characterization of a *Dehalococcoides* mixed culture grown on *cis*-1,2-dichloroethene, *J. Biosci. Bioeng.*, 2015, **120**, 69–77.

63 D. Cheng and J. He, Isolation and characterization of “*Dehalococcoides*” sp. strain MB which dechlorinates tetrachloroethene to *trans*-1,2-dichloroethene, *Appl. Environ. Microbiol.*, 2009, **75**, 5910–5918.

64 S. Wang, K. R. Chng, A. Wilm, S. Zhao, K.-L. Yang, N. Nagarajan and J. He, Genomic characterization of three unique *Dehalococcoides* that respire on persistent polychlorinated biphenyls, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 12103–12108.

65 O. Molenda, S. Tang, L. Lomheim, G. Vasu, S. Lemak, A. F. Yakunin and E. A. Edwards, Extrachromosomal circular elements targeted by CRISPR-Cas in *Dehalococcoides mccartyi* are linked to mobilization of reductive dehalogenase genes, *ISME J.*, 2019, **13**, 24–38.

66 A. Low, Z. Shen, D. Cheng, M. J. Rogers, P. K. H. Lee and J. He, A comparative genomics and reductive dehalogenase gene transcription study of two chloroethene-respiring bacteria, *Dehalococcoides mccartyi* strains MB and 11a, *Sci. Rep.*, 2015, **5**, 15204.

67 P. K. H. Lee, J. He, S. H. Zinder and L. Alvarez-Cohen, Evidence for nitrogen fixation by “*Dehalococcoides ethenogenes*” strain 195, *Appl. Environ. Microbiol.*, 2009, **75**, 7551–7555.

68 T. Kruse, J. Maillard, L. Goodwin, T. Woyke, H. Teshima, D. Bruce, C. Detter, R. Tapia, C. Han, M. Huntemann, C. L. Wei, J. Han, A. Chen, N. Kyrpides, E. Szeto, V. Markowitz, N. Ivanova, I. Pagani, A. Pati, S. Pitluck, M. Nolan, C. Holliger and H. Smidt, Complete genome sequence of *Dehalobacter restrictus* PER-K23(T), *Stand. Genomic Sci.*, 2013, **8**, 375–388.

69 L. A. Hug, R. G. Beiko, A. R. Rowe, R. E. Richardson and E. A. Edwards, Comparative metagenomics of three *Dehalococcoides*-containing enrichment cultures: the role of the non-dechlorinating community, *BMC Genomics*, 2012, **13**, 327.

70 A. S. Waller, R. Krajmalnik-Brown, F. E. Löffler and E. A. Edwards, Multiple reductive-dehalogenase-homologous genes are simultaneously transcribed during dechlorination by *Dehalococcoides*-containing cultures, *Appl. Environ. Microbiol.*, 2005, **71**, 8257–8264.

71 A. Wagner, L. Segler, S. Kleinsteuber, G. Sawers, H. Smidt and U. Lechner, Regulation of reductive dehalogenase gene transcription in *Dehalococcoides mccartyi*, *Philos. Trans. R. Soc., B*, 2013, **368**, 20120317.

72 O. Molenda, S. Tang, L. Lomheim and E. A. Edwards, Eight new genomes of organohalide-respiring *Dehalococcoides mccartyi* reveal evolutionary trends in reductive dehalogenase enzymes, *bioRxiv, Bioinf.*, 2018, 345173.

73 L. Krasper, H. Lilie, A. Kublik, L. Adrian, R. Golbik and U. Lechner, The MarR-Type Regulator Rdh2R Regulates *rdh* Gene Transcription in *Dehalococcoides mccartyi* strain CBDB1, *J. Bacteriol.*, 2016, **198**, 3130–3141.

74 C. B. Mansfeldt, A. R. Rowe, G. L. W. Heavner, S. H. Zinder and R. E. Richardson, Meta-analyses of *Dehalococcoides mccartyi* strain 195 transcriptomic profiles identify a respiration rate-related gene expression transition point and interoperator recruitment of a key oxidoreductase subunit, *Appl. Environ. Microbiol.*, 2014, **80**, 6062–6072.

75 R. M. Morris, J. M. Fung, B. G. Rahm, S. Zhang, D. L. Freedman, S. H. Zinder and R. E. Richardson, Comparative proteomics of *Dehalococcoides* spp. reveals strain-specific peptides associated with activity, *Appl. Environ. Microbiol.*, 2007, **73**, 320–326.

76 J. M. Fung, R. M. Morris, L. Adrian and S. H. Zinder, Expression of reductive dehalogenase genes in *Dehalococcoides ethenogenes* strain 195 growing on tetrachloroethene, trichloroethene, or 2,3-dichlorophenol, *Appl. Environ. Microbiol.*, 2007, **73**, 4439–4445.

77 D. J. Martínez-Cano, M. Reyes-Prieto, E. Martínez-Romero, L. P. Partida-Martínez, A. Latorre, A. Moya and L. Delaey, Evolution of small prokaryotic genomes, *Front. Microbiol.*, 2015, **5**.

78 Y. I. Wolf and E. V. Koonin, Genome reduction as the dominant mode of evolution, *Bioessays*, 2013, **35**, 829–837.

79 H. K. Allen, J. Donato, H. H. Wang, K. A. Cloud-Hansen, J. Davies and J. Handelsman, Call of the wild: antibiotic resistance genes in natural environments, *Nat. Rev. Microbiol.*, 2010, **8**, 251–259.

80 H. Ochman, J. G. Lawrence and E. A. Groisman, Lateral gene transfer and the nature of bacterial innovation, *Nature*, 2000, **405**, 299–304.

81 J. Davies and D. Davies, Origins and evolution of antibiotic resistance, *Microbiol. Mol. Biol. Rev.*, 2010, **74**, 417–433.

82 A.-M. Hanssen and J. U. Ericson Sollid, *SCCmec* in staphylococci: genes on the move, *FEMS Immunol. Med. Microbiol.*, 2006, **46**, 8–20.



83 J. Maillard and C. Holliger, in *Organohalide-Respiring Bacteria*, ed. L. Adrian and F. E. Löffler, Springer, Berlin, 2016, pp. 153–168.

84 P. H. Wang, S. Q. Tang, K. Nemr, R. Flick, J. Yan, R. Mahadevan, A. F. Yakunin, F. E. Löffler and E. A. Edwards, Refined experimental annotation reveals conserved corrinoid autotrophy in chloroform-respiring *Dehalobacter* isolates, *ISME J.*, 2017, **11**, 626–640.

85 Y. K. Wong, S. I. Holland, H. Ertan, M. Manefield and M. Lee, Isolation and characterization of *Dehalobacter* sp. strain UNSWDHB capable of chloroform and chlorinated ethane respiration, *Environ. Microbiol.*, 2016, **18**, 3092–3105.

86 S. Tang and E. A. Edwards, Identification of *Dehalobacter* reductive dehalogenases that catalyse dechlorination of chloroform, 1,1,1-trichloroethane and 1,1-dichloroethane, *Philos. Trans. R. Soc., B*, 2013, **368**, 20120318.

87 A. Grostern, W. W. Chan and E. A. Edwards, 1,1,1-trichloroethane and 1,1-dichloroethane reductive dechlorination kinetics and co-contaminant effects in a *Dehalobacter*-containing mixed culture, *Environ. Sci. Technol.*, 2009, **43**, 6799–6807.

88 B. Sun, B. M. Griffin, L. A.-d.-R. Héctor, S. A. Hashsham and J. M. Tiedje, Microbial Dehalorespiration with 1, 1, 1-Trichloroethane, *Science*, 2002, **298**, 1023–1025.

89 C. Holliger, D. Hahn, H. Harmsen, W. Ludwig, W. Schumacher, B. Tindall, F. Vazquez, N. Weiss and A. J. B. Zehnder, *Dehalobacter restrictus* gen. nov. and sp. nov., a strictly anaerobic bacterium that reductively dechlorinates tetra- and trichloroethene in an anaerobic respiration, *Arch. Microbiol.*, 1998, **169**, 313–321.

90 F. Maphosa, M. van Passel, W. De Vos and H. Smidt, Metagenome analysis reveals yet unexplored reductive dechlorinating potential of *Dehalobacter* sp. E1 growing in co-culture with *Sedimentibacter* sp., *Environ. Microbiol. Rep.*, 2012, **4**, 604–616.

91 W. Qiao, F. Luo, L. Lomheim, E. E. Mack, S. Ye, J. Wu and E. A. Edwards, A *Dehalogenimonas* population respire 1,2,4-trichlorobenzene and dichlorobenzenes, *Environ. Sci. Technol.*, 2018, **52**, 13391–13398.

92 J. L. Nelson, J. M. Fung, H. Cadillo-Quiroz, X. Cheng and S. H. Zinder, A Role for *Dehalobacter* spp. in the Reductive Dehalogenation of Dichlorobenzenes and Monochlorobenzene, *Environ. Sci. Technol.*, 2011, **45**, 6806–6813.

93 L. A. Puentes Jacome, P.-H. Wang, Y. X. Li, O. Molenda, K. Krivushin, M. Ahsanul Islam and E. A. Edwards, Sustained dechlorination of vinyl chloride to ethene in *Dehalococcoides*-enriched cultures grown without addition of exogenous vitamins at low pH, *Environ. Sci. Technol.*, 2019, **53**, 11364–11374.

94 S. Wang, W. Zhang, K. L. Yang and J. He, Isolation and characterization of a novel *Dehalobacter* species strain TCP1 that reductively dechlorinates 2,4,6-trichlorophenol, *Biodegradation*, 2014, **25**, 313–323.

95 N. Yoshida, L. Ye, D. Baba and A. Katayama, A novel *Dehalobacter* species is involved in extensive 4,5,6,7-tetrachlorophthalide dechlorination, *Appl. Environ. Microbiol.*, 2009, **75**, 2400.

96 S. Siddaramappa, J. F. Challacombe, S. F. Delano, L. D. Green, H. Daligault, D. Bruce, C. Detter, R. Tapia, S. Han, L. Goodwin, J. Han, T. Woyke, S. Pitluck, L. Pennacchio, M. Nolan, M. Land, Y. J. Chang, N. C. Kyripies, G. Ovchinnikova, L. Hauser, A. Lapidus, J. Yan, K. S. Bowman, M. S. da Costa, F. A. Rainey and W. M. Moe, Complete genome sequence of *Dehalogenimonas lykanthroporepellens* type strain (BL-DC-9(T)) and comparison to “*Dehalococcoides*” strains, *Stand. Genomic Sci.*, 2012, **6**, 251–264.

97 T. A. Key, D. P. Richmond, K. S. Bowman, Y. J. Cho, J. Chun, M. S. da Costa, F. A. Rainey and W. M. Moe, Genome sequence of the organohalide-respiring *Dehalogenimonas alkenigignens* type strain (IP3-3(T)), *Stand. Genomic Sci.*, 2016, **11**, 44.

98 T. A. Key, K. S. Bowman, I. Lee, J. Chun, L. Albuquerque, M. S. da Costa, F. A. Rainey and W. M. Moe, *Dehalogenimonas formicexedens* sp. nov., a chlorinated alkane-respiring bacterium isolated from contaminated groundwater, *Int. J. Syst. Evol. Microbiol.*, 2017, **67**, 1366–1373.

99 Y. Yang, S. A. Higgins, J. Yan, B. Simsir, K. Chourey, R. Iyer, R. L. Hettich, B. Baldwin, D. M. Ogles and F. E. Löffler, Grape pomace compost harbors organohalide-respiring *Dehalogenimonas* species with novel reductive dehalogenase genes, *ISME J.*, 2017, **11**, 2767–2780.

100 J. A. Müller, B. M. Rosner, G. Von Abendroth, G. Meshulam-Simon, P. L. McCarty and A. M. Spormann, Molecular identification of the catabolic vinyl chloride reductase from *Dehalococcoides* sp. strain VS and its environmental distribution, *Appl. Environ. Microbiol.*, 2004, **70**, 4880–4888.

101 X. Liang, O. Molenda, S. Tang and E. A. Edwards, Identity and substrate-specificity of reductive dehalogenases expressed in *Dehalococcoides*-containing enrichment cultures maintained on different chlorinated ethenes, *Appl. Environ. Microbiol.*, 2015, **81**, 4626–4633.

102 D. R. Johnson, E. L. Brodie, A. E. Hubbard, G. L. Andersen, S. H. Zinder and L. Alvarez-Cohen, Temporal transcriptomic microarray analysis of “*Dehalococcoides ethenogenes*” strain 195 during the transition into stationary phase, *Appl. Environ. Microbiol.*, 2008, **74**, 2864–2872.

103 J. K. Magnuson, M. F. Romine, D. R. Burris and M. T. Kingsley, Trichloroethene reductive dehalogenase from *Dehalococcoides ethenogenes*: sequence of *tceA* and substrate range characterization, *Appl. Environ. Microbiol.*, 2000, **66**, 5141–5147.

104 S. Tang, W. M. W. Chan, K. E. Fletcher, J. Seifert, X. Liang, F. E. Löffler, E. A. Edwards and L. Adrian, Functional characterization of reductive dehalogenases by using blue native polyacrylamide gel electrophoresis, *Appl. Environ. Microbiol.*, 2013, **79**, 974–981.

105 L. Adrian, J. Rahnenführer, J. Gobom and T. Hölscher, Identification of a chlorobenzene reductive dehalogenase in *Dehalococcoides* sp. strain CBDB1, *Appl. Environ. Microbiol.*, 2007, **73**, 7717–7724.

