

Cite this: *Chem. Sci.*, 2020, **11**, 3441

All publication charges for this article have been paid for by the Royal Society of Chemistry

Probing the electronic and mechanistic roles of the μ_4 -sulfur atom in a synthetic Cu_z model system†

Suresh C. Rathnayaka,^a Shahidul M. Islam,^a Ida M. DiMucci,^b Samantha N. MacMillan,^b Kyle M. Lancaster^b*^b and Neal P. Mankad^a*

Nitrous oxide (N_2O) contributes significantly to ozone layer depletion and is a potent greenhouse agent, motivating interest in the chemical details of biological N_2O fixation by nitrous oxide reductase (N_2OR) during bacterial denitrification. In this study, we report a combined experimental/computational study of a synthetic $[\text{4Cu:1S}]$ cluster supported by N-donor ligands that can be considered the closest structural and functional mimic of the Cu_z catalytic site in N_2OR reported to date. Quantitative N_2 measurements during synthetic N_2O reduction were used to determine reaction stoichiometry, which in turn was used as the basis for density functional theory (DFT) modeling of hypothetical reaction intermediates. The mechanism for N_2O reduction emerging from this computational modeling involves cooperative activation of N_2O across a Cu/S cluster edge. Direct interaction of the μ_4 -S ligand with the N_2O substrate during coordination and N–O bond cleavage represents an unconventional mechanistic paradigm to be considered for the chemistry of Cu_z and related metal–sulfur clusters. Consistent with hypothetical participation of the μ_4 -S unit in two-electron reduction of N_2O , Cu K-edge and S K-edge X-ray absorption spectroscopy (XAS) reveal a high degree of participation by the μ_4 -S in redox changes, with approximately 21% S 3p contribution to the redox-active molecular orbital in the highly covalent $[\text{4Cu:1S}]$ core, compared to approximately 14% Cu 3d contribution per copper. The XAS data included in this study represent the first spectroscopic interrogation of multiple redox levels of a $[\text{4Cu:1S}]$ cluster and show high fidelity to the biological Cu_z site.

Received 10th December 2019
Accepted 14th February 2020

DOI: 10.1039/c9sc06251c

rsc.li/chemical-science

Introduction

Metal–sulfide clusters represent a common motif in bio-inorganic chemistry. The most studied examples are iron–sulfur clusters (*e.g.* $[\text{2Fe:2S}]$, $[\text{4Fe:4S}]$, *etc.*) that serve as ubiquitous electron transfer sites in a wide range of metalloproteins.¹ Other scenarios such as the $[\text{NiFe}]$ and $[\text{MoCu}]$ catalytic sites of carbon monoxide dehydrogenases (CODHs),^{2,3} the H-cluster found in $[\text{FeFe}]$ hydrogenases, and the $[\text{FeMo}]$ -cofactor of nitrogenases⁴ involve multinuclear metal–sulfide clusters facilitating multielectron/multiproton catalytic transformations. Typically, the bridging sulfido (S^{2-}) ligands in these clusters are thought to be crucial for electronically coupling the transition metal sites, thereby facilitating electron delocalization and lowering barriers towards electron transfer either to/from a catalytic site or along an electron transport chain. However, only in rare cases are the sulfur centers proposed to play a direct rather than spectator role with regard to bond activation and/or bond

formation. In the case of the $[\text{FeMo}]$ -cofactor of nitrogenase, various hypotheses have been put forward in which sulfide ligands might act as redox-active proton relays⁵ or even that sulfide(s) may serve in a gating mechanism to nitrogenase activity.⁶ In the case of the $[\text{MoCu}]$ catalytic site of aerobic CODH, one mechanistic hypothesis based on crystallographic studies with substrate analogues involves the μ_2 -sulfide ligand actively participating in CO activation *via* transient S–C bond formation.²

During bacterial denitrification, nitrous oxide (N_2O) is converted to $\text{N}_2 + \text{H}_2\text{O}$ in a $2\text{e}^-/2\text{H}^+$ reaction catalyzed by the metalloenzyme, nitrous oxide reductase (N_2OR).⁷ The catalytic site of N_2OR is a tetranuclear copper–sulfur cluster, Cu_z , which has been structurally characterized in both $[\text{4Cu:1S}]$ and $[\text{4Cu:2S}]$ forms.^{8,9} Both forms show N_2O reductase activity to some extent, and both require physiological reduction to their most reduced redox states to activate N_2O : the 4Cu^{I} (“fully reduced”) state for the $[\text{4Cu:1S}]$ cluster and the $3\text{Cu}^{\text{I}}:1\text{Cu}^{\text{II}}$ (“1-hole”) state for the $[\text{4Cu:2S}]$ cluster.¹⁰ For the $[\text{4Cu:1S}]$ cluster, Solomon has proposed N_2O binding across a dicopper(I) cluster edge, with the N_2O molecule occupying a μ -1,3 binding mode, based on computational modeling (Fig. 1a).¹¹ For the $[\text{4Cu:2S}]$ form, Einsle has reported crystallographic data on N_2O -pressurized crystals of N_2OR showing a N_2O molecule within van der

^aDepartment of Chemistry, University of Illinois at Chicago, 845 W. Taylor St., Chicago, IL 60607, USA. E-mail: npm@uic.edu

^bDepartment of Chemistry & Chemical Biology, Cornell University, Baker Laboratory, Ithaca, NY 14853, USA. E-mail: kml236@cornell.edu

† Electronic supplementary information (ESI) available: Spectral, analytical, and computational data. See DOI: 10.1039/c9sc06251c





Fig. 1 N_2O interactions with the Cu_Z site of nitrous oxide reductase proposed by (a) Solomon and (b) Einsle. Distances are given in Å.

Waals contact of Cu_Z , but the N_2O molecule was not found within coordination distance of Cu_Z and had not undergone significant activation (Fig. 1b).⁹ In neither case has experimental data emerged to probe the nature of N_2O activation by the copper-sulfur clusters.

Studying synthetic model systems can aid understanding of how these unusual inorganic copper-sulfur functional groups behave,¹² which is particularly crucial knowledge in the context of N_2O 's significant impact as a greenhouse gas and an ozone layer depleting agent.^{13,14} Among the synthetic copper compounds and materials known to activate N_2O ,^{15–18} one of our groups has reported the only examples of N_2O activation by copper sulfide clusters. In one case, a dicuprous $[\text{Cu}_2\text{S}]$ cluster with an unsupported μ_2 -sulfide bridge¹⁹ was found to reduce multiple N_2O equivalents to N_2 , resulting in exhaustive oxidation of the sulfur center to a μ_2 -sulfate ligand (Scheme 1a).²⁰ Here, the copper centers remained redox inactive while the μ_2 -sulfide ligand was not only the redox-active center but also acted as an oxygen atom acceptor. In another case, a phosphine-supported tetranuclear

$[\text{Cu}_4\text{S}]$ cluster in its 4Cu^{I} state showed reactivity towards N_2O reduction,²¹ but the cluster lost structural integrity during the reaction, losing the sulfur atom to unknown products in the reaction medium and thus limiting insight that can be gained about its role. Finally, a formamidinate-supported $[\text{Cu}_4\text{S}]$ cluster in its formally $3\text{Cu}^{\text{I}}:1\text{Cu}^{\text{II}}$ ($[\text{4Cu:1S}]^{1-}$) state was found to reduce $^{15}\text{N}_2\text{O}$ to $^{15}\text{N}_2$ (Scheme 1b).^{22,23} Here the μ_4 -sulfide bridge remained intact during a formal oxidation to the $2\text{Cu}^{\text{I}}:2\text{Cu}^{\text{II}}$ ($[\text{4Cu:1S}]^0$) redox state of the cluster, allowing us to establish a closed cycle for N_2O reduction. Based on these results, the potential role (or lack thereof) of the bridging sulfide ligand in copper-sulfur clusters merits further investigation.

In this report, we disclose a combined experimental/computational study of the latter system that collectively implicates the μ_4 -sulfide ligand as participating in redox changes and directly interacting with N_2O during its activation (Scheme 1c). Our data includes the first spectroscopic interrogation of multiple $[\text{4Cu:1S}]$ redox levels, which has proven challenging in the metalloenzyme system,^{7,24} and highlights the fidelity of our synthetic model to the biological Cu_Z site. Additionally, the direct interaction of N_2O with the bridging sulfur atom(s) in Cu_Z has not been proposed before. Such reaction pathways should be considered for the chemistry of Cu_Z and related metal-sulfide clusters in light of the synthetic model studies reported herein.

Results and discussion

In our previous report of N_2O reduction by the anionic complex $[\text{Cu}_4(\mu_4\text{-S})(\mu_2\text{-NCN})_4]^{1-}$ (the 1-hole cluster, referred to here as $[\text{4Cu:1S}]^{1-}$) as its $[\text{K}(18\text{-crown-6})]^+$ salt ($\text{NCN} = [\text{MesNC}(\text{H})\text{NMe}_2]^{1-}$),²² we were able to use NMR spectroscopy, isotopic labeling experiments, and *post situ* electrophilic trapping to establish the presence of three products: neutral $[\text{Cu}_4(\mu_4\text{-S})(\mu_2\text{-NCN})_4]$ (the 2-hole cluster, referred to here as $[\text{4Cu:1S}]^0$), N_2 , and O^{2-} . However, we were unable to definitively establish the reaction stoichiometry at that time. Since then, we have undertaken quantitative GC-MS analysis of the reaction headspace to determine the yield of N_2 . According to this analysis (see ESI[†]), 0.53 ± 0.06 mol of N_2 are produced per mol of the $[\text{4Cu:1S}]^{1-}$ complex. When combining this result with our previous observations, we can now confidently propose the balanced reaction shown in Scheme 2 as the dominant pathway. Based on this reaction stoichiometry, we proceeded with the working hypothesis that one equivalent of $[\text{4Cu:1S}]^{1-}$ is responsible for N_2O activation while a second equivalent is acting as a sacrificial reductant, thus accounting for the overall two-electron redox reaction.



Scheme 1 Active participation of bridging sulfide ligands in N_2O activation by (a) binuclear and (b) and (c) tetranuclear synthetic copper sulfide models.



Scheme 2 Overall balanced reaction being studied.



Next, because we have been unable to detect any intermediates experimentally, we sought to examine the binding mode of N_2O using DFT modeling at the B3LYP/6-31G(d) level in the gas phase. To save computational time, the mesityl groups on the supporting NCN ligands were replaced with methyl groups. After attempting to simulate several types of adducts between the $[\text{4Cu:1S}]^{1-}$ model complex and N_2O , we were able to locate minima associated with N_2O coordination to both the $[\text{4Cu:1S}]^{1-}$ model (intermediate $[\text{A}]^{1-}$) and to its closed-shell, fully reduced $[\text{4Cu:1S}]^{2-}$ analogue (intermediate $[\text{A}]^{2-}$). In both cases N_2O occupied a μ -1,3 binding mode, but to our surprise the N_2O molecule was found to bridge one of the Cu centers and the S atom (Fig. 2a). In each case, one of the other Cu centers has moved away from the S atom to facilitate its direct interaction with N_2O . An alternative, μ -3,1,2 binding mode in which the N_2O molecule bridges two Cu centers as well as the S atom also was located but was determined to be significantly higher in energy by $+11.8 \text{ kcal mol}^{-1}$ on the Gibbs free energy surface (see ESI†). The preferred binding mode for this model system is distinct from the μ -1,3 bridging between two Cu centers that is proposed for Cu_z (see Fig. 1a), where the $\mu_4\text{-S}^{2-}$ ligand is not proposed to interact directly with N_2O . It should be noted that a mononuclear intermediate in which N_2O bridges across a terminal nickel-sulfide bond has been isolated and crystallographically characterized by Hayton and coworkers.^{25,26} The accord between the metrical parameters of the activated N_2O in our computational model with Hayton's experimental data (Fig. 2b) lends further support to the intermediacy of $[\text{A}]^{1-}$. Binding of N_2O to the 1-hole $[\text{4Cu:1S}]^{1-}$ model to form $[\text{A}]^{1-}$ was calculated to be endothermic by $+18.5 \text{ kcal mol}^{-1}$, consistent with our inability to observe an N_2O -bound intermediate experimentally.




Fig. 2 (a) Optimized structures of N_2O -activated intermediates $[\text{A}]^{1-}$ and $[\text{A}]^{2-}$ (color code: Cu, orange; S, yellow; O, red; N, blue; C, gray). (b) Comparison of the cyclic core structures of $[\text{A}]^{n-}$ and a related mononuclear Ni complex characterized by Hayton;²⁵ distances are given in Å.

Assuming that small equilibrium concentrations of an N_2O -bound intermediate akin to $[\text{A}]^{1-}$ form under N_2O atmosphere, we next considered the potential reaction pathway to $\text{N}_2 + \text{O}^{2-}$ (Scheme 3). We expect that N_2O binding to the $[\text{4Cu:1S}]^{1-}$ complex would raise its reduction potential, due to the π -accepting nature of N_2O .²⁷ Thus, there would be a driving force for $[\text{A}]^{1-}$ to undergo reduction by a sacrificial 1-hole complex to provide $[\text{A}]^{2-}$. As in the $[\text{4Cu:1S}]$ form of Cu_z ,¹¹ $[\text{A}]^{2-}$ is in the fully-reduced 4Cu^{I} state and thus is expected to π -backdonate sufficient electron density into the N_2O π^* manifold to induce N-O bond cleavage. Conversion to the resulting intermediate $[\text{B}]^{2-}$ from N_2 loss was calculated to be exothermic relative to $[\text{A}]^{2-}$. Further energy lowering was found by shifting the terminal O^{2-} ligand in $[\text{B}]^{2-}$ to a μ_2 -bridging position in $[\text{C}]^{2-}$. In the case of Cu_z , Solomon has reported that the on-cycle intermediate Cu_z° formed after N_2 loss features a terminal oxygen ligand stabilized by hydrogen bonding with a nearby lysine residue, and has found that disruption of hydrogen bonding produces the off-cycle intermediate Cu_z^* in which the oxygen ligand occupies its thermodynamically preferred bridging position.¹¹ Because we propose O^{2-} to be a stoichiometric product of our aprotic model reaction, we assume that O^{2-} dissociates from either $[\text{B}]^{2-}$ or $[\text{C}]^{2-}$.

An alternative pathway (Scheme 3, dotted arrows) would involve N_2 loss directly from 1-hole $[\text{A}]^{1-}$ prior to reduction, producing intermediate $[\text{B}]^{1-}$. Reduction of $[\text{B}]^{1-}$ by a sacrificial $[\text{4Cu:1S}]^{1-}$ complex would then produce intermediate $[\text{B}]^{2-}$ that is common to both pathways. However, because O^{2-} is expected to lower the reduction potential of the tetracopper cluster due to its π -donor character, it should be unfavorable for $[\text{B}]^{1-}$ to undergo reduction by the sacrificial 1-hole species. Indeed, $[\text{A}]^{1-}$ was calculated to be more oxidizing than $[\text{B}]^{1-}$ by 0.21 V. Thus, we consider this alternative pathway to be unlikely, but we cannot rule it out definitively.

Because the μ_4 -sulfide ligand seems to play a crucial and direct role in N_2O activation according to our DFT modeling, we wondered whether the frontier orbitals of these synthetic $[\text{4Cu:1S}]$ complexes have notable sulfur character. In order to validate our mechanistic model, we thus undertook multi-edge X-ray absorption spectroscopy (XAS) combined with higher-level computational modeling to interrogate the electronic structural changes underpinning the $[\text{4Cu:1S}]^{0/1-}$ redox process.

Cu K-edge XAS data obtained for $[\text{4Cu:1S}]^{1-}$ and $[\text{4Cu:1S}]^0$ are shown in Fig. 3a. Spectral subtraction was carried out to remove a minor contribution of $[\text{4Cu:1S}]^0$ in the spectrum of the monoanion (*vide infra*). Neither spectrum presents a resolved pre-edge ($1s \rightarrow 3d$) feature, although both spectra feature a shoulder that gives a peak in the second derivative spectrum at 8979.8 eV, consistent with the presence of Cu 3d vacancies (Fig. 3b). The rising edges of the two spectra have qualitatively similar fine structure including maxima at *ca.* 8983 eV suggesting the presence of Cu^{I} centers,²⁸ although the spectrum of the $[\text{4Cu:1S}]^{1-}$ cluster is shifted, with inflection points occurring at 0.8 to 1.1 eV lower energy relative to $[\text{4Cu:1S}]^0$. Given the effectively identical coordination environments between the two species, the shift in rising edge position largely reflects some Cu participation in the redox process. Moreover, the lack





Scheme 3 Reaction pathways modeled by DFT (B3LYP/6-31G(d)). Energies at 298 K are shown in kcal mol⁻¹. The favored pathway is shown with solid arrows, and the disfavored pathway with dotted arrows.

of dramatic intensity changes for the rising edge features suggests a delocalized redox process, *i.e.* that a localized [2Cu^I:2Cu^{II}]/[3Cu^I:1Cu^{II}] description is not appropriate.

Quantitative estimates of S participation in the redox-active molecular orbital (RAMO) can be gleaned through analysis of S K-edge XAS data²⁹ obtained for the two clusters, which are




Fig. 3 (a) Cu K-edge XAS spectra obtained for the [4Cu:1S]^{0/1-} redox couple. (b) First (top) and second (bottom) derivative Cu K-edge XAS spectra. Isoenergetic pre-edge (1s → 3d) excitations are evident in the second-derivative spectra at 8979.8 eV. Rising edge inflection points occur at 8982.3 and 8985.9 eV for [4Cu:1S]⁰ and 8981.5 and 8984.8 eV for [4Cu:1S]¹⁻.



- 13 A. R. Ravishankara, J. S. Daniel and R. W. Portmann, *Science*, 2009, **326**, 123–125.
- 14 A. J. Thomson, G. Giannopoulos, J. Pretty, E. M. Baggs and D. J. Richardson, *Philos. Trans. R. Soc., B*, 2012, **367**, 1157–1168.
- 15 I. Bar-Nahum, A. K. Gupta, S. M. Huber, M. Z. Ertem, C. J. Cramer and W. B. Tolman, *J. Am. Chem. Soc.*, 2009, **131**, 2812–2814.
- 16 C. Esmieu, M. Orio, S. Torelli, L. Le Pape, J. Pécaut, C. Lebrun and S. Ménage, *Chem. Sci.*, 2014, **5**, 4774–4784.
- 17 M.-L. Tsai, R. G. Hadt, P. Vanelderen, B. F. Sels, R. A. Schoonheydt and E. I. Solomon, *J. Am. Chem. Soc.*, 2014, **136**, 3522–3529.
- 18 V. Zhuravlev and P. J. Malinowski, *Angew. Chem., Int. Ed.*, 2018, **57**, 11697–11700.
- 19 J. Zhai, A. S. Filatov, G. L. Hillhouse and M. D. Hopkins, *Chem. Sci.*, 2016, **7**, 589–595.
- 20 S. Bagherzadeh and N. P. Mankad, *Chem. Commun.*, 2018, **54**, 1097–1100.
- 21 C.-W. Hsu, S. C. Rathnayaka, S. M. Islam, S. N. MacMillan and N. P. Mankad, *Angew. Chem., Int. Ed.*, 2020, **59**, 627–631.
- 22 B. J. Johnson, W. E. Antholine, S. V. Lindeman, M. J. Graham and N. P. Mankad, *J. Am. Chem. Soc.*, 2016, **138**, 13107–13110.
- 23 B. J. Johnson, W. E. Antholine, S. V. Lindeman and N. P. Mankad, *Chem. Commun.*, 2015, **51**, 11860–11863.
- 24 E. I. Solomon, D. E. Heppner, E. M. Johnston, J. W. Ginsbach, J. Cirera, M. Qayyum, M. T. Kieber-Emmons, C. H. Kjaergaard, R. G. Hadt and L. Tian, *Chem. Rev.*, 2014, **114**, 3659–3853.
- 25 N. J. Hartmann, G. Wu and T. W. Hayton, *Angew. Chem., Int. Ed.*, 2015, **54**, 1–5.
- 26 N. J. Hartmann, G. Wu and T. W. Hayton, *Chem. Sci.*, 2018, **326**, 123–129.
- 27 W. C. Trogler, *Coord. Chem. Rev.*, 1999, **187**, 303–327.
- 28 L. S. Kau, D. J. Spira-Solomon, J. E. Penner-Hahn, K. O. Hodgson and E. I. Solomon, *J. Am. Chem. Soc.*, 2001, **109**, 6433–6442.
- 29 T. Glaser, B. Hedman, K. O. Hodgson and E. I. Solomon, *Acc. Chem. Res.*, 2000, **33**, 859–868.
- 30 P. Chen, S. I. Gorelsky, S. Ghosh and E. I. Solomon, *Angew. Chem., Int. Ed.*, 2004, **43**, 4132–4140.
- 31 S. Ghosh, S. I. Gorelsky, S. D. George, J. M. Chan, I. Cabrito, D. M. Dooley, J. J. G. Moura, I. Moura and E. I. Solomon, *J. Am. Chem. Soc.*, 2007, **129**, 3955–3965.
- 32 R. Sarangi, S. DeBeer George, D. Jackson Rudd, R. K. Szilagyi, X. Ribas, C. Rovira, M. Almeida, K. O. Hodgson, B. Hedman and E. I. Solomon, *J. Am. Chem. Soc.*, 2007, **129**, 2316–2326.
- 33 F. Neese, *Inorg. Chem.*, 2005, **44**, 2245–2254.
- 34 F. Neese, *Inorg. Chim. Acta*, 2002, **337**, 181–192.
- 35 S. Sinnecker, L. D. Slep, E. Bill and F. Neese, *J. Chem. Theory Comput.*, 2008, **4**, 908–919.
- 36 D. A. Pantazis, X.-Y. Chen, C. R. Landis and F. Neese, *J. Chem. Theory Comput.*, 2008, **4**, 908–919.
- 37 S. DeBeer George and F. Neese, *Inorg. Chem.*, 2010, **49**, 1849–1853.
- 38 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297.

