



Cite this: *Org. Biomol. Chem.*, 2017, **15**, 7841

Why can a gold salt react as a base?†

Mariarosa Anania, , Lucie Jašíková, , Juraj Jašík  and Jana Roithová  *

This study shows that gold salts $[(L)AuX]$ ($L = PMe_3, PPh_3, JohnPhos, IPr$; $X = SbF_6, PF_6, BF_4, TfO, Tf_2N$) act as bases in aqueous solutions and can transform acetone to digold acetonol complexes $[(L)_2Au_2(CH_2COCH_3)]^+$ without any additional base present in solution. The key step is the formation of digold hydroxide complexes $[(L)_2Au_2(OH)]^+$. The kinetics of the formation of the digold complexes and their mutual transformation is studied by electrospray ionization mass spectrometry and the delayed reactant labelling method. We show that the formation of digold hydroxide is the essential first step towards the formation of the digold acetonol complex, the reaction is favoured by more polar solvents, and the effect of counter ions is negligible. DFT calculations suggest that digold hydroxide and digold acetonol complexes can exist in solution only due to the stabilization by the interaction with two gold atoms. The reaction between the digold hydroxide and acetone proceeds towards the dimer $\{[(L)Au(OH)] \cdot [(L)Au(CH_2COCH_3)]^+\}$. The monomeric units interact at the gold atoms in the perpendicular arrangement typical of the gold clusters bound by the aurophilic interaction. The hydrogen is transferred within the dimer and the reaction continues towards the digold acetonol complex and water.

Received 31st July 2017,
Accepted 25th August 2017

DOI: 10.1039/c7ob01905j

rsc.li/obc

Introduction

Nowadays, homogenous gold catalysis represents a broad research field.^{1–5} Unique properties of gold(I) complexes consist of their strong interactions with multiple bonds^{6–9} and the formation of rather stable, long-lived intermediates with carbon–gold bonds.^{10–12} In addition, the aurophilic interaction makes the formation of complexes with a gold–gold bond favourable which is probably in the background of cooperative effects in gold catalysis.^{13–16} These points make gold(I) catalysis an attractive topic for mechanistic research.^{17–20}

Mass spectrometry is a valuable tool in the research of gold(I) complexes, because they are easily detectable by electrospray ionization (ESI). The classical mechanistic tools such as the determination of reaction kinetics by NMR or UV-vis spectroscopy are related to the kinetics of the overall reaction. Mass spectrometry, on the contrary, provides data related directly to the gold-containing intermediates. It can be therefore extremely useful, for example, in answering the questions about the

role of monoaurated and diaurated intermediates in a reaction mechanism, the topic of an ongoing discussion.^{12,17,21,22}

The disadvantage of electrospray ionization mass spectrometry in the research of reactions proceeding in solutions is that the molecules or ions have to be transferred from the reaction mixture to the gas phase. This process can be accompanied by the formation of various ions in the gas phase during the electrospray ionization process. These ions are then not related to the actual reaction mixture and can be misleading in the interpretation of the results. We have developed a method – Delayed Reactant Labelling – that enables us to eliminate these artefacts and moreover to investigate the reaction kinetics of the ESI-MS detected ions in the reaction mixture.^{12,23}

Delayed Reactant Labelling consists of relative quantification of the detected signals with respect to their isotopically labelled analogues. Clearly, isotopically labelled intermediates cannot be just added to the reaction mixture, therefore we use a trick of adding an isotopically labelled reactant with a time delay to the reaction mixture. It means that we can observe pairs of the ESI-MS signals for all complexes containing the reactant molecule (*i.e.* reactant complexes, intermediates, product complexes, products). As the labelled reactant is added with a time delay, the labelled intermediates/products are initially present at a lower concentration. With time, the concentrations of unlabelled and labelled intermediates become equal. The time evolution is linked to the kinetics of their formation and degradation. As the ESI-MS response for isotopically labelled ions/molecules is the same, the whole

Department of Organic Chemistry, Faculty of Science, Charles University, Hlavova 2030/8, Prague 2, 12843, Czech Republic. E-mail: roithova@natur.cuni.cz

† Electronic supplementary information (ESI) available: Preparation of reaction mixtures; photodissociation theoretical IR spectra of gold complexes; details of the determination of kinetic isotope effects; data of the delayed reactant labelling experiments; ESI-MS spectra of reaction mixtures and the collision induced dissociation spectra of gold complexes; results of DFT calculations. See DOI: 10.1039/c7ob01905j



process can be quantitatively observed by ESI-MS. The ions that are formed in the gas-phase during the electrospray ionization (or in a fast equilibrium in solution) do not show any time evolution of the labelled *vs.* unlabelled signals and can thus be easily identified.

Here, we will address the formation of gold-acetonyl complexes, compounds that have a C–Au bond and are stable in aqueous solutions. Gold-ketonyl complexes can be prepared from ketones and a gold salt in the presence of a base.^{24,25} These complexes can be then used as precatalysts by releasing the gold complex in the presence of an acid.^{23,26} Gold-ketonyl complexes were also predicted as intermediates in a coupling reaction.²⁷ We have observed the formation of gold acetonyl complexes even in the absence of a base and could detect the signals of the monogold acetonyl as well as the digold acetonyl complexes. This raises interesting questions about the mechanism of the formation of these complexes and a possible role of the gold–gold cooperative effect in it.

Results

Generation of the digold acetonyl complex

$[(L)_2Au_2(CH_2COCH_3)]^+$

Previous studies reported the preparation of gold(i) acetonyl complexes from an acetone solution of a gold catalyst in the presence of a base. We have detected digold acetonyl complexes by electrospray ionization (ESI) from the solutions containing acetone even without any additional base. Importantly, however, the presence of water in the solution was essential. We note that this reaction is not exclusive for acetone, but other ketones react as well. An example with cyclohexanone can be found in the ESI (Fig. S1†).

We set out to determine the mechanism of the formation of the digold acetonyl complexes and the role of water molecules in it. All experiments were done in acetone, water and THF (tetrahydrofuran) or dioxane mixture of solvents (the exact ratios will be given for each experiment). THF and dioxane do not participate in the reaction and were used for making the solutions of gold complexes to which acetone and water were added.

The initial experiments were done with $(PMe_3)Au(SbF_6)$. The electrospray ionization mass spectrum (ESI-MS) of the solution of $(PMe_3)Au(SbF_6)$ in dry acetone and THF shows that the trimethylphosphino gold cation can be coordinated to acetone (m/z 331), THF (m/z 345), or another trimethylphosphine ligand (m/z 349) (Fig. 1a). We can also see two digold complexes: $[(PMe_3)_2Au_2OH]^+$ (m/z 563) and $[(PMe_3)_2Au_2Cl]^+$ (m/z 581); the former is a product of a reaction between the gold complex and a trace amount of water; the latter is present due to the chloride residual after the counter ion exchange (see the Experimental details). Addition of water to the solution results in an increase of the ESI-MS signal of $[(PMe_3)_2Au_2OH]^+$ and the appearance of the signal of the digold acetonyl complex $[(PMe_3)_2Au_2(CH_2COCH_3)]^+$ (m/z 603) (Fig. 1b).




Fig. 1 ESI-MS spectrum of a 0.3 mM solution of $[(PMe_3)Au(SbF_6)]$ in THF and acetone (1 : 1, v/v) (a) under dry conditions and (b) after the addition of 5% v/v water. (c) ESI-MS spectrum of a solution of 0.3 mM $[(PMe_3)Au(SbF_6)]$ in THF, acetone, acetone- d_6 , and water (2 : 1 : 1 : 0.2 v/v).

The signal of the digold acetonyl complex can indicate its presence in the solution or it can be due to a neutral monogold acetonyl complex that is tagged by another gold cation during the ESI process. The structure of the detected complex was probed by helium tagging IRPD (infrared photo-dissociation) spectroscopy (Fig. 2a) that provides the infrared spectra of mass-selected ions.²⁸ The IRPD spectrum of $[(PMe_3)_2Au_2(CH_2COCH_3)]^+$ is consistent with the structure in which one $(PMe_3)Au^+$ group replaced a hydrogen atom of a methyl group and the other $(PMe_3)Au^+$ unit is bound to the oxygen atom (Fig. 2b shows a theoretical IR spectrum of this complex). We can exclude all variants of the complex with the enol form of acetone. Such structures would be characterized by the stretching bands of the single C–O bond expected at about 1200 cm^{-1} and the C=C double bond expected above 1600 cm^{-1} (see Fig. S2 in the ESI†). Neither of these signatures is present in the experimental spectrum.

We have also obtained the IRPD spectrum of the complex between acetone and $(PMe_3)Au^+$ (Fig. 2c). It nicely corresponds to the expected coordination of the gold cation to the carbonyl group (Fig. 2d shows the corresponding theoretical IR spectrum; Fig. S3 in the ESI† shows the theoretical IR spectrum of






Fig. 2 Helium tagging IRPD spectra of mass-selected ions (a) $[(\text{PMe}_3)_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ (m/z 603) and (c) $[(\text{PMe}_3)\text{Au}(\text{CH}_3\text{COCH}_3)]^+$ (m/z 331). Theoretical IR spectra (B3LYP-D3/6-311+G*(SDD: Au); scaling factor: 0.975) of the most stable structures of (b) $[(\text{PMe}_3)_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ and (d) $[(\text{PMe}_3)\text{Au}(\text{CH}_3\text{COCH}_3)]^+$.

the enol form of acetone). The stretching mode of the gold-coordinated carbonyl group is at 1663 cm^{-1} in the IRPD spectrum of the monogold complex and at 1537 cm^{-1} in the spectrum of the digold complex. The large red-shift that occurs in the digold complex is a result of a partial electron delocalization between the gold atoms.

Kinetic isotope effect (KIE)

Knowing the structure of the digold acetonyl ions, we wanted to know how these ions are formed. One of the parameters that should be closely related to the reaction mechanism is the kinetic isotope effect for the C–H activation of acetone. We compared the formation of the digold acetonyl complexes from the reaction of $[(\text{PMe}_3)_3\text{AuSbF}_6]$ with a 1 : 1 mixture of CH_3COCH_3 and CD_3COCD_3 . If the reaction was irreversible, then the KIE should be simply the ratio between the concentrations of $[(\text{PMe}_3)_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ and $[(\text{PMe}_3)_2\text{Au}_2(\text{CD}_2\text{COCD}_3)]^+$. The ESI-MS response to isotopically labelled compounds is the same; therefore, the concentration ratio is equal to the ratio of the intensities of the corresponding peaks in the ESI-MS spectrum (Fig. 1c).^{29,30}

For reproducible conditions, we prepared first a solution of the gold complex in THF, then added a given amount of water, and a 1 : 1 mixture of acetone and acetone- d_6 and monitored the final mixture by ESI-MS. We found out that the ratio of the signals

of $[(\text{PMe}_3)_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ and $[(\text{PMe}_3)_2\text{Au}_2(\text{CD}_2\text{COCD}_3)]^+$ strongly depended on the water content in the solution and it was changing with time. At the beginning of the experiment the apparent KIE was large (above 10) and it converged to a smaller value in about 1 hour (see Fig. S4–S11 in the ESI†). The converged KIEs ranged from 3.8 (with 50% addition of water) to 6.2 (with 5% addition of water). We repeated the experiment with D_2O instead of H_2O and obtained the same results within the experimental error (see Table 1 and Fig. S12 in the ESI†).

These results suggest that the digold acetonyl complexes are not formed in a simple irreversible reaction, but probably in equilibrium reactions with different species in solution.

Counter ion effect


The formation of gold acetonyl complexes has to be mediated by a basic reactant. Possible basic species in solution are counter ions from the gold complex and hydroxide anions. As described above, we detect $[(\text{PMe}_3)_2\text{Au}_2\text{OH}]^+$ and $[(\text{PMe}_3)_2\text{Au}_2\text{Cl}]^+$ and these ions are in equilibrium with their neutral variants and SbF_6^- (see Scheme 1).

The requirement of the presence of water in solution for the formation of digold acetonyl can mean that the formation

Table 1 Kinetic isotope effects for the C–H activation of acetone by $[(\text{L})\text{AuX}]$ (L = ligand, X = counter ion) in the presence of water^a

Gold complex	KIE	
	5% water ^a	50% water ^a
$(\text{PMe}_3)_3\text{AuSbF}_6$	6.2 ± 0.2	3.8 ± 0.4
$(\text{PMe}_3)_3\text{AuSbF}_6$	6.0 ± 0.3^b	3.3 ± 0.2^b
$(\text{PMe}_3)_3\text{AuPF}_6$	6.9 ± 0.4	3.9 ± 0.3
$(\text{PMe}_3)_3\text{AuOTf}$	6.2 ± 0.2	3.1 ± 0.3
$(\text{PMe}_3)_3\text{AuNTf}_2$	6.7 ± 0.7	3.7 ± 0.5
$(\text{PPh}_3)_3\text{AuNTf}_2$	7.0 ± 0.3	5.9 ± 0.4
$(\text{PPh}_3)_3\text{AuSbF}_6$	6.8 ± 0.4	6.6 ± 0.2
$(\text{JohnPhos})\text{Au}(\text{CH}_3\text{CN})\text{SbF}_6$	5.1 ± 0.5	3.5 ± 0.2
$(\text{IPr})\text{Au}(\text{CH}_3\text{CN})\text{BF}_4^c$	6.5 ± 0.3	5.2 ± 0.3
$(\text{IPr})\text{AuOTf}^c$	3.5 ± 0.8	5.3 ± 0.3
$(\text{IPr})\text{AuNTf}_2^c$	4.7 ± 0.1	5.4 ± 0.1
$(\text{IPr})_2\text{Au}_2(\text{OH})\text{BF}_4^{c,d}$	6.0 ± 0.1	4.6 ± 0.2

^a The experiments were performed in the solutions of $[(\text{L})\text{AuX}]$ (0.3 mM) in THF, acetone- d_6 in a 2 : 1 : 1 ratio with additional 5% or 50% of water. ^b In the presence of D_2O instead of H_2O . ^c The experiments were performed in the solutions of $[(\text{IPr})\text{AuX}]$ (0.3 mM) in dioxane, acetone, acetone- d_6 in a 2 : 7 : 7 ratio with additional 5% or 50% of water. ^d $[(\text{IPr})_2\text{Au}_2\text{OH}]\text{BF}_4$ also reacts under dry conditions, the kinetic isotope effect is 5.5 ± 0.1 .



Scheme 1 Equilibrium between different complexes in the solution of $[(\text{L})\text{AuX}]$ complex (L = ligand, X = counter ion) in THF, water and acetone.



This experiment once more confirms that the ratio of the concentrations of $[(\text{PMe}_3)_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ and $[(\text{PMe}_3)_2\text{Au}_2(\text{CD}_2\text{COCd}_3)]^+$ depends on the water content in solution (the dashed lines in Fig. 3 are the interpolations of the ratios of the experimental points). It also shows that the rate of the conversion of $[(\text{PMe}_3)_2\text{Au}_2\text{OH}]^+$ and $[(\text{PMe}_3)_2\text{Au}_2\text{Cl}]^+$ to digold




Fig. 4 Time evolution of the relative intensities of the signals at m/z 331 ($[(\text{PMe}_3)\text{Au}(\text{CH}_3\text{COCH}_3)]^+$ + $[(\text{PMe}_3)\text{Au}(\text{CH}_2\text{COCH}_3)]\text{H}^+$ in red), m/z 336 ($[(\text{PMe}_3)\text{Au}(\text{CD}_2\text{COCD}_3)]\text{H}^+$ in violet), and m/z 337 ($[(\text{PMe}_3)\text{Au}(\text{CD}_3\text{COCD}_3)]^+$ in blue – this intensity was corrected for the ^{13}C contribution of the signal at m/z 336). Under the assumption that the complexes with intact acetone molecules are formed in the 1 : 1 ratio during the whole experiment, the evolution of the signal intensity of $[(\text{PMe}_3)\text{Au}(\text{CH}_2\text{COCH}_3)]\text{H}^+$ was extracted as a difference between the intensities of m/z 331 and m/z 337. Experiment: the same as described in the caption of Fig. 3. The experiment was performed with 41% water content in the solution of the gold complex in THF and water; evolution of $[(\text{PMe}_3)\text{Au}(\text{CD}_2\text{COCD}_3)]\text{H}^+$ in the experiments with 33% and 9% of water is shown as well, the rest of the data are in Fig. S30 and S31 in the ESI.†

Next, we repeated the same experiments with the $[(\text{IPr})\text{AuBF}_4]$ catalyst. The experiments are analogous, but we used dioxane instead of THF for the preparation of the gold complex solutions. The results are evaluated precisely in the same way as above. The time evolution of the formation of the

Org. Biomol. Chem., 2017, 15, 7841–7852 | 7845

digold acetonyl complexes is rather similar to that obtained with the phosphine ligand (Fig. S34 in the ESI†) and will not be discussed further.

The time evolution in the range of monogold complexes with acetone again reveals the formation of neutral gold acetonyl complexes (Fig. 5). The evaluation of the signals shows that more of the cationic gold is transformed to these neutral complexes than we found in the experiment with the trimethylphosphino gold catalyst. Repetition of the experiments with D₂O confirmed that there is no effect and molecules of water are not directly involved in the formation of the gold acetonyl complexes (Fig. S35 in the ESI†).

Stability of $[(L)_2Au_2(CH_2COCH_3)]^+$

Finally, we have probed the reversibility of the gold acetonyl complex formation by preparing a solution of $[(\text{PMe}_3)\text{AuSbF}_6]$ in THF, acetone and 33% water, let it react overnight and then adding D_6 -labeled acetone (or *vice versa*). In the experiment with overnight activation of CH_3COCH_3 in the presence of D_2O , we can see a significant amount of the $[(\text{PMe}_3)\text{Au}(\text{CH}_2\text{COCH}_3)]\text{D}^+$ complexes (m/z 332, see Fig. 6a). We can exclude that the signal would be due to an H/D scrambling between acetone and D_2O by examining the signals of other complexes with acetone (see Fig. 6a). We note in passing that the signal at m/z 224 and 226 corresponds to silver complexes with the product of the aldol reaction – $\text{CH}_3\text{-C}(\text{CH}_3)(\text{OD})\text{-CH}_2\text{-(CO-CH}_3\text{)}$. This is clear because there is almost no




Fig. 5 Time evolution of the relative intensities of the signals at m/z 643 ($[(\text{IPr})\text{Au}(\text{CH}_3\text{COCH}_3)]^+$ + $[(\text{IPr})\text{Au}(\text{CH}_2\text{COCH}_3)]\text{H}^+$ in red), m/z 648 ($[(\text{IPr})\text{Au}(\text{CD}_2\text{COCOD}_3)]\text{H}^+$ in violet), and m/z 649 ($[(\text{IPr})\text{Au}(\text{CD}_3\text{COCOD}_3)]$ in blue – this intensity was corrected for the ^{13}C contribution of the signal at m/z 648). Under the assumption that the complexes with intact acetone molecules are formed in the 1 : 1 ratio during the whole experiment, the evolution of the signal intensity of $[(\text{IPr})\text{Au}(\text{CH}_2\text{COCH}_3)]\text{H}^+$ was extracted as a difference between the intensities of m/z 643 and m/z 649. Experiment: $[(\text{IPr})\text{AuBF}_4]$ (358 μg) was dissolved in dioxane (0.2 ml) and H_2O (0.14 ml – 41% v/v) and left to react for 15 hours. Then, a 1 : 1 mixture of CH_3COCH_3 and $\text{CD}_3\text{COCOD}_3$ (1.4 ml) was added and the solution was immediately monitored by ESI-MS.




Fig. 6 (a) ESI-MS spectra of a solution prepared by the addition of 0.4 ml of CD_3COCD_3 to an overnight reaction mixture of $[(\text{PMe}_3)_4\text{AuSbF}_6]$ (77 μg) in THF (0.8 ml), D_2O (0.6 ml) and CH_3COCH_3 (0.4 ml). (b) ESI-MS spectra of a solution prepared by the addition of 0.4 ml of CH_3COCH_3 to an overnight reaction mixture of $[(\text{PMe}_3)_4\text{AuSbF}_6]$ (77 μg) in THF (0.8 ml), H_2O (0.6 ml) and CD_3COCD_3 (0.4 ml). The black spectra were collected during the first 5 minutes and the gray spectra were collected during 55–60 minutes after the addition of (a) CD_3COCD_3 or (b) CH_3COCH_3 to the reaction mixture.

H/D-scrambling in complexes with only one molecule of acetone (*e.g.* $[\text{Ag}(\text{CH}_3\text{COCH}_3)(\text{THF})]^+$).

In the experiment with overnight activation of CD_3COCD_3 in the presence of H_2O we detect $[(\text{PMe}_3)\text{Au}(\text{CD}_2\text{COCD}_3)]\text{H}^+$ (m/z 336, see Fig. 6b). In both experiments shown in Fig. 6, the ratio between the gold complexes of CH_3COCH_3 (m/z 331) and CD_3COCD_3 (m/z 337) right at the beginning of the experiment is close to 1. It perfectly reflects the relative concentration of CH_3COCH_3 and CD_3COCD_3 in solution and indicates that these signals are only due to the gold complexes with intact solvent molecules. On the other hand, $[(\text{PMe}_3)\text{Au}(\text{CH}_2\text{COCH}_3)]\text{D}^+$ (in Fig. 6a) and $[(\text{PMe}_3)\text{Au}(\text{CD}_2\text{COCD}_3)]\text{H}^+$ (in Fig. 6b) are formed by protonation of neutral gold acetylonyl complexes formed overnight.

The same experiments with the $[(\text{IPr})\text{AuBF}_4]$ complex led to analogous results. The spectra again showed a large build-up of the neutral gold-acetonyl complexes in an overnight experiment (see Fig. 7). All spectra of the reaction mixtures in which the gold acetonyl complexes were formed overnight reveal only small intensities of digold hydroxide and digold chloride complexes (Fig. S36–S43†). It can be assumed that most of the gold cations are incorporated into the gold acetonyl complexes. Also, the relative abundances of the other than gold




Fig. 7 (a) ESI-MS spectra of a solution prepared by addition of 0.7 ml of CD_3COCD_3 to an overnight reaction mixture of $[(\text{IPr})\text{AuBF}_4]$ (358 μg) in dioxane (0.2 ml), D_2O (0.45 ml) and CH_3COCH_3 (0.7 ml). (b) ESI-MS spectra of a solution prepared by addition of 0.7 ml of CH_3COCH_3 to an overnight reaction mixture of $[(\text{IPr})\text{AuBF}_4]$ (358 μg) in dioxane (0.2 ml), H_2O (0.45 ml) and CD_3COCD_3 (0.7 ml). The black spectra were collected during the first 5 minutes and the gray spectra were collected during 55–60 minutes after the addition of (a) CD_3COCD_3 or (b) CH_3COCH_3 to the reaction mixture.

acetonyl complexes do not change significantly during these experiments (Fig. S36–S43†).

Next we plotted the time evolution of the relative ESI-MS signal intensities of the unlabelled and labelled digold acetonyl and monogold acetonyl complexes. The latter is possible by using the same strategy as we applied in the experiments shown in Fig. 4 and 5. The advantage of the mutual evaluation of the signals corresponding to the very same ions differing only in the isotopic labelling is that it is quantitative and directly corresponds to their relative concentrations in solution. Hence, it allows us to study the kinetics of the evolution of their concentrations. Note that the time evolutions of the signals of the unlabelled and labelled complexes are mirror images because they are evaluated with respect to each other. Therefore, we show only the unlabelled signals in Fig. 8.

We will discuss the experiments, where the deuterated gold acetonyl complexes were formed in the overnight experiment and then let to react with unlabelled acetone (Fig. 8; the opposite variant is a much slower reaction). The black points show the formation of $[(\text{L})_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ ($\text{L} = \text{PMe}_3$ in Fig. 8a and IPr in Fig. 8b) in the solution containing H_2O , whereas the red points refer to the same experiment but in the presence of D_2O . Clearly, H_2O in the solution makes the reaction faster than D_2O .




Fig. 8 Time evolution of the relative concentration of $[(\text{L})_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ (L is (a) PMe_3 or (b) IPr) with respect to the sum of $[(\text{L})_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$ and $[(\text{L})_2\text{Au}_2(\text{CD}_2\text{COCD}_3)]^+$ in a solution with 33% H_2O (in black) or 33% D_2O (in red). The blue points show the corrected evolution of D^+ tagged neutral $[(\text{L})\text{Au}(\text{CH}_2\text{COCH}_3)]$ with respect to D^+ tagged $[(\text{L})\text{Au}(\text{CD}_2\text{COCD}_3)]$ under the assumption that the concentrations of $[(\text{L})\text{Au}(\text{CH}_3\text{COCH}_3)]^+$ and $[(\text{L})\text{Au}(\text{CD}_3\text{COCD}_3)]^+$ are equal (see the text). Experiments: (a) $[(\text{PMe}_3)\text{AuSbF}_6]$ (77 μg) was dissolved in THF (0.8 ml), CD_3COCD_3 (0.4 ml), and H_2O or D_2O (0.6 ml) and left to react for 15 hours. Then, 0.4 ml of CH_3COCH_3 was added and the solution was immediately monitored by ESI-MS. (b) $[(\text{IPr})\text{AuBF}_4]$ (358 μg) was dissolved in dioxane (0.2 ml), CD_3COCD_3 (0.7 ml) and H_2O or D_2O (0.45 ml) and left to react for 15 hours. Then, 0.7 ml of CH_3COCH_3 was added and the solution was immediately monitored by ESI-MS. The lines correspond to 10 points smoothing.

We know from the previous experiments that the equilibrium KIE is around 3.5 for $\text{L} = \text{PMe}_3$ and around 5 for $\text{L} = \text{IPr}$ (see Table 1, 50% water content). Here, after about one hour of the experiment in the presence of H_2O we achieve a ratio of only slightly higher than 1 for $\text{L} = \text{PMe}_3$ and we stay below 1 for $\text{L} = \text{IPr}$. The reaction is even slower in the presence of D_2O . We can thus conclude that the formation of the digold acetonyl complexes is reversible, but the reaction is very slow. The mechanism most probably involves the protodeauration of



Finally, we extracted the evolution of the concentrations of monogold complexes $[(L)Au(CH_2COCH_3)]$ and $[(L)Au(CD_2COCD_3)]$ in D_2O with respect to each other (blue curves in Fig. 8; the extraction procedure was the same as described for Fig. 4 and 5). The results show that the rate of the equilibration of the monogold acetonyl complexes is similar to that of the digold acetonyl complexes. The better signal-to-noise ratio of the results obtained with the IPr ligand allows us to conclude that the equilibration of the monogold acetonyl complexes is somewhat slower than the digold acetonyl complexes.

Thermochemistry

The reaction between $[(\text{PMe}_3)\text{Au}(\text{H}_2\text{O})]^+$ and a water molecule to form neutral $[(\text{PMe}_3)\text{Au}(\text{OH})]$ and the hydroxonium ion

Our computational treatment of the hydroxonium cation is probably not sufficient and further stabilization by interactions with more molecules of water is expected.³⁶ If we assume that the hydroxonium ion can be further stabilized by interactions with more basic molecules present in the solution (such as acetone or anions) it makes the overall energy balance as about thermoneutral. We can thus conclude that the stabilization of the hydroxide anion by interaction with two gold cations drives the formation of a base in solution. Furthermore, more polar solvents (here water) will shift the reaction more towards the formation of digold hydroxide than less polar solvents. This is mainly because of the stabilization of the hydroxonium ions.

The next step is the reaction of gold hydroxides with acetone. Both monogold hydroxide and digold hydroxide react exothermically with acetone in water to yield the acetonyl complexes. The less polar solvent THF makes the reaction to form digold acetonyl less favoured and we found it to be thermoneutral. These results are in agreement with the findings that a lower content of water led to a smaller conversion of digold hydroxide to digold acetonyl (see Fig. 3). Scheme 3 once more demonstrates that the formation of digold complexes is thermodynamically favoured and should largely prevail in solution.



The experiments suggest that the equilibration of the monogold acetonyl complexes is somewhat slower than that of the digold acetonyl complexes. It cannot be caused by a larger stability of the monogold complexes (because the opposite is true). If the formation of the digold acetonyl complexes proceeds *via* the monogold acetonyl complexes then the monogold complexes would be formed either faster or with the same rate as the digold complexes. This is again opposed to the experimental results. Hence, the only explanation for the observed results is that primarily the digold acetonyl complexes are formed and they only slowly decompose into the monogold acetonyl complexes.

The remaining question is how the digold hydroxide complex activates the C–H bond of acetone and is transformed into the digold acetylonyl complex. DFT calculations suggest that the gold–gold interaction plays an essential role in this transformation (Fig. 9). The initial complex between digold hydroxide

Scheme 4 Degradation of the gold acetonyl complexes.

and acetone (**A**) undergoes a rearrangement to the dimeric complex $\{[(\text{PMe}_3)\text{Au}(\text{CH}_3\text{COCH}_3)]\cdot[(\text{PMe}_3)\text{Au}(\text{OH})]\}^+$ (**B**). The monomeric units are oriented perpendicular to each other and interact *via* the gold atoms (see the optimized structure **B** in Fig. 9).^{37,38} The relative energy of **B** is only 7 kJ mol⁻¹ higher than the initial complex **A**.

The proton transfer occurs within the dimeric complex (TSB/C) and leads to another dimer composed of the $[(\text{PMe}_3)\text{Au}(\text{H}_2\text{O})]^+$ and $[(\text{PMe}_3)\text{Au}(\text{CH}_2\text{C}(\text{O})\text{CH}_3)]$ units. This step is endothermic (46 kJ mol⁻¹), because it contains energetically disfavoured gold enolate. The final step is a simple slip of the gold atom to the methylene group (TSC/D) which leads to the formation of a complex between digold acetonyl and water. The overall reaction is endothermic by 3 kJ mol⁻¹. As mentioned above, if isolated digold complexes are assumed as the starting and final points, then the reaction is exothermic by 9 kJ mol⁻¹.

Kinetic isotope effect

A decrease of the kinetic isotope effect for the C–H activation of acetone with the increasing amount of water (Table 1) can be explained by the participation of another molecule of water in the proton transfer reaction.³⁹ This explanation is in agreement with the fact that the most significant drop of KIEs is observed for the trimethylphosphino gold complex. The small ligand does not sterically hinder the participation of other water molecules in the transition state. On the other hand, for the complexes with large ligands, the KIE drop is much smaller (see the results for the complexes with the PPh₃ ligand or IPr together with non-coordinating anion BF₄⁻). The small change of the kinetic isotope effect indicates that the reaction mechanism stayed probably the same and the C–H activation proceeds in a “hydrophobic pocket” formed by the large ligands.⁴⁰

Conclusions

We have shown experimentally that a gold complex $[(\text{L})\text{AuX}]$, L is a ligand, X is a counter ion) in a solution containing water can act as a strong base. It deprotonates acetone to form stable gold–acetonyl complexes. The reaction proceeds firstly *via* the formation of a digold hydroxide complex $[(\text{L})_2\text{Au}_2\text{OH}]^+$ and this complex can deprotonate acetone and form the digold acetonyl complex $[(\text{L})_2\text{Au}_2(\text{CH}_2\text{COCH}_3)]^+$. We can also observe the formation of monogold acetonyl complexes $[(\text{L})\text{Au}(\text{CH}_2\text{COCH}_3)]$, but the kinetics of their formation suggests that they are formed from the primary digold acetonyl complexes. The conversion of digold hydroxide to digold acetonyl is larger, if the solution contains more water.

In order to evaluate the effects of ligands and counter ions in the reaction of gold complexes, we have measured the kinetic isotope effects of the C–H activation step. We showed that there is a negligible effect of counter ions with the exceptions of $[(\text{IPr})\text{Au}(\text{OTf})]$ and $[(\text{IPr})\text{Au}(\text{NTf}_2)]$ in a solution with a low content of water. Otherwise, counter ions are most prob-

ably not involved in the C–H activation reaction. Ligands influence the reaction which is consistent with the scenario that the gold complex is directly present in the transition structure of the reaction.

DFT calculations suggest that the interaction of two gold cations with the anion is essential and makes the formation of digold hydroxide and digold acetonyl complexes thermodynamically feasible. They also show that the monogold acetonyl complexes are prone to protodeauration with hydroxonium (or other protonated ions) in solution, whereas the cationic digold acetonyl complexes should be much more stable. The suggested reaction mechanism of the C–H activation reaction involves a dimer of gold complexes bound *via* the gold–gold interaction.

Experimental

Preparation of $[(\text{L})\text{AuX}]$

Solutions of $[(\text{L})\text{AuCl}]$ with AgX in THF or dioxane (only for L = IPr) were sonicated for 10 minutes. The AgCl precipitate was filtered off. The details of the reaction mixture preparations are in the figure captions and in the ESI.† We use a convention that lists v/v amount of water in the reaction mixture before it is “isotopically labelled” (*e.g.* before CD₃COCD₃ is added with a time delay). We were looking at relative changes with increasing amount of water and slight variations in the preparations of the reaction mixtures did not influence the results.

ESI-MS experiments

The spectra were acquired with a triple quadrupole mass spectrometer TSQ 7000 or a linear ion trap instrument LTQ, both equipped with an electrospray ionization source.⁴¹ Only a simple mode of the mass spectra acquisition was used here. The results are reproducible irrespective of the mass spectrometer used.

Ion spectroscopy

IRPD spectra measurements were conducted with the ISORI (Ion Spectroscopy Of Reaction Intermediates) instrument based on the TSQ 7000 platform.⁴² The ionization and mass-selection is performed in the very same way as for the classical experiments with TSQ 7000. Mass-selected ions are then trapped with helium buffer gas in a wire quadrupole trap operating at 3 K. The ions relax in collisions with helium and attach a helium atom. Helium complexes are then used for the determination of the absorption of IR photons (from a Nd:YAG-pumped IR optical parametric oscillator/amplifier). The absorption leads to the helium detachment and therefore can be monitored as a mass change. The number of helium complexes with (N_i) and without (N_{i0}) laser irradiation is counted using a Daly-type detector. The IRPD spectra are constructed as $(1 - N_i/N_{i0})$. More details can be found in ref. 43 and 44.



- 35 A. Kütt, T. Rodima, J. Saame, E. Raamat, V. Mäemets, I. Kaljurand, I. A. Koppel, R. Y. Garlyauskayte, Y. L. Yagupolskii, L. M. Yagupolskii, E. Bernhardt, H. Willner and I. Leito, *J. Org. Chem.*, 2011, **76**, 391.
- 36 A. P. Kelly, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2006, **110**, 16066.
- 37 F. Mendizabal and P. Pyykko, *Phys. Chem. Chem. Phys.*, 2004, **6**, 900.
- 38 H. Schmidbaur and A. Schier, *Chem. Soc. Rev.*, 2012, **41**, 370.
- 39 S. Cucinotta, A. Ruini, A. Catellani and A. Stirling, *ChemPhysChem*, 2006, **7**, 1229.
- 40 H. Clavier and S. P. Nolan, *Chem. Commun.*, 2010, **46**, 841.
- 41 L. Ducháčková and J. Roithová, *Chem. – Eur. J.*, 2009, **15**, 13399.
- 42 J. Jašík, J. Žabka, J. Roithová and D. Gerlich, *Int. J. Mass Spectrom.*, 2013, **354–355**, 204.
- 43 D. Gerlich, J. Jašík, E. Andris, R. Navrátil and J. Roithová, *ChemPhysChem*, 2016, **17**, 3723.
- 44 J. Jašík, D. Gerlich and J. Roithová, *J. Phys. Chem. A*, 2015, **119**, 2532.
- 45 Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215.
- 46 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 47 M. J. Frisch, *et al.*, *Gaussian 09, Revision D.01*, Gaussian, Inc., Wallingford CT, 2013.
- 48 A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 6378.
- 49 D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648.
- 50 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter*, 1988, **37**, 785.
- 51 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623.
- 52 A. P. Scott and L. Radom, *J. Phys. Chem.*, 1996, **100**, 16502.

