

PAPER

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *RSC Chem. Biol.*, 2020,
1, 281

Enhancing the enthalpic contribution of hydrogen bonds by solvent shielding†

Jonathan Cramer, ^a Xiaohua Jiang,^a Wojciech Schönemann,^a
Marleen Silbermann,^a Pascal Zihlmann,^a Stefan Siegrist,^a Brigitte Fiege,^a
Roman Peter Jakob, ^b Said Rabbani,^a Timm Maier ^b and Beat Ernst ^{*a}

In biological systems, polar interactions are heavily burdened by high desolvation penalties resulting from strong solute–solvent interactions. As a consequence thereof, enthalpic contributions of hydrogen bonds to the free energy of binding are severely diminished. However, this effect is strongly attenuated for interactions within solvent-shielded areas of proteins. In microcalorimetric experiments, we show that the bacterial lectin FimH utilizes conformational adaptations to effectively shield its binding site from solvent. The transition into a lower dielectric environment results in an enthalpic benefit of approximately -13 kJ mol^{-1} for mannoside binding. However, this effect can be abrogated, if the hydrogen bond network within the binding site is disturbed by deoxygenation of the ligand. Conformational adaptation leading to reduced local dielectric constants could represent a general mechanism for proteins to enable enthalpy-driven recognition of polar ligands.

Received 26th June 2020,
Accepted 19th August 2020

DOI: 10.1039/d0cb00108b

rsc.li/rsc-chembio

Introduction


The energetics of hydrogen bonds in biomolecular recognition are highly affected by severe desolvation penalties imposed by strong interactions with the aqueous solvent in the unbound state. A common conception is that strongly directional hydrogen bond interactions provide selectivity, but do not necessarily augment the overall binding affinity of a ligand because favorable contributions are often canceled out by significant desolvation penalties.^{1–3} The free energy penalty for the transfer of a hydroxyl group from aqueous solvent to the gas phase was estimated as 26 kJ mol^{-1} , with an associated enthalpy penalty of 36 kJ mol^{-1} and the transfer of an alcohol function from bulk solvent into a hydrophobic pocket was associated with a free energy penalty of $18\text{--}21 \text{ kJ mol}^{-1}$.^{4,5} Furthermore, in an aqueous system, the contribution of a single hydrogen bond to the free energy of binding has been estimated to equally amount to $18\text{--}21 \text{ kJ mol}^{-1}$.^{6–10} In summary, a hydrogen bond hardly contributes to the binding process, unless additional conditions are fulfilled. It has been proposed that hydrophobic occlusion and solvent shielding are able to advance the strength of hydrogen bonds in biological systems. Thus, enthalpies of

formation for solvent-exposed hydrogen bonds on protein surfaces amount to -6 kJ mol^{-1} , whereas buried interactions contribute with up to -25 kJ mol^{-1} .^{11,12} For the arabinose-binding protein, removal of buried hydrogen bond interactions in deoxy D-galactose derivatives was associated with an enthalpy-driven loss in binding free energy of 30 kJ mol^{-1} .¹³ Similarly, hydrophobic solvent-shielding has been shown to impact the activity of enzymes, as well as the pK_a of buried titratable groups.^{14–18} The exclusion of polar water molecules effectively creates a low dielectric environment that supports electrostatic interactions and accelerates the reaction of charged intermediates. Thus, solvent shielding of hydrophilic binding sites may contribute to enthalpy-driven recognition of highly polar molecules such as carbohydrates. In this study, we aim to demonstrate that proteins actively harness the modulation of local dielectric properties by conformational transitions and to quantify the thermodynamic consequences on polar hydrogen bond interactions.

As a model system to study the impact of the dielectric constant ϵ_r on the thermodynamics of binding, we employed FimH, a bacterial adhesin located on type I pili of uropathogenic *E. coli*.^{19–21} The functional lectin domain of FimH features a carbohydrate recognition site tailor-made to accommodate mannose epitopes. The mannose binding affinity of the lectin domain is allosterically controlled by the adjacent regulatory pilin domain.^{22,23} When mechanical forces, such as shear forces during micturition, cause the separation of lectin and pilin domain,^{23,24} a switch from the low-affinity conformation to the high-affinity conformation of FimH is induced and mannoside ligands are bound with ~ 100 -fold stronger affinity


^a Institute of Molecular Pharmacy, University of Basel, Klingelbergstrasse 50,
4056 Basel, Switzerland. E-mail: beat.ernst@unibas.ch^b Institute of Structural Biology, University of Basel, Klingelbergstrasse 70,
4056 Basel, Switzerland

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0cb00108b



mannose hydroxyl groups, pre-organizing the binding site for carbohydrate recognition ($\text{RMSD}_{\text{apo-bound}} = 0.35 \text{ \AA}$). The binding event is a single-step mechanism that proceeds without any major conformational adjustment of the residues in the mannose binding site.^{26,27} On the other hand, the domain associated full-length FimH construct (FimH_{FL}) has been shown to bind mannoside ligands by a more complex mechanism caused by conformational distortions of the protein due to the presence of the regulatory subunit.^{22,23} In the absence of a ligand, the protein exists in an ensemble of conformational states that are characterized by a shallow, solvent-exposed binding site, a low affinity to carbohydrate ligands, and a high degree of flexibility ($\text{FimH}_{\text{FL-open}}$; Fig. 1).^{23,28} In the presence of

The isolated FimH lectin domain (FimH_{LD}) is a common *in vitro* model for the domain-separated, high-affinity conformation of the protein.²⁵ FimH_{LD} features a rigid structure and a well-defined carbohydrate binding site (Fig. 1). Mannoside ligands engage in an impressive network of ten hydrogen bonds. In the absence of a ligand, water molecules assume the positions of



a ligand, the conformational equilibrium is shifted toward a closed state (FimH_{FL-closed}), in which the clamp loop is accommodated close to the carbohydrate binding site and shields the mannose moiety from the solvent. The atomistic features of this mechanism, namely clamp loop mobility in the open and solvent shielding of the binding site in the closed conformation, have been thoroughly characterized in MD simulations.²⁹

Although the overall structure of this state differs from the high-affinity conformation, the binding site regions are virtually identical (RMSD_{lectin_domain} = 2.0 Å, RMSD_{binding_site} = 0.2 Å). Fig. 2 depicts a thermodynamic model of ligand binding to FimH_{LD} and FimH_{FL}. The experimentally observed free energy of binding to FimH_{LD} (ΔG_{LD-obs}°) is composed of terms for desolvation ($\Delta G_{desolv-closed}^\circ$), protein–ligand interaction ($\Delta G_{bind-closed}^\circ$), and solvation of the assembled complex (ΔG_{solv}°) (Fig. 2, pathway A). For FimH_{FL}, the conformational rearrangement associated with ligand binding theoretically proceeds *via* a conformational selection (Fig. 2, pathway B) or an induced-fit (Fig. 2, pathway C) mechanism. In case of conformational selection, the experimentally observed thermodynamics (ΔG_{FL-obs}°) contain contributions from a conformational transition to the closed state ($\Delta G_{conf-sel}^\circ$), desolvation ($\Delta G_{desolv-closed}^\circ$), protein–ligand interaction ($\Delta G_{bind-closed}^\circ$), and solvation of the assembled complex (ΔG_{solv}°). In the induced-fit pathway C, ΔG_{FL-obs}° is composed of desolvation ($\Delta G_{desolv-open}^\circ$), low-affinity binding ($\Delta G_{bind-open}^\circ$), induced-fit transition ($\Delta G_{ind-fit}^\circ$), and solvation of the assembled complex (ΔG_{solv}°). Thus, the difference in the thermodynamic profiles of ligand binding to FimH_{FL} and FimH_{LD} ($\Delta\Delta G_{FL-LD}^\circ$) gives access to an experimental approximation of clamp loop rearrangement and hydrophobic occlusion of the polar mannose binding site. In the conformational selection model (Fig. 2, pathway B), $\Delta G_{conf-sel}^\circ$ is a constant term, and ligand binding affinity scales entirely with binding affinity to the closed state. In the induced-fit model (Fig. 2, pathway C), however, differences between ligands can scale with the affinity of a low-affinity encounter complex ($\Delta G_{bind-open}^\circ$) and the effect of the conformational transition ($\Delta G_{ind-fit}^\circ$).

As a result of an extensive network of hydrogen bonds between the pre-organized binding site and the mannose moiety, the interaction between mannositides 1–4 and FimH_{LD} (high-affinity state) is characterized by a strongly enthalpic contribution (Fig. 3A). The weak entropic penalty has been linked to the rigidification of the tyrosine gate, an ensemble of hydrophobic residues that control access to the mannose binding site and interact with the aglycones of ligands 1–4.^{26,27} An improved π – π -stacking and hydrophobic interaction between aglycone and tyrosine gate further accentuates their thermodynamic profile, as the electrostatic nature of the aglycones gives rise to progressively tighter, enthalpy-driven binding from 4 $\rightarrow$ 3 $\rightarrow$ 2 $\rightarrow$ 1.

The thermodynamic profiles of mannoside interactions with the FimH_{FL} construct (low-affinity state) show similar characteristics (Fig. 3B). The binding process is equally driven by

enthalpy, with a counteracting entropic penalty. However, the magnitude of this enthalpy–entropy compensation is much more substantial. Overall, the binding affinity of the mannositides 1–4 is reduced roughly by 12 kJ mol^{−1}, approximately two orders of magnitude in K_D , compared with the same ligands binding to FimH_{LD}. This value represents the experimental approximation of the thermodynamics related to the conformational rearrangement from the open (FimH_{FL-open}) to the closed (FimH_{FL-closed}) state.

The difference in the thermodynamics (Fig. 3C) shows that this fairly constant affinity reduction mainly originates from a





Fig. 3 Thermodynamic profiles (see Fig. S1–S4, ESI†) for the interaction of mannositides 1–4 with FimH_{LD} (A) and FimH_{FL} (B). Data for the interactions FimH_{LD}-2/3/4 and FimH_{FL}-2/4 have been published before.^{24,30,31} (C) Difference in the thermodynamic profiles between ligand binding to FimH_{FL} and FimH_{LD}. Error bars represent (propagated) 68% confidence intervals. Numeric data and further information can be found in the ESI.†



In order to investigate the outcome of a perturbation of the hydrogen bond network, several deoxygenated derivatives of *n*-heptyl α -D-mannoside (**4**) were subjected to thermodynamic and structural analyses. The X-ray crystal structures of **4**–**7** in

Similarly, CSP of the side chain signals of Asn135 indicate the interaction with 4-OH. The signal is shifted downfield for **4**, **5**, and **6**, whereas a relative upfield shift for the 4-deoxy derivative **7** signifies the absence of the interaction. Importantly, none of the observed residues show a marked shift in the

^{15}N -dimension of the spectra, reporting on changes in dihedral angles of protein backbone and side chains.

In summary, the structural information obtained by NMR and X-ray is in good agreement, and the conformational changes of protein and ligand are remarkably small upon the loss of relevant interactions. In both experiments, low binding affinity prevented a characterization of compound **8**.

The thermodynamics of **5–7** binding to FimH_{LD} (Fig. 5A) illustrate that removal of the hydroxyl groups in the 2-, 3-, and 4-position of *n*-heptyl α -D-mannoside **4** coincide with a sizeable reduction of the enthalpic contribution to binding.

In addition, the negative value for $-T\Delta S^\circ$ indicates an increased degree of disorder, probably as a consequence of the impaired hydrogen bond network. This is in line with the reduced downfield shifts in NMR experiments. The interaction of **5–7** with FimH_{FL} shows an equivalent trend in the thermodynamic profiles (Fig. 5B). Yet, the magnitude of the enthalpy–entropy compensation effect differs substantially. As a consequence, the fairly constant trend we observed in the differential profiles for **1–4** (Fig. 3C) is lost for **5–7** (Fig. 5C). Here, the enthalpic contribution that was attributed to the modulation of binding site permittivity ϵ_r in FimH_{FL} is considerably diminished and even reversed for **6** and **7**. Instead, a progressive shift toward entropic binding is observed.

When the thermodynamic profiles of **5–7** are compared with the parent compound **4**, notable dissimilarities between the two protein constructs, FimH_{LD} (Fig. 6A) and FimH_{FL} (Fig. 6B) can be observed. For FimH_{LD}, the enthalpic penalty for removal of a hydroxyl group amounts to 26–32 kJ mol^{−1}. The same modification in FimH_{FL}, however, results in an enthalpic penalty that ranges from 42 kJ mol^{−1} up to 65 kJ mol^{−1}. The discrepancy suggests that the loss of a hydrogen bond in the interaction of **5–7** and FimH_{FL} is superimposed by global




Fig. 5 Thermodynamic profiles (see Fig. S5–S8, ESI†) for the interaction of deoxygenated *n*-heptyl α -D-mannoside derivatives **5–8** with FimH_{LD} (A) and FimH_{FL} (B). Data for the interactions of FimH_{LD} with compounds **5–7** have been published elsewhere.³⁰ (C) Difference in the thermodynamic profiles between ligand binding to FimH_{FL} and FimH_{LD}. Error bars represent (propagated) 68% confidence intervals.




Fig. 6 Difference in thermodynamic profiles between **4** and deoxygenated *n*-heptyl α -D-mannoside derivatives **5–7** with FimH_{LD} (A) or FimH_{FL} (B). Error bars represent propagated 68% confidence intervals.



binding site permittivity could represent a general mechanism for carbohydrate binding proteins to enable enthalpy-driven recognition of polar ligands in aqueous solution.

Thus, the observed thermodynamics are not indicative of the protein–ligand interaction itself, but instead reflect a shift in the conformational equilibrium of the protein. This mechanism has been termed entropy–enthalpy transduction.³⁶ The existence of distinctive enthalpy–entropy compensation profiles for different ligands might also indicate that binding does not proceed *via* selection of a preexisting FimH_{FL} conformation (Fig. 2, pathway B), but through a ligand-induced fit (Fig. 2, pathway C). The conformational equilibrium in a conformational selection model is an inherent property of the protein and does not depend on the nature of a ligand. Thus, $\Delta G_{\text{conf-sel}}^{\circ}$ must be a constant term, and differences in ligand thermodynamics originate solely from interaction with the binding-competent conformation ($\Delta G_{\text{bind-closed}}^{\circ}$). In an induced-fit model, however, formation of an encounter complex ($\Delta G_{\text{bind-copen}}^{\circ}$) and contrasting consequences of the conformational transition $\Delta G_{\text{ind-fit}}^{\circ}$ (e.g. loss of enthalpic benefit from solvent shielding) can account for differences between ligands. For ligands with an intact mannoside core (**1–4**), the induced-fit transition reaches its full effect so that the differential thermodynamic profiles (Fig. 3C) appear constant.

Intriguingly, X-ray and NMR experiments yielded no information about the binding mode of **8** to FimH_{LD}. ITC data suggests that this compound actually binds with very low, but similar affinity to the high- and low-affinity construct of the protein (Fig. 5). These observations could signify a different binding mode that does not proceed *via* a conformational rearrangement in FimH_{FL}.

Conclusions

In conclusion, we have investigated the binding thermodynamics of two FimH constructs that share a similar binding mode but explore a very different conformational landscape. We could demonstrate that an observed enthalpic contribution of roughly -13 kJ mol^{-1} for mannoside ligands is a direct consequence of solvent shielding enabled by the rearrangement of the clamp loop in FimH_{FL} and the associated modulation of local dielectric properties. When the hydrogen bond network in the binding site is disrupted, the enthalpic benefit is completely abolished (Fig. 5C). In the case of FimH_{FL}, the effect of the solvent shielding mechanism is superimposed by an entropic penalty for the arrest of the clamp loop dynamics.

A similar transition of the binding site into a lower dielectric environment was observed for a number of other carbohydrate binding proteins, such as arabinose-binding protein (ABP), glucose/galactose-binding protein (GGBP) and sialic acid-binding periplasmic protein (SiaP). Their conformational transition upon ligand binding involves a twist of rigid protein domains around a fixed axis (“hinge-bending” motions).^{37–40} These transitions do not alter the flexibility of individual protein domains to an extent as observed in FimH and can be considered entropically neutral. Thus, modulation of

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

B. E. gratefully acknowledges funding from the Swiss National Science Foundation (PZ) and German Academic Exchange Service (BF).

Notes and references

- 1 H.-J. Böhm and G. Klebe, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 2588–2614.
- 2 R. E. Babine and S. L. Bender, *Chem. Rev.*, 1997, **97**, 1359–1472.
- 3 G. M. Keserü and D. C. Swinney, *Thermodynamics and Kinetics of Drug Binding*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2015.
- 4 S. Cabani, P. Gianni, V. Mollica and L. Lepori, *J. Solution Chem.*, 1981, **10**, 563–595.
- 5 E. Barratt, A. Bronowska, J. Vondrášek, J. Černý, R. Bingham, S. Phillips and S. W. Homans, *J. Mol. Biol.*, 2006, **362**, 994–1003.
- 6 G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford University Press, Oxford, 1997.
- 7 T. Steiner, *Angew. Chem., Int. Ed.*, 2002, **41**, 48–76.
- 8 C. P. Sager, D. Eriş, M. Smieško, R. Hevey and B. Ernst, *Beilstein J. Org. Chem.*, 2017, **13**, 2584–2595.
- 9 A. Vedani and J. D. Dunitz, *J. Am. Chem. Soc.*, 1985, **107**, 7653–7658.
- 10 J. Cramer, C. P. Sager and B. Ernst, *J. Med. Chem.*, 2019, **62**, 8915–8930.
- 11 A. V. Efimov and E. V. Brazhnikov, *FEBS Lett.*, 2003, **554**, 389–393.
- 12 G. Némethy, I. Z. Steinberg and H. A. Scheraga, *Biopolymers*, 1963, **1**, 43–69.
- 13 A. H. Daranas, H. Shimizu and S. W. Homans, *J. Am. Chem. Soc.*, 2004, **126**, 11870–11876.
- 14 S.-O. Shan and D. Herschlag, *Proc. Natl. Acad. Sci. U. S. A.*, 1996, **93**, 14474–14479.
- 15 M. M. Malabanan, T. L. Amyes and J. P. Richard, *Curr. Opin. Struct. Biol.*, 2010, **20**, 702–710.
- 16 D. G. Isom, C. A. Castañeda, B. R. Cannon, P. D. Velu and E. B. García-Moreno, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 16096–16100.
- 17 C. A. Fitch, D. A. Karp, K. K. Lee, W. E. Stites, E. E. Lattman and E. B. García-Moreno, *Biophys. J.*, 2002, **82**, 3289–3304.
- 18 E. L. Mehler, M. Fuxreiter, I. Simon and E. B. Garcia-Moreno, *Proteins: Struct., Funct., Genet.*, 2002, **48**, 283–292.
- 19 M. A. Mulvey, J. D. Schilling, J. J. Martinez and S. J. Hultgren, *Proc. Natl. Acad. Sci. U. S. A.*, 2000, **97**, 8829–8835.



- 20 J. D. Schilling, M. A. Mulvey and S. J. Hultgren, *J. Infect. Dis.*, 2001, **183**, S36–S40.
- 21 C. H. Jones, J. S. Pinkner, R. Roth, J. Heuser, A. V. Nicholes, S. N. Abraham and S. J. Hultgren, *Proc. Natl. Acad. Sci. U. S. A.*, 1995, **92**, 2081–2085.
- 22 I. Le Trong, P. Aprikian, B. A. Kidd, M. Forero-Shelton, V. Tchesnokova, P. Rajagopal, V. Rodriguez, G. Interlandi, R. Klevit, V. Vogel, R. E. Stenkamp, E. V. Sokurenko and W. E. Thomas, *Cell*, 2010, **141**, 645–655.
- 23 M. M. Sauer, R. P. Jakob, J. Eras, S. Baday, D. Eriş, G. Navarra, S. Bernèche, B. Ernst, T. Maier and R. Glockshuber, *Nat. Commun.*, 2016, **7**, 10738.
- 24 K. Mayer, D. Eris, O. Schwardt, C. P. Sager, S. Rabbani, S. Kleeb and B. Ernst, *J. Med. Chem.*, 2017, **60**, 5646–5662.
- 25 J. Bouckaert, J. Berglund, M. Schembri, E. De Genst, L. Cools, M. Wuhler, C.-S. Hung, J. Pinkner, R. Slättegård, A. Zavialov, D. Choudhury, S. Langermann, S. J. Hultgren, L. Wyns, P. Klemm, S. Oscarson, S. D. Knight and H. De Greve, *Mol. Microbiol.*, 2004, **55**, 441–455.
- 26 B. Fiege, S. Rabbani, R. C. Preston, R. P. Jakob, P. Zihlmann, O. Schwardt, X. Jiang, T. Maier and B. Ernst, *ChemBioChem*, 2015, **16**, 1235–1246.
- 27 W. Schönemann, J. Cramer, T. Mühlethaler, B. Fiege, M. Silbermann, S. Rabbani, P. Dätwyler, P. Zihlmann, R. P. Jakob, C. P. Sager, M. Smiesko, O. Schwardt, T. Maier and B. Ernst, *ChemMedChem*, 2019, **14**, 749–757.
- 28 V. Kalas, J. S. Pinkner, T. J. Hannan, M. E. Hibbing, K. W. Dodson, A. S. Holehouse, H. Zhang, N. H. Tolia, M. L. Gross, R. V. Pappu, J. Janetka and S. J. Hultgren, *Sci. Adv.*, 2017, **3**, e1601944.
- 29 G. Interlandi and W. E. Thomas, *Proteins: Struct., Funct., Bioinf.*, 2016, **84**, 990–1008.
- 30 P. Zihlmann, M. Silbermann, T. Sharpe, X. Jiang, T. Mühlethaler, R. P. Jakob, S. Rabbani, C. P. Sager, P. Frei, L. Pang, T. Maier and B. Ernst, *Chem. – Eur. J.*, 2018, **24**, 13049–13057.
- 31 S. Kleeb, L. Pang, K. Mayer, D. Eris, A. Sigl, R. C. Preston, P. Zihlmann, T. Sharpe, R. P. Jakob, D. Abgottspon, A. S. Hutter, M. Scharenberg, X. Jiang, G. Navarra, S. Rabbani, M. Smiesko, N. Lüdin, J. Bezençon, O. Schwardt, T. Maier and B. Ernst, *J. Med. Chem.*, 2015, **58**, 2221–2239.
- 32 J. Lameira, R. P. Bora, Z. T. Chu and A. Warshel, *Proteins: Struct., Funct., Bioinf.*, 2015, **83**, 318–330.
- 33 G. Jindal and A. Warshel, *Proteins: Struct., Funct., Bioinf.*, 2017, **85**, 2157–2161.
- 34 C. N. Schutz and A. Warshel, *Proteins: Struct., Funct., Genet.*, 2001, **44**, 400–417.
- 35 A. Warshel, P. K. Sharma, M. Kato and W. W. Parson, *Biochim. Biophys. Acta, Proteins Proteomics*, 2006, **1764**, 1647–1676.
- 36 A. T. Fenley, H. S. Muddana and M. K. Gilson, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 20006–20011.
- 37 M. J. Borrok, L. L. Kiessling and K. T. Forest, *Protein Sci.*, 2007, **16**, 1032–1041.
- 38 L. Unione, G. Ortega, A. Mallagaray, F. Corzana, J. Pérez-Castells, A. Canales, J. Jiménez-Barbero and O. Millet, *ACS Chem. Biol.*, 2016, **11**, 2149–2157.
- 39 A. Müller, E. Severi, C. Mulligan, A. G. Watts, D. J. Kelly, K. S. Wilson, A. J. Wilkinson and G. H. Thomas, *J. Biol. Chem.*, 2006, **281**, 22212–22222.
- 40 J. F. Darby, A. P. Hopkins, S. Shimizu, S. M. Roberts, J. A. Brannigan, J. P. Turkenburg, G. H. Thomas, R. E. Hubbard and M. Fischer, *J. Am. Chem. Soc.*, 2019, **141**, 15818–15826.

