

rsc.li/obc



ISSN 1477-0520

REVIEW

View Article Online
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2021, **19**, 4380

Received 9th February 2021,
Accepted 29th March 2021

DOI: 10.1039/d1ob00257k

rscl.li/obc

Chemical probes targeting the kinase CK2: a journey outside the catalytic box

Jessica Iegre, ^{†a} Eleanor L. Atkinson, ^{†a} Paul D. Brear, ^b Bethany M. Cooper, ^a Marko Hyvönen ^b and David R. Spring ^{*a}

CK2 is a protein kinase that plays important roles in many physio-pathological cellular processes. As such, the development of chemical probes for CK2 has received increasing attention in the past decade with more than 40 lead compounds developed. In this review, we aim to provide the reader with a comprehensive overview of the chemical probes acting outside the highly-conserved ATP-site developed to date. Such probes belong to different classes of molecules spanning from small molecules to peptides, act with a range of mechanisms of action and some of them present themselves as promising tools to investigate the biology of CK2 and therefore develop therapeutics for many disease areas including cancer and COVID-19.

Introduction

Protein kinases are considered the master regulators of the cell machinery. They exert their functions through the phosphorylation of other proteins with consequent activation or deactivation of crucially important cellular signaling pathways.¹

The essential cellular processes that kinases regulate comprise cell growth, proliferation, differentiation, and death. Therefore, it is unsurprising that gene mutations that alter the function or expression levels of protein kinases lead to the insurgence of several diseases including neurological, immunological, hematological, endocrine, and skeletal disorders, as well as cancer.² The implication of protein kinases in disease generation and progression has been widely recognized by the pharmaceutical community, and, as of today, there are 75 FDA approved drugs targeting this class of proteins.³

^aDepartment of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK. E-mail: spring@ch.cam.ac.uk

^bDepartment of Biochemistry, University of Cambridge, 80 Tennis Court Road, Cambridge, CB2 1GA, UK

[†]These authors contributed equally to the work.



Jessica Iegre

Jessica Iegre was born in Italy and obtained her MSci in Medicinal Chemistry and Pharmaceutical Technology at the University of Pisa, Italy in 2013. The same year, she joined the AstraZeneca IMED Graduate programme in Göteborg, Sweden where she spent 2 years working across three different departments: medicinal chemistry, DMPK and computational chemistry. In 2015 she joined the Spring group at the University of

Cambridge, and she obtained her PhD in chemical biology in 2019. Jessica is currently a Postdoctoral Research Associate in the group, and she is developing novel stapled peptides to inhibit medically relevant protein-protein interactions.



Eleanor L. Atkinson

Eleanor L. Atkinson received her MChem degree from the University of Oxford in 2018 after completing her Masters' project under the supervision of Professor Angela Russell, investigating the metabolism of novel drug candidates. In 2018, she started her PhD studies at the University of Cambridge under the supervision of Professor David Spring, where her research focuses on the development of novel stapled peptides for the inhibition of protein-protein interactions.



Among the >500 protein kinases identified in the human proteome, there is CK2s (formerly referred to as Casein Kinase 2).⁴ CK2 α and its very similar paralog CK2 α' are a serine/threonine kinases belonging to the CMCG family. Unlike other eukaryotic kinases, CK2 α is constitutively active and as such, does not require upstream phosphorylation to exert its function.⁵ They form a tetrameric complex with a non-catalytic scaffolding subunit CK2 β . CK2 phosphorylates more than 300 substrates and is thus one of the most promiscuous kinase we know.⁶ CK2 is found in both healthy and cancerous cells – whilst the protein levels in healthy cells increase during proliferation only, cancerous cells are characterized by overexpression of the different subunits which contributes to their rapid growth, proliferation, and their ability to escape apoptosis.⁷ Breast, lung, prostate, colorectal, renal cancer, leukemia, and glioblastoma are among the cancerous tissues featuring particularly high CK2 content, and hence, CK2 inhibitors could prove particularly useful as therapeutic agents for the treatment of these types of cancer.⁸ Indeed, an orally available CK2 α inhibitor, named **CX4945** (silmitasertib), is currently undergoing clinical studies for the treatment of cholangiocarcinoma paving the way for CK2 to be validated as an anticancer target.⁹

Although the role of CK2 in cancer progression is the most studied, it should be noted that the kinase is also involved in other pathological processes. For example, recently CK2 has been found to be interacting with the nucleocapsid of SARS-CoV-2, possibly contributing to COVID-19.^{10,11} Studies probing this relationship are in their infancy, however, it is possible that CK2 may represent a valid target for the treatment of coronaviruses in the future.

Many are the mechanisms by which CK2 can be inhibited, and therefore, there is a large number of inhibitors and chemical probes developed to date: from small molecules to peptide and peptidomimetics, to polyoxometalates.¹² Similar to other kinases, the most studied mode of inhibition is ATP-competitive, where a ligand competes with ATP and hence prevents substrate phosphorylation. Though effective, one of the draw-

backs of this strategy is that the ATP binding site is highly conserved across the kinome, leading to numerous challenges when developing a selective chemical probe for the target kinase.^{13,14} Most CK2 α inhibitors serve as an example to this problem, including the clinical trial candidate **CX4945**.⁹ In addition, cellular concentration of ATP is high and therefore high-affinity ATP-competitive inhibitors are needed to effectively reduce the activity of the target. To overcome this issue, more selective strategies of CK2 inhibition have been explored including targeting of sites outside the ATP pocket, inhibition of the formation of the holoenzyme, and displacement of endogenous substrates.

In this review, we aim to provide a comprehensive overview of the molecular and structural biology of CK2, mechanisms of inhibition, and chemical probes for CK2 developed to date with emphasis on those acting outside the catalytic box (Fig. 1). In addition to describing their discovery and applications, this review provides a critical view of the challenges and opportunities of this promising class of chemical probes.

CK2 kinase

Protein structure and characteristics

The CK2 holoenzyme is a tetramer constituted by two catalytic subunits, α and/or α' connected by a dimer of the two regulatory subunits, β (Fig. 1). The α subunit presents 20 additional amino acids at the C-terminus that are absent in α' . The catalytic domain (~48 kDa) is responsible for the phosphorylation of protein substrates and comprises the ATP binding site between the N- and C-terminal domains, linked *via* the so-called α D region. The N-terminal domain comprises of one α -helix, named α C and five β -sheets, β 1– β 5. The α C helix contacts the kinase substrates and, due to its lysine-rich content, imparts to the kinase a preference for acidic substrates. The C-terminal domain of CK2 α comprises seven α -helices and two β -sheets that form the floor of the ATP binding site and the



Paul D. Brear

Paul Brear obtained his MChem degree from the University of Durham in 2008 and obtained his PhD from the University of St Andrews in 2012 under the supervision of Prof. Nicholas Westwood and Dr Rupert J. Russell. He joined the group of Dr Marko Hyvönen at the University of Cambridge in 2012 and from 2018 is also the X-ray crystallography facility manager at the biochemistry department of the University of Cambridge.



Bethany M. Cooper

Bethany M. Cooper received her MChem degree from the University of Leeds in 2018, having completed her 3rd year of study at Lubrizol Ltd, Hazelwood. On return to the University of Leeds her final year was spent under the supervision of Professor Steve Marsden. In 2019, she started her PhD studies at the University of Cambridge under the supervision of Professor David Spring and industrial supervisor

Dr Maria Öhwegård-Halvarsson, where her research has focused on peptide stapling methodologies for use within therapeutics.






Fig. 1 The structure of the CK2 holoenzyme is shown on the top right (PDB: 1JWH). The surfaces of two catalytic subunits are shown in grey and green; the surfaces of the two regulatory subunits are shown in pink and purple. Key binding sites and inhibitors of CK2 α are shown on the left (ATP-site in white, α/β interface site in purple, substrate channel in black and α D pocket in light blue). Key binding site and inhibitor of CK2 β is shown on the bottom right.

activation loop. The α D pocket, a pocket that has been recently discovered and that can be exploited to achieve selective CK2 inhibition, is located near the α D helix of the C-terminal domain.^{15,16} It is the conformation of the α C helix and the DWG motif in the activation segment that make the kinase constitutively active. In particular, the protein's N-terminus makes extensive contact with the α C helix which remains locked in an open conformation and allows for substrate

binding. Unlike other closely related kinases, CK2 features a DWG motif (Asp175–Trp176–Gly177) in the activation loop rather than a DFG motif. The Phe to Trp mutation allows the creation of an additional H-bond between the indole of Trp176 and the Leu173 which maintains the kinase in an active state. This constitutive activity means that, notably, the majority of CK2 substrates do not require the presence of CK2 β to be phosphorylated.^{16,17}



Marko Hyvönen

Marko Hyvönen is a research group head at the University of Cambridge. He got his undergraduate degree from University of Helsinki and did his PhD work at EMBL in Heidelberg, Germany, graduating in 1997. He moved to Cambridge in 1998 as a postdoctoral fellow in the group of Tom Blundell. In 2001 he was awarded a BBSRC David Phillips Fellowship, appointed to a University Lectureship in 2008 and became a Reader in 2018. Dr Hyvönen and his group use structural biology techniques to study the structure and function of proteins and for structure-guided and fragment-based drug discovery.



David R. Spring

David Spring is currently Professor of Chemistry and Chemical Biology at the University of Cambridge within the Chemistry Department. He received his DPhil (1998) at Oxford University under Sir Jack Baldwin. He then worked as a Wellcome Trust Postdoctoral Fellow at Harvard University with Stuart Schreiber (1999–2001), after which he joined the faculty at the University of Cambridge. His research programme is focused on the use of chemistry to explore biology.



Org. Biomol. Chem., 2021, 19, 4380–4396 | 4383




Fig. 3 Schematic of the six hallmarks in cancer and the involvement of CK2 in each of them. Where specific proteins are known to be affected by CK2 with respect to each of these hallmarks, examples are given: proteins attached to a P are phosphorylated by CK2, proteins in green are stabilised by CK2 and proteins in yellow have increased expression due to CK2. Despite evidence that CK2 induces angiogenesis and enables replicative immortality, the specific pathways affected by CK2 are not currently known.^{24,32–39} Created with BioRender.com.

bition of cell growth, proliferation and an increase in apoptotic activity with no redundant pathways able to compensate for such downregulation.

Here, we summarise key findings that support the relevance of CK2 in different types of cancer. The reader is directed to several reviews for more information on the importance of CK2 in other cancer types.^{7,40}

Lung cancer. A study by O-Charoenrat *et al.* in 2004, demonstrated the correlation between overexpression of the CK2 α transcript and the poor prognosis outcome determined through relapse timeframes and overall survival of patients with lung cancer.⁴¹ The CK2 response differed depending on the subtype of lung cancer: for non-small cell lung cancer (NSCLC) and adenocarcinoma cell lines, MMP-2 transcript expression and ERK pathway activity was downregulated as a result of CK2 inhibition, resulting in decreased cell migration and invasion.⁴² Furthermore, Benavent *et al.* demonstrated that CIBG-300, an anti-CK2 peptide, inhibited lung cell adhesion, invasion and migration through *in vitro* studies.⁴³

Colorectal cancer. It is well documented that within colorectal cancer (CRC) cells there is an overexpression of the CK2 gene transcript. Lin *et al.* found a correlation between the overexpression of the CK2 β subunit and the prognosis outcome and hence survival rate.⁴⁴ In addition, Zou *et al.* showed that CK2 α was overexpressed in tissues from 144 patients with colorectal abnormalities.⁴⁵ Suppression of CK2 α resulted in G₀/G₁ phase arrest, enhanced expression of p53/p21, reduced expression of C-Myc and induced cell senescence. In addition, knockdown of CK2 α resulted in the inhibition of cell invasion and motility.

Cholangiocarcinoma. A study by Zhou *et al.* determined that the CK2 β subunit is overexpressed in cholangiocarcinoma (CAA) tumorigenesis in comparison to normal epithelial liver cells, and that there is a correlation between the level of CK2 β and the tumour progression.⁴⁶ CK2 α was also identified as a potential biomarker for CAA.⁴⁷ In 2016 the US FDA granted

CX4945, a CK2 inhibitor, Orphan Drug Designation for the treatment of cholangiocarcinoma, highlighting the potential of CK2 as an anticancer target.³¹

Breast cancer. Studies have determined that CK2 α and CK2 β are overexpressed in many different types of breast cancer.⁷ Whilst in all breast cancer types the CK2 α' subunit is under-expressed in both invasive and non-invasive cells, CK2 β was found to be over-expressed only in invasive breast cancer cells. It was determined that the two subunits which influenced survival rate were CK2 α and CK2 β , with CK2 α' having no significant effect.⁷

Importance of CK2 in COVID-19

CK2 is emerging within the infectious disease field as an attractive target. There are validated studies regarding CK2 phosphorylation and modulation of viral proteins' function in human immunodeficiency virus (HIV), vesicular stomatitis virus (VSV), hepatitis C virus (HCV), herpes simplex-1 (HSV-1) and human papilloma virus (HPV).³⁰ CK2 also plays a key role in a number of signalling cascades that viruses hijack including JAK/STAT, PTEN/PI3K/Akt-PKB and NF- κ B.³⁰ It is thought that if CK2 was inhibited, viral replication would be impacted. A recent study by Bouhaddou *et al.* used a combination of 87 kinase inhibitors to identify those with antiviral efficacy. Kinases p38, CK2, CDK, PIKFYVE and AXL were identified as having antiviral potential. The conducted experiments demonstrated the instrumental role of CK2 inhibition for the SARS-CoV2 infection *in vitro*.^{30,48} SARS-CoV2 nucleocapsid protein has been shown to directly interact with CK2. By enabling co-localization along the filopodia protrusions, this interaction promotes several functions including virus egress and cell-to-cell spread through infected cells' epithelial monolayers.^{30,48} Particularly for CK2, the virus-host PPI was associated with increased activity of CK2 as documented by the enhanced phosphorylation of CK2 substrates such as STAT1, HMGN 1, HMGA 1, HDAC 2 and CTNNA 1.⁴⁸

The potential of CK2 as an anti-COVID-19 target is documented by two small clinical trials using previously developed CK2 inhibitors: CX4945 and CIGB-325. In particular, it was found that CX4945 was able to improve the oxygen level of patients with COVID-19-induced pneumonia within 24 hours of starting treatment, and a further five days later the first patient was discharged.³¹ Similarly, a small study was conducted on twenty SARS-CoV-2 positive patients using the anti-CK2 peptide CIGB-325 (formerly CIGB-325). It was found that 50% of COVID-19 positive patients that were treated with CIGB-325 had a reduction in the number of pulmonary lesions by day 7 in comparison to only 20% in the standard-of-care only treatment.³⁰

Inhibition mechanism and CK2 inhibitors

Approaches to the discovery of CK2 inhibitors used so far

The first CK2 inhibitors were identified by the investigation into biologically active compounds and natural products, such



as heparin and emodin.^{49,50} However, with the rapid growth in the field of computational chemistry and structural biology, more recently inhibitors have been designed specifically for CK2 using structural-based approaches such as fragment-based drug discovery and phage display. The rapid growth in technologies facilitating the guided and rational design of protein modulators has had a drastic impact upon the development of CK2 inhibitors, as is self-evident from the rapid growth in the field over the past 20 years.

From the many inhibitor development studies targeting CK2, multiple different binding sites have been identified on CK2 α . These sites all have different advantages and disadvantages which have led to them being utilised with varying success for the inhibition of CK2. These sites can be initially grouped into four broad categories: the ATP site, CK2 β protein-protein interaction site, the substrate binding channel and potential allosteric sites outside of the classic ATP site (Fig. 4a).

In the next sections, we will focus on the approaches used to discover CK2 inhibitors targeting sites outside the catalytic box and we aim to provide a comprehensive overview of the molecules reported to date. A brief overview of the ATP competitive inhibitors is also provided.

Overview of ATP competitive inhibitors

The site where the most development has happened is the ATP site. This site is the where the protein naturally binds the small molecule cofactor, and as such, is seen as the easiest site to target due to its characteristics. Firstly, the site is pre-formed in the apo structure meaning that there is no barrier to ligand binding. Secondly, the site is composed of a deep hydrophobic cleft sandwiched by Lys68 and the hinge region at opposite ends. This layout is ideally set up for sandwiching an aromatic ring system with hydrogen bonding substituents protruding off either side (Fig. 4b).

In a recent crystallographic fragment screen conducted by Brear *et al.*, the ATP site was the only site identified as binding fragments.⁵¹ This is significant as fragment screens are often used to probe the surface of proteins for small-molecule

binding sites. The ATP site has been used with great success in identifying ligands that bind with high affinity and displace the endogenous ATP hence preventing the phosphorylation of the substrates. Indeed, a number of small molecules with picomolar affinities have been identified for this site.⁵²

CK2 ATP-competitive inhibitors can broadly be divided into four main classes: halogenated compounds such as **DRB** and its derivatives,^{53,54} condensed polyphenolic compounds such as emodin and its derivatives,^{50,55} pyrazolo-triazines and pyrazolo-pyrimidines^{56,57} and indoloquinazolines such as **CX4945**.⁹ The chronology of the most influential inhibitors developed thus far is detailed in Fig. 5.

The reader is redirected to a recent review for an up to date comprehensive review on ATP-competitive inhibitors for CK2.⁵⁸

Unfortunately, there are a number of drawbacks associated with the use of this site, with the primary issue being selectivity. The ATP site has evolved to bind ATP which is a very common cofactor amongst the genome. Indeed, Uniprot annotates 1391 human genes as binding ATP. Therefore, any small molecule that targets just the ATP site is likely to bind and inhibit any number of these other ATP-binding proteins. An example highlighting the selectivity issue of this class of compounds is given by the small molecule clinical candidate **CX4945** which is frequently described in the literature as highly selective for CK2 with an IC₅₀ of 1 nM. However, when screened in a selectivity panel of 238 kinases, **CX4945** also inhibits another seven kinases more than 90% (tested at 500 nM). The most significant is the effect seen with Dyrk1A and 1B where **CX4945** showed IC₅₀ values of 6.8 and 6.4 nM, respectively.⁹ A number of efforts are currently ongoing to identify new selective series of CK2 ATP-competitive inhibitors.

In 2016, Dowling *et al.* identified compound **7h** as the most promising CK2 α inhibitors out of a pyrazolepyrimidine series. The compound showed picomolar activity in an SPR assay and was found to inhibit the Wnt pathways with an IC₅₀ of 50 nM. Despite its limited oral bioavailability, **7h** presented promising pharmacokinetic properties after IV administration, and it was effective as monotherapy in both HCT116 cells and




Fig. 4 Crystal structure of CK2 α in complex with ADP (PDB: 6YPN). (a) The small-molecule binding sites on CK2 α . The ATP site is shown in red and the sites outside the ATP pocket are shown in blue. (b) and (c) The ATP site of CK2 α . The hinge residues are shown in cyan, the polar phosphate binding residues are shown in green and the hydrophobic sandwiching residues are shown in grey.





Fig. 5 Structures of the most influential ATP-competitive inhibitors of CK2 reported in chronological order. The X-ray co-crystal structure of CX4945 and CK2 is shown on the bottom left (PDB: 3NGA). Residues which hydrogen bond with CX4945 are shown in cyan alongside their hydrogen bond lengths.

SW-620 xenografts. The compound also showed limited off-target activity when screened in a panel of 402 kinases with 12 members of the CMGC family being inhibited more than 50%.⁵²

Oshima *et al.* identified GO289 in 2019 as having excellent activity *in vitro* ($IC_{50} = 7$ nM) as well as high selectivity for CK2; the second most inhibited protein, PIM2, is much more weakly inhibited by GO289 than CK2 (PIM2 $IC_{50} = 13$ μ M).⁵⁹ In 2020, SRPIN803-rev-derivatives were found to have moderate inhibitory activity against CK2 ($IC_{50} = 0.28$ μ M for the lead compound) as well as good selectivity (only CK2 α inhibited over 50% at 1 μ M in a 320 kinase panel) and cellular activity ($DC_{50} = 12.8$ μ M, Jurkat cells).⁶⁰

Following on from Dowling's work on pyrazolopyrimidines, in 2020 Wells *et al.* developed compound 24.⁶¹ The lead compound proved to be selective against a panel of 402 kinases and showed an IC_{50} of 36 nM and 16 nM against CK2 α and CK2 α' respectively in a nanoBRET assay carried out in Hek-293 cells. While compound 24 showed target-specific inhibition of Akt Ser129 phosphorylation, it did not activate Caspase 3/7 as been seen with other CK2 α inhibitors and had variable effects on proliferation in a panel of cancer cell lines.⁶¹

Most of the newly identified series of inhibitors require further optimisation, characterisation, and evaluation before highly potent, selective and fully understood cell-permeable CK2 inhibitors are obtained, but all classes represent promising starting points for this to take place. Importantly, whilst minor off target effects could be beneficial in cancer treatment, they are not desirable while investigating signalling pathways related to a targeted protein.

In order to overcome selectivity issues associated with the use of ATP competitive inhibitors, developers of kinase inhibitors are looking for alternative mechanisms of inhibition. This

is also true for CK2 inhibition. Within CK2 α alternative binding sites can be split into four categories: the substrate-binding site, the CK2 β interface site, the α D site, and possible allosteric sites. As of yet, only the α D site, the CK2 β interface site and the substrate-binding channel have been successfully utilised and validated for the development of inhibitors (Fig. 4a). As of today, no allosteric, non-competitive inhibitors are known for CK2.

Substrate channel inhibitors

An alternative strategy for CK2 inhibition is through blocking of the substrate-binding site only. One of CK2's distinctive features is its substrate preferences – it phosphorylates only acidophilic substrates.⁶² Therefore, greater selectivity of CK2 modulation may be achieved through the development of substrate-competitive inhibitors.

Polyglutamyl peptides were identified as CK2 substrate-competitive inhibitors in 1983 by Meggio *et al.* It was found that (Glu)₇₀ was able to inhibit CK2 *in vitro* ($K_i = 0.11$ μ M) but was inactive towards the closely related CK1 and cAMP-dependent protein kinase.⁶³ They subsequently identified the hexapeptide RSEEEVE as a very weak substrate-competitive inhibitor of CK2 ($K_i = 3.98$ M)⁶⁴ as well as the random heteropolymers Glu/Tyr (4 : 1) and Glu/Tyr (1 : 1) which had good activities against CK2 ($K_i = 25$ nM, 50 nM respectively).⁶⁵ Successive work by Tellez *et al.* in the early 1990s further reinforced the notion that polymers incorporating Glu and Tyr were able to inhibit CK2.^{66,67}

Through screening of a random cyclic peptide phage display library, Perea *et al.* identified peptide P15 (WMSPRHLGT) which prevented CK2 phosphorylation *in vitro* in a substrate-competitive manner.⁶⁸ Addition of two cysteine residues to either end of the peptide, as well as attachment of



Org. Biomol. Chem., 2021, **19**, 4380–4396 | 4387



Fig. 7 Bi-specific ATP/substrate competitive inhibitors discovered to date. The co-crystal structure of **ARC1502** and CK2 α is shown in the bottom left (PDB: 6SPX). Residues which hydrogen bond to the ligand are shown in cyan alongside the hydrogen bond lengths. Portion of the molecule binding in the ATP site is shown in red, the motif binding to the substrate channel is shown in black. For **ARC-1513-5O** the fluorophore is shown in blue.

whereas **K137-E4** inhibited CK2 the most strongly when tested against 140 other enzymes. However, **K137-E4** is not cell permeable and therefore exhibits negligible biological effects in cellular assays.⁷³

In 2017, Vahter *et al.* developed a series of bi-substrate inhibitors using a fragment derived from **CX4945** as the ATP-competitive inhibiting portion of the compound.⁷⁶ **ARC-1513-5O** showed exceptional binding ability to CK2 ($K_d = 16\text{--}37$ pM, Fig. 7). Similarly to **CX4945**, **ARC-1424-5O** showed moderate selectivity when tested in a panel of 140 protein kinases; when tested at 1 μM , 23 kinases were inhibited more than 50%, including 6 which were strongly inhibited (>90%).⁷⁶ Despite its promiscuity, **ARC-1424-5O** remains one of the best candidates for the measurement of binding affinities of other CK2 inhibitors in competition experiments due to its picomolar activity.

The field of bi-substrate inhibitors of CK2 is relatively new compared to ATP-competitive inhibitors. However, the compounds identified so far highlight that it is a promising strategy of inhibition for the development of both potent and selective inhibitors of CK2, but pharmacological properties need to be improved before these can be considered as viable developmental candidates.

Inhibitors of the holoenzyme assembly

An alternative method to modulate the activity of CK2 is *via* disruption of the PPI at the CK2 α/β interface. CK2 α is rare for a kinase in that it is constitutively active.⁶ However, it forms a hetero-tetrameric complex with CK2 β which is believed to regulate the activity and substrate selectivity of CK2 α . CK2 substrates can be divided into three classes: Class I, II and III.⁷⁷ Class I can be phosphorylated by the CK2 α /CK2 β complex or by CK2 α alone; Class II can be phosphorylated by CK2 α alone; Class III can be phosphorylated only by the CK2 α /CK2 β

complex. It has been shown that preventing the formation of the complex prevents the phosphorylation of the Class III substrates, affects the localisation of the complex and destabilises CK2 α .⁷⁷ There are some reports of weak inhibition of the phosphorylation of Class I and II substrates by ligands binding at the CK2 β interface site.⁷⁸ However, higher affinity compounds that bind more selectively in the interface site and have been more reliably validated do not show this inhibition.⁷⁹ Therefore, it seems unlikely that this site can be used for the direct inhibition of all CK2 substrates. The dominant feature of the interface site is a pocket in which Phe190 binds, a shallow groove in which Tyr125 binds and a flexible loop (Val101-Pro109 loop) that opens and closes the interface site (Fig. 8a).⁸⁰ The size of this site varies depending on which ligands are bound in the site. For example, the Val101-Pro109 loop can be seen in a closed conformation in the structure (Fig. 8b grey cartoon, c grey surface). Whereas, in the CK2 β -bound structure the loop folds back to give a significantly more open pocket (Fig. 8b cyan loop, d grey surface). This range of movement allows the pocket to accommodate a large range of ligands and provides more potential for optimisation. The main advantage of the interface site over the ATP site is that this site is less conserved amongst kinases. This means there is a greater possibility of generating selective inhibitors using this site. However, there are also a number of drawbacks to this site. Firstly, generating high-affinity small molecule inhibitors using this site is less facile than the ATP site for which it is possible to access small molecules with a pico-molar affinity.^{52,81} This has a lot to do with the depth of the ATP site where inhibitors can make extensive contacts with the target, shielded from the solvent. In contrast, the interface site is a shallow pocket, or rather a dent, on the surface of the protein, which offers little three-dimensionality and contact.





Fig. 8 (a) CK2 β interface loop (purple) binding in the CK2 α interface site (grey) (PDB: 4MD7). (b) Flexibility of the interface site. Open/peptide bound structure in cyan (PDB: 6Q4Q), closed/small molecule bound structure in grey (PDB: 5CU6). (c) Surface of the closed/small molecule bound structure (PDB: 5CU6) and (d) surface of the open/peptide bound structure (PDB: 6Q4Q). Ligand is shown in green and the surface in grey.

The highest affinity reported so far for a small molecule binding at the interface site is 30 μ M (ref. 82) and higher affinity inhibitors for this site are peptides or stapled peptides. This is not surprising if we consider that bigger molecules like peptides better interact with the big and shallow pocket found at the interface.⁸³ The indirect nature of the inhibition from this pocket may be an advantage or a disadvantage to this site; however, more advanced chemical tools and testing are required to confirm this.

Small molecules. The first identified modulator of CK2 holoenzyme assembly was the small molecule 5,6-dichloro-1- β -D-ribofuranosyl-benzimidazole (**DRB**).^{78,84} As well as being an ATP-competitive inhibitor, X-ray structures showed that **DRB** was also binding weakly to CK2 α at the α/β interface. This was the first identification of the CK2 PPI as a druggable site. Although it disrupted the CK2 α/β assembly, **DRB** did not cause a biological response.⁷⁸ Due to its binding promiscuity, **DRB** now has limited use as a CK2 inhibitor. However, its role in identification of the PPI as a potential site of inhibition was key for the subsequent development of small molecule CK2 PPI inhibitors, the best of which are outlined in Fig. 9.^{78,82,85,86}




Fig. 9 CK2 inhibitors reported in chronological order and acting at the protein interface between the catalytic and regulatory subunit with the co-crystal structure of CK2 α and **CAM7117** shown on the left (PDB: 6Q4Q). The hydrophobic Tyr and Phe pockets key to binding are highlighted, alongside the π - π stacking interactions of the ligand. sC18 = H₂N-GLRKRLKFRNKIEK-NH₂,⁹³ TAT = GRKKRRQRRRPPQ.⁹⁶ Compounds have been labelled as in the related papers. r = D-Arg.

The second small molecule modulator of the PPI was the podophyllotoxin indole-analogue **W16**, identified as having a modest IC_{50} of 30–40 μM in enzymatic assays (Fig. 9).⁸⁶ Kinetic analysis showed that **W16** is a non-competitive inhibitor of CK2 α and that this inhibition was reversed upon the addition of CK2 β or the CK2 β -mimicking peptide **Pc**, suggesting that **W16** binds in or close to the CK2 β -binding pocket on CK2 α . The inhibition of CK2 α may result from small conformational changes of the protein upon binding of **W16**.⁸⁶

Recently, Kröger *et al.* found that the enantiomer of **W16** was six times more potent than its analogue ($K_i = 4.9 \mu M$ vs. 31 μM respectively).⁸⁷ They also replaced the labile anhydride group of **W16** with a more stable imide and *N*-methylimide, increasing stability without the loss of binding affinity ($K_i = 3.6 \mu M$ vs. 2.8 μM respectively). However, the increased inhibition of the PPI did not result in a greater biological effect as the new analogues of **W16** were not seen to inhibit CK2. This clearly highlights that perturbation to the equilibrium of the CK2 holoenzyme complex formation does not necessarily lead to enzyme inhibition.⁸⁷ Due to the large aromatic structures of the podophyllotoxin indole-analogues, the compounds suffer from poor solubility making the likelihood of developing therapeutics derived from **W16** low.^{86,87}

A high-concentration X-ray crystallography fragment screen identified **1** (in Brear *et al.*) binding at both the CK2 α/β interface and in the ATP-pocket of CK2 with a weak IC_{50} (900 μM).⁸⁸ Iterative design-test cycles resulted in the development of the small molecule **CAM187** which had greatly improved bioactivity ($IC_{50} = 44 \mu M$) and did not show promiscuity (Fig. 9). Due to its significantly smaller size and, as such, more favourable physical properties than **W16**, the fragment **CAM187** is a much-improved starting point for the development of selective small molecule drug-like compounds to target the CK2 PPI. However, further analyses of its activity and selectivity against other kinases, including evaluation of its biological effect in cells are required.

With a virtual screening of a compound library followed by surface-plasmon resonance (SPR), NMR and crystallography, Kufareva *et al.* identified compound **6** ($K_d = 30 \mu M$) a novel CK2 PPI inhibiting molecule (Fig. 9).⁸² Compound **6** (in Kufareva *et al.*) successfully inhibited the phosphorylation of CK2 β -dependent substrates, suppressed MDA-MB231 triple-negative breast cancer cell growth and induced apoptosis in the μM range.⁸² As such, compound **6** is the first example of a small molecule inhibitor of the CK2 α/β interaction which has been shown to be effective in cells. Alongside **CAM187**, compound **6** represents an excellent starting point for the development of selective and potent small molecule therapeutics for the inhibition of CK2 *via* the perturbation of the CK2 PPI.

Peptides. In 2008, Laudet *et al.* developed a peptide to bind to CK2 α at the site of the CK2 β binding pocket.¹⁰⁷ Peptides represent good candidates for PPI modulators as their flexibility allows them to adapt to the large surfaces involved in PPIs.⁸⁹ They can be designed to mimic the section of the partner protein binding to the desired target region, hence overcoming the difficulties associated with *de novo* design.⁹⁰

Using the prior knowledge of residues Tyr188, Gly189 and Phe190 being the main source of affinity between the β -hairpin loop and CK2 α , in combination with the crystal structure of CK2 α bound to CK2 β , Laudet *et al.* designed a handful of CK2 β -derived peptides, the most promising of which was **Pc** (Fig. 9).^{15,91} **Pc** consists of the central interacting segment of CK2 β C-terminal loop cyclised by a disulfide bridge between two cysteine residues. The peptide was shown to antagonise the assembly of the holoenzyme complex ($IC_{50} = 3.0 \mu M$) as well as altering its substrate specificity.⁹¹ **Pc** represented the first peptide antagonist of CK2 which exerted its effect by binding to CK2 α and preventing the formation of the holoenzyme.⁹¹ However, **Pc** did not successfully exhibit cellular activity. To improve the cellular activity of the peptide, predicted to suffer from poor stability of disulfides in reducing intracellular environment, the disulfide bond was replaced with a head-to-tail cyclisation.⁹² Subsequently, the cyclic peptide was fused with the TAT cell-penetrating peptide which is known to facilitate transport across cell membranes.⁸⁹ The resulting compound, **TAT-Pc**, was shown to enter cells, inhibit the phosphorylation of CK2 β -dependent substrates and lead to caspase independent cell death.⁹² In 2019, three simultaneous but separate optimisations of the **Pc** and **TAT-Pc** peptides occurred leading to the development of **sC18-I-Pc**,⁹³ **I192F**⁹⁴ and **CAM7117**.⁷⁹

In 2015, Hochscherf *et al.* altered the Phe residue in the structure of **Pc** to include *para*-halogens. The iodinated-**Pc** peptide was the most effective, showing a slight improvement in K_d compared to **Pc** (0.24 vs. 0.56 μM).⁹⁵ In an attempt to make **I-Pc** cell-permeable, in 2019, the same research group introduced a cell-penetrating tag, namely, sC18⁹⁶ to translocate the peptide into cells. Although the introduction of sC18 resulted in a 4-time increase of the K_i , **sC18-I-Pc** successfully inhibited the phosphorylation of CK2 β -dependent substrates as well as evoking cell death in the mid- μM range (Fig. 9).⁹³

Tang *et al.* used structure-based computational design in combination with experimental evaluation to develop point mutated **Pc** derivatives.¹⁰⁸ The most promising mutant, **I192F** exhibited a 10-fold improvement in the predicted binding affinity to CK2 α compared to **Pc** (0.5 vs. 8.9 μM). When the TAT-conjugated cell-permeable derivative of the peptide was synthesised, **TAT-I192F** showed comparable anti-proliferative effects against HepG2 cells to **TAT-Pc** (20.1 μM vs. 30.4 μM).^{96,108}

The development of **CAM7117** took a different approach to the aforementioned peptides in that it was the cyclic constraint that was the main site of optimisation due to the lability of the disulfide bond in the **Pc** derivatives. Using a structure-based approach, a variety of covalent constraints were designed and tested to determine which held the peptide in the optimal conformation. In addition, molecular modelling and X-ray crystallography of the peptide sequence directed a point mutation in the central sequence of **Pc**, namely Ile 192 to Trp. Movement of one of the cyclising residues by one position and the use of CuAAC chemistry gave **CAM7117** (Fig. 9).⁷⁹ The peptide had an improved K_d value compared to **Pc** (0.2 vs. 1 μM respectively) in



an isothermal titration calorimetry (ITC) assay, was stable in human serum and cell-permeable. CAM7117 was able to inhibit cell growth with a GI_{50} of 33 μ M in U2OS cells. It was postulated that the difference between the enzymatic and the cellular activity was due to partial trapping in the Golgi apparatus.⁷⁹

In just over a decade, there has been a movement from the identification of CK2 α /CK2 β interaction site as a potential target for small molecule-directed inhibition of CK2 to the development of a handful of potent CK2 PPI inhibitors. Both the peptide and small molecule inhibitors have their limitations and require further optimisation before a sufficiently stable, extremely potent and highly selective cellular chemical probe or therapeutic is obtained for this site. However, the most promising inhibitors currently known are an excellent place for optimisation to continue from.

Inhibitors of CK2 α acting outside the ATP pocket

Other strategies to inhibit CK2 target binding sites outside of the ATP pocket. Upon ligand binding, conformational modifications take place in the catalytic subunits resulting in the inhibition of its phosphorylating activity. Today, the only pockets outside the ATP site identified within CK2 are the so-called α D pocket and a pocket located at the interface between the α C helix and the glycine-rich loop, although recent findings question the existence of the latter.

Inhibitors utilising the α D pocket. The α D site is unique to CK2 α amongst all known kinase structures, and therefore, provides exciting possibilities for the development of incredibly selective CK2 α inhibitors (Fig. 10). The α D site is not formed in the majority of the apo crystal structures as it requires a large reorganisation of the α D helix for the site to be accessible. In the unligated structures, the main pocket of the α D site is filled by Phe121 and the hydrophilic water channel off the main pocket is filled by Tyr125. The α D site was first discovered in a high concentration fragment screen carried out by Brear *et al.*⁵¹ It is possible for the α D site to form uniquely in CK2 α due to the increased flexibility of this region compared to other kinases: in the published structures of CK2 α , the loop is observed in a large spread of conformations. These range from the loop closed structures with Phe121 in the centre of the α D site (PDB: 6YPK), to the partially open apo structure (PDB: 5CU6), to the open fragment bound structure (PDB: 5CLP) and finally to the apo/inactive conformation (PDB: 5CVG). Interestingly, in all other kinase structures, only the closed conformation has been observed for this helix. The binding of small molecules in the α D site differentially affects the α D loop conformation depending on the ligand. Some ligands interact with the α D site and partially stabilise the loop, others do not engage the α D loop and the loop is not well defined in the crystal structure. It is unclear if the different type of interaction has any effect on the activity of the ligands. The majority of the reported small molecules binding in the α D site inhibit CK2 α by binding in the α D site and linking or growing into the ATP site.⁹⁷ However, there has been one report of an inhibitor that is proposed to bind in the



Fig. 10 (a) Different conformations of the α D loop (cyan = 6YPK, green = 5CU6, purple = 5CLP, yellow = 5CVG, magenta = 5CS6); (b) cross-section of the α D pocket in the apo conformation showing the surface of the pocket and loop in cartoon format with Phe121 and Tyr125 shown (PDB: 6YPK); (c) cross-section of the α D pocket in the apo conformation showing the surface of the pocket and loop in cartoon format with Tyr125 shown in the water channel. (PDB: 6YPK).

α D site allosterically inhibiting CK2 α .⁹⁸ Unfortunately, there is currently no structural data confirming that this compound binds in the α D site. Therefore, no reliable conclusions can be drawn from this data as to whether the α D site alone can be used as an allosteric site.

Using a fragment-based drug discovery approach, a series of benzylamine biaryl molecules binding in the α D pocket have been developed in a collaboration between our groups.^{51,97,99} Starting from high-concentration X-ray crystallography of commercially available fragments, compound **1** (in Brear *et al.*) was found to bind to the cryptic pocket in addition to other binding sites, namely the ATP binding site and in a hydrophobic pocket located at the interface with CK2 β .⁹⁷ Interestingly, compound **1** was the first ligand to reveal the full size of the α D pocket. Despite the lack of inhibitory effect on the catalytic activity of CK2, **1** set the basis for the development of molecules with improved binding affinity and potentially inhibitory activity (Fig. 11).

Initially, it was decided to optimise the fragment by growing towards vacant parts of the pocket. This led to com-





Recently, Bestgen *et al.* reported aryl aminothiazole derivatives as allosteric modulators of CK2 by targeting the interface between the α C helix and the glycine-rich loop (G-loop).^{100,101} Starting from a virtual ligand screening campaign of around 2 million compounds, compound **1** (in Bestgen *et al.*) was identified as a hit and it was able to inhibit CK2 with an IC_{50} of 28 μ M. SAR studies initially led to the discovery of compound **2** (IC_{50} of 7 μ M) and later to the lead compound **27** (in Bestgen *et al.*) with an IC_{50} of 0.6 μ M, K_d of 0.3 μ M, and EC_{50} of 5 μ M (Fig. 11). Enzymatic kinetic studies, native mass spectroscopy, circular dichroism experiments together with STD NMR studies hinted that compound **i** was acting as an allosteric inhibitor with a non-ATP-competitive mechanism of action able to stabilise the inactive conformation of CK2.^{100,101} However, Brear *et al.* used a combination of crystal structures, competitive ITC and NMR, hydrogen-deuterium exchange

Despite the numerous efforts gone into developing chemical tools to inhibit such a crucial kinase, clinical candidates are yet to reach the market. Only one molecule, **CX4945**, is currently undergoing clinical studies; however, due to its limited selectivity, even if approved, there will still be a long way to go before CK2 can be considered a truly validated target in oncology. For this reason, scientists have recently shifted their attention to the development of CK2 inhibitors that interact selectively with CK2. Strategies have seen the development of chemical probes acting at sites located outside the ATP-pocket on CK2 α , with the site at the interface with CK2 β and the α D site being the most validated. With the increasing amount of information around these sites and advances in biochemical techniques, it is reasonable to believe that potent and selective CK2 inhibitors with suitable pharmacokinetic and pharmacodynamic properties will soon be discovered. In addition,

recent advances in chemistry make it possible to foresee that different types of chemical probes will soon be developed including PROTACs acting outside the catalytic box, covalent inhibitors and small molecule/peptide hybrids to target multiple sites simultaneously.

Based upon the explosion of interest in the field of CK2 inhibition over the past two decades, it is not then unrealistic to imagine that CK2 inhibition will remain a key and active area of research over the coming years, as the search for novel therapeutics for cancer, COVID-19 and a multitude of other diseases in which CK2 is implicated continues.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 F. Ardito, M. Giuliani, D. Perrone, G. Troiano and L. Lo Muzio, *Int. J. Mol. Med.*, 2017, **40**, 271–280.
- 2 P. Lahiry, A. Torkamani, N. J. Schork and R. A. Hegele, *Nat. Rev. Genet.*, 2010, **1**, 60–74.
- 3 <http://www.kinase-screen.mrc.ac.uk/phosphorylation-ubiquitylation-drug-discovery> Accessed January 2021.
- 4 G. Manning, D. B. Whyte, R. Martinez, T. Hunter and S. Sudarsanam, *Science*, 2002, **298**, 1912–1934.
- 5 N. N. Singh and D. P. Ramji, *J. Mol. Med.*, 2008, **86**, 227–897.
- 6 L. A. Pinna, *J. Cell Sci.*, 2002, **115**, 3873–3878.
- 7 C. E. Ortega, Y. Seidner and I. Dominguez, *PLoS One*, 2014, **9**, e115609.
- 8 J. Zhang, P. L. Yang and N. S. Gray, *Nat. Rev. Cancer*, 2009, **9**, 28–39.
- 9 A. Siddiqui-Jain, D. Drygin, N. Streiner, P. Chua, F. Pierre, S. E. O' Brien, J. Bliesath, M. Omori, N. Huser, C. Ho, *et al.*, *Cancer Res.*, 2010, **70**, 10288–10299.
- 10 M. Bouhaddou, D. Memon, B. Meyer, K. M. White, V. V. Rezelj, M. Correa Marrero, B. J. Polacco, J. E. Melnyk, S. Ulferts, R. M. Kaake, *et al.*, *Cell*, 2020, **182**, 685–712.
- 11 D. E. Gordon, G. M. Jang, M. Bouhaddou, J. Xu, K. Obernier, K. M. White, M. J. O'Meara, V. V. Rezelj, J. Z. Guo, D. L. Swaney, *et al.*, *Nature*, 2020, **583**, 459–468.
- 12 R. Prudent and C. Cochet, *Chem. Biol.*, 2009, **16**, 112–120.
- 13 Z. A. Knight and K. M. Shokat, *Chem. Biol.*, 2005, **12**, 621–637.
- 14 M. I. Davis, J. P. Hunt, S. Herrgard, P. Ciceri, L. M. Wodicka, G. Pallares, M. Hocker, D. K. Treiber and P. P. Zarrinkar, *Nat. Biotechnol.*, 2011, **29**, 1046–1051.
- 15 K. Niefind, B. Guerra, I. Ermakowa and O. G. Issinger, *EMBO J.*, 2001, **20**, 5320–5331.
- 16 K. Niefind, J. Raaf and O. G. Issinger, *Cell. Mol. Life Sci.*, 2009, **66**, 1800–1816.
- 17 E. Papinutto, A. Ranchio, G. Lolli, L. A. Pinna and R. Battistutta, *J. Struct. Biol.*, 2012, **177**, 382–391.
- 18 A. C. Bibby and D. W. Litchfield, *Int. J. Biol. Sci.*, 2005, **1**, 67–79.
- 19 D. W. Litchfield, *Biochem. J.*, 2003, **369**, 1–15.
- 20 O. Filhol, J. L. Martiel and C. Cochet, *EMBO Rep.*, 2004, **5**, 351–355.
- 21 G. Poletto, J. Vilardell, O. Marin, M. A. Pagano, G. Cozza, S. Sarno, A. Falqués, E. Itarte, L. A. Pinna and F. Meggio, *Biochemistry*, 2008, **32**, 8317–8325.
- 22 J. Raaf, N. Bischoff, K. Klopffleisch, E. Brunstein, B. B. Olsen, G. Vilck, D. W. Litchfield, O.-G. Issinger and K. Niefind, *Biochemistry*, 2011, **50**, 512–522.
- 23 J. Raaf, E. Brunstein, O.-G. Issinger and K. Niefind, *Protein Sci.*, 2008, **12**, 2180–2186.
- 24 J. H. Trembley, Z. Chen, G. Unger, J. Slaton, B. T. Kren, C. Van Waes and K. Ahmed, *BioFactors*, 2010, **36**, 187–195.
- 25 S. A. Gibson and E. N. Benveniste, *Trends Immunol.*, 2018, **38**, 82–85.
- 26 J. Castello, A. Ragnauth, E. Friedman and H. Rebholz, *Pharmaceuticals*, 2017, **10**(1), 7.
- 27 S. Koch, C. T. Capaldo, R. S. Hilgarth, B. Fournier, C. A. Parkos and A. Nusrat, *Mucosal Immunol.*, 2013, **6**, 136–145.
- 28 E. Ampofo, L. Nalbach, M. D. Menger, M. Montenarh and C. Götz, *Int. J. Mol. Sci.*, 2019, **18**, 4398.
- 29 K. J. Treharne, R. M. Crawford, Z. Xu, J. H. Chen, O. G. Best, E. A. Schulte, D. C. Gruenert, S. M. Wilson, D. N. Sheppard, K. Kunzelmann and A. Mehta, *J. Biol. Chem.*, 2007, **282**, 10804–10813.
- 30 L. R. Cruz, I. Baladrón, A. Rittoles, P. A. Díaz, R. Santana, M. M. Vázquez, A. García, D. Chacón, G. Perera, A. González, *et al.*, *ACS Pharmacol. Transl. Sci.*, 2021, **4**(1), 206–212.
- 31 R. Turner, *BeingWell Online*, <https://medium.com/being-well/is-silmitasertib-the-covid-19-treatment-breakthrough-weve-been-waiting-for-263cf9c696d5> (accessed January 2021).
- 32 D. Hanahan and R. A. Weinberg, *Cell*, 2011, **144**, 646–674.
- 33 D. Hanahan and R. A. Weinberg, *Cell*, 2000, **100**, 57–70.
- 34 R. A. Faust, G. Niehans, M. Gapany, D. Hoistad, D. Knapp, D. Cherwitz, A. Davis, G. L. Adams and K. Ahmed, *Int. J. Biochem. Cell Biol.*, 1999, **31**, 941–949.
- 35 J. H. Trembley, G. Wang, G. Unger, J. Slaton and K. Ahmed, *Cell. Mol. Life Sci.*, 2009, **66**, 1858–1867.
- 36 R. A. Faust, M. Gapany, P. Tristani, A. Davis, G. L. Adams and K. Ahmed, *Cancer Lett.*, 1996, **101**, 31–35.
- 37 S. Yenice, A. T. Davis, S. A. Goueli, A. Akdas, C. Limas and K. Ahmed, *Prostate*, 1994, **24**, 11–16.
- 38 G. Seitz, U. Münstermann, H. R. Schneider and O. G. Issinger, *Biochem. Biophys. Res. Commun.*, 1989, **163**, 635–641.
- 39 M. Gapany, R. A. Faust, S. Tawfic, A. Davis, G. L. Adams and K. Ahmed, *Mol. Med.*, 1995, **1**, 659–666.
- 40 M. Chua, C. Ortega, A. Sheikh, M. Lee, H. Abdul-Rassoul, K. Hartshorn and I. Dominguez, *Pharmaceuticals*, 2017, **10**, 18.



- 41 P. O-Charoenrat, V. Rusch, S. G. Talbot, I. Sarkaria, A. Viale, N. Socci, I. Ngai, P. Rao and B. Singh, *Clin. Cancer Res.*, 2004, **10**, 5792–5803.
- 42 M. J. Ku, J. W. Park, B. J. Ryu, Y. J. Son, S. H. Kim and S. Y. Lee, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 5609–5613.
- 43 F. Benavent Acero, C. S. Capobianco, J. Garona, S. M. Cirigliano, Y. Perera, A. J. Urtreger, S. E. Perea, D. F. Alonso and H. G. Farina, *Lung Cancer*, 2017, **107**, 14–21.
- 44 K. Y. Lin, C. L. Fang, Y. Chen, C. F. Li, S. H. Chen, C. Y. Kuo, C. Tai and Y. H. Uen, *Ann. Surg. Oncol.*, 2010, **17**, 1695–1702.
- 45 J. Zou, H. Luo, Q. Zeng, Z. Dong, D. Wu and L. Liu, *J. Transl. Med.*, 2011, **9**, 97.
- 46 F. Zhou, J. Xu, G. Ding and L. Cao, *Pathol. Oncol. Res.*, 2014, **20**, 73–79.
- 47 G. Di Maira, A. Gentilini, M. Pastore, A. Caligiuri, B. Piombanti, C. Raggi, E. Rovida, M. Lewinska, J. B. Andersen, C. Borgo, *et al.*, *Oncogenesis*, 2019, **8**, 61.
- 48 M. Bouhaddou, D. Memon, B. Meyer, K. M. White, V. V. Rezeli, M. C. Marrero, B. J. Polacco, J. E. Melnyk, S. Ulferts, R. M. Kaake, *et al.*, *Cell*, 2020, **182**, 685–712.
- 49 G. M. Hathaway, T. H. Lubben and J. A. Traugh, *J. Biol. Chem.*, 1980, **255**, 8038–8041.
- 50 H. Yim, Y. H. Lee, C. H. Lee and S. K. Lee, *Planta Med.*, 1999, **65**, 009–013.
- 51 C. De Fusco, P. Brear, J. Iegre, K. H. Georgiou, H. F. Sore, M. Hyvönen and D. R. Spring, *Bioorg. Med. Chem.*, 2017, **13**, 3471–3482.
- 52 J. E. Dowling, M. Alimzhanov, L. Bao, C. Chuaqui, C. R. Denz, E. Jenkins, N. A. Larsen, P. D. Lyne, T. Pontz, Q. Ye, G. A. Holdgate, L. Snow, N. O'Connell and A. D. Ferguson, *ACS Med. Chem. Lett.*, 2016, **7**, 300–305.
- 53 R. Zandomeni, M. C. Zandomeni, D. Shugar and R. Weinmann, *J. Biol. Chem.*, 1986, **261**, 3414–3419.
- 54 S. Sarno, E. de Moliner, M. Ruzzene, M. A. Pagano, R. Battistutta, J. Bain, D. Fabbro, J. Schoepfer, M. Elliott, P. Furet, F. Meggio, G. Zanotti and L. A. Pinna, *Biochem. J.*, 2003, **374**, 639–646.
- 55 F. Meggio, M. A. Pagano, S. Moro, G. Zagotto, M. Ruzzene, S. Sarno, G. Cozza, J. Bain, M. Elliott, A. D. Deana, A. M. Brunati and L. A. Pinna, *Biochemistry*, 2004, **43**, 12931–12936.
- 56 Z. Nie, C. Perretta, P. Erickson, S. Margosiak, R. Almassy, J. Lu, A. Averill, K. M. Yager and S. Chu, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 4191–4195.
- 57 J. E. Dowling, C. Chuaqui, T. W. Pontz, P. D. Lyne, N. A. Larsen, M. H. Block, H. Chen, N. Su, A. Wu, D. Russell, *et al.*, *ACS Med. Chem. Lett.*, 2012, **3**, 278–283.
- 58 E. L. Atkinson, J. Iegre, P. D. Brear, E. A. Zhabina, M. Hyvönen and D. R. Spring, *Molecules*, 2021, **26**, 1977.
- 59 T. Oshima, Y. Niwa, K. Kuwata, A. Srivastava, T. Hyoda, Y. Tsuchiya, M. Kumagai, M. Tsuyuguchi, T. Tamaru, A. Sugiyama, N. Ono, N. Zolboot, Y. Aikawa, S. Oishi, A. Nonami, F. Arai, S. Hagihara, J. Yamaguchi, F. Tama, Y. Kunisaki, K. Yagita, M. Ikeda, T. Kinoshita, S. A. Kay, K. Itami and T. Hirota, *Sci. Adv.*, 2019, **5**, 9060–9083.
- 60 A. Dalle Vedove, F. Zonta, E. Zanforlin, N. Demitri, G. Ribaudo, G. Cazzanelli, A. Ongaro, S. Sarno, G. Zagotto, R. Battistutta, M. Ruzzene and G. Lolli, *Eur. J. Med. Chem.*, 2020, **195**, 112267.
- 61 C. Wells, D. H. Drewry, J. E. Pickett, D. W. Litchfield, S. Knapp and A. D. Axtman, *Cell Chem. Biol.*, 2021, DOI: 10.1016/j.chembiol.2020.12.013.
- 62 B. Kobe, T. Kampmann, J. K. Forwood, P. Listwan and R. I. Brinkworth, *Biochim. Biophys. Acta, Proteins Proteomics*, 2005, **1754**, 200–209.
- 63 F. Meggio, L. A. Pinna, F. Marchiori and G. Borin, *FEBS Lett.*, 1983, **162**, 235–238.
- 64 F. Meggio, F. Marchiori, G. Borin, G. Chessa and L. A. Pinna, *J. Biol. Chem.*, 1984, **259**, 14576–14579.
- 65 F. Meggio and L. A. Pinna, *Biochim. Biophys. Acta, Mol. Cell Res.*, 1989, **1010**, 128–130.
- 66 R. Tellez, M. Gatica, C. C. Allende and J. E. Allende, *FEBS Lett.*, 1990, **265**, 113–116.
- 67 R. Tellez, C. C. Allende and J. E. Allende, *FEBS Lett.*, 1992, **308**, 113–115.
- 68 Y. Perera, H. G. Farina, I. Hernandez, O. Mendoza, J. M. Serrano, O. Reyer, D. E. Gomez, R. E. Gomez, B. E. Acevedo, D. F. Alfonso and S. E. Perea, *Int. J. Cancer*, 2008, **122**, 57–62.
- 69 S. E. Perea, O. Reyes, Y. Puchades, O. Mendoza, N. S. Vispo, I. Torrens, A. Santos, R. Silva, B. Acevedo, E. López, V. Falcón and D. F. Alonso, *Cancer Res.*, 2004, **64**, 7127–7129.
- 70 M. F. Gottardo, C. S. Capobianco, J. E. Sidabra, J. Garona, Y. Perera, S. E. Perea, D. F. Alonso and H. G. Farina, *Sci. Rep.*, 2020, **10**, 14689.
- 71 M. Winiewska-Szajewska, D. Płonka, I. Zhukov and J. Poznański, *Sci. Rep.*, 2019, **9**, 11018.
- 72 G. Cozza, S. Zanin, S. Sarno, E. Costa, C. Girardi, G. Ribaudo, M. Salvi, G. Zagotto, M. Ruzzene and L. A. Pinna, *Biochem. J.*, 2015, **471**, 415–430.
- 73 E. Enkvist, K. Viht, N. Bischoff, J. Vahter, S. Saaver, G. Raidaru, O. G. Issinger, K. Niefind and A. Uri, *Org. Biomol. Chem.*, 2012, **10**, 8645–8653.
- 74 K. Viht, S. Saaver, J. Vahter, E. Enkvist, D. Lavogina, H. Sinijärvi, G. Raidaru, B. Guerra, O. G. Issinger and A. Uri, *Bioconjugate Chem.*, 2015, **26**, 2324–2335.
- 75 R. Y. Tsien, *Nature*, 1981, **290**, 527–528.
- 76 J. Vahter, K. Viht, A. Uri and E. Enkvist, *Bioorg. Med. Chem.*, 2017, **25**, 2277–2284.
- 77 T. N. de Villavicencio-Diaz, Y. Mazola, Y. P. Negrín, Y. C. García, O. G. Cruz and S. E. Perea Rodríguez, *Biochem. Biophys. Rep.*, 2015, **4**, 20–27.
- 78 F. Meggio, D. Shugar and L. A. Pinna, *Eur. J. Biochem.*, 1990, **187**, 89–94.
- 79 J. Iegre, P. Brear, D. J. Baker, Y. S. Tan, E. L. Atkinson, H. F. Sore, D. H. O'Donovan, C. S. Verma, M. Hyvönen and D. R. Spring, *Chem. Sci.*, 2019, **10**, 5056–5063.



- 80 G. Lolli, A. Ranchio and R. Battistutta, *ACS Chem. Biol.*, 2014, **9**, 366–371.
- 81 S. Zanin, C. Borgo, C. Girardi, S. E. O'Brien, Y. Miyata, L. A. Pinna, A. Donella-Deana and M. Ruzzene, *PLoS One*, 2012, **11**, e49193.
- 82 I. Kufareva, B. Bestgen, P. Brear, R. Prudent, B. Laudet, V. Moucadet, M. Ettaoussi, C. F. Sautel, I. Krimm, M. Engel, O. Filhol, M. Le Borgne, T. Lomberget, C. Cochet and R. Abagyan, *Sci. Rep.*, 2019, **9**, 15893.
- 83 J. C. Fuller, N. J. Burgoyene and R. M. Jackson, *Drug Discovery Today*, 2009, **14**, 155–161.
- 84 J. Raaf, E. Brunstein, O. G. Issinger and K. Niefind, *Chem. Biol.*, 2008, **2**, 111–117.
- 85 B. Laudet, V. Moucadet, R. Prudent, O. Filhol, Y. S. Wong, D. Royer and C. Cochet, *Mol. Cell. Biochem.*, 2008, **316**, 63–69.
- 86 V. Moucadet, R. Prudent, C. F. Sautel, F. Teillet, C. Barette, L. Lafanechere, V. Receveur-Brechot and C. Cochet, *Oncotarget*, 2011, **2**, 997–1010.
- 87 L. Kröger, C. G. Daniliuc, D. Ensan, S. Borgert, C. Nienberg, M. Lauwers, M. Steinkrüger, J. Jose, M. Pietsch and B. Wünsch, *ChemMedChem*, 2020, **15**, 871–881.
- 88 P. Brear, A. North, J. Iegre, K. Hadje Georgiou, A. Lubin, L. Carro, W. Green, H. F. Sore, M. Hyvonen and D. R. Spring, *Bioorg. Med. Chem.*, 2018, **26**, 3016–3020.
- 89 L. Nevola and E. Giralt, *Chem. Commun.*, 2015, **51**, 3302–3315.
- 90 J. A. Wells and C. L. McClendon, *Nature*, 2007, **450**, 1001–1009.
- 91 B. Laudet, C. Barette, V. Dulery, O. Renaudet, P. Dumy, A. Metz, R. Prudent, A. Deshiere, O. Dideberg, O. Filhol and C. Cochet, *Biochem. J.*, 2007, **408**, 363–373.
- 92 B. Bestgen, Z. Belaid-Choucair, T. Lomberget, M. Le Borgne, O. Filhol and C. Cochet, *Pharmaceuticals*, 2017, **1**, 16.
- 93 D. Lindenblatt, M. Horn, C. Götz, K. Niefind, I. Neundorf and M. Pietsch, *ChemMedChem*, 2019, **14**, 833–841.
- 94 S. Tang, N. Zhang, Y. Zhou, W. A. Cortopassi, M. P. Jacobson, L. Zhao and R. Zhong, *Mol. Inf.*, 2019, **38**, 1800089.
- 95 J. Hochscherf, D. Lindenblatt, M. Steinkrüger, E. Yoo, Ö. Ulucan, S. Herzig, O.-G. G. Issinger, V. Helms, C. Götz, I. Neundorf, K. Niefind and M. Pietsch, *Anal. Biochem.*, 2015, **468**, 4–14.
- 96 I. Neundorf, R. Rennert, J. Hoyer, F. Schramm, K. Löbner, I. Kitanovic and S. Wölfl, *Pharmaceuticals*, 2009, **2**, 49–65.
- 97 P. Brear, C. De Fusco, K. Hadje-Georgiou, N. J. Francis-Newton, C. J. Stubbs, H. F. Sore, A. R. Venkitaraman, C. Abell, D. R. Spring and M. Hyvönen, *Chem. Sci.*, 2016, **7**, 6839–6845.
- 98 C. Li, X. Zhang, N. Zhang, Y. Zhou, G. Sun, L. Zhao and R. Zhong, *Molecules*, 2020, **25**, 237.
- 99 J. Iegre, P. Brear, C. De Fusco, M. Yoshida, S. L. Mitchell, M. Rossmann, L. Carro, H. F. Sore, M. Hyvönen and D. R. Spring, *Chem. Sci.*, 2018, **9**, 3041–3049.
- 100 B. Bestgen, I. Krimm, I. Kufareva, A. A. M. Kamal, W. G. Seetoh, C. Abell, R. W. Hartmann, R. Abagyan, C. Cochet, M. Le Borgne, M. Engel and T. Lomberget, *J. Med. Chem.*, 2019, **62**, 1803–1816.
- 101 B. Bestgen, I. Kufareva, W. Seetoh, C. Abell, R. W. Hartmann, R. Abagyan, M. Le Borgne, O. Filhol, C. Cochet, T. Lomberget and M. Engel, *J. Med. Chem.*, 2019, **62**, 1817–1836.
- 102 X. Chen, C. Li, D. Wang, Y. Chen and N. Zhang, *Molecules*, 2020, **25**, 870.
- 103 R. Prudent, V. Moucadet, B. Laudet, C. Barette, L. Lafanechère, B. Hasenknopf, J. Li, S. Bareyt, E. Lacôte, S. Thorimbert, M. Malacria, P. Gouzerh and C. Cochet, *Chem. Biol.*, 2008, **15**, 683–692.
- 104 P. Brear, D. Ball, K. Stott, S. D'Arcy and M. Hyvönen, *J. Med. Chem.*, 2020, **21**, 12786–12798.
- 105 D. Lindenblatt, A. Nickelsen, V. M. Applegate, J. Jose and K. Niefind, *J. Med. Chem.*, 2020, **14**, 7766–7772.
- 106 V. Martel, O. Filhol, P. Colas and C. Cochet, *Oncogene*, 2006, **25**, 7343–7353.
- 107 B. Laudet, C. Barette, V. Dulery, O. Renaudet, P. Dumy, A. Metz, R. Prudent, A. Deshiere, O. Dideberg, O. Filhol and C. Cochet, *Biochem. J.*, 2007, **408**, 363–373, 17714077.
- 108 S. Tang, N. Zhang, Y. Zhou, W. A. Cortopassi and M. P. Jacobson, *Mol. Inf.*, 2018, **38**, 1800089.

