



Cite this: *Chem. Commun.*, 2014, 50, 14290

Metal–ligand bifunctional reactivity and catalysis of protic N-heterocyclic carbene and pyrazole complexes featuring β -NH units

Shigeki Kuwata* and Takao Ikariya*

Received 11th June 2014,
Accepted 22nd July 2014

DOI: 10.1039/c4cc04457f

www.rsc.org/chemcomm

Metal–ligand bifunctional cooperation has attracted much attention because it offers a powerful methodology to realize a number of highly efficient and selective catalysts. In this article, recent developments in the metal–ligand cooperative reactions of protic N-heterocyclic carbene (NHC) and pyrazole complexes bearing an acidic NH group at the position β to the metal are surveyed. Protic 2-pyridylidenes as related cooperating non-innocent ligands are also described.

1. Introduction

Metal–ligand bifunctional cooperation has been implicated in a number of catalysts including metalloenzymes in nature. The cooperating ligands therein impart superior substrate-activation and transformation ability as well as selectivity to the catalyst through non-covalent interactions such as hydrogen bonding. In [FeFe] hydrogenases, which catalyze reversible conversion of hydrogen gas into two protons and electrons, the bridging

azadithiolato ligand serves as a Brønsted acid–base catalytic group to promote heterolytic cleavage of dihydrogen and proton relay involved in the catalysis (Fig. 1a).¹ Inspired by the prominent role of the amino group in the second coordination sphere, DuBois and co-workers developed metal–ligand bifunctional electrocatalysts for H₂ production and oxidation featuring uncoordinated amino groups, typified by the nickel complex shown in Fig. 1b.²

Provided that the remote cooperating ligand can tautomerize, at least formally, to an “ α -protic” form where the Brønsted acidic hydrogen attaches to the donor atom in the position α to the metal, protonation/deprotonation of the cooperating unit in the second coordination sphere would be coupled with coordination/dissociation events on the metal center. For example, Milstein and

Department of Applied Chemistry, Graduate School of Science and Engineering,
Tokyo Institute of Technology, O-okayama 2-12-1 E4-1, Meguro-ku, Tokyo, Japan.
E-mail: skuwata@apc.titech.ac.jp, tikariya@apc.titech.ac.jp



Shigeki Kuwata

Shigeki Kuwata was born in 1969 in Osaka, Japan. He received the Dr Eng. degree from Department of Chemistry and Biotechnology at the University of Tokyo in 1997 under the supervision of Professors Masanobu Hidai and Yasushi Mizobe. Just after that, he was appointed as an Assistant Professor in the department. In 2002, he joined Professor Ikariya's group at Tokyo Tech as an Associate Professor. He obtained the Award for Young Scientists in

Coordination Chemistry, Japan, in 2002. His current research interests focus on the chemistry of “azametallics” featuring reactive metal–nitrogen units.



Takao Ikariya

Takao Ikariya completed his PhD degree in 1976 at Tokyo Institute of Technology and then he was appointed as an assistant professor in the Department of Synthetic Chemistry at the University of Tokyo. He spent one and a half years in 1979–1981 as a postdoctoral fellow in Professor Robert H. Grubbs' group at Caltech. In 1985 he moved to the Central Research Center of NKK Corp. In 1991, he joined the ERATO Molecular Cat-

alysis Project of Japan Science and Technology Corporation, which was directed by Professor Ryoji Noyori. Ikariya was then promoted to professor at Tokyo Institute of Technology in 1997. In 2014, he retired from Tokyo Institute of Technology and now he is an Emeritus Professor.






Fig. 1 (a) Active site of the [FeFe] hydrogenase enzyme. (b) Metal-ligand bifunctional nickel complex that catalyzes electrochemical H_2 formation.


co-workers demonstrated that a series of lutidine-based pincer-type complexes catalyze various hydrogenations and dehydrogenative transformations.³ These reactions are believed to proceed through addition and elimination of the substrates at the bifunctional platform containing the methylene group remote from the metal (Scheme 1), as observed in the prototypical α -protic metal-amine bifunctional catalysts.⁴ The hydroxycyclopentadienyl-hydrido Ru^{II} complex **2** derived from Shvo's catalyst **1** also undergoes reversible proton-coupled hydride transfer to the substrates, giving the coordinatively unsaturated cyclopentadienone Ru^0 species **3** as shown in Scheme 2.⁵ It is to be noted that the interconversion, which leads to various transfer hydrogenation catalysis, involves formal redox of the metal and cooperating ligand in contrast to the corresponding transformation of the conventional metal-amine bifunctional catalysts as well as Milstein's catalyst.



Scheme 1 Reversible deprotonation of the methylene group and metal-ligand bifunctional activation of substrates in Milstein's catalysts.



Scheme 2 Fragmentation of Shvo's catalyst **1** into a hydroxycyclopentadienyl-hydrido complex **2** and coordinatively unsaturated cyclopentadienone complex **3**, which are interconvertible with a hydrogen acceptor (A) and donor ($\text{A}'\text{H}_2$).



Scheme 3 β -Protic metal-ligand bifunctional catalysts and their reversible deprotonation.

The bifunctional cooperation of a metal center and conjugated proton-responsive units placed at appropriate positions for substrate recognition, activation and transformation thus results in efficient and tunable catalysis owing to the structural diversity and electronic flexibility of the cooperating ligands.⁶


Among such bifunctional catalysts containing remote cooperating ligands, we focus here on the protic N-heterocyclic carbene (NHC)^{7,8} and pyrazole^{9,10} complexes featuring an NH group in the position β to the metal. The reasonable stability of the conjugate bases facilitates transfer of the nucleophilic group X to external substrates as well as the bifunctional activation of various pronucleophiles HX (Scheme 3). The metal-ligand cooperating reactivities and catalysis of these β -protic bifunctional complexes with different structures and Brønsted acidity will be described herein.

2. Protic NHC complexes

The NHCs have been used as auxiliary ligands in numerous homogeneous catalysts owing to their accessibility, robustness and σ -donating ability.¹¹ However, NHCs that have an NH-wingtip (protic NHC; also designated as NH-NHC, NR,NH-stabilized carbene, NH-functionalized NHC and so on) have been explored to a much less extent, at least partly because the protic NHCs are not generally obtained by simple deprotonation of the corresponding imidazolium salt with an acidic NH group and tend to isomerize to the corresponding imidazoles.¹² Protic NHC complexes thus have been synthesized as follows: (i) cyclization of isocyanides and amines on metal templates,¹³ (ii) base/acid-promoted tautomerization of imidazole ligands, (iii) chelation-assisted formal tautomerization of imidazoles, (iv) oxidative addition of 2-haloimidazoles followed by N-protonation¹⁴ and (v) deprotection of NH-masked NHC complexes. Readers can find the details in recent reviews.^{7,8} In this article, we emphasize the metal-ligand bifunctional reactivity of protic NHC complexes.

2.1 Monodentate NHC complexes


2.1.1 Substrate recognition and addition reaction. Hahn and co-workers demonstrated that the protic NHC complex **4a** catalyzes hydrogenation of a 3-butenic acid ester more than two times faster than 1-dodecene.¹⁵ The acceleration may be rationalized by the substrate recognition through a hydrogen bond between the protic NHC ligand and the carbonyl group of the alkenic ester (Scheme 4). The ability of the NH group to




Scheme 4 Competitive hydrogenation of alkenes with/without a carbonyl group catalyzed by a protic NHC complex.

form a hydrogen bond was evaluated by ^1H NMR titration experiments using the tungsten analogue **4b** and a urea derivative.


Chloride abstraction of the protic NHC complex **5** with KPF_6 in acetonitrile leads to *anti*-addition of the N–H bond to acetonitrile, giving the chelate imine complex **6** as shown in eqn (1).¹⁶ This transformation most likely occurs *via* a cationic acetonitrile complex, wherein the Lewis acidic iridium center activates the nitrile toward the intramolecular addition of the protic NHC ligand to the $\text{C}\equiv\text{N}$ bond.¹⁷



2.1.2 C–C bond formation. In the extensive work on Rh-catalyzed C–H bond functionalization of N-heterocycles,¹⁸ Ellman, Bergman and co-workers described catalytic arylation¹⁹ and alkylation²⁰ of imidazoles; typical examples are shown in Scheme 5. Scheme 6 illustrates a proposed mechanism of the catalytic arylation. Tautomerization of benzimidazole on the rhodium catalyst affords the protic NHC complex **7**, which has been unambiguously characterized by an X-ray analysis (for $\text{R} = \text{Me}$). Complex **7** with the electron-donating NHC ligand then undergoes



Scheme 5 Rh-catalyzed functionalization of benzimidazoles. coe = cyclooctene, LutHCl = 2,6-lutidinium chloride.




Scheme 6 A proposed mechanism for Rh-catalyzed arylation of benzimidazoles. $\text{Rh} = \text{Rh}(\text{PCy}_3)_n$.

oxidative addition of iodobenzene. The Lewis acidic Rh^{III} center in the resultant complex **8** promotes deprotonation of the protic NHC ligand to give **9**, which yields the C-arylation product through reductive elimination. A related reaction of a six-membered dihydroquinazoline that proceeds *via* a protic NHC complex is also known.²¹


2.2 Chelate-stabilized NHC complexes

The protic NHC ligand implicated in the catalysis shown in Scheme 5 originates in the substrate and is incorporated into the product. To expand the scope of the metal–ligand bifunctional catalysis of the protic NHC complexes, the cooperating ligand needs to be installed in the catalyst in advance and to be protected from dissociation. Chelate-stabilization of the protic NHC ligand offers a rational strategy for this purpose since such ligands are easily accessed by cyclometalation of imidazoles with appropriate anchoring groups.

2.2.1 C–C bond formation. Tan, Bergman and Ellman synthesized the chelation-stabilized protic NHC complex **11** through formal tautomerization of **10** on a rhodium complex as shown in eqn (2).²² They also demonstrated that **11** catalyzes cyclization of alkenyl-substituted imidazoles such as **10**, an intramolecular version of the reaction shown in Scheme 5,^{22,23} in which the protic NHC ligand in **11** still comes from the substrate imidazole.



We have revealed the metal–ligand bifunctional cooperating reactivities of the C–N chelate protic NHC complex **12** prepared by cyclometalation of *N*-(2-pyridyl)benzimidazole. Complex **12**




Scheme 7 A proposed mechanism for dehydrative coupling of pyridylbenzimidazole and allyl alcohol catalyzed by protic NHC complex **12**. $Ru = Cp^*Ru$.

catalyzes dehydrative coupling of the benzimidazole and allyl alcohol.²⁴ The proposed mechanism is illustrated in Scheme 7. A two-point interaction between the protic NHC complex and the substrate would enable facile C–O bond cleavage of allyl alcohol to afford the π -allyl-imidazolyl Ru^{IV} species **13** under neutral conditions. In this dehydration step, the β -protic bifunctional complex **12** formally provides one proton and two electrons to the substrate. Reductive elimination easily occurs despite the presence of the anchoring pyridyl group to yield the crystallographically characterized 2-allylimidazole complex **14**. The involvement of **14** in the catalysis has been established by the facile formation of **14** in the reaction of **12** and allyl alcohol at room temperature. Replacement of the allylimidazole ligand in **14** by the pyridylimidazole substrate leads to tautomerization of the incoming imidazole to regenerate the protic NHC complex **12**.

Proton migration from the NH unit in **12** to ligated substrates is also observed in the reaction of **12** and silver nitrite.²⁴ The resultant nitrosyl-imidazolyl complex undergoes reversible N-protonation to regenerate the protic NHC ligand. Structural changes of the imidazolyl ligand upon protonation have been scrutinized.^{7,24}


2.2.2 H–H and H–X (X = O, C) bond cleavage. Grotjahn and co-workers reported the reactivities of the phosphine-tethered protic NHC ruthenium complexes as summarized in Scheme 8.²⁵ The protic NHC ligand in **15** is reversibly deprotonated with *n*-butyllithium or lithium diisopropylamide to afford the anionic imidazolyl complex **16**, which has been unambiguously

characterized by an X-ray analysis as well as the 7Li and ^{13}C NMR spectroscopy. Complex **16** can be regarded as a LiCl-masked coordinatively unsaturated imidazolyl complex. In fact, even ethylene instantaneously reacts with **16** to provide the imidazolyl–ethylene complex **17** ($L = C_2H_4$). The latent metal–ligand bifunctionality in **16** becomes more evident in the reactions with dihydrogen and 2-propanol, which result in heterolysis of H_2 (ref. 26) and dehydrogenative oxidation of the alcohol to give the hydrido complex **18**. The hydrido complex is also obtainable in a one-pot reaction from **15**. In addition, the metal–ligand bifunctional hydrido complex **18** catalyzes transfer hydrogenation of acetophenone with 2-propanol in the absence of a base (eqn (3)).




The reactions of **15** with protic amines as well as ethylene in the presence of the hexafluorophosphate anion lead to the formation of a series of cationic complexes **19** (Scheme 8).²⁵ The protic NHC complexes **19** are converted to the imidazolyl complexes **17** upon treatment with a base. It is to be noted that the deprotonation occurs at the carbene site rather than the coordinated amine, indicating the stronger Brønsted acidity of the protic NHC ligand. The ^{15}N NMR study of the natural abundance samples using the gHSQC technique revealed that deprotonation of the NH-wingtip results in a significant downfield shift of the ^{15}N NMR signal.²⁵

Stepwise dehydrochlorination of the isoelectronic iridium complex **20** shown in Scheme 9 also offers a bifunctional platform for heterolytic cleavage of dihydrogen as well as addition of acetylene.²⁷ The hydrogen bond between a related protic NHC




Scheme 8 Reactions of C–P chelate protic NHC ruthenium complexes. $Ru = CpRu$.



Scheme 9 Metal–ligand bifunctional reactions of protic NHC iridium complex **20**. Ir = Cp*Ir, BAR₄ = B(C₆F₅)₄.

ruthenium complex **21** and an external urea has been assessed by Hahn and co-workers.²⁸ Esteruelas and co-workers revealed the intramolecular dihydrogen bond between the protic NHC and hydrido ligands in the osmium complex **22** by X-ray analysis.²⁹




3. Pyridine-derived protic carbene complexes


Protic 2-pyridylidene complexes³⁰ represent a subclass of β -protic NHC complexes. The synthesis has been achieved mainly through (i) formal tautomerization of pyridines on the metal center^{31–37} and (ii) oxidative addition of 2-haloimidazoles followed by N-protonation.^{38,39}

3.1 Reactions with H₂ and unsaturated molecules

Poveda, Carmona and co-workers demonstrated that the reaction of the dinitrogen complex [Tp*IrPh₂(N₂)] (Tp* = hydrotris-(3,5-dimethylpyrazolyl)borato) with 2-substituted pyridines affords a series of protic 2-pyridylidene complexes **23** as shown in eqn (4).^{32,33} Theoretical calculations revealed that the tautomerization of pyridines most likely occurs through the σ -bond complex-assisted metathesis involving the phenyl ligand.³² The parent 2-pyridylidene complex (R = H) can be obtained by alkaline hydrolysis of the trimethylsilyl derivative (R = SiMe₃).³²




When **23** is heated at 90 °C in C₆D₆, replacement of the phenyl ligands by C₆D₅ groups and deuteration of the NH unit take place.³² This observation indicates that thermal elimination of benzene from **23** generates a coordinatively unsaturated



Scheme 10 Reactions of protic 2-pyridylidene iridium complex **23**. The 6-substituents on the pyridine ring are omitted. Ir = Tp*Ir.


pyridyl complex **24** (Scheme 10), which cleaves the C–D bond in the solvent C₆D₆ to give the deuterated **23**. In fact, **24** can be trapped as the carbonyl complex **26**.⁴⁰ Thermal liberation of water from the aqua (or hydroxo–pyridylidene) complex **25** also generates **24**, which subsequently reacts with benzene to give **23**.³² The metal–ligand bifunctional platform in **24** undergoes additions of carbon dioxide and nonpolar alkenes to afford the iridacycles **28** and **29**, respectively.⁴⁰ The reaction of **23** with dihydrogen leads to heterolytic cleavage of H₂ in **24**,²⁶ giving the hydrido–carbene complex **27**.⁴⁰ Upon treatment with CO, this hydrido complex is converted to the carbonyl complex **26**, which may indicate the intermediacy of an η -H₂ complex in the H₂ heterolysis. Under strictly anhydrous conditions, **23** reacts with acetylene to provide an addition product **31** via intramolecular nucleophilic attack of the pyridyl ligand on the vinylidene moiety in the intermediate **30**, whilst attack of adventitious water on the vinylidene ligand yields the acyl complex **32**.⁴⁰ Similar hydration occurs in the reaction of **23** with acetonitrile and water, resulting in the formation of the acetamidato complex **33**.⁴⁰ The hydration appears to be aided by the neighboring Brønsted basic pyridyl ligand⁴¹ because [Tp*IrPh₂(NCMe)] without the cooperating ligand does not react with water under the same conditions. The protic 2-pyridylidene ligand in related Tp*Ir complexes also undergoes intramolecular C–N bond formation with the supporting phenyl,⁴² alkyl⁴³ and diene⁴⁴ ligands.

As part of their intensive study on protic 2-pyridylidene complexes,³⁴ Esteruelas and co-workers described C–C bond forming reactions of the protic carbene complex **34** obtained by the tautomerization of 2-methylpyridine on [TpOs(PiPr₃)(acetone)₂][BF₄]




Scheme 11 C–C bond formation between protic 2-pyridylidene and acetylide ligands. Os = TpOs.

(Tp = hydrotris(pyrazolyl)borato) and a subsequent reaction with terminal alkynes (Scheme 11).³⁵ Deprotonation of **34** leads to the formation of an equilibrium mixture of the alkynyl-pyridylidene complex **35** and pyridyl-vinylidene complex **36**, the latter of which is converted to the 1,1-insertion product **37** instead of a C–N bond formation product as observed in a related Tp*Ir system (**31** in Scheme 10). Such a C–C bond forming step may be involved in the catalytic C–H functionalization of pyridines with alkynes exemplified by eqn (5).⁴⁵ Berman, Bergman and Ellman proposed the involvement of a 2-pyridylidene complex in an Rh-catalyzed arylation of pyridines as the protic NHC complexes in the catalytic functionalization of imidazoles.⁴⁶




3.2 Reversible deprotonation

Chelation-assisted tautomerization of 2,3'-bipyridines and 1,9-phenanthrolines on [Ir(cod)₂]BF₄ yields a series of protic 2-pyridylidene complexes such as **38**.^{36,37} These iridium(i) complexes are in equilibrium with the hydrido-pyridyl Ir^{III} complexes, represented by **39** (eqn (6)). The equilibrium is significantly shifted to the protic 2-pyridylidene side by the counteranion X and solvent that can form a hydrogen bond with the NH group and thus stabilize the protic carbene ligand.^{37,47} Similarly, Brønsted basic substituents at an appropriate position on the pyridine ring advantage the protic carbene isomer.^{36,47} The 1,3-shift associated with formal redox of the metal center and umpolung of the hydrido ligand is accelerated by water, which participates in the proton transfer through a cyclic transition state.



Oxidative addition of 2-bromopyridines bearing nitrogen and sulfur donor tethers to [Pd(dba)₂] (dba = dibenzylideneacetone)

affords a series of dinuclear complexes **40**.³⁹ Subsequent protonation with HBr leads to the reversible formation of the protic 2-pyridylidene complexes **41** as shown in eqn (7). Complexes **40** catalyze the Mizoroki–Heck reaction, although the catalysis appears to be heterogeneous on the basis of mechanistic studies.




4. Protic pyrazole complexes

Having the structural similarity with the protic NHCs, protic pyrazoles provide the scaffolds for metal–ligand bifunctional cooperation.^{48,49} In comparison with the protic NHCs, protic pyrazoles have an advantage of having no stable aprotic tautomer. The ease of the ligand synthesis and complexation allows more flexible design of the metal–ligand bifunctional platforms. In this section, we describe the metal–ligand cooperative reactions of the protic pyrazole complexes including those bearing two NH units in the second coordination sphere.


4.1 Mono(pyrazole) complexes

4.1.1 Deprotonation and hydrogen transfer. Given the proton-responsive nature of the N-unsubstituted pyrazole ligand,^{7,50} coordinatively unsaturated pyrazolato complexes derived from the protic pyrazole complexes are expected to undergo hydrogenation and transfer hydrogenation through metal–ligand bifunctional cooperation as described in Schemes 8–10. We have demonstrated that the dehydrochlorination of the protic pyrazole complex **42** affords the pyrazolato-bridged dinuclear complex **43** in a reversible manner (Scheme 12).⁵¹ The dimeric complex **43** is functionally equivalent to the expected coordinatively unsaturated mononuclear complex owing to the significant distortion of the pyrazolato bridge. In fact, transfer hydrogenation of **43** with 2-propanol smoothly takes place to produce the hydrido-bridged pyrazole–pyrazolato complex **44** whose structure is topologically related to Shvo's catalyst. This partially hydrogenated complex **44** is also obtained by dehydrochlorination of **42** in 2-propanol.

Yu and co-workers described catalytic transfer hydrogenation of ketones with pincer-type ruthenium complexes bearing a protic pyrazole arm such as **45**.⁵² In this highly efficient catalysis, however, the NH unit does not appear to participate directly in substrate recognition and transformation because related imidazole complexes typified by **46** exhibit comparable catalytic activity despite that the NH unit lies at a more remote position γ to the metal.^{53,54} The facile hydrogen transfer promoted by the protic pyrazole ligand **47** has also been applied to α -alkylation of amides with primary alcohols shown in eqn (8).⁵⁵ Thiel and co-workers disclosed that the N–N chelate pyridylpyrazolato complexes **48** catalyze transfer hydrogenation of acetophenone


Scheme 12 Formation and transfer hydrogenation of pyrazolato complex **43**.

and hydrogenation of CO₂.⁵⁶ A mechanism without direct cooperation of the pyrazolato ligand has been proposed.




Gimeno, Lledós and co-workers have demonstrated that isomerization of allylic alcohols to carbonyl compounds is efficiently catalyzed by protic pyrazole Ru^{IV} complexes (Scheme 13).⁵⁷ The reaction occurs *via* bidirectional hydrogen transfer between the substrate and catalyst, which has been analyzed by DFT calculations. In the first half of the catalytic cycle, the hydroxo ligand derived from the water solvent participates in the hydrogen transfer. This step is also assisted by the pyrazolato ligand when the reaction is carried out in THF. On the other hand, the hydrogen transfer back to the substrate is achieved by the metal-pyrazole cooperation in both solvents. The same authors, however, also reported that a related γ -protic imidazole complex exhibits higher catalytic activity in water.⁵⁸

4.1.2 C–C and C–X (X = N, O, Si) bond formation. Some C–C bond forming reactions catalyzed by protic pyrazole complexes have been known. Satake and co-workers reported catalytic cyclopropanation of ketene silyl acetals with a cationic




Scheme 13 A proposed mechanism for redox isomerization of allylic alcohols catalyzed by protic pyrazole complexes. Substituents on the pyrazole ligand and allylic alcohol are omitted. Ru = {bis(allyl)}Ru.


allyl-pyridylpyrazole Pd^{II} complex as shown in eqn (9).⁵⁹ The active species is a deprotonated pyrazolato complex, and the secondary interactions of the β -NH unit with the substrates are not involved in the proposed mechanism. Protic pyrazole complexes and conjugate bases that catalyze the Mizoroki–Heck reaction,⁶⁰ Sonogashira reaction,⁶¹ hydrosilylation of phenylacetylene⁶² and oxidation of alkanes⁶³ are also reported.



Nitriles undergo nucleophilic addition of pyrazoles in the coordination sphere to afford pyrazolylamidino complexes.⁶⁴ In addition, catalytic hydration of nitriles has been achieved by protic pyrazole complexes under neutral conditions (Scheme 14).⁶⁵ Although the detailed mechanism remains unclear, the catalysis may be explained by the metal-ligand bifunctional cooperation, wherein the Brønsted basic pyrazolato ligand activates the nucleophilic water molecule through hydrogen bonding.⁴¹


The bifunctional nature of the protic pyrazole complexes would also be beneficial for simultaneous activation of two functional groups in one substrate molecule. We have disclosed that the iridium pyrazolato complex **43** catalyzes intramolecular

Scheme 14 Catalytic hydration of nitriles with a protic pyrazole complex. L = Me₂SO.



Scheme 15 A proposed mechanism for intramolecular hydroamination of aminoalkenes catalyzed by pyrazolato iridium complex **43**. The phenyl group in the pyrazole ring is omitted. Ir = Cp*Ir.

cyclization of aminoalkenes as shown in eqn (10).^{51,66} This hydroamination reaction is tolerant of various functional groups such as ester, bromo, cyano and hydroxy groups. The catalyst is also applicable to less reactive aminoalkenes bearing a primary amino group or without *gem*-substituents on the linker chain. The low activity of the N-methylated pyrazole complex as well as a related γ -protic imidazole complex indicates that the β -NH group is pivotal for efficient catalysis. Scheme 15 illustrates a possible mechanism of the catalytic hydroamination. The olefin bond is activated by the Lewis acidic metal center toward nucleophilic attack of the amino group, which may be assisted by the hydrogen bond with the Brønsted basic pyrazolato ligand. The two-point interaction between the aminoalkenes and catalyst in **49** may rationalize the excellent catalytic activity and diastereoselectivity. Subsequent proton transfer from the pyrazole ligand would afford the cyclization product. An alternative mechanism, which features *anti*-addition of the amino group (**50**) and rate-determining Ir–C bond protonolysis aided by a hydrogen bond network involving external protic amine molecules $R''NH_2$ (**51**), has also been proposed.⁶⁷




4.2 Bis(pyrazole) chelate complexes

The intriguing bifunctional reactivities of the protic pyrazole complexes described above prompted the exploration of poly(protic pyrazole) complexes bearing two or more NH groups, which would


efficiently work in multipoint recognition of substrates and multiple proton transfer. In this context, the 2,6-di(1*H*-pyrazol-3-yl)pyridines are attractive components since this class of pincer-type ligands provides a unique metal–ligand multifunctional platform wherein the rigid chelate framework fixes the two coplanar β -NH units adjacent to a coordination site.^{9,68}

4.2.1 Hydrogenation and polymerization catalysis. In their pioneering work, Thiel and co-workers revealed that the protic pincer-type bis(pyrazolyl)pyridine complex **52** along with the bis(pyrazole) complex **53** catalyzes hydrogenation and transfer hydrogenation of acetophenone.⁶⁹ As in the transfer hydrogenation with the related mono(pyrazole) complexes as well as bis(pyrazolyl)pyridine complexes,⁷⁰ however, cooperation of these NH units in the second coordination sphere has not been substantiated because an N-allylated derivative of **52** also exhibits comparable catalytic activity.⁷¹ Lang and co-workers described polymerization of methyl methacrylate⁷² and ring-opening polymerization of ϵ -caprolactone⁷³ catalyzed by protic bis(pyrazolyl)pyridine copper and iron complexes. Application to Suzuki–Miyaura cross-coupling has also been reported.⁷⁴




4.2.2 Multiple proton–electron transfer. We have recently described catalytic disproportionation of hydrazine to ammonia and dinitrogen promoted by the protic bis(pyrazolyl)pyridine iron complex **54**.⁷⁵ Masking even one of the two β -NH units with a methyl group leads to a significant decrease in the catalytic activity, indicating that both of the two cooperating groups are indispensable. The proposed mechanism shown in Scheme 16 features multiple and bidirectional proton-coupled electron transfer (PCET) between the metal–ligand bifunctional platform and the substrate. A hydrogen bond between one of the β -NH units and the distal nitrogen atom in the ligated hydrazine in **55** assists reductive N–N bond cleavage of hydrazine to give the $Fe^{IV}(NH_2)$ complex **56**. This step formally involves PCET from the bifunctional iron complex to the hydrazine. The remaining β -NH unit facilitates the substitution of the amido ligand by the second molecule of hydrazine to afford the hydrazido(1–) Fe^{IV} complex **57**. Subsequent PCET from the hydrazido ligand to the pyrazolato Fe^{IV} fragment leads to the formation of the Fe^{II} diazene complexes **58**. An X-ray analysis of the isolable phenyl-diazene complex **58b** has revealed that the diazene ligand, a key catalytic intermediate, is stabilized by a hydrogen bonding network involving the two β -NH units and counteranions. Finally, the diazene ligand in **58a** liberates and disproportionates to dinitrogen and hydrazine.

We have also synthesized a pincer-type ruthenium complex **59** bearing both protic NHC and pyrazole arms.⁷⁶ Deprotonation of **59** afforded an NHC–pyrazolato complex, indicating that the pyrazole group is more acidic. The C_3 -symmetric iron complex **60**



Scheme 16 A proposed mechanism for N–N bond cleavage of hydrazine catalyzed by protic bis(pyrazolyl)pyridine iron complex **54**. The total charges of the iron complexes (2+), triflate counteranions and *tert*-butyl groups in the pincer ligand have been omitted. Fe = {Fe(PMe₃)₂}²⁺.

has three protic pyrazole groups fixed by a hexadentate chelate ligand.⁷⁷ The NH units in the second coordination sphere are engaged in hydrogen bonds with the perchlorate counteranions.



5. Conclusion and perspectives

Cooperation of the metals and NH functionality in the coordination sphere has been recognized as an attractive strategy for efficient catalysis. In α -protic amine complexes⁴ the metal and NH group are in close proximity, whilst in β -protic NHC and pyrazole complexes the two functional groups are more separated yet electronically conjugated *via* π -electron systems, which allow the interplay between coordination and hydrogen bonds. A number of intriguing stoichiometric reactions of these β -protic complexes thus have been reported over the last decade. These findings also led to the

development of the catalysis such as hydration of nitrile and intramolecular hydroamination along with simple hydrogenation and transfer hydrogenation. The direct cooperation of the β -NH unit to these catalysts is, however, not always evident. In some cases, N-alkylated derivatives and γ -protic imidazole analogues exhibit comparable or even better catalytic activity. More detailed mechanistic studies are obviously required to establish the bifunctional cooperative catalysis of the β -protic complexes. These studies should include quantitative assessment of the acidity of the NH units as well as the electron-donating ability of the proton-responsive N-heterocyclic ligands. The multiproton-responsive complex based on two or more β -protic moieties would also provide a promising research area. In particular, electron transfer triggered by multiple proton transfer from the β -NH units may facilitate transformation of chemically inert molecules such as carbon dioxide and dinitrogen.

Acknowledgements

This work was supported by a Grant-in-Aid for Scientific Research (S) (No. 22225004) from the Ministry of Education, Culture, Sports, Science and Technology, Japan, The Naito Foundation (S.K.) and The Sumitomo Foundation (S.K.).

Notes and references

- 1 Y. Nicolet, A. L. de Lacey, X. Vernède, V. M. Fernandez, E. C. Hatchikian and J. C. Fontecilla-Camps, *J. Am. Chem. Soc.*, 2001, **123**, 1596; N. Wang, M. Wang, L. Chen and L. Sun, *Dalton Trans.*, 2013, **42**, 12059; D. Schilter and T. B. Rauchfuss, *Angew. Chem., Int. Ed.*, 2013, **52**, 13518.
- 2 A. D. Wilson, R. H. Newell, M. J. McNeven, J. T. Muckerman, M. Rakowski DuBois and D. L. DuBois, *J. Am. Chem. Soc.*, 2006, **128**, 358; M. Rakowski DuBois and D. L. DuBois, *Chem. Soc. Rev.*, 2009, **38**, 62; M. Rakowski DuBois and D. L. DuBois, *Acc. Chem. Res.*, 2009, **42**, 1974; D. L. DuBois and R. M. Bullock, *Eur. J. Inorg. Chem.*, 2011, 1017; J. Y. Yang, S. E. Smith, T. Liu, W. G. Dougherty, W. A. Hoffert, W. S. Kassel, M. Rakowski DuBois, D. L. DuBois and R. M. Bullock, *J. Am. Chem. Soc.*, 2013, **135**, 9700.
- 3 C. Gunanathan and D. Milstein, *Acc. Chem. Res.*, 2011, **44**, 588; T. Zell, P. Milko, K. L. Fillman, Y. Diskin-Posner, T. Bendikov, M. A. Iron, G. Leituss, Y. Ben-David, M. L. Neidig and D. Milstein, *Chem. – Eur. J.*, 2014, **20**, 4403; M. Vogt, A. Nerush, M. A. Iron, G. Leituss, Y. Diskin-Posner, L. J. W. Shimon, Y. Ben-David and D. Milstein, *J. Am. Chem. Soc.*, 2013, **135**, 17004; D. Srimani, Y. Ben-David and D. Milstein, *Chem. Commun.*, 2013, **49**, 6632.
- 4 T. Ikariya, *Bull. Chem. Soc. Jpn.*, 2011, **84**, 1; P. A. Dub and T. Ikariya, *ACS Catal.*, 2012, **2**, 1718; B. Zhao, Z. Han and K. Ding, *Angew. Chem., Int. Ed.*, 2013, **52**, 4744; P. E. Sues, K. Z. Demmans and R. H. Morris, *Dalton Trans.*, 2014, **43**, 7650; M. L. Clarke, *Synlett*, 2014, 1371.
- 5 M. C. Warner and J.-E. Bäckvall, *Acc. Chem. Res.*, 2013, **46**, 2545; B. L. Conley, M. K. Pennington-Boggio, E. Boz and T. J. Williams, *Chem. Rev.*, 2010, **110**, 2294; R. Karvembu, R. Prabhakaran and K. Natarajan, *Coord. Chem. Rev.*, 2005, **249**, 911.
- 6 Recent reviews on cooperating ligands: M. Raynal, P. Ballester, A. Vidal-Ferran and P. W. N. M. van Leeuwen, *Chem. Soc. Rev.*, 2014, **43**, 1660; S. R. Flynn and D. F. Wass, *ACS Catal.*, 2013, **3**, 2574; J. Reedijk, *Chem. Soc. Rev.*, 2013, **42**, 1776; O. R. Luca and R. H. Crabtree, *Chem. Soc. Rev.*, 2013, **42**, 1440; O. Eisenstein and R. H. Crabtree, *New J. Chem.*, 2013, **37**, 21; V. K. K. Praneeth, M. R. Ringenberg and T. R. Ward, *Angew. Chem., Int. Ed.*, 2012, **51**, 11228; D. Gelman and S. Musa, *ACS Catal.*, 2012, **2**, 2456; B. Askevold, H. W. Roesky and S. Schneider, *ChemCatChem*, 2012, **4**, 307; J. I. van der Vlugt, *Eur. J. Inorg. Chem.*, 2012, 363; G. Bauer

- and K. A. Kirchner, *Angew. Chem., Int. Ed.*, 2011, **50**, 5798; D. B. Grotjahn, *Chem. Lett.*, 2010, **39**, 908.
- 7 S. Kuwata and T. Ikariya, *Chem. – Eur. J.*, 2011, **17**, 3542.
 - 8 F. E. Hahn, *ChemCatChem*, 2013, **5**, 419; S. J. Houk, L.-A. Schaper, W. A. Herrmann and F. E. Kühn, *Chem. Soc. Rev.*, 2013, **42**, 5073; P. G. Edwards and F. E. Hahn, *Dalton Trans.*, 2011, **40**, 10278.
 - 9 M. A. Halcrow, *New J. Chem.*, 2014, **38**, 1868; M. A. Halcrow, *Coord. Chem. Rev.*, 2005, **249**, 2880.
 - 10 M. A. Halcrow, *Dalton Trans.*, 2009, 2059; J. Pérez and L. Riera, *Eur. J. Inorg. Chem.*, 2009, 4913.
 - 11 Recent reviews: K. Riener, S. Haslinger, A. Raba, M. P. Högerl, M. Cokoja, W. A. Herrmann and F. E. Kühn, *Chem. Rev.*, 2014, **114**, 5215; C. Valente, M. Pompeo, M. Sayah and M. G. Organ, *Org. Process Res. Dev.*, 2014, **18**, 180; O. Songis and C. S. J. Cazin, *Synlett*, 2013, 1877; L. Zhang and Z. Hou, *Chem. Sci.*, 2013, **4**, 3395; M. Yus and I. M. Pastor, *Chem. Lett.*, 2013, **42**, 94; H. D. Velazquez and F. Verpoort, *Chem. Soc. Rev.*, 2012, **41**, 7032; A. Kumar and P. Ghosh, *Eur. J. Inorg. Chem.*, 2012, 3955; L. Delaude and A. Demonceau, *Dalton Trans.*, 2012, **41**, 9257; J. Tornatzky, A. Kannenberg and S. Blechert, *Dalton Trans.*, 2012, **41**, 8215.
 - 12 G. A. McGibbon, C. Heinemann, D. J. Lavorato and H. Schwarz, *Angew. Chem., Int. Ed.*, 1997, **36**, 1478; G. Sini, O. Eisenstein and R. H. Crabtree, *Inorg. Chem.*, 2002, **41**, 602; L. J. L. Häller and S. A. Macgregor, *Eur. J. Inorg. Chem.*, 2009, 2000.
 - 13 Recent examples: C.-O. Ng, S.-M. Yiu and C.-C. Ko, *Inorg. Chem.*, 2014, **53**, 3022; J. Ruiz, L. García, C. Mejuto, M. Vivanco, M. R. Diaz and S. García-Granda, *Chem. Commun.*, 2014, **50**, 2129; C.-H. Chen, Y.-H. Liu, S.-M. Peng, J.-T. Chen and S.-T. Liu, *Dalton Trans.*, 2012, **41**, 2747.
 - 14 T. Kösterke, T. Pape and F. E. Hahn, *J. Am. Chem. Soc.*, 2011, **133**, 2112; recent examples: R. Das, C. G. Daniliuc and F. E. Hahn, *Angew. Chem., Int. Ed.*, 2014, **53**, 1163; D. Brackmeyer, A. Hervé, C. Shulte to Brinke, M. C. Jahnke and F. E. Hahn, *J. Am. Chem. Soc.*, 2014, **136**, 7841.
 - 15 N. Meier, F. E. Hahn, T. Pape, C. Siering and S. R. Waldvogel, *Eur. J. Inorg. Chem.*, 2007, 1210.
 - 16 X. Wang, H. Chen and X. Li, *Organometallics*, 2007, **26**, 4684.
 - 17 For addition reactions to coordinated nitriles, see: V. Yu Kukushkin and A. J. L. Pombeiro, *Chem. Rev.*, 2002, **102**, 1771.
 - 18 D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; J. C. Lewis, R. G. Bergman and J. A. Ellman, *Acc. Chem. Res.*, 2008, **41**, 1013.
 - 19 J. C. Lewis, S. H. Wiedermann, R. G. Bergman and J. A. Ellman, *Org. Lett.*, 2004, **6**, 35; J. C. Lewis, A. M. Berman, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2008, **130**, 2493.
 - 20 K. L. Tan, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2002, **124**, 13964; K. L. Tan, S. Park, J. A. Ellman and R. G. Bergman, *J. Org. Chem.*, 2004, **69**, 7329.
 - 21 S. H. Wiedermann, J. C. Lewis, J. A. Ellman and R. G. Bergman, *J. Am. Chem. Soc.*, 2006, **128**, 2452.
 - 22 K. L. Tan, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2002, **124**, 3202.
 - 23 K. L. Tan, R. G. Bergman and J. A. Ellman, *J. Am. Chem. Soc.*, 2001, **123**, 2685.
 - 24 K. Araki, S. Kuwata and T. Ikariya, *Organometallics*, 2008, **27**, 2176.
 - 25 V. Miranda-Soto, D. B. Grotjahn, A. L. Cooksy, J. A. Golen, C. E. Moore and A. L. Rheingold, *Angew. Chem., Int. Ed.*, 2011, **50**, 631.
 - 26 For heterolytic cleavage of dihydrogen on metal–ligand cooperating bifunctional complexes, see: G. J. Kubas, *J. Organomet. Chem.*, 2014, **751**, 33; R. H. Crabtree, *New J. Chem.*, 2011, **35**, 18; J. C. Gordon and G. J. Kubas, *Organometallics*, 2010, **29**, 4682; M. Ito and T. Ikariya, *Chem. Commun.*, 2007, 5134.
 - 27 V. Miranda-Soto, D. B. Grotjahn, A. G. DiPasquale and A. L. Rheingold, *J. Am. Chem. Soc.*, 2008, **130**, 13200.
 - 28 F. E. Hahn, A. R. Naziruddin, A. Hepp and T. Pape, *Organometallics*, 2010, **29**, 5283.
 - 29 B. Eguillor, M. A. Esteruelas, J. García-Raboso, M. Oliván, E. Oñate, I. M. Pastor, I. Peñaflor and M. Yus, *Organometallics*, 2011, **30**, 1658.
 - 30 O. Schuster, L. Yang, H. G. Raubenheimer and M. Albrecht, *Chem. Rev.*, 2009, **109**, 3445.
 - 31 D. Kunz, *Angew. Chem., Int. Ed.*, 2007, **46**, 3405; E. Álvarez, S. Conejero, P. Lara, J. A. López, M. Paneque, A. Petronilho, M. L. Poveda, D. del Río, O. Serrano and E. Carmona, *J. Am. Chem. Soc.*, 2007, **129**, 14130.
 - 32 S. Conejero, J. López-Serrano, M. Paneque, A. Petronilho, M. L. Poveda, F. Vattier, E. Alvarez and E. Carmona, *Chem. – Eur. J.*, 2012, **18**, 4644.
 - 33 E. Alvarez, S. Conejero, M. Paneque, A. Petronilho, M. L. Poveda, O. Serrano and E. Carmona, *J. Am. Chem. Soc.*, 2006, **128**, 13060; M. Paneque, M. L. Poveda, F. Vattier, E. Alvarez and E. Carmona, *Chem. Commun.*, 2009, 5561; S. Conejero, C. Maya, M. Paneque, A. Petronilho, M. L. Poveda, F. Vattier, E. Álvarez, E. Carmona, A. Laguna and O. Crespo, *Dalton Trans.*, 2012, **41**, 14126.
 - 34 M. A. Esteruelas, F. J. Fernández-Alvarez and E. Oñate, *J. Am. Chem. Soc.*, 2006, **128**, 13044; M. A. Esteruelas, F. J. Fernández-Alvarez and E. Oñate, *Organometallics*, 2007, **26**, 5239; M. L. Buil, M. A. Esteruelas, K. Garcés, M. Oliván and E. Oñate, *J. Am. Chem. Soc.*, 2007, **129**, 10998; M. L. Buil, M. A. Esteruelas, K. Garcés, M. Oliván and E. Oñate, *Organometallics*, 2008, **27**, 4680; M. A. Esteruelas, F. J. Fernández-Alvarez and E. Oñate, *Organometallics*, 2008, **27**, 6236; M. A. Esteruelas, F. J. Fernández-Alvarez, M. Oliván and E. Oñate, *Organometallics*, 2009, **28**, 2276.
 - 35 S. Bajo, M. A. Esteruelas, A. M. López and E. Oñate, *Organometallics*, 2012, **31**, 8618.
 - 36 G. Song, Y. Li, S. Chen and X. Li, *Chem. Commun.*, 2008, 3558.
 - 37 G. Song, Y. Su, R. A. Periana, R. H. Crabtree, K. Han, H. Zhang and X. Li, *Angew. Chem., Int. Ed.*, 2010, **49**, 912.
 - 38 B. Crociani, F. Di Bianca, A. Gioenco and A. Scriveri, *J. Organomet. Chem.*, 1983, **251**, 393; K. Isobe, Y. Nakamura, T. Miwa and S. Kawaguchi, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 149; H. G. Raubenheimer, J. G. Toerien, G. J. Kruger, R. Otte, W. van Zyl and P. Olivier, *J. Organomet. Chem.*, 1994, **466**, 291.
 - 39 A. Poulain, A. Neels and M. Albrecht, *Eur. J. Inorg. Chem.*, 2009, 1871.
 - 40 E. Álvarez, Y. A. Hernández, J. López-Serrano, C. Maya, M. Paneque, A. Petronilho, M. L. Poveda, V. Salazar, F. Vattier and E. Carmona, *Angew. Chem., Int. Ed.*, 2010, **49**, 3496; C. Cristóbal, Y. A. Hernández, J. López-Serrano, M. Paneque, A. Petronilho, M. L. Poveda, V. Salazar, F. Vattier, E. Álvarez, C. Maya and E. Carmona, *Chem. – Eur. J.*, 2013, **19**, 4003.
 - 41 T. J. Ahmed, S. M. M. Knapp and D. R. Tyler, *Coord. Chem. Rev.*, 2011, **255**, 949; R. García-Álvarez, P. Crochet and V. Cadierno, *Green Chem.*, 2013, **15**, 46.
 - 42 Y. A. Hernández, J. López-Serrano, M. Paneque, M. L. Poveda, F. Vattier, V. Salazar, E. Álvarez and E. Carmona, *Chem. – Eur. J.*, 2011, **17**, 9302.
 - 43 C. Cristóbal, E. Álvarez, M. Paneque and M. L. Poveda, *Organometallics*, 2013, **32**, 714.
 - 44 F. Vattier, V. Salazar, M. Paneque, M. L. Poveda and E. Álvarez, *Organometallics*, 2014, **33**, 498.
 - 45 D. G. Johnson, J. M. Lynam, N. S. Mistry, J. M. Slattery, R. J. Thatcher and A. C. Whitwood, *J. Am. Chem. Soc.*, 2013, **135**, 2222; M. Murakami and S. Hori, *J. Am. Chem. Soc.*, 2003, **125**, 4720.
 - 46 A. M. Berman, R. G. Bergman and J. A. Ellman, *J. Org. Chem.*, 2010, **75**, 7863.
 - 47 Y. Su, G. Song, K. Han and X. Li, *J. Organomet. Chem.*, 2011, **696**, 1640.
 - 48 D. B. Grotjahn, *Dalton Trans.*, 2008, 6497.
 - 49 Even a nonmetallic, pyrazole-based frustrated Lewis pair that reacts with H₂, CO₂ and some organic molecules is reported: E. Theuergarten, J. Schlösser, D. Schlüns, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, *Dalton Trans.*, 2012, **41**, 9101 and references therein.
 - 50 Y. Kashiwame, M. Watanabe, K. Araki, S. Kuwata and T. Ikariya, *Bull. Chem. Soc. Jpn.*, 2011, **84**, 251.
 - 51 Y. Kashiwame, S. Kuwata and T. Ikariya, *Chem. – Eur. J.*, 2010, **16**, 766.
 - 52 F. Zeng and Z. Yu, *Organometallics*, 2008, **27**, 6025; W. Jin, L. Wang and Z. Yu, *Organometallics*, 2012, **31**, 5664; W. Ye, M. Zhao and Z. Yu, *Chem. – Eur. J.*, 2012, **18**, 10843.
 - 53 F. Zeng and Z. Yu, *Organometallics*, 2008, **27**, 2898; F. Zeng and Z. Yu, *Organometallics*, 2009, **28**, 1855.
 - 54 W. Du, Q. Wang, L. Wang and Z. Yu, *Organometallics*, 2014, **33**, 974 and references therein.
 - 55 T. Kuwahara, T. Fukuyama and I. Ryu, *RSC Adv.*, 2013, **3**, 13702.
 - 56 K. Muller, Y. Sun, A. Heimermann, F. Menges, G. Niedner-Schatteburg, C. van Wüllen and W. R. Thiel, *Chem. – Eur. J.*, 2013, **19**, 7825.
 - 57 L. Bellarosa, J. Díez, J. Gimeno, A. Lledós, F. J. Suárez, G. Ujaque and C. Vicent, *Chem. – Eur. J.*, 2012, **18**, 7749.
 - 58 J. Díez, J. Gimeno, A. Lledós, F. J. Suárez and C. Vicent, *ACS Catal.*, 2012, **2**, 2087.

- 59 A. Satake and T. Nakata, *J. Am. Chem. Soc.*, 1998, **120**, 10391.
- 60 Y. Sun and W. R. Thiel, *Inorg. Chim. Acta*, 2006, **359**, 4807; V. Montoya, J. Pons, V. Branchadell, J. Garcia-Antón, X. Solans, M. Font-Bardia and J. Ros, *Organometallics*, 2008, **27**, 1084.
- 61 C. Chen, H. Qin and W. Chen, *Inorg. Chem.*, 2011, **50**, 8671.
- 62 A. P. Martínez, M. J. Fabra, M. P. García, F. J. Lahoz, L. A. Oro and S. J. Teat, *Inorg. Chim. Acta*, 2005, **358**, 1635.
- 63 I. Gryca, B. Machura, J. G. Matecki, L. S. Shul'pina, A. J. L. Pombeiro and G. B. Shul'pin, *Dalton Trans.*, 2014, **43**, 5759.
- 64 C.-C. Hsieh, C.-J. Lee and Y.-C. Horng, *Organometallics*, 2009, **28**, 4923; M. Arroyo, P. Gómez-Iglesias, J. M. Martín-Alvarez, C. M. Alvarez, D. Miguel and F. Villafañe, *Inorg. Chem.*, 2012, **51**, 6070 and references therein.
- 65 Í. Ferrer, J. Rich, X. Fontrodona, M. Rodríguez and I. Romero, *Dalton Trans.*, 2013, **42**, 13461.
- 66 Y. Kashiwame, S. Kuwata and T. Ikariya, *Organometallics*, 2012, **31**, 8444.
- 67 S. Tobisch, *Chem. – Eur. J.*, 2012, **18**, 7248.
- 68 K. Umehara, S. Kuwata and T. Ikariya, *Inorg. Chim. Acta*, 2014, **413**, 136; J. S. Costa, S. Rodríguez-Jiménez, G. A. Craig, B. Barth, C. M. Beavers, S. J. Teat and G. Aromí, *J. Am. Chem. Soc.*, 2014, **136**, 3869; I.-R. Jeon, J. G. Park, C. R. Haney and T. D. Harris, *Chem. Sci.*, 2014, **5**, 2461; X. Wang, F. Y. Bai, Y. H. Xing, L. J. Wan, Q. L. Guan, Y. N. Hou and R. Zhang, *Inorg. Chim. Acta*, 2014, **416**, 171; T. D. Roberts, M. A. Little, L. J. K. Cook and M. A. Halcrow, *Dalton Trans.*, 2014, **43**, 7577; A. K. Hui, B. J. Cook, D. J. Mindiola and K. G. Caulton, *Inorg. Chem.*, 2014, **53**, 3307; G. A. Craig, M. Schütze, D. Aguila, O. Roubeau, J. Ribas-Ariño, S. Vela, S. J. Teat and G. Aromí, *Inorg. Chem.*, 2014, **53**, 3290; T. Shiga, M. Noguchi, H. Sato, T. Matsumoto, G. N. Newton and H. Oshio, *Dalton Trans.*, 2013, **42**, 16185; X. Zhang, N. Xing, F. Bai, L. Wan, H. Shan, Y. Hou, Y. Xing and Z. Shi, *CrystEngComm*, 2013, **15**, 9135; C. Chen, Q. Sun, D.-X. Ren, R. Zhang, F.-Y. Bai, Y.-H. Xing and Z. Shi, *CrystEngComm*, 2013, **15**, 5561; A. Yoshinari, A. Tazawa, S. Kuwata and T. Ikariya, *Chem. – Asian J.*, 2012, **7**, 1417.
- 69 T. Jozak, D. Zabel, A. Schubert, Y. Sun and W. R. Thiel, *Eur. J. Inorg. Chem.*, 2010, 5135.
- 70 L. Wang, Q. Yang, H. Chen and R.-X. Li, *Inorg. Chem. Commun.*, 2011, **14**, 1884; S. Günnaz, N. Özdemir, S. Dayan, O. Dayan and B. Çetinkaya, *Organometallics*, 2011, **30**, 6145; O. Dayan, N. Özdemir, Z. Şerbetci, M. Dinçer, B. Çetinkaya and O. Büyükgüngör, *Inorg. Chim. Acta*, 2012, **392**, 246; O. Dayan, S. Dayan, İ. Kani and B. Çetinkaya, *Adv. Organomet. Chem.*, 2012, **26**, 663.
- 71 L. T. Ghoochany, S. Farsadpour, Y. Sun and W. R. Thiel, *Eur. J. Inorg. Chem.*, 2011, 3431.
- 72 L.-L. Miao, H.-X. Li, M. Yu, W. Zhao, W.-J. Gong, J. Gao, Z.-G. Ren, H.-F. Wang and J.-P. Lang, *Dalton Trans.*, 2012, **41**, 3424.
- 73 Y.-Y. Fang, W.-J. Gong, X.-J. Shang, H.-X. Li, J. Gao and J.-P. Lang, *Dalton Trans.*, 2014, **43**, 8282.
- 74 Q. Yang, L. Wang, L. Lei, X.-L. Zheng, H.-Y. Fu, M.-L. Yuan, H. Chen and R.-X. Li, *Catal. Commun.*, 2012, **29**, 194.
- 75 K. Umehara, S. Kuwata and T. Ikariya, *J. Am. Chem. Soc.*, 2013, **135**, 6754.
- 76 T. Toda, S. Kuwata and T. Ikariya, *Chem. – Eur. J.*, 2014, **20**, 9539.
- 77 S. A. Diener, A. Santoro, C. A. Kilner, J. J. Loughrey and M. A. Halcrow, *Dalton Trans.*, 2012, **41**, 3731.