

Cite this: *RSC Adv.*, 2014, 4, 48191

Ball milling for the quantitative and specific solvent-free Knoevenagel condensation + Michael addition cascade in the synthesis of various 2-amino-4-aryl-3-cyano-4*H*-chromenes without heating†

Omid Hosseini Qareaghaj,^a Sara Mashkouri,^a M. Reza Naimi-Jamal^{*a} and Gerd Kaupp^b

Ball milling is used as a facile, efficient, cheap, and environmentally friendly procedure for the solvent-free three-component reaction of aromatic aldehydes with malononitrile and dimesone, or 1,3-cyclohexanedione, resorcinol, and α - and β -naphthol. The reactions proceed quantitatively at room temperature by milling stoichiometric mixtures of the reagents in the presence of 10 mol% Na_2CO_3 . This method offers the advantages of short reaction times, low cost, quantitative yields and simple work-up with no need of any organic solvent. It is thus enviro-economic without producing dangerous wastes. The orientation selectivity with β -naphthol and resorcinol is discussed. Existing structural misattributions are pointed out and clarified by ^1H NMR spin coupling analysis.

Received 3rd July 2014
Accepted 2nd September 2014

DOI: 10.1039/c4ra06603k

www.rsc.org/advances

Introduction

The versatile synthesis of 2-amino-3-cyano-4-aryl-4*H*-chromenes is presently one of the most studied vehicles for the presentation of new catalysts with mostly weak acids and bases, even though uncatalyzed reactions of aldehydes with malononitrile and 1,3-diketones provide quantitative wasteless yields in stoichiometric solventless melt-reactions at still moderate temperatures (125–150 °C). Clearly, the uncatalyzed syntheses provide the best possible scale-up possibilities,¹ but also enviro-economic catalysis by milling with soda at room temperature is of interest for reactions with less reactive reagents.² The general importance of chromene derivatives or amino-4*H*-chromenes has been amply reported in ref. 3 and in the numerous papers cited therein, but we focus here on the importance of 2-amino-3-cyano-4-aryl-4*H*-chromenes. These compounds have been identified as having the capacity to prevent various types of human diseases.^{4,5} These include antiallergic,⁶ antitumor^{7–9} and antibacterial activities.¹⁰ Most recent interest has focused on these compounds with additional substituents for anti-microbial activities in

pharmaceutical applications and biological studies.^{11,12} Furthermore, the 2-amino-4-aryl-3-cyano-substituted 4*H*-chromenes are being used as building blocks for the synthesis of human drugs for many kinds of neurodegenerative and other serious diseases.^{13,14} For example, most 4*H*-chromene derivatives with particular substituents show antibacterial activity even against highly resistant *Staphylococcus aureus* and others.¹⁵ Several of these compounds are thus already commercially available. Importantly, also both the 2-amino and the 3-cyano substituents have been variously converted with standard transformations to create a multitude of compounds with useful applications. But unfortunately there is some confusion as to the correct chemical structure of these very important commercial compounds, because the necessary corrections of unduly claimed orientation selectivities in some of the multiple component reactions (MCRs) were never performed, and the reasons for the selectivities were not discussed.

The synthesis by Knoevenagel condensation + Michael addition of the highly substituted 4*H*-chromenes has been varied by using many methods in recent years in an attempt to improve ease and efficiency by catalysis. The two-step approach by cyclization of preformed arylidenemalononitriles with β -dicarbonyl compounds is accelerated by the presence of an organic base such as piperidine,¹⁶ morpholine, pyridine,¹⁷ and triethylamine.^{18,19} Furthermore, despite our report of an early quantitative wasteless procedure for stoichiometric melt reactions without catalyst,¹ further two-step procedures have been tried to decrease the temperature by use of a catalyst at the expense of lower yield and the need for chromatographic

^aResearch Laboratory of Green Organic Synthesis and Polymers, Department of Chemistry, Iran University of Science and Technology, 16846 Tehran, Iran. E-mail: naimi@iust.ac.ir

^bFaculty 5, Department of Chemistry, University of Oldenburg, 26188 Edewecht, Germany

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ra06603k

workup.^{20–37} This is also true for the later tried MCRs for the preparation of 4*H*-chromenes. The catalytic processes for the one-pot synthesis of 2-amino-4-aryl-3-cyano-substituted 4*H*-chromenes have been recently complemented by the application of potassium phthalimido-*N*-hydroxylate in water as another base though by use of 10% excess of the reagent malononitrile,³ which however creates additional wastes when recycling the catalyst. Other approaches utilized for example hexadecyltrimethylammonium bromide (HTMAB),²⁰ sodium stearate,²¹ 4-dodecylbenzenesulfonic acid (DBSA),²⁰ $\text{KF-Al}_2\text{O}_3$,²² the rare earth perfluorooctanoate (Re(PFO)_3),²³ (*S*)-proline,²⁴ $(\text{NH}_4)_2\text{HPO}_4$,²⁵ NMe_4OH ,²⁶ MgO ,^{27,28} tetrabutylammonium fluoride (TBAF),²⁹ $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$,³⁰ $\text{ZnO-}\beta$ -zeolite,³¹ 1,4-diazabicyclo [2.2.2]octane (DABCO),³² silica gel-supported polyphosphoric acid (PPA-SiO_2),³³ $\gamma\text{-Fe}_2\text{O}_3$,³⁴ Caro's acid silica gel,³⁵ and amino-functionalized ionic liquids.³⁶ Some of these methods applied organic solvent such as ethanol, dimethylformamide (DMF) or acetic acid,^{24,37} others water with propylpiperazine-*N*-sulfonic acid,³⁸ or hydrocalcite,³⁹ or 1-deoxy-1-methylaminosorbitol.⁴⁰ Also, heating by microwave⁴¹ or ultrasonic irradiation⁴² was applied. However, most of these procedures have drawbacks of producing dangerous wastes or providing less than quantitative yield of pure product. These are as yet only overcome upon stoichiometric heating of the components at 150 °C and 100–130 °C without catalyst.¹ Some catalysis protocols still require elevated temperatures, often hazardous organic solvents, and organic bases. Furthermore, the catalysts are often expensive or not commercially available, and tedious workup procedures are the rule with many wastes, because of only moderate product yields, requiring chromatographic workup and recrystallization. Thus, quantitative yield of the target products at room temperature omitting wasteful workup is highly desired. We therefore applied the usefulness of the solventless (solvent-free when choosing aqueous workup) stoichiometric MCR technique with the easily removable environmentally safe catalyst sodium carbonate in a ball-mill,^{2,43–47} and keep in mind the possibility that ball milling is scalable to hundreds of grams,^{48–52} which contrasts with recent reviews on milling in organic synthesis with complaints of lack of scale-up developments.⁵³ But the existence of the reviews^{48–52} on such scale-up (further applications in ref. 48–52) has been pointed out.⁵⁴ We have already reported that combining MCRs and ball milling allows for the synthesis of a wide variety of interesting organic building blocks.^{43–52,55,56} We also reported that sodium carbonate can be a useful basic catalyst in kneading milling.^{44–47} This technique benefits from both the economical and the synthetic point of view by not only omitting the unnecessary use of organic solvents, but also enhancing the rate of many organic reactions.


The ongoing hype with multiple repetitions of the MCR syntheses in question, providing high melting-point products, bears the risk that the characterization of the products rests solely on melting point comparison rather than on interpretation of ¹H NMR coupling schemes that must fit with the claimed orientation specificities in these reactions. Unfortunately, there are never-resolved discrepancies in the claimed molecular structures when for example β -naphthol or resorcinol was the

reagent for Michael additions followed by cyclization, and one must not invoke melting points of published unassigned or merely claimed molecular structures without considering a safe structure elucidation.

With the desire to understand the surprising orientation specificities in these MCR syntheses, we continued our preliminary grinding and heating experiments² by ball milling with sodium carbonate catalysis in order to obtain quantitative yields without heating and without producing dangerous wastes. We assess the correctness of the product structures and report on the selectivity of the weakly basic catalyst towards phenols. These one-pot solvent-free syntheses react various benzaldehydes with malononitrile and dimedone, or 1,3-cyclohexandione, or resorcinol, or α - or β -naphthol. Quantitative yields are indeed obtained by kneading ball milling with sodium carbonate as an efficient, selective, cheap, environmentally economic, and easily separated catalyst without the need for external heating.

Results

It was previously reported that the catalyst-free ball-milling reaction of 4-chlorobenzaldehyde (**1**, Ar = 4-ClC₆H₄) with malononitrile (**2**) to give the Knoevenagel condensate could not be completed in the solid state after 10 min at room temperature but afforded only 7% of the corresponding benzylidene malononitrile **9** (Scheme 1).⁵⁷ However, the uncatalyzed melt reactions occurred quantitatively at 150 °C with 4-H-, 4-OH-, 4-Cl- and 4-NO₂-benzaldehyde and the stoichiometric follow-up Michael additions with dimedone (**3a**) occurred in melts at 100–130 °C to quantitatively give the 7,7-dimethyl-5-oxo-5,6,7,8-tetrahydro-4*H*-chromenes **4a**.¹ The one-pot MCR of 4-chlorobenzaldehyde (**1a**) with malononitrile (**2**) and less reactive β -



Scheme 1 Possible mechanism for the preparation of 4–8.



naphthol (**3e**) was incomplete (78% **8a**) at 150 °C (4 h), but when 10 mol% of ground Na₂CO₃ was admixed, the reaction temperature could be decreased to 125 °C (1 h) and quantitative or near-quantitative yields were obtained also with a series of further benzaldehydes (Scheme 1).²

We now found that a quantitative yield of the product **8a** (Table 1) was obtained at room temperature when the above mixture was milled for 10 min. 100% conversion was achieved because the product becomes an insoluble powder. After sublimation in a vacuum at 180 °C the product was completely separated from the recycled catalyst and this process was thus “solventless”. But such high workup temperature could be avoided by simple extraction of the catalyst from the insoluble product crystals with water in a then “solvent-free” process. No extra purifying workup was required in both techniques.

Kneading in the presence of sodium carbonate can now be used to provide the completion of both the Knoevenagel condensations by addition of aldehydes **1** with malononitrile **2** (m.p. 32 °C) followed by dehydration to give **9** and the Michael addition with 1,3-diketones **3a** and **3b** (as their enols) or with the reactive phenols **3c–e** in the 25–30 °C range. Mechanistically, the intermediates **10**, which are in equilibrium with their enols **11**, undergo intramolecular cyclization to give **12**, the tautomers of the products **4–8**.¹ Apparently, the basicity of ball-milled Na₂CO₃ and the acidity of NaHCO₃ are improved by continued creation of new surfaces of these solids (Scheme 1).

The corresponding results with dimedone (**3a**), 1,3-cyclohexanedione (**3b**), resorcinol (**3c**), and α - (**3d**) and β -naphthol (**3b**) (Scheme 2) are reported in Table 1. The times for quantitative reaction were determined by thin layer chromatography (TLC) tests. Table 1 reveals that electron-withdrawing groups accelerate the reaction. For example, all reactions of 4-nitro- and 4-cyanobenzaldehyde and **2** with **3a–3e** were completed after 10 minutes milling, but 4-methoxy- and 4-hydroxybenzaldehyde needed up to 40 minutes for completion. Also the ball milling of 4-chloro- and 4-bromobenzaldehydes proceeded rapidly and quantitatively without external heating. Only in the case of **8c** was the subsequent heating of the reaction mixture to 125 °C for 1 h necessary to afford the product quantitatively, probably due to the presence of the electron-releasing phenolic OH group and the least reactivity of β -naphthol among the other reagents **3**. Conversely, all uncatalyzed quantitative corresponding reactions required heating to 125 °C for 1 h.² Solid Na₂CO₃ has a clear advantage over K₂CO₃. In similar work, Zhou *et al.*⁵⁸ reported lower yields of 67–74% when grinding with stoichiometric amounts of K₂CO₃ and could not avoid the formation of deeply colored by-products. A major difference appears to be the different qualities of the continuously hydrated catalysts when the water of reaction is liberated. It is well known that sodium carbonate takes up 10H₂O to form a stable solid hydrate, whereas the stable solid hydrate of potassium carbonate contains only 1.5H₂O. Also stoichiometric grinding with the weaker basic NaHCO₃ was inferior in that it gave 74–93% yield upon similar grinding for 90 min, here without coloration.⁵⁸

The surprising orientation specificities giving only one product in all cases were determined by purity checks with sharp melting points, TLC and the physical data that also

correspond to the available published ones. The NMR spectroscopic data reveal only one series of resonance signals.

The structures of all products were obtained on the basis of their spectral data. The typical NH₂ and C≡N FT-IR bands are of course present in the spectra of all of the corresponding derivatives. The ¹H NMR spectra of all described products show one characteristic singlet signal in the range of 4.1–5.6 ppm for the 4-H and the ¹³C NMR spectra exhibit one specific signal in the range of 54–60 ppm for the carbon at position 4. The reaction orientation of the products **6** and **8** is still controversial. Thus, in a series of papers, Kumar^{59–61} published the reactions of β -naphthol with aromatic aldehydes and malononitrile using various catalysts and heating devices including microwaves as giving straight “2-amino-4-aryl-3-cyano-4H-benzo[*g*]chromenes” (from claimed reaction at C-3 of β -naphthol), rather than the angular 2-amino-4-aryl-3-cyano-4H-benzo[*f*]chromenes (reaction at C-1 of β -naphthol), as we do and many other research groups in the field. The same discrepancy exists when cyanoacetic ethylester is used instead of malononitrile for providing the corresponding products with 3-ethoxycarbonyl substitution. Furthermore, the CAS Registry Database also interprets the poorly defined products with β -naphthol in the papers of Pasha⁶² and Jin⁶³ with the straight ring condensation. A clear discussion of the correct structure of the products of these MCRs is of utmost importance, as several of these products are commercially available. For example the product from benzaldehyde, malononitrile and β -naphthol is still sold and advertised (December 2013 according to CAS Registry Database) by 4 companies (American Custom Chemical Corp.; ChemPur GmbH; Fluorochem Ltd; Gencore Bio PharmaPVT Ltd) as straight “4H-naphtho[2,3-*b*]pyran” (a synonym for “4H-benzo[*g*]chromene”) (CAS registry number 924708-55-4). Similar advertisements continue to exist with substituted products with claimed straight ring condensation. Conversely, 23 suppliers sell these products as angular “4H-naphtho[2,1-*b*]pyran” (a synonym for “4H-benzo[*f*]chromene”) (CAS registry number 111861-46-2 for Ar = C₆H₅). As the research groups that published the angular structure (reaction at C-1 of β -naphthol) also used various catalysts (*e.g.* TiCl₄ or bases or phase-transfer catalysis) at room temperature or with heating (including microwave), it appears clear that one of these orientation claims must be wrong. Clearly, high-resolution ¹H NMR provides a clear decision. Unfortunately, the claims of the straight structure were never substantiated by a crystal structure determination or by an analysis or discussion of the aromatic proton resonance coupling schemes by ¹H NMR. Thus these papers^{59–63} do not account for the Ar–H resonances that are only published as multiplet containing 10 or 11 Ar–H. However, our ¹H NMR spectra² exhibit the Ar–H signal at highest field for all substituted derivatives of the β -naphthol products with large splitting (7–9 Hz range of an AB spin system) evidently by the coupling with the vicinal Ar–H, and so do our here studied products (see the examples **8b** and **8c** in Fig. 1 with the analysis in the Experimental section).

Also when checking the corresponding ¹H NMR data of others, who disclosed the spectra (though without discussion), one finds the same coupling scheme. There cannot be any



Table 1 Synthesis of 2-amino-4-aryl-3-cyano-4*H*-chromenes under solvent-free kneading ball-milling conditions catalyzed by 10 mol% sodium carbonate










Entry	Aldehyde (1)	Enol-naphthol	Product	Time (min)	Yield ^a (%)	M.p. (obsd.) (°C)	M.p. (lit.) (°C)
1	4-Cl-C ₆ H ₄ CHO	3a	 4a	10	>99	213–214	(208–210) ¹⁷
2	PhCHO	3a	 4b	25	>99	219–221	(226–228) ¹⁰
3	2-Cl-C ₆ H ₄ CHO	3a	 4c	15	>99	196–197	(191–192) ²⁵
4	4-Me-C ₆ H ₄ CHO	3a	 4d	35	>99	211–213	(210–213) ²³
5	4-NO ₂ -C ₆ H ₄ CHO	3a	 4e	10	>99	177–178	(177–179) ²³
6	3-NO ₂ -C ₆ H ₄ CHO	3a	 4f	20	>99	206–209	(204–205) ²³
7	4-CN-C ₆ H ₄ CHO	3a	 4g	10	>99	221–224	(225–228) ²³
8	4-Br-C ₆ H ₄ CHO	3a	 4h	35	>99	196–198	(196–198) ⁶⁹
9	4-MeO-C ₆ H ₄ CHO	3a	 4i	35	>99	197–199	(201–202) ²⁵



Table 1 (Contd.)


















Entry	Aldehyde (1)	Enol-naphthol	Product	Time (min)	Yield ^a (%)	M.p. (obsd.) (°C)	M.p. (lit.) (°C)
10	4-HO-C ₆ H ₄ CHO	3a	 4j	40	>99	205–206	(204–205) ²⁵
11	2-NO ₂ -C ₆ H ₄ CHO	3a	 4k	15	>99	220–221	(233–234) ²⁵
12	4-Cl-C ₆ H ₄ CHO	3b	 5a	20	>99	241–244	(241–243) ⁷⁰
13	3-NO ₂ -C ₆ H ₄ CHO	3b	 5b	20	>99	229–230	(221–223) ⁷¹
14	4-NO ₂ -C ₆ H ₄ CHO	3b	 5c	10	>99	223–225	(234–236) ⁷²
15	4-Br-C ₆ H ₄ CHO	3b	 5d	35	>99	238–240	(241–242) ⁷³
16	4-HO-C ₆ H ₄ CHO	3b	 5e	40	>99	244–246	(234–236) ⁷²
17	2-Cl-C ₆ H ₄ CHO	3b	 5f	15	>99	205–208	(208–210) ⁷⁴
18	4-MeO-C ₆ H ₄ CHO	3b	 5g	35	>99	199–202	(198–200) ⁷⁵



Table 1 (Contd.)


Entry	Aldehyde (1)	Enol-naphthol	Product	Time (min)	Yield ^a (%)	M.p. (obsd.) (°C)	M.p. (lit.) (°C)
19	PhCHO	3c	 6a	10	>99	230–231	(230–232) ³
20	4-Cl-C ₆ H ₄ CHO	3c	 6b	5	>99	228–230	(230–231) ⁷⁶
21	4-Br-C ₆ H ₄ CHO	3c	 6c	5	>99	240–241	(240–241) ⁷⁷
22	4-Cl-C ₆ H ₄ CHO	3d	 7a	10	>99	233–235	(233–235) ³
23	4-Br-C ₆ H ₄ CHO	3d	 7b	10	>99	232–234	(234–236) ³
24	4-Cl-C ₆ H ₄ CHO	3e	 8a	12	>99	207–209	(207–208) ³
25	3-NO ₂ -C ₆ H ₄ CHO	3e	 8b	12	>99	237–239	(232–235) ²
26	4-HO-C ₆ H ₄ CHO	3e	 8c	30 ^b	>99	260–262	(260–262) ²

^a Isolated yields after extraction of Na₂CO₃ from the insoluble crystals with water and drying. ^b For the quantitative yield for this case, subsequent heating of the reaction mixture at 125 °C was necessary.


doubt that the Ar–H signal at highest field belongs to the hydrogen that is next to the heterocyclic oxygen. Furthermore, two slightly broadened singlet signals would have to be expected from the Ar–H moieties of the central ring in the straight

structure. But such singlets cannot be found in any of these spectra. This now safely validates the angular -4*H*-benzo[*f*]-chromene structures of the products from the reactions with β-naphthol, independent of the reaction conditions and aryl





Scheme 2 Preparation of different 2-amino-4H-pyran derivatives.

Fig. 1 500 MHz ^1H NMR spectra of **8b** and **8c** in $\text{DMSO}-d_6$ for the aromatic proton resonances.

substitution. Also, published X-ray diffraction analyses of related compounds (**8a** with H,⁶⁴ CH_3 ,⁶⁵ or Br (ref. 66) instead of Cl, the latter in the (*S*)-configuration) support the angular structure. Clearly, all chemical, biological, and medicinal results that were based on the still commercially advertised and sold products with the unsupported nominal straight “-4H-”benzo[*g*]-chromene structure must be urgently reinterpreted.

Another structural question to be settled is the surprising orientation specificity of the reactions with resorcinol. While

a reaction at its C-2 position would be electronically more probable, the greater crowding of the aryl group with the remaining OH group would disfavor that possibility. Thus, reaction at C-4 of resorcinol comes to the fore. Experimentally, one obtains the 7-hydroxy-4H-chromene structures **6** but not the 5-OH-4H-chromene isomers, not even as detectable side-products. Unfortunately, we cannot argue with product stabilities because quantum chemical calculations predict the actually obtained **6a** and the not obtained isomeric structure are energetically very close. The semi-empirical PM3 method calculates the obtained 7-OH structure as being $0.28 \text{ kcal mol}^{-1}$ more stable, but DFT B3LYP 6-31G* calculates the not obtained 5-OH structure as being $0.015 \text{ kcal mol}^{-1}$ more stable. The ^1H NMR spectra of **6a–6c** provide a clear structural decision. They exhibit the high-field Ar-H (the 8-H that is next to the heterocyclic oxygen atom) as slightly broadened singlet signal (with unresolved 4J or 5J long-range couplings below 1.8 Hz or smaller), and a 6-H/5-H AB system (the 6-H signal with more long-range broadening), in addition to the deeper field Ar-H signals of the 4-substituent (AA'BB' for **6b** and **6c**). Thus, the structure **6** (reaction at C-4 of resorcinol as proposed in ref. 67) is confirmed, and the suggestion in ref. 68 (claiming reaction at C-2 of resorcinol) is unsupported. Importantly, our ^1H NMR spectra of **6a–6c** concur with the ones of ref. 13 also with the singlet signal from 8-H without vicinal coupling partner. These products were synthesized in solution with a “tungstic acid functionalized mesoporous catalyst”. Furthermore, the melting points of the corresponding products from refluxing water with potassium phthalimido-*N*-hydroxylate³ and those in ref. 76 and 77 correspond with ours. Thus, the structure **6** is found independent of the reaction conditions.

The remaining 7-OH group of **6a–6c** was not able to undergo the Michael addition with excess arylidenemalononitriles under our reaction conditions. The electronic conditions would allow for it, but the acidity of 7-OH in the compounds **6** is apparently not high enough. Stronger conditions should be helpful for the corresponding second Michael addition.

The advantage of the present enviro-economic methodology may for example be demonstrated by comparison of the efficiencies for the syntheses of compound **4e** in various previous publications (Table 2). The reported results clearly show that the present protocol at room temperature and with short reaction times is the only one giving quantitative yields and not requiring the use of organic solvents *etc.* for workup. The previous catalytic techniques sometimes at elevated temperatures and with long reaction times exhibit lower and never quantitative yields. They require costly purification workup with serious waste formation by removal/recovery of catalysts and organic non-product materials in organic solvents. It is also noteworthy that our easily separated catalyst, Na_2CO_3 , is the cheapest and greenest one among those being compared. Our work-up procedure is extremely simple. The obtained solid product powder is quantitatively flushed out with water from the milling vessel, filtered and dried for quantitatively isolated pure product yields and a soda solution. No purification workup was necessary. These solvent-free syntheses are thus enviro-

Table 2 Comparison of the present method with some reported methods for the preparation of **4e**

Catalyst	Temp.	Time	Catalyst loading	Yield (%)
This work (Na_2CO_3)	rt	10 min	10 mol%	100
$\text{Re}(\text{PFO})_3$ (ref. 23)	60 °C	4 h	5 mol%	90
Diammonium hydrogen phosphate ²⁵	rt	0.5–2 h	10 mol%	78
NMe_4OH (ref. 26)	rt	0.5–2 h	10 mol%	80
CeCl_3 (ref. 30)	Reflux	1 h	10 mol%	93
ZnO-beta zeolite ³¹	Reflux	50 min	100 mg	87
DABCO ³²	Reflux	2 h	10 mol%	94
Silica gel-supported polyphosphoric acid ³³	Reflux	8 min	100 mg	85
Caro's acid-silica gel ³⁵	Reflux	15 min	16 mol%	93
D-L-Proline ²⁴	rt	>30 min	20 mol%	92

economic. When desired, the powders can also be directly sublimed off from the catalyst.

Discussion

The selectivity of the milled solid catalyst Na_2CO_3 between different phenols is remarkable. The reactions go to completion despite the increasing amounts of water of crystallization from the Knoevenagel condensation. Since the solid decahydrate of soda (m.p. 34 °C) might have easily melted in its environment, it is almost certain that this hydration state was not reached and that at least part of the water of reaction was also included in the solid products with many functional groups. While resorcinol and the naphthols react smoothly, phenol and the phenolic groups in **4j** or **6a–6c** do not react with the Knoevenagel reagents **9**. This may be roughly related to the well-known pK_a values for aqueous solution even though the acidities in the present medium would be different. Thus, the reactive phenols such as resorcinol ($\text{pK}_{a1} = 9.15$), α -naphthol ($\text{pK}_a = 9.3$), and β -naphthol ($\text{pK}_a = 9.57$) are considerably stronger acids than the weaker alkyl- and alkoxyphenols. We may refer here to phenol ($\text{pK}_a = 9.99$) and reference compounds such as 4-methylphenol ($\text{pK}_a = 10.26$) and 2-methoxyphenol ($\text{pK}_a = 9.98$). It thus appears that our solid catalyst cannot deprotonate the weaker phenols **4j** and **6a–6c** and this would explain its selection between phenols.

Also the exclusive Michael addition of β -naphthol at C-1 is highly remarkable, because reaction at C-3 would be facilitated by less steric crowding. It appears, however, that the much higher electron density of the naphthol and of the naphtholate at C-1 is the prevailing factor. Density functional calculations at the B3LYP 6-31G* level calculate the electrostatic charges at C-1 as -0.505 and at C-3 as -0.224 for β -naphthol or as -0.605 at C-1 and as -0.326 at C-3 for the β -naphtholate anion. The large electrostatic differences substantiate the experimentally found orientation specificities to C-1 but not to C-3 of β -naphthol in its reactions with the Knoevenagel intermediate **9**. Kumar's group unduly claimed the latter choice without analysis of their ^1H NMR data.^{59–61}

To exclude the possibility that their experimental technique would have drastically changed the orientation specificity of β -naphthol, the reported melting points in ref. 60–62 are invoked. The now challenged “2-amino-4-aryl-3-cyano-4*H*-benzo[*g*]

chromenes” are reported with melting points of 274, 187, 191, and 260 °C for Ar = 4-H, 4- NO_2 , 4- CH_3O , and 2-Cl-phenyl.^{59–61} These values concur with those of the ^1H NMR spectroscopically validated 2-amino-4-aryl-3-cyano-4*H*-benzo[*g*]chromenes. Furthermore, the melting points correspond with those of all other published -4*H*-benzo[*g*]chromenes that consistently exhibit the splitting of the highest field Ar-H signal in the ^1H NMR spectrum from the cited authors in Table 1 and also in Table 2 of ref. 2. Thus β -naphthol reacts at C-1 but not at C-3 independent of the experimental conditions, because the electron density is almost twice as high at the experimentally chosen position.


Conclusions and outlook

We have demonstrated a “green” atom-economic and environmental method for the solid-state preparation of several 2-amino-4-aryl-3-cyano-4*H*-chromenes and clarified still existing detrimental controversies concerning the product structures. The solvent-free technique with the eco-efficient catalyst is superior to the less productive solution techniques that use expensive and harmful catalysts, solid supports, ultrasound or microwave, which require purifying workup by using organic solvents for removal of the support, of unreacted reagents, and of side products. The here reported milling protocol without heating has the advantages of operational simplicity, mild reaction conditions, quantitative product yields, and no need of organic solvents for workup and purification. The soda catalyst is cheap and non-toxic. These benefits make our methodology a valid contribution to the existing processes in the field of 2-amino-4-aryl-3-cyano-4*H*-chromene derivative synthesis.

Experimental section

Material and methods

The ball mill was a Retsch MM 400 swing mill. A 10 mL stainless steel vessel was used. Two stainless steel balls with 7 mm diameter were used, and the milling frequency was 30–35 Hz. The temperature within the milling beaker was close to room temperature of about 25 °C. It never reached or exceeded 30 °C by the milling, but malononitrile (m.p. 32 °C) melts in the presence of the other reactants, so this is actually kneading.



Melting points were determined with a Barnstead electro-thermal melting point apparatus and are uncorrected. Infrared (IR) spectra were recorded with a Shimadzu 8400s FT-IR spectrometer using potassium bromide pellets. 500 MHz ^1H NMR spectra were recorded with a Bruker DRX-500 Avance spectrometer and 250 MHz spectra with a Bruker Avance DPX-250. The chemical shifts are reported in ppm (δ -scale) relative to internal TMS and coupling constants are reported in DMSO- d_6 . Density functional calculations were performed with the TITAN program of Wave Function Inc., Irvine, USA and Schrödinger Inc., Portland, USA. All solid reagents were purchased from commercial sources and used without further purification when uniformity was tested by TLC (silica gel with ethyl acetate and hexane in a 1 : 5 ratio). The liquid aldehydes were distilled prior to use. All products are previously described compounds and they were identified and structurally characterized. Their physical and spectral data corresponded with the cited literature data. Product purity was checked by melting point, by TLC on silica gel with ethyl acetate and hexane in a 1 : 5 ratio, and by high-resolution ^1H NMR spectra.

Typical procedure for the synthesis of the 4H-chromenes

In a typical experiment, a stoichiometric mixture of an aromatic aldehyde (**1**) (1.0 mmol), malononitrile (**2**) (66 mg, 1.0 mmol), dimedone (**3a**) (140 mg, 1.0 mmol) or **3b–3e** (112 mg, 110 mg, 144 mg, and 144 mg, respectively, 1.0 mmol each) and anhydrous sodium carbonate (11 mg, 0.1 mmol) were ball-milled for the appropriate time (Table 1). When the reaction was completed, as indicated by the FTIR spectrum of the solid in KBr pellets and TLC to detect uniformity, the reaction product (from repeat runs without taking material for IR and TLC) was scraped out on a filter and the holdup flushed out with cold washing water (3 times, 5 mL each) that was shaken in the mill for about 1 min prior to pouring on the solid product for extracting the inorganic catalyst. After drying the product from the filter in a vacuum, the pure product was directly obtained with no need for extra purification. The products could also be quantitatively obtained in pure form by complete collection from the ball-milling vessel (using the holdup from a previous run) and sublimation in a good vacuum in the temperature range of 150–190 °C. FTIR spectra of the washed or recrystallized or sublimed products corresponded to the raw spectra still containing the catalyst.

Representative spectral data

2-Amino-7,7-dimethyl-4-(4-nitrophenyl)-5-oxo-5,6,7,8-tetrahydro-4H-chromene-3-carbonitrile (4e). IR (KBr), $\nu(\text{cm}^{-1})$: 3406, 3325, 3182, 3080, 2964, 2190, 1681, 1519, 1367; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 0.96 (s, 3H), 1.04 (s, 3H), 2.11 (AB, $J = 16.0$ Hz, 1H), 2.25 (BA, $J = 16.0$ Hz, 1H), 2.50 (s, 2H), 4.36 (s, 1H), 7.17 (s, 2H, NH_2), 7.45 (AA'BB', " J " = 8.2 Hz, 2H), 8.17 (BB'AA', " J " = 8.2 Hz, 2H); ^{13}C NMR (125 MHz, DMSO- d_6): 27.79, 29.11, 32.66, 36.56, 50.74, 56.91, 57.91, 112.63, 120.17 (2C), 124.51, 129.49 (2C), 147.14, 153.15, 159.48, 163.96, 196.53.

2-Amino-7,7-dimethyl-4-(4-methoxyphenyl)-5-oxo-5,6,7,8-tetrahydro-4H-chromene-3-carbonitrile (4i). IR (KBr), $\nu(\text{cm}^{-1})$:

3375, 3321, 3186, 2962, 2192, 1683, 1654, 1510, 1369; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 0.95 (s, 3H), 1.04 (s, 3H), 2.09 (AB, $J = 16$ Hz, 1H), 2.23 (BA, $J = 16$ Hz, 1H), 2.50 (s, 2H), 3.72 (s, 3H), 4.12 (s, 1H), 6.84 (AA'BB', " J " = 8.05 Hz, 2H), 6.94 (s, 2H, NH_2), 7.05 (BB'AA', " J " = 8.05 Hz, 2H); ^{13}C NMR (125 MHz, DMSO- d_6) δ (ppm): 27.65, 29.28, 32.64, 35.63, 50.70, 50.89, 55.87, 59.46, 113.87, 114.55 (2C), 120.65, 129.31 (2C), 137.72, 158.79, 159.29, 162.99, 196.51.

2-Amino-7-hydroxy-4-phenyl-4H-chromene-3-carbonitrile (6a). IR (KBr), $\nu(\text{cm}^{-1})$: 3495, 3425, 3337, 2189, 1649, 1589, 1504, 1456, 1404, 1153; ^1H NMR (DMSO- d_6 , 250 MHz), δ (ppm): 4.64 (s, 4-H), 6.43 (s, 8-H), 6.49 (AB, $J = 8.25$ Hz, 6-H), 6.83 (BA, $J = 8.25$ Hz, 5-H), 6.91 (s, 2NH), 7.17–7.25 (m, 3ArH), 7.30–7.36 (m, 2ArH), 9.74 (br, s, 1OH).

2-Amino-4-(4-chlorophenyl)-7-hydroxy-4H-chromene-3-carbonitrile (6b). IR (KBr), $\nu(\text{cm}^{-1})$: 3470, 3340, 3250, 2191, 1641, 1587, 1502, 1460, 1406, 1155; ^1H NMR (DMSO- d_6 , 250 MHz), δ (ppm): 4.67 (s, 4-H), 6.41 (s, 8-H), 6.48 (AB, $J = 7.5$ Hz, 6-H), 6.78 (BA, $J = 7.5$ Hz, 5-H), 6.94 (s, 2NH), 7.19 (AA'BB', " J " = 7.0 Hz, 2ArH), 7.37 (BB'AA', " J " = 7.0 Hz, 2ArH), 9.89 (br, s, 1OH).

2-Amino-4-(4-bromophenyl)-7-hydroxy-4H-chromene-3-carbonitrile (6c). IR (KBr), $\nu(\text{cm}^{-1})$: 3472, 3340, 3259, 2189, 1641, 1585, 1502, 1460, 1404, 1155; ^1H NMR (DMSO- d_6 , 250 MHz), δ (ppm): 4.66 (s, 4-H), 6.41 (s, 8-H), 6.49 (AB, $J = 8.25$ Hz, 6-H), 6.79 (BA, $J = 8.25$ Hz, 5-H), 6.95 (s, 2NH), 7.13 (AA'BB', " J " = 8.2 Hz, 2ArH), 7.51 (BB'AA', $J = 8.2$ Hz, 2ArH), 9.76 (br, s, 1OH).

2-Amino-4-(4-chlorophenyl)-4H-benzo[h]chromene-3-carbonitrile (7a). IR (KBr), $\nu(\text{cm}^{-1})$: 3452, 3334, 2190, 1666, 1590, 1409; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 4.96 (s, 1H, CH), 7.10 (AB, 1H, " J " = 8.5 Hz ArH), 7.20 (s, 2H, NH_2), 7.28 (AA'BB', 2H, " J " = 8.4 Hz, ArH), 7.38 (BB'AA', 2H, " J " = 8.4 Hz, ArH), 7.57–7.66 (m, 3H, ArH), 7.90 (d, 1H, $J = 8.0$ Hz, ArH), 8.24 (BA, 1H, " J " = 8.5 Hz, ArH).

2-Amino-4-(4-bromophenyl)-4H-benzo[h]chromene-3-carbonitrile (7b). IR (KBr), $\nu(\text{cm}^{-1})$: 3452, 3337, 2189, 1664, 1600, 1570, 1408; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 4.94 (s, 1H, CH), 7.10 (AB, 1H, $J = 7.9$ Hz, ArH), 7.2–7.23 (m, 4H, NH_2 + ArH), 7.52 (AA'BB', 2H, " J " = 7.3 Hz, ArH), 7.59–7.65 (m, 3H, ArH), 7.90 (d, 1H, " J " = 7.7 Hz, ArH), 8.24 (BA, 1H, $J = 7.9$ Hz, ArH); ^{13}C NMR (DMSO- d_6 , 125 MHz): 39.84, 56.62, 118.18, 120.95, 121.24, 121.58, 123.59, 124.89, 126.94, 127.62, 127.75, 128.57, 130.82 (2C), 132.48 (2C), 133.62, 143.61, 145.95, 161.02.

2-Amino-4-(4-chlorophenyl)-4H-benzo[f]chromene-3-carbonitrile (8a). IR (KBr), $\nu(\text{cm}^{-1})$: 3452, 3334, 2190, 1666, 1590, 1409; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 4.95 (s, 1H, CH), 7.09 (AB, 1H, $J = 8.5$ Hz ArH), 7.19 (s, 2H, NH_2), 7.27 (AA'BB', 2H, " J " = 8.35 Hz, ArH), 7.36 (BB'AA', 2H, " J " = 8.35 Hz, ArH), 7.57–7.66 (m, 3H, ArH), 7.89 (br, d, 1H, $J = 8.0$ Hz, ArH), 8.24 (BA, 1H, $J = 8.5$ Hz, ArH).

2-Amino-4-(3-nitrophenyl)-4H-benzo[f]chromene-3-carbonitrile (8b). IR (KBr), $\nu(\text{cm}^{-1})$: 3462, 3357, 3206, 2190, 1658, 1589, 1410; ^1H NMR (DMSO- d_6 , 500 MHz), δ (ppm): 5.63 (s, 1H, CH), 7.17 (s, 2H, NH_2), 7.41 (AB, 1H, " J " = 8.9 Hz, ArH), 7.42–7.48 (m, 2H, ArH), 7.57–7.60 (m, 1H, ArH), 7.67 (d, 1H, " J " = 7.7 Hz, ArH), 7.87 (d, 1H, " J " = 8.2 Hz, ArH), 7.94 (d, 1H, " J " = 7.6 Hz, ArH),



7.99 (d, 1H, “J” = 7.9 Hz, ArH), 8.04 (d, 1H, “J” = 8.1 Hz, ArH), 8.08 (s, 1H, ArH); ¹³C NMR (DMSO-*d*₆, 125 MHz): 39.87, 57.82 (=C–CN), 115.47, 117.74, 121.04, 122.18, 122.72, 124.36, 128.28, 129.48, 130.78, 130.95, 131.33 (2C), 131.73 (2C), 134.59, 148.61, 148.86, 160.84 (=C–O, –N).

2-Amino-4-(4-hydroxyphenyl)-4H-benzof[chromene-3-carbonitrile (8c). IR (KBr), $\nu(\text{cm}^{-1})$: 3452, 3308, 3175, 2195, 1653, 1589, 1510, 1412; ¹H NMR (DMSO-*d*₆, 500 MHz), δ (ppm): 5.17 (s, 1H, CH), 6.63 (d, 2H, *J* = 7.8 Hz, AA'BB'), 6.89 (s, 2H, NH₂), 6.98 (d, 2H, *J* = 7.8 Hz, BB'AA'), 7.32 (d, 1H, “J” = 8.8 Hz, ArH), 7.40–7.46 (m, 2H, ArH), 7.85 (br, d, “J” = 7.7 Hz, 1H, ArH), 7.905–7.92 (br, d, 2H, ArH), 9.26 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆, 125 MHz): 40.89, 59.28 (=C–CN), 116.21 (2C), 117.04, 117.65, 121.51, 124.59, 125.71, 127.81, 128.88 (2C), 129.28, 130.10, 131.09, 131.68, 137.11, 147.54, 156.82, 160.38 (=C–O, –N).


The ESI⁺ contains typical prints of the recorded NMR and IR spectra. It can be accessed free of charge.

Acknowledgements

The financial support from The Research Council of Iran University of Science and Technology (IUST), Tehran, Iran is gratefully acknowledged.

Notes and references

- G. Kaupp, M. R. Naimi-Jamal and J. Schmeyer, *Tetrahedron*, 2003, **59**, 3753–3760.
- M. R. Naimi-Jamal, S. Mashkouri and A. Sharifi, *Mol. Diversity*, 2010, **14**, 473–477.
- M. G. Dekamin, M. Eslami and A. Maleki, *Tetrahedron*, 2013, **69**, 1074–1085.
- L. Bonsignore, G. Loy, D. Secci and A. Calignano, *Eur. J. Med. Chem.*, 1993, **28**, 517–520.
- Y. Gao, W. Yang and D.-M. Du, *Tetrahedron: Asymmetry*, 2012, **23**, 339–344.
- R. Ling, M. Yoshida and P. S. Mariano, *J. Org. Chem.*, 1996, **61**, 4439–4449.
- L. R. Morgan, B. S. Jursic, C. L. Hooper, D. M. Neumann, K. Thangaraj and B. LeBlanc, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 3407–3411.
- A. Bolognese, G. Correale, M. Manfra, A. Lavecchia, O. Mazzoni, E. Novellino, P. La Colla, G. Sanna and R. Loddo, *J. Med. Chem.*, 2004, **47**, 849–858.
- H. Gourdeau, L. Leblond, B. Hamelin, C. Desputeau, K. Dong, I. Kianicka, D. Custeau, C. Boudreau, L. Geerts, S.-X. Cai, J. Drewe, D. Labrecque, S. Kasibhatla and B. Tseng, *Mol. Cancer Ther.*, 2004, **3**, 1375–1384.
- N. Fokialakis, P. Magiatis, I. Chinou, S. Mitaku and F. Tillequin, *Chem. Pharm. Bull.*, 2002, **50**, 413–414.
- A. Afantitis, G. Melagraki, H. Sarimveis, P. A. Koutentis, J. Markopoulos and O. Iggleksi-Markopoulou, *Bioorg. Med. Chem.*, 2006, **14**, 6686–6694.
- M. H. Fatemi and S. Gharaghani, *Bioorg. Med. Chem.*, 2007, **15**, 7746–7754.
- S. K. Kundu, J. Mondal and A. Bhaumik, *Dalton Trans.*, 2013, **42**, 10515–10524.
- C. S. Konkoy, D. B. Fick, S. X. Cai, N. C. Lan and J. F. W. Keana, *Pct Int. Appl.*, WO 0075123A1, 2000 p. 1214; *Chem. Abstr.* 2001, 134, 29313a.
- M. Kidwai, S. Saxena, M. K. Rahman Khan and S. S. Thukral, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 4295–4298.
- N. Martin, C. Pascual, C. Seoane and J. Soto, *Heterocycles*, 1987, **26**, 2811–2816.
- A.-F. A. Harb, A.-H. M. Hesien, S. A. Metwally and M. H. Elnagdi, *Liebigs Ann. Chem.*, 1989, 585–588.
- S. E. Zayed, E. I. Abou Elmaged, S. A. Metwally and M. H. Elnagdi, *Collect. Czech. Chem. Commun.*, 1991, **56**, 2175–2182.
- M. H. Elnagdi, R. M. Abdel-Motaleb, M. Mustafa, M. F. Zayed and E. M. Kamel, *J. Heterocycl. Chem.*, 1987, **24**, 1677–1681.
- T. Jin, A. Wang, J. Zhang, F. Zhang and T. Li, *Chin. J. Org. Chem.*, 2004, **24**, 1598–1600.
- L.-M. Wang, N. Jiao, J. Qiu, J.-J. Yu, J.-Q. Liu, F.-L. Guo and Y. Liu, *Tetrahedron*, 2010, **66**, 339–343.
- X.-S. Wang, D.-Q. Shi, S.-J. Tu and C.-S. Yao, *Synth. Commun.*, 2003, **33**, 119–126.
- L.-M. Wang, J.-H. Shao, H. Tian, Y.-H. Wang and B. Liu, *J. Fluorine Chem.*, 2006, **127**, 97–100.
- S. Balalaie, M. Bararjanian, A. M. Amani and B. Movassagh, *Synlett*, 2006, 263–266.
- S. Balalaie, M. Bararjanian, M. Sheikh-Ahmadi, S. Hekmat and P. Salehi, *Synth. Commun.*, 2007, **37**, 1097–1108.
- S. Balalaie, M. Sheikh-Ahmadi and M. Bararjanian, *Catal. Commun.*, 2007, **8**, 1724–1728.
- D. Kumar, V. B. Reddy, B. G. Mishra, R. K. Rana, M. N. Nadagouda and R. S. Varma, *Tetrahedron*, 2007, **63**, 3093–3097.
- M. Seifi and H. Sheibani, *Catal. Lett.*, 2008, **126**, 275–279.
- S. Gao, C. Tseng, C. H. Tsai and C.-F. Yao, *Tetrahedron*, 2008, **64**, 1955–1961.
- G. Sabitha, K. Arundhathi, K. Sudhakar, B. Sastry and J. Yadav, *Synth. Commun.*, 2009, **39**, 2843–2851.
- S. S. Katkar, M. K. Lande, B. R. Arbad and S. T. Gaikwad, *Chin. J. Chem.*, 2011, **29**, 199–202.
- D. Tahmassebi, J. A. Bryson and S. I. Binz, *Synth. Commun.*, 2011, **41**, 2701–2711.
- A. Davoodnia, S. Allameh, S. Fazli and N. Tavakoli-Hoseini, *Chem. Pap.*, 2011, **65**, 714–720.
- S. Rostamnia, A. Nuri, H. Xin, A. Pourjavadi and S. H. Hosseini, *Tetrahedron Lett.*, 2013, **54**, 3344–3347.
- H. A. Oskooie, M. M. Heravi, N. Karimi and M. E. Zadeh, *Synth. Commun.*, 2011, **41**, 436–440.
- P. P. Salvi, A. M. Mandhare, A. S. Sartape, D. K. Pawar, S. H. Han and S. S. Kolekar, *C. R. Chim.*, 2011, **14**, 878–882.
- K. Singh, J. Singh and H. Singh, *Tetrahedron*, 1996, **52**, 14273–14280.
- M. Ghashang, S. S. Mansoor and K. Aswin, *Res. Chem. Intermed.*, 2013, 1–17, DOI: 10.1007/s11164-013-1419-2.
- S. R. Kale, S. S. Kahandal, A. S. Burange, M. B. Gawande and R. V. Jayaram, *Catal. Sci. Technol.*, 2013, **3**, 2050–2056.
- R.-Y. Guo, Z.-M. An, L.-P. Mo, R.-Z. Wang, H.-X. Liu, S.-X. Wang and Z.-H. Zhang, *ACS Comb. Sci.*, 2013, **15**, 557–563.



- 41 J. Zhou, S. Tu, Y. Gao and M. Ji, *Chin. J. Org. Chem.*, 2001, **21**, 742–745.
- 42 S. Tu, H. Jiang, Q. Zhuang, C. Miao, D. Shi, X. Wang and Y. Gao, *Chin. J. Org. Chem.*, 2003, **23**, 488–490.
- 43 J. Mokhtari, M. R. Naimi-Jamal, H. Hamzeali, M. G. Dekamin and G. Kaupp, *ChemSusChem*, 2009, **2**, 248–254.
- 44 S. Mashkouri and M. R. Naimi-Jamal, *Molecules*, 2009, **14**, 474–479.
- 45 M. R. Naimi-Jamal, M. Mirzaei, M. Bolourtchian and A. Sharifi, *Synth. Commun.*, 2006, **36**, 2711–2717.
- 46 A. Sharifi, M. Mirzaei and M. R. Naimi-Jamal, *J. Chem. Res.*, 2007, 129–132.
- 47 A. Sharifi, M. Mirzaei and M. R. Naimi-Jamal, *Synth. Commun.*, 2005, **35**, 1039–1044.
- 48 G. Kaupp, *CrystEngComm*, 2006, **8**, 794–804.
- 49 G. Kaupp, *CrystEngComm*, 2011, **13**, 3108–3121.
- 50 G. Kaupp, M. R. Naimi-Jamal and J. Schmeyers, *Chem.–Eur. J.*, 2002, **8**, 594–600.
- 51 G. Kaupp, in *Encyclopedia of Life Support Systems (EOLSS), developed under the Auspices of the UNESCO*, EOLSS Publishers, Oxford, UK, 2012, pp. 38, <http://www.eolss.net>.
- 52 G. Kaupp, in *Kirk-Othmer Encyclopedia of Chemical Technology*, John Wiley & Sons, Inc., 2012, DOI: 10.1002/0471238961.solvkaup.a01.
- 53 A. Stolle, T. Szuppa, S. E. S. Leonhardt and B. Ondruschka, *Chem. Soc. Rev.*, 2011, **40**, 2317–2329.
- 54 G.-W. Wang, *Chem. Soc. Rev.*, 2013, **42**, 7668–7700.
- 55 G. Kaupp and M. R. Naimi-Jamal, *Eur. J. Org. Chem.*, 2002, 1368–1373.
- 56 G. Kaupp, M. R. Naimi-Jamal and V. Stepanenko, *Chem.–Eur. J.*, 2003, **9**, 4156–4161.
- 57 R. Trotzki, M. M. Hoffmann and B. Ondruschka, *Green Chem.*, 2008, **10**, 767–772.
- 58 D. Zhou, Z. Ren, J. Chen, W. Cao and H. Deng, *J. Heterocycl. Chem.*, 2008, **45**, 1865–1867.
- 59 B. S. Kumar, N. Srinivasulu, R. H. Udipi, B. Rajitha, Y. T. Reddy, P. N. Reddy and P. S. Kumar, *Russ. J. Org. Chem.*, 2006, **42**, 1813–1815.
- 60 S. K. Kumar, M. S. Khalid and R. H. Udipi, *Asian J. Chem.*, 2010, **22**, 2685–2688.
- 61 B. S. Kumar, N. Srinivasulu, R. H. Udipi, B. Rajitha, Y. T. Reddy, P. N. Reddy and P. S. Kumar, *J. Heterocycl. Chem.*, 2006, **43**, 1691–1693.
- 62 M. A. Pasha and V. Jayashankara, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 2007, **46**, 1328–1331.
- 63 T.-S. Jin, J.-C. Xiao, S.-J. Wang, T.-S. Li and X.-R. Song, *Synlett*, 2003, **2003**, 2001–2004.
- 64 M. Akkurt, A. R. Kennedy, S. K. Mohamed, S. H. H. Younes and G. J. Miller, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2013, **69**, o401.
- 65 P. Das, A. Dutta, A. Bhaumik and C. Mukhopadhyay, *Green Chem.*, 2014, **16**, 1426–1435.
- 66 X.-S. Wang, G.-S. Yang and G. Zhao, *Tetrahedron: Asymmetry*, 2008, **19**, 709–714.
- 67 V. J. Ram and M. Verma, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 1994, **33**, 908–911.
- 68 F. F. Abdel-Latif, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 1990, **29**, 664–666.
- 69 S. B. Guo, S. X. Wang and J. T. Li, *Synth. Commun.*, 2007, **37**, 2111–2120.
- 70 A. T. Khan, M. Lal, S. Ali and M. M. Khan, *Tetrahedron Lett.*, 2011, **52**, 5327–5332.
- 71 C. Revanna, T. Swaroop, G. Raghavendra, D. Bhadregowda, K. Mantelingu and K. Rangappa, *J. Heterocycl. Chem.*, 2012, **49**, 851–855.
- 72 J.-C. Xu, W.-M. Li, H. Zheng, Y.-F. Lai and P.-F. Zhang, *Tetrahedron*, 2011, **67**, 9582–9587.
- 73 G. Zhang, Y. Zhang, J. Yan, R. Chen, S. Wang, Y. Ma and R. Wang, *J. Org. Chem.*, 2011, **77**, 878–888.
- 74 H. R. Safaei, M. Shekouhy, S. Rahmanpur and A. Shirinfeshan, *Green Chem.*, 2012, **14**, 1696–1704.
- 75 S. Nemouchi, R. Boulcina, B. Carboni and A. Debache, *C. R. Chim.*, 2012, **15**, 394–397.
- 76 A. M. El-Agrody, A. H. Bedair, T. C. Corkery, A. Ata and A. S. Abd-El-Aziz, *Heterocycles*, 2004, **63**, 1793–1812.
- 77 G. Klokol, L. Sharanina, V. Nesterov, V. Shklover, Y. Sharanin and Y. Struchkov, *Zh. Org. Khim.*, 1987, **23**, 412–421.

