


Photochemical & Photobiological Sciences

An international journal

rsc.li/pps

Volume 19
Number 5
May 2020
Pages 533-738



ISSN 1474-905X



Cite this: *Photochem. Photobiol. Sci.*, 2020, **19**, 585

Revisiting cyanobacterial state transitions

Pablo I. Calzadilla and Diana Kirilovsky *

Photosynthetic organisms are exposed to a fluctuating environment in which light intensity and quality change continuously. Specific illumination of either photosystem (PSI or PSII) creates an energy imbalance, leading to the reduction or oxidation of the intersystem electron transport chain. This redox imbalance could trigger the formation of dangerous reactive oxygen species. Cyanobacteria, like plants and algae, have developed a mechanism to re-balance this preferential excitation of either reaction center, called state transitions. State transitions are triggered by changes in the redox state of the membrane-soluble plastoquinone (PQ) pool. In plants and green algae, these changes in redox potential are sensed by Cytochrome *b₆f*, which interacts with a specific kinase that triggers the movement of the main PSII antenna (the light-harvesting complex II). By contrast, although cyanobacterial state transitions have been studied extensively, there is still no agreement about the molecular mechanism, the PQ redox state sensor and the signaling pathways involved. In this review, we aimed to critically evaluate the results published on cyanobacterial state transitions, and discuss the "new" and "old" models in the subject. The phycobilisome and membrane contributions to this physiological process were addressed and the current hypotheses regarding its signaling transduction pathway were discussed.

Received 18th November 2019,
Accepted 20th February 2020

DOI: 10.1039/c9pp00451c


rsc.li/pps

1. Introduction

Light is essential for photosynthetic organisms that harvest solar energy and convert it into chemical energy (ATP) and

reducing power (NADPH). These molecules are used in the assimilation of CO₂ and synthesis of organic carbon molecules, in a process called photosynthesis, in which molecular oxygen is produced as a sub-product. However, light can become dangerous in certain cases. For example, when the light energy reaching the photosystems exceeds the energy consumption by multiple cellular processes, the entire photo-

Université Paris-Saclay, CNRS, CEA, Institute for Integrative Biology of the Cell (I2BC), 91198 Gif sur Yvette, France. E-mail: diana.kirilovsky@cea.fr



Pablo I. Calzadilla

Pablo Calzadilla was born in Quilmes (Argentina) and obtained his Ph.D. in Biological Sciences at the University of Buenos Aires (Argentina). He did his first post-doc in the Laboratory of Regulatory Mechanisms in Photosynthetic Organisms (I2BC, CNRS) in Gif sur Yvette, France, headed by Dr Diana Kirilovsky. Presently, he is a Research Associate in Dr Giles Johnson's lab in the University of Manchester

(Manchester, UK). His main research interests are how photosynthetic organisms acclimates their photosynthetic capacity to cope with a changing environment. He has worked in cyanobacteria and different plant species (models and crops) along his short career.



Diana Kirilovsky

Diana Kirilovsky was born in Buenos Aires (Argentina) and obtained her Ph.D. in Biochemistry at the Hebrew University of Jerusalem (Israel). Presently, she is a Research Director (DR1, CNRS) and Group Leader in the Institute of Integral Biology of the Cell (I2BC, CNRS) in Gif sur Yvette, France. Diana Kirilovsky has studied the molecular biology and physiology of cyanobacteria for over 30 years focusing on light induced pro-

cesses, in particular, the role of light as a source of stress and as a regulator. She has principally studied the mechanisms of photoinhibition, photoprotection and light acclimation with more of 100 publications.



synthetic chain is reduced. Under these conditions, secondary reactions are induced, and the accumulation of high energy reactive intermediates is promoted, subsequently leading to the production of singlet oxygen and oxygen radicals (ROS). These molecules cause severe cell damage, especially at the level of the two photosystems: photosystem I (PSI) and photosystem II (PSII). In addition, changes in light quality can imbalance the photosystems' activity, decreasing their photosynthetic efficiency, and/or altering the redox state of the electron transport chain, which again leads to ROS production.

In order to efficiently use the incoming light and to avoid damage, sensing and adapting to environmental changes becomes crucial. Thus, photo-protective mechanisms were evolutionarily developed by photosynthetic organisms. These mechanisms induce a rapid reorganization of the photosynthetic apparatus, changing the interaction between the antennae and reaction centers, and adjusting light absorption and utilization. Particularly in cyanobacteria, two photo-protective mechanisms, the Orange Carotenoid Protein (OCP)-related Non-Photochemical-Quenching (NPQ) (see reviews¹⁻⁵) and state transitions (see reviews⁶⁻⁸), regulate the transfer of energy arriving at the photochemical reaction centers, sensing the intensity and the quality of light. While OCP-related NPQ simultaneously decreases the energy arriving at both the photosystems under high light conditions,^{9,10} state transitions decrease (or increase) the activity of only one of the two photosystems under fluctuating light quality.

The state transitions mechanism was first described fifty years ago in green and red algae.^{11,12} Two states were characterized, one induced by light preferentially absorbed by PSII (State II), and the other triggered by light preferentially absorbed by PSI (State I). Specific illumination of PSI or PSII creates an imbalance in the electron transport chain, rendering the plastoquinone (PQ) pool more oxidized or more reduced. This effect is rebalanced by state transitions. Each state differs in its fluorescence characteristics, *i.e.*, in State II, the PSII to PSI fluorescence ratio is low, while in State I this ratio is higher.

During the last few decades, state transitions have been well characterized in plants and green algae (see review¹³). In these organisms, the differences in the fluorescence of PSI and PSII are related to a partial migration of the Light-Harvesting Complex II (LHCII), which is the main membrane chlorophyll-carotenoid PSII antenna. This migration is triggered by redox changes in the PQ pool, in a process involving Cytochrome *b₆f* (Cyt *b₆f*) and phosphorylation of LHCII (see reviews¹³⁻¹⁵). When the PQ pool gets reduced, due to preferential illumination of PSII, a specific kinase (STN7/STT^{16,17}) is activated by the binding of plastoquinol (PQH₂) to the Cyt *b₆f* Qo site.¹⁸⁻²⁰ This binding leads to conformational changes in the Cyt *b₆f*, particularly in the Rieske protein,²⁰⁻²³ and in specific amino acids on the peripheral stromal structure of subunit IV which are in contact with the kinase.²⁴ The activated kinase phosphorylates the mobile trimers of LHCII, resulting in their detachment from PSII and attachment to PSI, triggering the transition from State I to State II.^{25,26} Conversely, when the PQ

pool gets oxidized, in darkness or due to preferential absorption of light by PSI, the kinase is deactivated, and a specific phosphatase (TAP38/PPH1^{27,28}) dephosphorylates LHCII, which migrates back to PSII, leading to State I transition. It must be noted that in plants and green algae, Cyt *b₆f* is the redox sensor of the PQ pool, and in its absence, state transitions are inhibited.²⁹

Cyanobacteria do not contain LHCII; instead, the PSII antenna is a huge membrane-extrinsic complex, the phycobilisome (PBS), which is bound to the stromal surface of the thylakoid membranes. The PBS is constituted of water soluble phycobiliproteins (which bind blue, violet and red bilins) and linker proteins. In most cyanobacteria, these proteins are organized as rods radiating from a central core (see reviews³⁰⁻³⁴) (Fig. 1). Each rod is formed by hexamers of phycocyanin (PC, $\lambda_{\text{max}} = 620$ nm) and, in some cases, phycoerythrin (PE, $\lambda_{\text{max}} = 577$ nm) and phycoerythrocyanin (PEC, $\lambda_{\text{max}} = 625$ nm).^{31,35} When these last two phycobiliproteins are present, PC is localized in the core proximal hexamers, and PEC and PE form the distal hexamers.³¹

The core structure varies between the different cyanobacterial strains, and can be formed by two (*Synechococcus elongatus* PCC 7942, hereafter *S. elongatus*), three (*Synechocystis* PCC 6803, hereafter *Synechocystis*), or five cylinders (*Nostoc* PCC 7120, *Thermosynechococcus elongatus*) of allophycocyanin (APC). Each cylinder is made of 4 APC trimers, composed of α APC- β APC heterodimers, which absorb at 650 nm and emit at 660 nm. Within each basal cylinder, one α APC subunit in an external trimer is replaced by an α APC-like subunit, called ApcD, and one β APC in an internal trimer is replaced by a β APC-like subunit, called ApcF. Adjacent to ApcF, another APC dimer is formed by a β APC subunit and the bilin-linker region of an α APC-like domain, called ApcE (or L_{CM}) (Fig. 1). ApcD and ApcE emit at 680 nm, and are terminal emitters of the PBS, which transfer the energy absorbed by this antenna to the photosystems.

The PBS architecture favors the funneling of absorbed photons from the peripheral rods to the central core, and then to the photosystems (ref. 36 and 37, see review³⁸). This energy transfer model is based on Förster Resonance Energy Transfer (FRET), where the excitation energy is transmitted from the donor to the acceptor molecule by resonance of excited states.³⁹ The down-hill energy transfer is possible due to the overlapping of the emission spectra of the different chromophores (ref. 36 and 37 see review³⁸). Mutations in the PBS terminal emitters affect the energy transfer to one or both photosystems in different cyanobacterial strains (see discussion in following sections).⁴⁰⁻⁴⁴

The entire scientific community agrees that in cyanobacteria, like in plants and green algae, state transitions are triggered by redox changes in the PQ pool.^{45,46} However, no agreement can be found in literature about the molecular mechanism, the sensor of the PQ redox state, or the signaling pathways involved. The present review critically evaluates the different propositions regarding the aforementioned knowledge gaps, and describes the most recent working models of






Fig. 1 Model of *Synechocystis* PCC 6803 phycobilisomes (A) and models of phycobilisome cores (B–D). The PBSs are constituted by a tricylindrical core in *Synechocystis* (A and B), a dicylindrical core in *S. elongatus* (C) or a pentacylindrical core in *Anabaena* PCC 7120. In *Synechocystis*, the core is surrounded by six rods (A). Each rod is formed by three hexamers of blue phycocyanin (PC). The rods are stabilized by non-chromophorylated linker proteins (in green). The rod-core linker (LR_c) protein (small cylinder in green) stabilizes the binding of the rods to the core. The linker-core (Lc) protein (in orange) present in the external trimers of each cylinder core stabilizes them. In the tricylindrical core (B), the upper cylinder is formed by 4 trimers of allophycocyanin (APC), composed by a α APC– β APC heterodimer which binds two phycocyanobilins (maximal fluorescence 660 nm). In the basal cylinders, in one of the external trimers (in dark blue) one α APC subunit is replaced by a special α APC-like subunit called ApcD. In the adjacent trimer (in violet), one β -subunit is replaced by a β APC-like subunit named ApcF. In the same trimer, another dimer is formed by a β -APC subunit and the bilin-linker domain of an α APC-like subunit (ApcE). Both ApcD and ApcE proteins were described as terminal energy acceptors of the PBS, presenting a maximal fluorescence at 680 nm.

cyanobacterial state transitions, based on the results obtained in the last few years.

2. Following photosystem fluorescence changes during state transitions

State transitions can be studied by fluorescence measurements at room temperature or at 77 K. At room temperature, changes in only PSII fluorescence can be measured, while at 77 K, both PSII and PSI fluorescence can be resolved. The fluorescence emission spectra of cyanobacterial cells depend on the wavelength of the excitation light (Fig. 2).

When cells are excited at 430 nm, which is preferentially absorbed by the Chl antennae, fluorescence emitted by Chl is principally observed. Thus, under these conditions, the changes observed are mainly related to processes occurring in the membrane Chl antennae and reaction centers. When cells are excited by light absorbed mainly by the PBS (550 to 620 nm, depending on the presence or absence of PE), fluorescence emitted by phycobiliproteins, phycobilisome final

emitters, and Chl is detected. In this case, the fluorescence changes observed are related to processes occurring in the PBS and the reaction center, and to the energy transfer from PBS to the photosystems. Independently of the excitation light, the peak at 695 nm is related to CP47 and Reaction Center II fluorescence, while the peak at 715–740 nm is related to PSI.^{47,48} Emissions at 575, 650, and 660 nm, related to PE, PC, and APC, respectively, contribute much more to the 77 K fluorescence spectrum when PBS are excited.^{49,50} In addition, with PBS excitation, the peak at 685 nm is mainly related to the PBS terminal emitter, with a contribution of CP43 emission.⁴⁸ In contrast, when Chl is excited, the 685 nm peak is mainly related to CP43 emission.

State I is characterized by a high PSII (685 and 695 nm) to PSI (715–720 nm) fluorescence ratio in the 77 K spectra. Conversely, State II is characterized by a low PSII to PSI fluorescence ratio. These differences are visible when the PBS or the Chl are excited (Fig. 2), suggesting that changes in both PBS and Chl-containing-complexes are involved in state transitions.^{51,52} It has been observed that PSII-related fluorescence is lower in State II compared to that in State I (see Tables 1 and 2). However, only one third of the realized studies showed that PSI-related fluorescence is lower in State I com-






Fig. 2 Following state transitions by fluorescence spectra at 77 K. 77 K fluorescence emission spectra of dark (black, State II) and blue-light (red, 85 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, State I) adapted *S. elongatus* (A and C) and *T. elongatus* (B and D) cells. The excitation was done at 430 nm (A and B) and 550 nm (C and D). Normalization was done at the peak of Rhodamine B (570 nm; 0.4 μM) in A and B; and at the peak of fluorescein (560 nm; 0.4 μM) in C and D. When Chls are mainly excited ($\lambda_{\text{exc}} = 430 \text{ nm}$), fluorescence changes are related principally to membrane processes. Meanwhile, when PBSs are mainly excited ($\lambda_{\text{exc}} = 550\text{--}620 \text{ nm}$), the observed fluorescence changes are related to changes in energy transfer inside the PBS and from the PBS to the photosystems. The peak at 650–660 nm is related to PC and APC; the peak at 685 nm is related to PBS terminal emitters and CP43; the peak at 695 nm is related to CP47 and RCII and PSI related peaks appear at 715–740 nm. While State I is characterized by a high PSII to PSI fluorescence ratio, State II is characterized by a low PSII to PSI fluorescence ratio. The spectra are the average of at least 3 independent biological replicates.

pared to that in State II (see Tables 1 and 2, and discussion about spillover below).

State transitions kinetics can be studied using modulated fluorimeters, which record the fluorescence resulting from excitation by non-actinic pulses (non-induced charge separation),⁵³ and measuring changes in fluorescence yield. The most frequently used fluorimeters are PAM101–103, dual-PAM-100, and multicolor PAM. The only fluorimeter used in state transition studies, *i.e.*, PAM101–103, uses a detecting light of 650 nm, which is absorbed by both PBS and Chl. Thus, changes in fluorescence yield could be a result of a change in the PBS and/or the antenna-Chl emission, or a change in the

excitation energy transfer from PBS to PSII. In multicolor PAM, several detecting pulses with different wavelengths (going from blue to red) can be used. Since fluorescence emission spectra of cyanobacterial cells depend markedly on the excitation wavelength, the contribution of PBS and Chl fluorescence will be different depending on the wavelength of the detecting pulses (see ref. 54 for a detailed explanation). Changes at the PBS level will mainly be observed when orange detecting light (590–630 nm) is used, while Chl emission changes will be more prominent when using blue detecting light. In dual-PAM, red, which is the most frequently used, and blue detecting pulses are also available.



Table 1 Evidence for PSII and PSI fluorescence changes in cyanobacterial state transitions at 77 K, when PBSs are excited

Article	Species	Changes in PSII fluorescence	Changes in PSI fluorescence	Internal standard used
Calzadilla <i>et al.</i> (2019) ⁵²	<i>S. elongatus</i> and <i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at 800 nm
McConnell <i>et al.</i> (2002) ⁵¹	<i>Synechococcus</i> 7002 and <i>Synechocystis</i> PCC 6803	Increase in State I	No change	No
Zlenko <i>et al.</i> (2019) ⁴⁴	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at 650 nm
Zhao <i>et al.</i> (2014) ⁹⁴	<i>Synechocystis</i> PCC 6803	Increase in State I	Decrease in State I	No – normalization at 707 nm
Ogawa <i>et al.</i> (2013) ⁷⁴	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak
Kaňa <i>et al.</i> (2012) ¹⁰⁵	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak
Kondo <i>et al.</i> (2009) ¹⁰⁷	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak
Yang <i>et al.</i> 2009 ¹⁵⁵	<i>Synechocystis</i> PCC 6803	Increase in State I	Decrease in State I	No – normalization at 709 nm
Fujimori <i>et al.</i> (2005) ¹⁵⁶	<i>Synechocystis</i> PCC 6803	No change	Increase in State I	No – normalization at 695 nm
El-Bissati <i>et al.</i> (2000) ⁶⁵	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	Yes – fluorescein
Emlyn-Jones <i>et al.</i> (1999) ⁷⁸	<i>Synechocystis</i> PCC 6803	Increase in State I	Slight increase in State I	No – normalization at phycocyanin fluorescence peak
Olive <i>et al.</i> (1997) ⁶¹	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at 665 nm
Joshua and Mullineaux, (2005) ⁹²	<i>S. elongatus</i>	Increase in State I	No change	No – normalization at 654 nm
Joshua and Mullineaux, (2004) ⁸⁹	<i>S. elongatus</i>	Increase in State I	No change	No – normalization at 654 nm
Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶	<i>S. elongatus</i>	Increase in State I	No change	Yes Rhodamine B
Aspinwall <i>et al.</i> (2004) ⁹⁹	<i>S. elongatus</i>	Increase in State I	No change	No
Deng <i>et al.</i> (2012) ⁹³	<i>Synechococcus</i> PCC 7002	Increase in State I	Slight decrease in State I	No
Huang <i>et al.</i> (2003) ⁷³	<i>Synechococcus</i> PCC 7002	Increase in State I	Slight decrease in State I	No – normalization at 640 nm
Schluchter <i>et al.</i> (1996) ⁹⁸	<i>Synechococcus</i> PCC 7002	Increase in State I	No change	No
Bruce <i>et al.</i> (1989) ⁸¹	<i>Synechococcus</i> PCC 7002	Increase in State I	Decrease in State I	No – normalization at 660 nm
Li <i>et al.</i> (2006) ⁶⁸	<i>Spirulina platensis</i>	Increase in State I	Decrease in State I	No – normalization at 710 nm
Li <i>et al.</i> (2004) ⁸³	<i>Spirulina platensis</i>	Increase in State I	Decrease in State I	No – normalization at 660 nm
Dong and Zhao, (2008) ⁸⁵	<i>Anabaena</i> PCC 7120	Increase in State I	Decrease in State I	No
Bruce and Biggins, (1985) ¹⁵⁷	<i>Anacystis nidulans</i>	Increase in State I	No change	No – normalization at 650 nm
Mullineaux and Allen, (1990) ⁴⁶	<i>Synechococcus</i> PCC 6301	Increase in State I	No change	No – normalization at 654 nm
Olive <i>et al.</i> (1986) ¹¹⁰	<i>Synechocystis</i> PCC 6714	Increase in State I	Decrease in State I	No

In PAM fluorometers, three levels of PSII fluorescence can be measured (Fig. 3). The minimal fluorescence level, F_0 , is determined using only the detecting modulated light in dark-adapted cells. No measurable charge separation is induced under these conditions, and all PSII are open. Strong pulses are applied to close all PSII centers and measure the maximal fluorescence level (F_m – in darkness and F'_m – in light) (Fig. 3). Under continuous illumination, the steady-state fluorescence level (F_s) is measured, which depends on the relative PSII and PSI activity influencing the PQ pool redox state. The contributions of PBS and Chl fluorescence in F_0 and F_m are different.⁵⁴

When State II adapted cells are illuminated with far-red light (absorbed by PSI) or low intensities of blue light (principally absorbed by Chl), an increase in all levels of fluorescence yield is observed, which is related to State II to State I transition (Fig. 3). The Chl antenna of PSI is almost three times larger than that of PSII, *i.e.*, 100 and 35 Chl molecules, respectively.^{55,56} Thus, illumination with light principally absorbed by Chl results in a higher activity of PSI compared to that of PSII, and oxidation of the PQ pool occurs.

When State I adapted cells are illuminated with orange or green light (principally absorbed by phycobiliproteins), a decrease in all levels of fluorescence related to State I to State II transition is induced (Fig. 3). Since PBS is the principal antenna of PSII,⁵⁷ this illumination increases PSII activity, resulting in reduction of the PQ pool. However, PSI is still active under orange and green light, because PBS can transfer

energy directly to both photosystems.^{58,59} Moreover, PSI can also receive energy from PSII (known as spillover), when the excess energy arriving at PSII is transferred to PSI *via* the Chl antennae.^{60–66} The percentage of energy arriving at one or the other photosystem depends on the cyanobacterial strain and the environmental conditions.^{59,67}

It is worth mentioning that upon illumination of dark-adapted cells with any kind of light (including orange light), PSI is activated and the PQ pool becomes more oxidized. Consequently, as mentioned previously, an increase in fluorescence yield related to State I transition is observed. However, blue light is more effective than orange light in inducing this transition.⁴⁶ Thus, both dark to light and orange to blue transitions induce an increase in fluorescence related to State I. Although majority of published results suggest that the mechanisms involved in these transitions are identical, one publication proposed that while changes in light quality induce only PBS movement, dark-light transition involves changes in concentration of PSI trimers and photosystem movements as well.⁶⁸

In the dark, in contrast to plants and green algae, cyanobacterial cells are in State II.^{69,70} In cyanobacteria, both photosynthesis and respiration take place in the thylakoid membranes, and PQ, Cyt *b₆f* and plastocyanin (or cytochrome *c₆*) are common to both electron transport chains (see review⁷¹). Homologs of mitochondrial Complex I (NDH-1) and Complex II (succinate dehydrogenase, SDH) reduce the PQ pool, and terminal oxidases oxidize it *via* the respiratory chain.⁷¹ Since



Table 2 Evidence of PSII and PSI fluorescence changes in cyanobacterial state transitions at 77 K, when Chls are excited

Article	Species	Changes in PSII fluorescence	Changes in PSI fluorescence	Internal standard used	Conclusions about spill-over
Calzadilla <i>et al.</i> (2019) ⁵²	<i>S. elongatus</i> and <i>Synechocystis</i> PCC 6803	Increase in State I	No change	Yes – Rhodamine B	No involvement
Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶	<i>S. elongatus</i> and <i>Synechocystis</i> PCC 6803	Increase in State I	No change	Yes – Rhodamine B	No involvement
Aspinwall <i>et al.</i> (2004) ⁹⁹	<i>S. elongatus</i>	Increase in State I	No change	No	Not addressed
Ogawa <i>et al.</i> (2013) ⁷⁴	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak	Not addressed
Stadnichuk <i>et al.</i> (2009) ⁸⁰	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	Yes – Rhodamine B	No involvement
Kondo <i>et al.</i> (2009) ¹⁰⁷	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak	Not addressed
Olive <i>et al.</i> (1997) ⁶¹	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at 665 nm	Spillover
Emlyn-Jones <i>et al.</i> (1999) ⁷⁸	<i>Synechocystis</i> PCC 6803	Increase in State I	No change	No – normalization at PSI peak	Minor role, if any, in cyanobacterial state transitions
McConnel <i>et al.</i> (2002) ⁵¹	<i>Synechococcus</i> 7002 and <i>Synechocystis</i> PCC 6803	Increase in State I	No change	No	Spillover
Deng <i>et al.</i> (2012) ⁹³	<i>Synechococcus</i> PCC 7002	No change	Increase in State I	No	Not addressed
Mullineaux and Allen, (1990) ⁴⁶	<i>Synechococcus</i> 6301	Increase in State I	No change	No – normalization at 654 nm	Not addressed
Salehian and Bruce, (1992) ¹⁵⁸	<i>Synechococcus</i> PCC 6301	Increase in State I	Slight increase in State I	No	Spillover
Li <i>et al.</i> (2006) ⁶⁸	<i>Spirulina platensis</i>	Increase in State I	Increase in State I	No – normalization at 708 nm	Spillover
Li <i>et al.</i> (2004) ⁸³	<i>Spirulina platensis</i>	Increase in State I	Increase in State I	No	Spillover
Bruce <i>et al.</i> (1989) ⁸¹	<i>Synechococcus</i> PCC 7002	Increase in State I	Increase in State I	No – normalization at 660 nm	Spillover

the PQ pool reduction is generally faster than its oxidation *via* respiration, the PQ pool is reduced in the dark.^{69,70} However, due to differences in this redox state balance between cyanobacterial strains, the level of PQ reduction state varies, leading to different levels of State II fluorescence in the dark.^{52,72} Mutants in the respiratory process also affect state transitions. For instance, NDH-1 and SDH mutants are in State I in the dark,^{73–76} while terminal oxidase mutants have a more pronounced dark State II.⁷⁵

The different “levels” of State II in the dark lead to different amplitudes of fluorescence changes during dark to light transitions. In *Synechocystis*,^{51,52,61,77–80} *Synechococcus* PCC 7002^{40,51,81} and *Spirulina platensis*,^{68,82,83} these changes are smaller than in *S. elongatus* or *T. elongatus* (ref. 52 and Fig. 3). Interestingly, both *S. elongatus* and *T. elongatus* lack the photo-protective mechanism OCP-related NPQ.⁸⁴ Thus, studying cyanobacterial state transitions in *S. elongatus* could lead to a better understanding of the PBS and membrane contributions to the process, which are further discussed in the following sections.

3. PBS contribution to state transitions

PBS contribution in cyanobacterial state transitions has been largely studied due to the role of LHCII in plant and green alga state transitions. First, it was shown that PBS mutations affect this process in different cyanobacterial strains.^{40,41,51,85} Then, FRAP experiments showing that PBS are capable of moving along the surface of the thylakoid membranes,^{86–88} and that the same chemicals that inhibit PBS movement inhibit state transitions,^{68,83,89} seemed to confirm that PBS movement from one photosystem to the other could be involved in state transitions. However, more recent experiments and results showed that the recovery of fluorescence after PBS bleaching is not only due to PBS diffusion but also to internal processes in this macromolecular complex.⁹⁰ The role of PBS in cyanobacterial state transitions remains a matter of discussion.

3.1 PBS mutants impaired in state transitions

Mutations in ApcD, ApcF^{40,41,51,85} and a particular domain of ApcE⁴⁴ affect state transitions in several cyanobacterial strains (Table 3). Dong *et al.* (2009)⁴⁰ suggested that in the ΔApcD *Synechococcus* PCC 7002 mutant, the inhibition of state transitions was due to a permanent decrease in energy transfer from PBS to PSI. However, it has recently been shown that a lack of ApcD also decreases the energy transfer from PBS to PSI in *S. elongatus*, without impairing state transitions.⁴³ Furthermore, in *Synechocystis*, the lack of ApcD does not decrease energy transfer from PBS to any of the photosystems, but it inhibits state transitions.^{41,43} In the ApcE-C190S and the PB-loop ApcE *Synechocystis* mutants, the energy transfer from PBS to the photosystems is significantly affected.^{43,44} However, state transitions in the former are not impaired,⁹¹ while they appear to be decreased in the latter.⁴⁴






Fig. 3 Following state transitions by fluorescence measurements at room temperature. The fluorescence changes of *S. elongatus* (A), *T. elongatus* (B) and *Synechocystis* (C) were followed by a PAM101 fluorometer. Dark adapted cells (Chl concentration $2.5 \mu\text{g mL}^{-1}$) were successively illuminated with blue light ($85 \mu\text{mol photons per m}^2 \text{ per s}^{-1}$) and orange light ($20 \mu\text{mol photons per m}^2 \text{ per s}^{-1}$). Saturating pulses ($400 \text{ ms} \times 12\,000 \mu\text{mol photons per m}^2 \text{ per s}$) were applied each 90 s. The PAM used in the measurements has a 650 nm detecting light, which is absorbed by PBS and Chl. Consequently, the changes in fluorescence yield observed under different conditions can be due to changes in the emission of PBS and/or Chl-antenna, or changes in excitation energy transfer between PBS and PSII. Three levels of fluorescence can be observed: F_0 , the minimal fluorescence level determined by detecting light in dark-adapted cells; F_m or F_m' , the maximal fluorescence obtained when saturating pulses are applied to the cells under darkness or light conditions, respectively; F_s ; steady-state fluorescence under continuous illumination. In the dark, cyanobacteria cells are in State II due to reduction of the PQ pool by respiration. Upon illumination of dark adapted cells with blue light (mainly absorbed by Chl), PSI is activated and the PQ pool becomes more oxidized, leading to State I. When State I adapted cells are illuminated with orange light (principally absorbed by phycobiliproteins), PSII activity is increased, resulting in reduction of the PQ pool. Thus, a decrease of F_m related to State I to State II transition is observed.

It has been proposed that the impairment of state transitions in the ΔApcD *Synechocystis* mutant is due to a permanent binding of PBS to PSII, which increases the PSII antenna even in darkness and PSII illumination (when the PQ pool is reduced).⁴¹ This hypothesis can explain the relatively high fluorescence observed in the dark-adapted ΔApcD *Synechocystis* cells. In addition, a lack of state transitions in the ΔApcD mutants was observed in *Synechocystis*, *Synechococcus* PCC 7002 and *Anabaena* PCC 7120,^{40,41,51,85} but not in *S. elongatus*⁴³ (Table 3). These results could be due to the different roles of ApcD in the PBS–photosystem interaction in different strains, which could also explain the different roles of ApcD in energy transfer to photosystems.^{40–43} As mentioned previously, absence of ApcD decreases energy transfer from PBS to PSI in *S. elongatus* and *Synechococcus* PCC 7002, but not in *Synechocystis*, while the ΔApcF mutation affects energy transfer to PSII and PSI only in *Synechocystis*.^{40–43}

A mutation in the *rpaC* gene also affects state transitions in *Synechocystis*, but not in *S. elongatus*.^{78,92} This gene, which is

strongly conserved among different cyanobacteria, codes for a 9 kDa protein, that has been proposed to be localized on the cytoplasmic side of the thylakoid membrane.⁷ The *rpaC* *Synechocystis* mutant was suggested to present a stronger PBS–PSII interaction, like the ΔApcD mutant, impairing state transitions.⁷ In contrast, a large decrease in the RpaC concentration did not affect this process in *S. elongatus*, which might again be due to different PBS–photosystem interaction mechanisms in different strains.⁹² A complete segregation of this mutation was not viable in *S. elongatus*, suggesting an essential role of this gene in this strain.

The different effects of the ApcD and RpaC mutants in *S. elongatus* and *Synechocystis* could also be explained by a different PBS contribution to state transitions in these strains. If this is the case, membrane processes could have a more important role in *S. elongatus* than in *Synechocystis*. It is worth mentioning that in the ΔRpaC *Synechocystis*, ΔApcD *Synechocystis*, *Synechococcus* 7002 and *Anabaena* PCC 7120 mutants, only the fluorescence changes related to PBS



Table 3 Alterations in state transitions and energy transfer in PBS mutants

Article	Species	Mutation	State transitions			Energy transfer PSII	Energy transfer PSI
			State transitions λ_{exc} PBS	λ_{exc} Chl	State transitions		
Calzadilla <i>et al.</i> (2019) ⁴³	<i>S. elongatus</i> and <i>Synechocystis</i> PCC 6803	ApcD, Apcf	Inhibited in <i>Synechocystis</i> Δ ApcD, impaired in <i>Synechocystis</i> Δ Apcf	Yes	Affected in <i>Synechocystis</i> Δ ApcF	Affected in <i>S. elongatus</i> Δ ApcD and <i>Synechocystis</i> Δ Apcf	
Ashby and Mullineaux, (1999) ⁴¹	<i>Synechocystis</i> PCC 6803	ApcD, Apcf	Inhibited in <i>Synechocystis</i> Δ ApcD, impaired in <i>Synechocystis</i> Δ Apcf	—	Affected in <i>Synechocystis</i> Δ ApcF	Affected in <i>Synechocystis</i> Δ Apcf	
Zlenko <i>et al.</i> (2019) ⁴⁴	<i>Synechocystis</i> PCC 6803	PB-loop in ApcE	Impaired	—	Affected	Not affected	
Deng <i>et al.</i> (2012) ⁹³	<i>Synechococcus</i> PCC 7002	ApcD	Inhibited	—	—	—	
Dong <i>et al.</i> (2009) ⁴⁰	<i>Synechococcus</i> PCC 7002	ApcD	Inhibited	—	—	Affected	
McConnel <i>et al.</i> (2002) ⁵¹	<i>Synechococcus</i> PCC 7002	ApcD	Inhibited	Yes	—	—	
Dong and Zhao (2008) ⁸⁵	<i>Anabaena</i> PCC 7120	ApcD	Inhibited	—	—	Affected	

were inhibited^{40,41,43,51,78,85,92,93} (Table 3). When Chl were preferentially excited at 77 K, the small fluorescence changes between State I and State II conditions were still observable.^{43,51,78,92} Thus, although the results related to PBS mutants are in favor of PBS participation in cyanobacterial state transitions, the relevance of this contribution is still not clear, and it might depend on the cyanobacterial strain studied and the prevailing growth conditions.

3.2 PBS long distance movement *versus* PBS detachment and small changes in PBS–photosystems interaction

The possible diffusion of PBS along the thylakoid membrane was first showed by fluorescence recovery after photobleaching (FRAP) experiments in *Dactylococcopsis salina* and *S. elongatus*, showing that PBS–photosystem association is a dynamic process which depends on weak interactions with membrane lipids.^{86,87} Based on FRAP experiments and to the possible association of PBSs with PSI and PSII,^{57,82,94} the hypothesis of a light-harvesting complex moving and modulating the energy reaching each photosystem, analogous to plant and green algae state transitions, has frequently been proposed (Fig. 4).⁹⁵ Within this hypothesis, it was suggested that in State II, the low PSII fluorescence is related to a smaller antenna, since most of the PBS are connected to PSI, while in State I, the high PSII fluorescence is related to a bigger PSII antenna, due to the migration of PBS from PSI to PSII.^{51,58,96,97}

However, FRAP results were suspected of some artificial artefacts due to the high energy laser used⁵¹ and possible effects induced in the PBS. Indeed, although PBS mobility was confirmed by FRAP and the fluorescence loss in photobleaching (FLIP) technique in *T. elongatus*, reversible fluorescence processes within the PBS were detected.⁸⁸ This led to the correction of PBS diffusion coefficient to a lower value ($1.7 \pm 0.4 \times 10^{-10} \text{ cm}^2 \text{ s}^{-1}$, which might differ depending on experimental conditions).⁸⁸ More recently, it was demonstrated by single molecule spectroscopy that PBS can lose (dark state) and recover their fluorescence spontaneously (blinking).⁹⁰ Under high light intensities, the “dark state” is formed more often and the PBSs remain longer time in this state.⁹⁰ Then, in darkness the PBSs recover their fluorescence. Because this blinking also affects FRAP experiments, the movement of PBS could be largely slower than it was supposed in the first studies.

Another argument for the involvement of PBS movement in state transitions was the fact that PBS mobility is inhibited by high osmotic buffers, which also inhibit state transitions in *S. elongatus* and *Spirulina platensis*.^{68,83,89} Incubation of cyanobacterial cells with phosphate (>0.5 M), betaine (1 M) or sucrose (1 M) suppresses the fluorescence changes related to state transitions observed at 77 K when PBS are excited ($\lambda_{exc} \approx 590 \text{ nm}$).^{68,83,89} These results seemed to support the hypothesis that the movement of PBS was the principal mechanism involved in cyanobacterial state transitions.⁸⁹ However, it was shown that these hyper-osmotic buffers also impair membrane processes, inhibiting the changes observed in 77 K fluorescence spectra when Chl is excited ($\lambda_{exc} \approx 430$).^{52,83} In






Fig. 4 Different models of phycobilisome contribution to cyanobacterial state transitions. (A) PBS-mobile model: During state transitions some PBSs move from one photosystem to the other changing the antenna size of both photosystems. In State I are more attached to PSII (high PSII fluorescence) and in State II to PSI (high PSI fluorescence); (B) PBS detachment model: the PBSs detach from both photosystems or from one of them in State I. The high “PSII” fluorescence in State I is related to the longer fluorescence life of detached PBSs.

addition, these chemical were also shown to inhibit quenching processes in the PBS.⁹⁰ Thus, it is not possible to conclude from these experiments that PBS long distance movements are involved in state transitions.

It was also shown that PSI oligomerization affects both state transitions^{98,99} and PBS diffusion.⁹⁹ In $\Delta PsaL$ strains of *Synechococcus* 7002 and *S. elongatus*, PSI trimerization is impaired, and State II to State I transition is faster than in the WT.^{98,99} In the *S. elongatus* mutant, it was also shown that the PBS diffusion rate was increased.⁹⁹ Although Aspinwall *et al.* (2004)⁹⁹ ascribed the faster kinetics of state transitions to the increase in PBS diffusion rate, the effect of differing distribution of protein complexes in the membrane cannot be discarded.

It has recently been shown that cyanobacterial thylakoid membranes are organized in a mosaic-like structure defined by domains.¹⁰⁰ Three types of domains were described, the first domain was enriched mainly by PSI, the second by PSII and PBS, and in the third domain, the abundance of PSII, PSI and PBS was balanced. These microdomains seem to be very stable and not change with light conditions. The fact that macrocomplexes containing PSII-PSI-PBS were isolated,⁵⁷ most probably from the third type of microdomain, suggests that very small position modification of the PBS (and/or photo-

systems) could induce changes in energy transfer from PBS to one or other photosystem. Thus, small PBS movement could be involved in state transitions. However, although the heterogeneous organization of cyanobacterial thylakoid membranes have been shown in several works,^{101–104} a conclusive model especially concerning the stability of the domains is still not available.

As an alternative to the long distance PBS movement model, recent publications have associated the high PSII fluorescence in State I to PBS detachment from thylakoid membranes (Fig. 4).^{79,105,106} Kaňa *et al.* (2012)¹⁰⁵ studied chlorophyll fluorescence induction curves and spectrally resolved fluorescence induction (SRFI) in different cyanobacterial strains, and proposed that transition from State II to State I involves PBS decoupling from PSII. Furthermore, Chukhutsina *et al.* (2015)⁷⁹ and Ranjbar Choubeh *et al.* (2018)¹⁰⁶ demonstrated, by measurements of fast time-resolved fluorescence, that the duration of PBS fluorescence was higher in State I than in State II, correlating with a small population of functionally disconnected PBS. However, in *Synechocystis*, PBS decoupling was estimated to be around 13% of the total PBS pool,⁷⁹ once again questioning the relevance of this light-harvesting complex in cyanobacterial state transitions.



As mentioned earlier, PBS contribution to state transitions can be observed by 77 K fluorescence emission spectra. All authors have reported an increase in PSII fluorescence (685 and 695 nm) during transition from State II to State I, on preferential excitation of the PBS ($\lambda_{\text{exc}} \approx 590$) (Table 1). Conversely, for PSI fluorescence (715–740 nm), some authors have reported no changes between State I and State II, while others have observed a decrease in its fluorescence during State I to State II transition, and still others have reported an increase during this transition (Table 1). The variability in the results can be ascribed at least partially to the different cyanobacterial strains studied. However different results were obtained with the same strains. Thus, different growth conditions and lights used to trigger state transitions could influence the obtained results. In addition, the normalization method employed in each case could significantly affect the interpretation of these results. In most cases, the fluorescence emission spectra were normalized within the spectra themselves (to PC, APC, PSII or even PSI fluorescence peaks) (Table 1), hindering the analysis of the fluorescence changes observed. For instance, some authors did not observe changes in PSI fluorescence between State II and State I in *Synechocystis*, but the PSI peak was used for normalization.^{74,105,107} In addition, several studies cited in Table 1 normalized their results to the APC peak, which was also found to change during state transitions.¹⁰⁸

In only two of the papers mentioned in Table 1,^{65,106} the fluorescence changes were normalized to an internal standard. When these chemicals, such as fluorescein or rhodamine B,^{65,106} are used, the fluorescence emission values are normalized to a wavelength outside the spectrum, allowing a better calibration of the fluorescence changes observed. Similar results can be obtained when normalization is done to wavelengths outside the spectrum (such as 800 nm), even in the absence of internal standards.⁵² In summary, the normalization of 77 K fluorescence emission spectra needs to be correctly addressed in order to facilitate the understanding of the PBS contribution to cyanobacterial state transitions. This subject is further discussed in section 4, where the contribution of membrane fluorescence to this physiological process is analyzed.

As a conclusion, although the analysis of PBS fluorescence changes during state transitions has been largely addressed in different cyanobacterial strains, it is still unclear how PBS participates in this process. As it is depicted in Table 1, the use of different experimental procedures to analyze the PBS-fluorescence related changes, hinders the possibility of finding general mechanisms for cyanobacterial state transitions. Nevertheless, based on the more recent results and our own interpretation of them, we propose that long distance movement of PBS is not involved in state transitions. Moreover, changes in the relative interaction of PBS and photosystems (involving slight PBS position changes) are most probably not involved neither. We suggest that only partial PBS detachment from one or both photosystems could contribute to the increase of PSII fluorescence in State I.

4. Membrane processes contribution to state transitions: spillover versus PSII quenching

Membrane participation in cyanobacterial state transitions was first proposed by Bruce *et al.* (1985) in the cyanobacteria *Anacystis nidulans*.¹⁰⁹ The involvement of the membrane was also suggested in *Synechocystis* cells, due to the changes in the thylakoid membrane topology of PSII distribution between the two states.^{61,110,111} In this cyanobacterium, a row disposition was observed for PSII in State I, and a random distribution (favoring spillover) was observed in State II. Further evidence supporting the involvement of the movement of membrane complexes, or changes in the oligomerization state of these complexes was presented by El-Bissati *et al.* (2000),⁶⁵ who demonstrated that this process is affected by alterations in membrane fluidity. However, more recently, Maksimov *et al.* (2017)¹¹² have suggested that the rate of oxidation of the PQ pool depends on membrane fluidity, and therefore on temperature. Thus, the changes observed by Bissati *et al.* could have been related to slower kinetics of reduction and oxidation of the PQ pool in less fluid membranes.⁶⁵ Finally, state transitions were also observed in mutants of *Synechocystis* and *Synechococcus* 7002 lacking PBS.^{81,113}

It was first proposed that the rearrangement of photosystem positions in the membrane induced changes in spillover (energy transfer from PSII to PSI) during state transitions.^{61,109} However, although changes in spillover have long been studied (Fig. 5),^{51,61,77} there are no clear results confirming the participation of this phenomenon in cyanobacterial state transitions. A key evidence supporting the spillover model would be a decrease in PSI fluorescence at 77 K (when Chl is preferentially excited) during State II to State I transition (when photosystems move apart), concomitant with an increase in PSII fluorescence under the same conditions. However, most studies on spillover did not show this result (Table 2). Of the articles listed in Table 2, only one work realized by Bruce *et al.* (1989) in *Synechococcus* PCC 7002,⁸¹ and two studies on *Spirulina platensis*,^{68,83} showed an apparent increase in PSI fluorescence in State II, which can be considered as an evidence for spillover. However, none of these works analyzed the 77 K steady-state fluorescence emission spectra using internal standards (such as fluorescein or rhodamine B). The utilization of these standards allows the study of the absolute variation in the different fluorescence contributions.^{52,79,80,106} Interestingly, when these internal standards were used, authors lead to the conclusion that due to the constant level of PSI emission between State I and II, spillover is an unlikely mechanism of cyanobacterial state transitions.^{52,80,106} Moreover, a recent study employing a streak camera to measure fast (ps to ms) fluorescence decays, strongly suggested that at least in *S. elongatus* (and most probably in *Synechocystis*) changes in spillover are not involved in state transitions.¹⁰⁶ These authors measured fast fluorescence decay kinetics (at 77 K) of cells adapted to State I and II, and subsequently obtained the






Fig. 5 Different models of membrane contribution to cyanobacterial state transitions. (A) Spillover: Movement of one or both photosystems induces changes in spillover. Larger spillover (energy transfer from PSII to PSI) in State II than in State I. (B) PSII quenching: In this model, the excitation energy arriving to PSI do not change between State I and II, and the fluorescence decrease observed during State I to State II transition is due to a specific PSII-quenching at the level of the reaction center and/or the inner antennae CP47 and CP43.

decay-associated spectra (DAS) after global analysis of the data. When 430 nm excitation wavelengths were used, PSII emission decreased in State II, however, the DAS showed that PSI emission was similar in State I and State II.

Ranjbar Choubeh *et al.* (2018) recently proposed that the decrease in PSII fluorescence is related to a reversible PSII-quenching in State II (Fig. 5), at least in *S. elongatus* and *Synechocystis* cells.¹⁰⁶ This quenching has been suggested to take place at the level of PSII, and is not related to the NPQ mechanism described for cyanobacteria, which occurs in the PBS.⁹ It is worth noting that higher PSII quenching should lead to extra heat emission in State II. However, optoacoustic studies did not show differences between the heat production in States I and II.^{114,115} As previously discussed, Van Amerongen *et al.* showed that in both *S. elongatus* and *Synechocystis*, a partial disconnection of PBS occurs in State I,^{79,106} which should increase the conversion of excitation energy into heat. This increase in heat could be adventitiously similar to the heat increase due to PSII quenching in State II, which might explain the absence of heat emission changes during state transitions. Overall, the results of Ranjbar Choubeh *et al.* (2018) demonstrated that PSII is not quenched by PSI in State II in *S. elongatus*.¹⁰⁶ The quenching mechanism related to PSII remains to be elucidated.

Stadnichuk *et al.* (2009) also suggested that state transitions do not involve PSI, and could be completely attributed to PSII.⁸⁰ In their study, fluorescence emission changes at 77 K were observed for PSII, but not for PSI, when Chl were excited in *Synechocystis* cells. Based on these results, they suggested that spillover is an unlikely mechanism implicated in cyanobacterial state transitions.⁸⁰ These observations were made in WT, CK (mutant without PC) and PAL (mutant depleted in PBS) strains, leading to the conclusion that thylakoid membranes participate in state transitions, and their contribution is not negligible.

5. The signal transduction pathways and the role of Cyt *b*₆*f*

Apart from the mechanism involved in cyanobacterial state transitions, the signal transduction pathways leading to State I/II transition remain to be deciphered. It is well established that changes in the redox state of the PQ pool trigger state transitions in cyanobacteria, like in plants and green algae.^{46,69,73,74,116,117} While a reduced PQ pool induces State II transition, State I transition is induced by its oxidation. As previously discussed, in the dark, cyanobacterial cells are in State II due to PQ pool reduction *via* respiration.^{69,70} In addition,



chemicals oxidizing or reducing the PQ pool also trigger state transitions⁴⁶ (Fig. 6). Among these chemicals, some quinones, such as *p*-benzoquinone (PBQ) or 2,6-dimethoxy-1,4-benzoquinone (DMBQ), induce State I transition through oxidation of the PQ pool in dark-adapted cells.^{73,117} In contrast, addition of 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone (DBMIB) induces State I to State II transition even under blue-light illumination.^{46,73,117} DBMIB binds close to the [2Fe–2S] cluster of the Cyt *b₆f* and blocks photosynthetic and respiratory electron transport,^{118,119} inhibiting the re-oxidation of the PQ pool by the complex. Thus, the PQ pool is reduced in its presence.

There are two DBMIB binding sites in Cyt *b₆f*, one with high affinity and the other with low affinity.¹¹⁹ The high affinity binding site is located at the periphery of the complex, outside the Q_o pocket. At low concentrations (0.5–1 μM), DBMIB binds only to the high affinity site, and in darkness, it does not inhibit electron transport *in vivo*, as reported in *Synechococcus* 7002.¹¹⁹ For inhibition of electron transport, the cells must be illuminated to induce the migration of DBMIB from its peripheral position to the low affinity site (close to the [2Fe–2S] cluster).¹¹⁹ However, at higher concentrations

(>2 μM), DBMIB fully inhibits Cyt *f* re-reduction also in darkness.

Given the relationship between the redox state of the PQ pool and state transitions in both cyanobacteria and plants,^{45,46} and the involvement of Cyt *b₆f* in the signal transduction pathway of plants and algae,²⁹ the participation of this complex in cyanobacterial state transitions has also been hypothesized. Results suggesting the participation of phosphorylation reactions in this signal transduction pathway strengthened this idea.^{120–125} Hence, Mao *et al.* (2002) and Huang *et al.* (2003) proposed Cyt *b₆f* as the redox sensor of the PQ pool in cyanobacterial state transitions.^{73,117} They based their conclusions on the effect of DBMIB on PBQ or DMBQ poisoned cells, suggesting that DBMIB induces transition to State II by binding to the Cyt *b₆f* complex, even when the PQ pool remains oxidized in the presence of the aforementioned quinones.^{73,117} However, they did not measure the redox state of the pool under these conditions.

It has recently been shown, using chlorophyll fluorescence induction curves, that the PQ pool is rapidly reduced by addition of DBMIB, even in the presence of DMBQ.⁵²




Fig. 6 Effect of different chemicals on the redox state of the PQ pool in cyanobacteria. Cyanobacterial state transitions is triggered by changes in the redox state of the PQ pool. While accumulation of oxidized PQ induces State I transition, increase in the concentration of its reduced form (PQH₂) leads to State II. Different chemicals affect the redox state of PQ pool. The quinones *p*-benzoquinone (PBQ) and 2,6-dimethoxy-1,4-benzoquinone (DMBQ) by accepting electrons from PQH₂ induces PQ pool oxidation triggering State I transition. On the contrary, addition of 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone (DBMIB) induces State II transition. DBMIB inhibits Cyt *b₆f* re-oxidation of the PQ pool, and as a consequence PQH₂ is accumulated. *N,N,N',N'*-Tetramethyl-*p*-phenylenediamine (TMPD) is capable of oxidizing the PQ pool under blue light illumination, leading to State I, even in the presence of DBMIB. In the figure, the arrows indicates the direction of electron transport between different protein complexes/chemical compounds, and its size is proportional to the quantity of electrons transfer under different conditions. Orange and blue arrows indicate electron transfer under orange or blue light illumination, respectively. Black arrows indicate electron transfer occurring under dark and light conditions. Dashed arrows indicate electrons taken/given by the added chemicals. Protein complexes: PSII, photosystem II; PSI, photosystem I; PQ, plastoquinone pool; Cyt *b₆f*, cytochrome *b₆f*. Chemical compounds: DCMU; 3-(3,4-dichlorophenyl)-1,1-dimethylurea; DMBQ, 2,6-dimethoxy-1,4-benzoquinone; DBMIB, 2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone; TMPD, *N,N,N',N'*-tetramethyl-*p*-phenylenediamine; PBQ, *p*-benzoquinone.



Moreover, the same concentration of DBMIB has different effects on state transitions under light-induced or DMBQ-induced State I-adapted cells. The concentration of DBMIB used were higher than 2 μM . These experiments strongly suggest that DBMIB induces State II transition due to the reduction of the PQ pool, and not due to its binding to Cyt b_6/f . In another set of experiments, the same authors showed that it was possible to induce State I transition when illuminating DBMIB-poisoned cells, using *N,N,N'*-tetramethyl-*p*-phenylenediamine (TMPD).⁵² This chemical was shown to restore photosynthetic electron transport, even when Cyt b_6/f was inhibited by DBMIB.¹²⁶ Thus, the oxidation of the PQ pool triggers State I transition, through a signal transduction pathway that does not involve Cyt b_6/f .

It has also been proposed that the Chl *a* molecule of the Cyt b_6/f complex is the signal transmitter in cyanobacterial state transitions.¹²⁷ In this model, the Chl *a* senses the changes in the Qo site, and participates in this signal transduction pathway by altering its volume. This hypothesis proposed by Vladkova (2016) was based on the analysis of various Cyt b_6/f X-ray structures, differing in the volume of Chl *a*.¹²⁷ Changes in the Chl *a* volume, along with changes in the redox states of the PQ pool, would affect the hydrophobic thickness of Cyt b_6/f , which would induce a hydrophobic mismatch in the membrane. This hydrophobic mismatch is the difference between the hydrophobic thickness of the protein and the hydrophobic thickness of the lipid bilayer, and would be the driving force behind membrane reorganization during the progression of state transitions.¹²⁷

If the Chl *a* volume is the signal transmitter in cyanobacterial state transitions, it would be expected that inhibition of Cyt b_6/f would have an irreversible effect on this process. The volume of Chl *a* must not change in DBMIB-poisoned cells, however, addition of TMPD was found to induce state transitions.⁵² This suggests that the involvement of this Chl *a* molecule is also unlikely. It is worth mentioning that under the conditions tested by Calzadilla *et al.* (2019) (DBMIB concentration >2 μM),⁵² the two DBMIB binding sites in Cyt b_6/f were occupied, fully inhibiting the complex and photosynthetic electron transport.¹¹⁹

The possible participation of phosphorylation reactions in cyanobacterial state transitions has also been studied. First reports showing a correlation between changes in thylakoid phosphorylation patterns, PQ pool redox changes and state transitions came from studies conducted on *Synechococcus* 6301.^{120,122,128} A model was proposed, in which protein phosphorylation and PBS migration were requisites for State I/II transition (ref. 121, see review¹²⁹). However, none of these studies showed a direct connection between the changes in phosphorylation and state transitions. Moreover, the results of Biggins *et al.* (1984)¹³⁰ suggest that state transitions in organisms with PBS do not involve reversible protein phosphorylation. This has been discussed in detail by Allen (1992).¹²⁹

In more recent works, it has been observed that PBS and photosystem proteins can undergo phosphorylation, indicating the participation of these reactions in cyanobacterial state

transitions.^{123–125} However, only Chen *et al.* (2015) directly addressed this possible relationship.¹²⁴ These authors showed that four residues of the β subunits of PC (Ser22, Ser49, Thr94 and Ser154) were phosphorylated. Mutation of these amino acids avoiding phosphorylation seemed to affect state transitions. A slower and smaller increase of fluorescence was observed upon illumination of dark-adapted mutant cells. However, these cells were not extensively characterized, for instance, alterations in the energy transfer within the PBS were suggested, which might affect PBS concentration and/or PSI/PSII ratio between strains.¹²⁴ Moreover, the dark PQ pool redox state was not measured; in addition, the differences between mutants cannot be ignored. Consequently, the observed impairment in state transitions could not conclusively be linked to the lack of PBS phosphorylation.

Ser/Thr kinases have been widely described as key factors in the signal transduction pathways of cyanobacteria (ref. 131 and 132, see review¹³³). *Synechocystis* presents 12 genes coding for Ser/Thr kinases (Spk),^{134,135} of which seven are classified in the PKN2 subfamily (spkA to spkG),¹³⁶ and the rest have been assigned to the ABC1 subfamily (spkH to spkL).¹³⁷ These kinases have been shown to participate in different cell functions,¹³⁸ such as cell motility,¹³⁹ oxidative stress response,¹⁴⁰ salt and low temperature stress,^{141,142} and regulation of carbon¹⁴³ and nitrogen metabolism.¹⁴⁴ In addition, the GroES chaperone protein and ferredoxin were also reported to be targets of some Spk.^{145,146}

As previously mentioned, Chen *et al.* (2015) suggested the possible participation of Ser/Thr kinases in cyanobacterial state transitions,¹²⁴ like in plants and green algae.^{16,17,124} Recently, Calzadilla *et al.* (2019) addressed this issue by constructing single mutants for all the Ser/Thr kinases and phosphatases identified in *Synechocystis*.⁵² None of them were impaired in state transitions. In addition, experiments with kinase (Staurosporine and K252a) and phosphatase inhibitors (NaF and Na_3VO_4) were also performed. Incubation with these chemicals had no effect on state transitions in *Synechocystis* and *S. elongatus*, although they affected other processes in the cells.⁵² Thus, these results lead to the conclusion that phosphorylation reactions are not essential in cyanobacterial state transitions.

6. Conclusions and perspectives

Despite the large number of studies conducted on cyanobacterial state transitions, the mechanism behind this physiological process has not yet been determined. Four different mechanisms (PBS movement, PBS detachment, spillover and PSII quenching), and combinations of them, were proposed to be related to the changes of fluorescence observed when PSII or PSI is specifically illuminated. Moreover, different models were proposed for the same cyanobacteria strain (Table 4).

Nevertheless, considering the literature included in the present review, it is possible to make some conclusions about the molecular mechanism and the signal transduction path-



Table 4 State transitions models proposed for the most common studied cyanobacteria

Organism	Model proposed	References
<i>S. elongatus</i>	PBS-mobile	Sarcina <i>et al.</i> (2001) ⁸⁷ ; Joshua and Mullineaux (2004) ⁸⁹ ; Aspinwall <i>et al.</i> (2004) ⁹⁹ ;
	PBS-detachment	Joshua and Mullineaux (2005) ⁹²
	PSII-quenching	Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶ Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶
<i>Synechocystis</i> PCC 6803	PBS-mobile	Ashby and Mullineaux (1999) ⁴¹ ; Joshua and Mullineaux (2005) ⁹²
	PBS-mobile/spillover	McConnell <i>et al.</i> (2002) ⁵¹
	PBS-detachment	Chukhutsina <i>et al.</i> (2015) ⁷⁹ ; Kaňa <i>et al.</i> (2012) ¹⁰⁵ ; Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶
	Spillover	Olive <i>et al.</i> (1997) ⁶¹ ; Folea <i>et al.</i> (2008) ¹¹¹ ; El-Bissati <i>et al.</i> (2000) ⁶⁵ ; Vernotte <i>et al.</i> (1990) ⁷⁷
	PSII-quenching	Ranjbar Choubeh <i>et al.</i> (2018) ¹⁰⁶ ; Stadnichuk <i>et al.</i> (2009) ⁸⁰
<i>Synechococcus</i> PCC 7002	PBS-mobile	Dong <i>et al.</i> (2009) ⁴⁰
	PBS-mobile/spillover	McConnell <i>et al.</i> (2002) ⁵¹
	Spillover	Bruce <i>et al.</i> (1989) ⁸¹
<i>Spirulina platensis</i>	PBS-mobile/spillover	Li <i>et al.</i> (2006) ⁶⁸ ; Li <i>et al.</i> (2004) ⁸³
<i>Dactylococcopsis salina</i>	PBS-mobile	Mullineaux <i>et al.</i> (1997) ⁸⁶

ways involved. Our proposed model is described in Fig. 7. First, it is clear that cyanobacterial state transitions cannot be reduced to changes in the energy transfer from PBS to the photosystems, as was proposed several years ago. Although it was demonstrated that PBS could move on the thylakoid surface, there is no direct evidence showing that this movement has a physiological role in state transitions. Furthermore, recent results suggest that only a partial detachment of PBS from one or both photosystems is involved in state transitions. However, we cannot totally discard small changes in their position favoring a better energy transfer to one or other photosystem, at least in some cyanobacteria strains. Finally, PBS contribution seems to depend on the cyanobacterial strain and growth conditions.

In contrast, membrane processes seem to be an intrinsic part of cyanobacterial state transitions. Chl fluorescence changes can be ascribed to the proposed PSII-quenching, and/or changes in spillover involvement. The different membrane topologies observed in State I and State II, favoring or hindering direct energy transfer from PSII to PSI, supported changes in spillover during state transitions. However, recent studies employing a streak camera to measure fast (ps to ms) fluorescence decays, strongly suggested that changes in spillover are not involved in state transitions.¹⁰⁶

Thus, many questions remain to be answered: How changes in membrane topology participate in cyanobacterial state transitions? Do these membrane changes also affect PBS energy transfer to photosystems? What is the mechanism involved in PSII-specific quenching? Regarding this last question, it has recently been shown that the *thermal phase* of the O–J–I–P curve can be ascribed to a light-induced conformational change in PSII, triggered by saturating multiple turnover flashes (ref. 147–149, see review¹⁵⁰). This increase in the fluorescence yield of Chl *a* is temperature dependent,¹⁴⁷ and its kinetics depends on the length, rather than the intensity, of the flashes.¹⁴⁸ This increase was further found to be related with changes in the interaction between the PSII reaction center and the antenna proteins CP43 and CP47.^{149,151}

Although it has been established that this phenomenon is not related with state transitions, because it takes place at the ms scale and can relax and be regenerated without changes in redox state of the PQ pool,¹⁴⁷ similar PSII conformational changes can be associated with the PSII-specific quenching suggested by Ranjbar Choubeh *et al.* (2018).¹⁰⁶ If the relevance of this quenching is confirmed, the definition of cyanobacterial state transitions as a re-balance of the excitation energy reaching the photosystems needs to be reconsidered.

The different contributions of PBS and membrane processes in cyanobacterial state transitions seem to be dependent on the cyanobacterial strain. Hence, comparing the same processes in different strains would lead to more conclusive results regarding state transitions. Along these lines, the utilization of *S. elongatus* and *T. elongatus* as model organisms can facilitate this work, due to the larger fluorescence differences observed between State I and II, compared to the other organisms used in the past (such as *Synechocystis* or *Synechococcus* 7002). It is also worth mentioning that state transitions could be dependent on growth conditions of the cultured strains. If this is the case, analyzing the fluorescence changes of cells acclimated to different conditions would allow a better understanding of the relative contributions of PBS and membrane processes to state transitions.

In case of the signal transduction pathways, showing that Cyt *b₆f* and phosphorylation reactions are not involved in cyanobacterial state transitions was a step forward. However, how changes in the redox state of the PQ pool could be transmitted to the PBS and/or photosystems remains to be discovered. If we consider that one of the main fluorescence changes between State I and II might come from the membrane contribution, and might involve a PSII-specific quenching, one tempting hypothesis would be that PSII itself is able to sense the redox state of the PQ pool. Accumulation of Q_A⁻ cannot be this redox sensor, because while its accumulation due to 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) addition induces State I, its accumulation due to the reduction of the PQ pool induces State II.






Fig. 7 A proposed model for cyanobacterial state transitions. Based on all the literature revised in the present article, we believed that the most probable mechanism involved in cyanobacterial state transitions involves a specific PSII-quenching during State I to State II transition (A), and a partial detachment of PBS during State II to State I transition (B). Nevertheless, small PBS movements that might affect the energy transfer to either of both photosystems cannot be ruled out. In addition, regarding the signal transduction pathways, it was demonstrated that neither Cyt b₆f nor phosphorylation reactions are involved in cyanobacterial state transitions. The participation of known or unknown proteins in this signaling pathway needs to be elucidated.

It was suggested that a Qc channel between Cyt b559 and PsbJ could exist.¹⁵² This channel, which is connected to the Q_B site, has been suggested to have a quinone-binding function, and can participate in PSII regulation. Thus, a possible hypothesis would be to consider the Qc hydrophobic tunnel as the redox sensor of the PQ pool. However, the existence of this Qc site is still unclear, since its presence was not confirmed in the most recent high resolution PSII structure model.⁵⁶ Notwithstanding, it was shown that mutations in Cyt b559, PsbJ, and near the putative Qc site, can alter the kinetics and amplitude of state transitions in *Synechocystis* cells.¹⁵³ These results present an interesting starting point for further studies,

although the participation of novel factors or known proteins in this signal transduction pathway cannot be discarded. New approaches, such as the utilization of mutant library screenings,¹⁵⁴ need to be developed to shed light on the mechanism connecting the redox state of the PQ pool and the fluorescence changes observed in cyanobacterial state transitions.

Funding

This work was supported by grants from the Agence Nationale de la Recherche (ANR projects RECYFUEL (ANR-16-CE05-



0026)) and by the European Union's Horizon 2020 Research and Innovation program under grant agreement no. 675006 (SE2B). P. C. salary is financed by RECYFUEL (ANR project). The research is also supported by the Centre National de la Recherche Scientifique (CNRS) and the Commissariat à l'Energie Atomique (CEA).

Conflicts of interest


There are no conflicts to declare.

References

- N. Sluchanko, Y. Slonimskiy and E. Maksimov, *Biochemistry*, 2017, **82**, 1592–1614.
- D. Kirilovsky and C. A. Kerfeld, *Nat. Plants*, 2016, **2**, 16180.
- D. Kirilovsky, *Photosynth. Res.*, 2007, **93**, 7.
- D. Kirilovsky and C. A. Kerfeld, *Biochim. Biophys. Acta, Bioenerg.*, 2012, **1817**, 158–166.
- C. A. Kerfeld, M. R. Melnicki, M. Sutter and M. A. Dominguez-Martin, *New Phytol.*, 2017, **215**, 937–951.
- J. Van Thor, C. Mullineaux, H. Matthijs and K. Hellingwerf, *Bot. Acta*, 1998, **111**, 430–443.
- C. W. Mullineaux and D. Emlyn-Jones, *J. Exp. Bot.*, 2004, **56**, 389–393.
- D. Kirilovsky, *Photosynth. Res.*, 2015, **126**, 3–17.
- A. Wilson, G. Ajlani, J.-M. Verbavatz, I. Vass, C. A. Kerfeld and D. Kirilovsky, *Plant Cell*, 2006, **18**, 992–1007.
- M. G. Rakhimberdieva, I. V. Elanskaya, W. F. Vermaas and N. V. Karapetyan, *Biochim. Biophys. Acta, Bioenerg.*, 2010, **1797**, 241–249.
- N. Murata, *Biochim. Biophys. Acta, Bioenerg.*, 1969, **172**, 242–251.
- C. Bonaventura and J. Myers, *Biochim. Biophys. Acta, Bioenerg.*, 1969, **189**, 366–383.
- J. Minagawa, *Biochim. Biophys. Acta, Bioenerg.*, 2011, **1807**, 897–905.
- S. Lemeille and J.-D. Rochaix, *Photosynth. Res.*, 2010, **106**, 33–46.
- P. Pesaresi, M. Pribil, T. Wunder and D. Leister, *Biochim. Biophys. Acta, Bioenerg.*, 2011, **1807**, 887–896.
- N. Depège, S. Bellafiore and J.-D. Rochaix, *Science*, 2003, **299**, 1572–1575.
- S. Bellafiore, F. Barneche, G. Peltier and J.-D. Rochaix, *Nature*, 2005, **433**, 892.
- A. V. Vener, P. J. van Kan, A. Gal, B. Andersson and I. Ohad, *J. Biol. Chem.*, 1995, **270**, 25225–25232.
- A. V. Vener, P. J. Van Kan, P. R. Rich, I. Ohad and B. Andersson, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 1585–1590.
- F. Zito, G. Finazzi, R. Delosme, W. Nitschke, D. Picot and F. A. Wollman, *EMBO J.*, 1999, **18**, 2961–2969.
- Z. Zhang, L. Huang, V. M. Shulmeister, Y.-I. Chi, K. K. Kim, L.-W. Hung, A. R. Crofts, E. A. Berry and S.-H. Kim, *Nature*, 1998, **392**, 677.
- C. Breyton, *J. Biol. Chem.*, 2000, **275**, 13195–13201.
- G. Finazzi, F. Zito, R. P. Barbagallo and F.-A. Wollman, *J. Biol. Chem.*, 2001, **276**, 9770–9774.
- L. Dumas, F. Zito, S. Blangy, P. Auroy, X. Johnson, G. Peltier and J. Alric, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 12063–12068.
- U. K. Larsson, B. Jergil and B. Andersson, *Eur. J. Biochem.*, 1983, **136**, 25–29.
- D. Kyle, L. Staehelin and C. Arntzen, *Arch. Biochem. Biophys.*, 1983, **222**, 527–541.
- M. Pribil, P. Pesaresi, A. Hertle, R. Barbato and D. Leister, *PLoS Biol.*, 2010, **8**, e1000288.
- A. Shapiguzov, B. Ingelsson, I. Samol, C. Andres, F. Kessler, J.-D. Rochaix, A. V. Vener and M. Goldschmidt-Clermont, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 4782–4787.
- F.-A. Wollman and C. Lemaire, *Biochim. Biophys. Acta, Bioenerg.*, 1988, **933**, 85–94.
- A. N. Glazer, *Biochim. Biophys. Acta*, 1984, **768**, 29–51.
- R. MacColl, *J. Struct. Biol.*, 1998, **124**, 311–334.
- N. Adir, *Photosynth. Protein Complexes*, 2008, 243–274.
- N. Adir, S. Bar-Zvi and D. Harris, *Biochim. Biophys. Acta, Bioenerg.*, 2019, DOI: 10.1016/j.bbabi.2019.07.002.
- M. Watanabe and M. Ikeuchi, *Photosynth. Res.*, 2013, **116**, 265–276.
- D. A. Bryant, *Microbiology*, 1982, **128**, 835–844.
- A. R. Holzwarth, *Physiol. Plant.*, 1991, **83**, 518–528.
- I. Eisenberg, F. Caycedo-Soler, D. Harris, S. Yochelis, S. F. Huelga, M. B. Plenio, N. Adir, N. Keren and Y. Paltiel, *J. Phys. Chem. B*, 2017, **121**, 1240–1247.
- T. Mirkovic, E. E. Ostroumov, J. M. Anna, R. Van Grondelle and G. D. Scholes, *Chem. Rev.*, 2016, **117**, 249–293.
- M. Şener, J. Strümpfer, J. Hsin, D. Chandler, S. Scheuring, C. N. Hunter and K. Schulten, *ChemPhysChem*, 2011, **12**, 518–531.
- C. Dong, A. Tang, J. Zhao, C. W. Mullineaux, G. Shen and D. A. Bryant, *Biochim. Biophys. Acta, Bioenerg.*, 2009, **1787**, 1122–1128.
- M. K. Ashby and C. W. Mullineaux, *Photosynth. Res.*, 1999, **61**, 169–179.
- J. Zhao, J. Zhou and D. Bryant, *Photosynth. Res.*, 1992, **34**, 83.
- P. I. Calzadilla, F. Muzzopappa, P. Sétif and D. Kirilovsky, *Biochim. Biophys. Acta, Bioenerg.*, 2019, **1860**, 488–498.
- D. V. Zlenko, I. V. Elanskaya, E. P. Lukashev, Y. V. Bolychevtseva, N. E. Suzina, E. S. Pojidaeva, I. A. Kononova, A. V. Loktyushkin and I. N. Stadnichuk, *Biochim. Biophys. Acta, Bioenerg.*, 2019, **1860**, 155–166.
- J. F. Allen, J. Bennett, K. E. Steinback and C. J. Arntzen, *Nature*, 1981, **291**, 25.
- C. W. Mullineaux and J. F. Allen, *Photosynth. Res.*, 1990, **23**, 297–311.



- 47 R. Van Dorssen, J. Breton, J. Plijter, K. Satoh, H. Van Gorkom and J. Ames, *Biochim. Biophys. Acta, Bioenerg.*, 1987, **893**, 267–274.
- 48 D. Siefermann-Harms, *J. Chromatogr. A*, 1988, **448**, 411–416.
- 49 A. N. Glazer and L. Stryer, *Biophys. J.*, 1983, **43**, 383–386.
- 50 M. Wu, P. M. Goodwin, W. P. Ambrose and R. A. Keller, *J. Phys. Chem.*, 1996, **100**, 17406–17409.
- 51 M. D. McConnell, R. Koop, S. Vasil'ev and D. Bruce, *Plant Physiol.*, 2002, **130**, 1201–1212.
- 52 P. I. Calzadilla, J. Zhan, P. Sétif, C. Lemaire, D. Solymosi, N. Battchikova, Q. Wang and D. Kirilovsky, *Plant Cell*, 2019, **31**, 911–931.
- 53 U. Schreiber, W. Bilger and C. Neubauer, in *Ecophysiology of photosynthesis*, Springer, 1995, pp. 49–70.
- 54 W. Remelli and S. Santabarbara, *Biochim. Biophys. Acta, Bioenerg.*, 2018, **1859**, 1207–1222.
- 55 P. Jordan, P. Fromme, H. T. Witt, O. Klukas, W. Saenger and N. Krauß, *Nature*, 2001, **411**, 909.
- 56 Y. Umena, K. Kawakami, J.-R. Shen and N. Kamiya, *Nature*, 2011, **473**, 55.
- 57 H. Liu, H. Zhang, D. M. Niedzwiedzki, M. Prado, G. He, M. L. Gross and R. E. Blankenship, *Science*, 2013, **342**, 1104–1107.
- 58 C. W. Mullineaux, *Biochim. Biophys. Acta, Bioenerg.*, 1992, **1100**, 285–292.
- 59 M. G. Rakhimberdieva, V. A. Boichenko, N. V. Karapetyan and I. N. Stadnichuk, *Biochemistry*, 2001, **40**, 15780–15788.
- 60 A. Ley and W. Butler, *Biochim. Biophys. Acta, Bioenerg.*, 1980, **592**, 349–363.
- 61 J. Olive, G. Ajlani, C. Astier, M. Recouvreur and C. Verrotte, *Biochim. Biophys. Acta, Bioenerg.*, 1997, **1319**, 275–282.
- 62 J. Biggins and D. Bruce, *Photosynth. Res.*, 1989, **20**, 1–34.
- 63 J. Biggins, N. A. Tanguay and H. A. Frank, *FEBS Lett.*, 1989, **250**, 271–274.
- 64 C. Verrotte, M. Picaud, D. Kirilovsky, J. Olive, G. Ajlani and C. Astier, *Photosynth. Res.*, 1992, **32**, 45–57.
- 65 K. El Bissati, E. Delphin, N. Murata, A.-L. Etienne and D. Kirilovsky, *Biochim. Biophys. Acta, Bioenerg.*, 2000, **1457**, 229–242.
- 66 S. Federman, S. Malkin and A. Scherz, *Photosynth. Res.*, 2000, **64**, 199.
- 67 L. Tian, I. H. van Stokkum, R. B. Koehorst, A. Jongerius, D. Kirilovsky and H. van Amerongen, *J. Am. Chem. Soc.*, 2011, **133**, 18304–18311.
- 68 H. Li, D. Li, S. Yang, J. Xie and J. Zhao, *Biochim. Biophys. Acta, Bioenerg.*, 2006, **1757**, 1512–1519.
- 69 C. W. Mullineaux and J. F. Allen, *FEBS Lett.*, 1986, **205**, 155–160.
- 70 M. Aoki and S. Katoh, *Biochim. Biophys. Acta, Bioenerg.*, 1982, **682**, 307–314.
- 71 C. W. Mullineaux, *Biochim. Biophys. Acta, Bioenerg.*, 2014, **1837**, 503–511.
- 72 M. Misumi, H. Katoh, T. Tomo and K. Sonoike, *Plant Cell Physiol.*, 2015, **57**, 1510–1517.
- 73 C. Huang, X. Yuan, J. Zhao and D. A. Bryant, *Biochim. Biophys. Acta, Bioenerg.*, 2003, **1607**, 121–130.
- 74 T. Ogawa, T. Harada, H. Ozaki and K. Sonoike, *Plant Cell Physiol.*, 2013, **54**, 1164–1171.
- 75 Y. V. Bolychevtseva, F. Kuzminov, I. Elanskaya, M. Y. Gorbunov and N. Karapetyan, *Biochemistry*, 2015, **80**, 50–60.
- 76 T. Ogawa and K. Sonoike, *J. Photochem. Photobiol., B*, 2015, **144**, 61–67.
- 77 C. Verrotte, C. Astier and J. Olive, *Photosynth. Res.*, 1990, **26**, 203–212.
- 78 D. Emlyn-Jones, M. K. Ashby and C. W. Mullineaux, *Mol. Microbiol.*, 1999, **33**, 1050–1058.
- 79 V. Chukhutsina, L. Bersanini, E.-M. Aro and H. Van Amerongen, *Sci. Rep.*, 2015, **5**, 14193.
- 80 I. N. Stadnichuk, E. P. Lukashev and I. V. Elanskaya, *Photosynth. Res.*, 2009, **99**, 227–241.
- 81 D. Bruce, S. Brimble and D. A. Bryant, *Biochim. Biophys. Acta, Bioenerg.*, 1989, **974**, 66–73.
- 82 D. Li, J. Xie, Y. Zhao and J. Zhao, *Biochim. Biophys. Acta, Bioenerg.*, 2003, **1557**, 35–40.
- 83 D. Li, J. Xie, J. Zhao, A. Xia, D. Li and Y. Gong, *Biochim. Biophys. Acta, Bioenerg.*, 2004, **1608**, 114–121.
- 84 C. Boulay, L. Abasova, C. Six, I. Vass and D. Kirilovsky, *Biochim. Biophys. Acta, Bioenerg.*, 2008, **1777**, 1344–1354.
- 85 C. Dong and J. Zhao, *Chin. Sci. Bull.*, 2008, **53**, 3422–3424.
- 86 C. W. Mullineaux, M. J. Tobin and G. R. Jones, *Nature*, 1997, **390**, 421.
- 87 M. Sarcina, M. J. Tobin and C. W. Mullineaux, *J. Biol. Chem.*, 2001, **276**, 46830–46834.
- 88 S. Yang, Z. Su, H. Li, J. Feng, J. Xie, A. Xia, Y. Gong and J. Zhao, *Biochim. Biophys. Acta, Bioenerg.*, 2007, **1767**, 15–21.
- 89 S. Joshua and C. W. Mullineaux, *Plant Physiol.*, 2004, **135**, 2112–2119.
- 90 M. Gwizdala, R. Berera, D. Kirilovsky, R. Van Grondelle and T. P. Krüger, *J. Am. Chem. Soc.*, 2016, **138**, 11616–11622.
- 91 D. Jallet, M. Gwizdala and D. Kirilovsky, *Biochim. Biophys. Acta, Bioenerg.*, 2012, **1817**, 1418–1427.
- 92 S. Joshua and C. W. Mullineaux, *Biochim. Biophys. Acta, Bioenerg.*, 2005, **1709**, 58–68.
- 93 G. Deng, F. Liu, X. Liu and J. Zhao, *FEBS Lett.*, 2012, **586**, 2342–2345.
- 94 J. Zhao, L. Chen, F. Gao, Q. Wang, Z. Qiu and W. Ma, *Acta Biochim. Biophys. Sin.*, 2014, **46**, 911–916.
- 95 C. W. Mullineaux, *Photosynth. Res.*, 2008, **95**, 175.
- 96 C. W. Mullineaux, *Biochim. Biophys. Acta, Bioenerg.*, 1994, **1184**, 71–77.
- 97 D. Bald, J. Kruip and M. Rögner, *Photosynth. Res.*, 1996, **49**, 103–118.
- 98 W. M. Schluchter, G. Shen, J. Zhao and D. A. Bryant, *Photochem. Photobiol.*, 1996, **64**, 53–66.
- 99 C. L. Aspinwall, M. Sarcina and C. W. Mullineaux, *Photosynth. Res.*, 2004, **79**, 179.



- 100 A. Strašková, G. Steinbach, G. Konert, E. Kotabová, J. Komenda, M. Tichý and R. Kaňa, *Biochim. Biophys. Acta, Bioenerg.*, 2019, DOI: 10.1016/j.bbabi.2019.07.008.
- 101 W. F. Vermaas, J. A. Timlin, H. D. Jones, M. B. Sinclair, L. T. Nieman, S. W. Hamad, D. K. Melgaard and D. M. Haaland, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 4050–4055.
- 102 D. M. Sherman, T. A. Troyan and L. A. Sherman, *Plant Physiol.*, 1994, **106**, 251–262.
- 103 S. Casella, F. Huang, D. Mason, G.-Y. Zhao, G. N. Johnson, C. W. Mullineaux and L.-N. Liu, *Mol. Plant*, 2017, **10**, 1434–1448.
- 104 G. Steinbach, F. Schubert and R. Kaňa, *J. Photochem. Photobiol., B*, 2015, **152**, 395–399.
- 105 R. Kaňa, E. Kotabová, O. Komárek, B. Šedivá, G. C. Papageorgiou and O. Prášil, *Biochim. Biophys. Acta, Bioenerg.*, 2012, **1817**, 1237–1247.
- 106 R. R. Choubeh, E. Wientjes, P. C. Struik, D. Kirilovsky and H. van Amerongen, *Biochim. Biophys. Acta, Bioenerg.*, 2018, **1859**, 1059–1066.
- 107 K. Kondo, C. W. Mullineaux and M. Ikeuchi, *Photosynth. Res.*, 2009, **99**, 217–225.
- 108 W. Zhao, J. Xie and J. Zhao, *Chin. Sci. Bull.*, 2014, **59**, 4712–4719.
- 109 D. Bruce, J. Biggins, T. Steiner and M. Thewalt, *Biochim. Biophys. Acta, Bioenerg.*, 1985, **806**, 237–246.
- 110 J. Olive, O. Vallon, F.-A. Wollman, M. Recouvreur and P. Bennoun, *Biochim. Biophys. Acta, Bioenerg.*, 1986, **851**, 239–248.
- 111 I. M. Folea, P. Zhang, E.-M. Aro and E. J. Boekema, *FEBS Lett.*, 2008, **582**, 1749–1754.
- 112 E. G. Maksimov, K. S. Mironov, M. S. Trofimova, N. L. Nechaeva, D. A. Todorenko, K. E. Klementiev, G. V. Tsoraev, E. V. Tyutyayev, A. A. Zorina and P. V. Feduraev, *Photosynth. Res.*, 2017, **133**, 215–223.
- 113 K. El Bissati and D. Kirilovsky, *Plant Physiol.*, 2001, **125**, 1988–2000.
- 114 C. W. Mullineaux, S. Griebenow and S. E. Braslavsky, *Biochim. Biophys. Acta*, 1991, **1060**, 315–318.
- 115 D. Bruce and O. Salehian, *Biochim. Biophys. Acta, Bioenerg.*, 1992, **1100**, 242–250.
- 116 W. P. Williams and P. J. Dominy, *Biochim. Biophys. Acta, Bioenerg.*, 1990, **1015**, 121–130.
- 117 H.-B. Mao, G.-F. Li, X. Ruan, Q.-Y. Wu, Y.-D. Gong, X.-F. Zhang and N.-M. Zhao, *FEBS Lett.*, 2002, **519**, 82–86.
- 118 A. G. Roberts, M. K. Bowman and D. M. Kramer, *Biochemistry*, 2004, **43**, 7707–7716.
- 119 J. Yan, G. Kurisu and W. A. Cramer, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 69–74.
- 120 J. F. Allen, C. E. Sanders and N. G. Holmes, *FEBS Lett.*, 1985, **193**, 271–275.
- 121 J. F. Allen and N. G. Holmes, *FEBS Lett.*, 1986, **202**, 175–181.
- 122 M. A. Harrison, N. F. Tsinoremas and J. F. Allen, *FEBS Lett.*, 1991, **282**, 295–299.
- 123 M.-k. Yang, Z.-x. Qiao, W.-y. Zhang, Q. Xiong, J. Zhang, T. Li, F. Ge and J.-d. Zhao, *J. Proteome Res.*, 2013, **12**, 1909–1923.
- 124 Z. Chen, J. Zhan, Y. Chen, M. Yang, C. He, F. Ge and Q. Wang, *Plant Cell Physiol.*, 2015, **56**, 1997–2013.
- 125 P. Spaet, B. Maček and K. Forchhammer, *Fron. Microbiol.*, 2015, **6**, 248.
- 126 M. Nanba and S. Katoh, *Biochim. Biophys. Acta, Bioenerg.*, 1985, **809**, 74–80.
- 127 R. Vladkova, *J. Biomol. Struct. Dyn.*, 2016, **34**, 824–854.
- 128 C. E. Sanders, A. Melis and J. F. Allen, *Biochim. Biophys. Acta, Bioenerg.*, 1989, **976**, 168–172.
- 129 J. F. Allen, *Biochim. Biophys. Acta, Bioenerg.*, 1992, **1098**, 275–335.
- 130 J. Biggins, C. L. Campbell and D. Bruce, *Biochim. Biophys. Acta, Bioenerg.*, 1984, **767**, 138–144.
- 131 A. Krupa and N. Srinivasan, *BMC Genomics*, 2005, **6**, 129.
- 132 J. Perez, A. Castaneda-Garcia, H. Jenke-Kodama, R. Müller and J. Munoz-Dorado, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 15950–15955.
- 133 N. H. Mann, *Microbiology*, 1994, **140**, 3207–3215.
- 134 T. Kaneko, S. Sato, H. Kotani, A. Tanaka, E. Asamizu, Y. Nakamura, N. Miyajima, M. Hirosawa, M. Sugiura and S. Sasamoto, *DNA Res.*, 1996, **3**, 109–136.
- 135 C.-C. Zhang, L. Gonzalez and V. Phalip, *Nucleic Acids Res.*, 1998, **26**, 3619–3625.
- 136 A. Kamei, T. Yuasa, X. Geng and M. Ikeuchi, *DNA Res.*, 2002, **9**, 71–78.
- 137 C. J. Leonard, L. Aravind and E. V. Koonin, *Genome Res.*, 1998, **8**, 1038–1047.
- 138 A. Zorina, *Russ. J. Plant Physiol.*, 2013, **60**, 589–596.
- 139 A. Kamei, T. Yuasa, K. Orikawa, X. X. Geng and M. Ikeuchi, *J. Bacteriol.*, 2001, **183**, 1505–1510.
- 140 A. Mata-Cabana, M. García-Domínguez, F. J. Florencio and M. Lindahl, *Antioxid. Redox Signaling*, 2012, **17**, 521–533.
- 141 C. Liang, X. Zhang, X. Chi, X. Guan, Y. Li, S. Qin and H. bo Shao, *PLoS One*, 2011, **6**, e18718.
- 142 A. Zorina, V. Bedbenov, G. Novikova and V. Panichkin, *Mol. Biol.*, 2014, **48**, 390–398.
- 143 S. Laurent, J. Jang, A. Janicki, C.-C. Zhang and S. Bedu, *Microbiology*, 2008, **154**, 2161–2167.
- 144 A. Galkin, L. Mikheeva and S. Shestakov, *Mikrobiologiya*, 2003, **72**, 64–69.
- 145 A. Zorina, N. Stepanchenko, G. V. Novikova, M. Sinetova, V. B. Panichkin, I. E. Moshkov, V. V. Zinchenko, S. V. Shestakov, I. Suzuki and N. Murata, *DNA Res.*, 2011, **18**, 137–151.
- 146 M. Angeleri, A. Zorina, E. M. Aro and N. Battchikova, *FEBS Lett.*, 2018, **592**, 411–421.
- 147 G. Schansker, S. Z. Tóth, L. Kovács, A. R. Holzwarth and G. Garab, *Biochim. Biophys. Acta, Bioenerg.*, 2011, **1807**, 1032–1043.
- 148 M. Magyar, G. Sipka, L. Kovács, B. Ughy, Q. Zhu, G. Han, V. Špunda, P. H. Lambrev, J.-R. Shen and G. Garab, *Sci. Rep.*, 2018, **8**, 2755.



- 149 G. Sipka, P. Müller, K. Brettel, M. Magyar, L. Kovács, Q. Zhu, Y. Xiao, G. Han, P. H. Lambrev and J. R. Shen, *Physiol. Plant.*, 2019, **166**, 22–32.
- 150 G. Schansker, S. Z. Tóth, A. R. Holzwarth and G. Garab, *Photosynth. Res.*, 2014, **120**, 43–58.
- 151 Y. Shibata, S. Nishi, K. Kawakami, J.-R. Shen and T. Renger, *J. Am. Chem. Soc.*, 2013, **135**, 6903–6914.
- 152 A. Guskov, J. Kern, A. Gabdulkhakov, M. Broser, A. Zouni and W. Saenger, *Nat. Struct. Mol. Biol.*, 2009, **16**, 334.
- 153 J.-Y. Huang, Y.-F. Chiu, J. M. Ortega, H.-T. Wang, T.-S. Tseng, S.-C. Ke, M. Roncel and H.-A. Chu, *Biochemistry*, 2016, **55**, 2214–2226.
- 154 B. E. Rubin, K. M. Wetmore, M. N. Price, S. Diamond, R. K. Shultzaberger, L. C. Lowe, G. Curtin, A. P. Arkin, A. Deutschbauer and S. S. Golden, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, E6634–E6643.
- 155 S. Yang, R. Zhang, C. Hu, J. Xie and J. Zhao, *Photosynth. Res.*, 2009, **99**, 99–106.
- 156 T. Fujimori, Y. Hihara and K. Sonoike, *J. Biol. Chem.*, 2005, **280**, 22191–22197.
- 157 D. Bruce and J. Biggins, *Biochim. Biophys. Acta, Bioenerg.*, 1985, **810**, 295–301.
- 158 O. Salehian and D. Bruce, *J. Lumin.*, 1992, **51**, 91–98.

