RSC Advances



PAPER View Article Online



Cite this: RSC Adv., 2019, 9, 39684

Microwave-assisted synthesis of 7-azaindoles *via* iron-catalyzed cyclization of an *o*-haloaromatic amine with terminal alkynes†

Yi Le, abc Zhisong Yang, Yumei Chen, Dongmei Chen, Longjia Yan, Longjia Yan, Uhanchao Wangbc and Guiping Ouyang Yabc

Received 24th October 2019 Accepted 26th November 2019

DOI: 10.1039/c9ra08742g

rsc.li/rsc-advances

An efficient and practical procedure was developed to prepare 7-azaindole, starting from an o-haloaromatic amine and corresponding terminal alkynes under microwave irradiation and the scope was demonstrated with a number of examples. The valuable features of this procedure included the iron-catalyzed cyclization, short reaction times and convenient operation. Furthermore, iron catalysis is an interesting alternative to homogeneous catalysis for the synthesis of heterocycles.

Introduction

In the recent years, microwave (MW) irradiation in synthetic organic chemistry has attracted considerable practical and theoretical attention as a very effective and non-polluting method of activation. Microwave irradiation often helps to reduce reaction times, minimize side products, increase yields and improve reproducibility. Therefore, the technique of microwave-assisted organic synthesis has been widely and successfully employed in dipolar cycloadditions, transition metal-catalyzed cross-coupling reactions and polymer formation.

1*H*-Pyrrolo[2,3-*b*]pyridines, often referred to as 7-azaindoles, are bioisosteres of the indole scaffold have been used in diverse areas such as materials and medicinal chemistry due to their physicochemical and pharmacological properties.⁵ For example, the natural product Variolin B⁵ in Fig. 1 isolated from an extremely rare antarctic sponge is a promising anti-cancer agent. PLX5622 in Fig. 1,⁶ a brain pentrant CSF1R inhibitor has been used in Alzheimer's disease (AD). Recently, the 7-azaindole derivatives were also shown potential anti-cancer activity, including AZD6738 a potent and selective ATR kinase inhibitor⁷ and GSK1070916 an aurora kinase inhibitor.⁸ Thus, development of synthetic methods to access these 7-azaindoles is of great importance to the drug discovery community.⁹

Our group has been focused on metal-catalyzed crosscoupling reactions for the preparation of bioactive heterocycles. 10 Based on the broad activities of 7-azaindole, we are interested in developing new strategies to prepare this novel heterocyclic scaffold. In the literature, the common synthetic methods to prepare azaindoles usually start from commercial aminopyridines, followed by Sonogashira alkynylation by subsequent base-mediated cyclization with various base for building up the pyrrole ring.11 As shown in Fig. 2, these methods were always used Pd-catalysts, such as Pd(PPh3)2Cl2,12a Pd2(dba)3,126 and Pd-NaY,126 and also have some disadvantages such as the high price of palladium and problem of pollution. Recently, the establishment of new catalytic methods using iron is attractive owing to the low cost, abundance, ready availability, and very low toxicity of iron.13 Furthermore, due to the availability of several catalytic cycles, iron catalysis offers a potential for orthogonal selectivity when compared to other metals,




Fig. 1 Bioactive skeletons containing 7-azaindole framework.

[&]quot;State Key Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering, Key Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, State-Local Joint Laboratory for Comprehensive Utilization of Biomass, Center for Research and Development of Fine Chemicals, Guizhou University, Guiyang 550025, China

^bSchool of Pharmaceutical Sciences, Guizhou University, Guiyang 550025, China. E-mail: ylj1089@163.com; oygp710@163.com

^cGuizhou Engineering Laboratory for Synthetic Drugs, Guiyang 550025, China

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra08742g

Fig. 2 Different strategies for synthesis of 7-azaindoles.

which allows for streamlined construction of complex molecules by modern cross-coupling chemistry.¹⁴ Encouraged by the development of iron-catalytic Sonogashira alkynylation¹⁵ and

the microwave-assisted synthesis of indole, ¹⁶ we wish to report the iron-catalyzed cyclization of *o*-haloaromatic amine with terminal alkynes under the microwave irradiation.

Results and discussion

Initially, we chose the commercially available 3-iodo-pyridin-2vlamine 1a and ethynyl-benzene 2a as model starting materials. We examined the reaction conditions using 10% Pd(PPh₃)₂Cl₂, 10% CuI, 1.5 equiv. of K₃PO₄ in DMF under microwave irradiation of 100 °C, the reaction proceeded for 30 min and product 3a was obtained in 12%, which was purified the intermediate 3aa for 35% yield (Table 1, entry 1). We subsequently investigated the effect of various catalysts including palladium, silver and iron (Table 1, entries 2-6). Fortunately, we found that Fe(acac)3 was given the best result (Table 1, entry 6). We also tested the amount of CuI, it was no product without CuI and the similar yield using 20% CuI (Table 1, entries 7 and 8). Another blank test was performed only with CuI and shown that 3aa was obtained in 8% yield (Table 1, entry 9). It was no reaction at the absence of Fe(acac)₃ and CuI (Table 1, entry 10). Replacement of K₃PO₄ by KOAc, or K₂CO₃ led to lower yields and KO^tBu gave a satisfactory yield (Table 1, entries 11-13). Next, we screened different organic solvents and

Table 1 Optimization of the metal-catalyzed cyclization 3a^a

Entry	Catalyst	CuI (%)	Base	Solvent	T (°C)	Yield ^b (%)
1	$Pd(PPh_3)_2Cl_2$	10	K_3PO_4	DMF	100	12
2	$Pd(dppf)Cl_2$	10	K_3PO_4	DMF	100	21
3	$AgNO_3$	10	K_3PO_4	DMF	100	Trace
4	AgOAc	10	K_3PO_4	DMF	100	Trace
5	FeCl_3	10	K_3PO_4	DMF	100	11
6	$Fe(acac)_3$	10	K_3PO_4	DMF	100	32
7	Fe(acac) ₃	_	K_3PO_4	DMF	100	0
8	Fe(acac) ₃	20	K_3PO_4	DMF	100	31
9		10	K_3PO_4	DMF	100	<u></u> c
10	_	_	K_3PO_4	DMF	100	0
11	$Fe(acac)_3$	10	KOAc	DMF	100	28
12	Fe(acac) ₃	10	K_2CO_3	DMF	100	25
13	Fe(acac) ₃	10	KO^t Bu	DMF	100	38
14	Fe(acac) ₃	10	KO^t Bu	DMSO	100	36
15	Fe(acac) ₃	10	KO^t Bu	NMP	100	42
16	Fe(acac) ₃	10	KO^t Bu	Dioxane	100	22
17	Fe(acac) ₃	10	KO^t Bu	NMP	110	58
18	Fe(acac) ₃	10	KO^t Bu	NMP	120	61
19	$Fe(acac)_3$	10	KO^t Bu	NMP	130	62
20	Fe(acac) ₃	10	KO^t Bu	NMP	140	58
21	$Pd(PPh_3)_2Cl_2$	10	KO^t Bu	NMP	130	44
22	$Pd(dppf)Cl_2$	10	KO^t Bu	NMP	130	48

^a Reagents and conditions: **1a** (1 mmol), **2a** (2 mmol), catalyst (0.1 mmol), CuI (0.1 mmol), base (1.5 mmol), solvent (2 mL), 30 min MW, 100 °C. ^b Isolated yields. ^c Only **3aa** was obtained with 8% yield.

RSC Advances

identified NMP as the most effective solvent (Table 1, entry 15), while dioxane resulted in a poor yield (Table 1, entry 16). The yield could be further improved by increasing the temperature to a maximum of 130 °C (Table 1, entry 19), whereas higher temperatures led to partial decomposition (Table 1, entry 20). Interestingly, when we used $Pd(PPh_3)_2Cl_2$ reported by Mueller^{12a} or $Pd(dppf)Cl_2$ under the optimized condition, the yield was 44% and 48%, respectively (Table 1, entries 21 and 22).

In order to optimize the reaction conditions, we speculated that the equivalent of alkyne and microwave time might also be efficient in promoting the microwave-assisted cyclized reaction. After exploring several reaction conditions (Table 2), **3a** was obtained in 72% yield by heating 3 equiv. **2a**, 0.1 equiv. Fe(acac)₃, 0.1 equiv. CuI, 1.5 equiv. KO^tBu in NMP for 60 min at 130 °C (Table 2, entry 4). We also tried the reaction under conventional thermal heating condition and the yield of the desired product was 33% (Table 2, entry 7).

Having determined the optimal reaction conditions (Table 2, entry 4), the scope of the reaction was explored with different substituted 3-iodo-pyridin-2-ylamine 1 (Table 3), which were prepared by the reaction of pyridin-2-ylamines with Ag₂SO₄ and I₂, to afford various 7-azaindoles 3. As illustrated in Table 3, a variety of substituted 7-azaindoles was readily prepared in good yield through this protocol, and assigned by spectroscopic methods (¹H NMR and ¹³C NMR) and HRMS. Both electrondonating group -CH3 (Table 3, entries 3 and 4) and electronwithdrawing group -CN (Table 3, entries 5-8) of R1 are tolerated at a variety of positions on the aromatic ring. Alkynes bearing benzene ring such as 4-OCH3 and 4-F were also tolerated under reaction conditions as 3b and 3d were afforded in 73% and 68% yields, respectively. Additionally, aliphatic groups afforded the corresponding products with decreased yields (3h). This result was consistent with the observation by Mueller et al. when Pd(PPh₃)₂Cl₂/(1-Ad)₂PBn was used as catalyst. 12a The introduction of trifluoromethyl group into organic compounds frequently renders remarkable changes in their special size,

Table 2 Optimization of the iron-catalyzed cyclization 3a^a

Entry	2a (equiv.)	Time (min)	Yield ^b (%)
1	2	30	62
2	3	30	68
3	4	30	66
4	3	60	72
5	3	90	71
6	3	120	58
7	3	120	58 33 ^c

^a Reagents and conditions: **1a** (1 mmol), **2a**, Fe(acac)₃ (0.1 mmol), CuI (0.1 mmol), KO^tBu (1.5 mmol), NMP (2 mL), MW, 130 °C. ^b Isolated yields. ^c Under conventional thermal heating condition.

bioactivity and metabolic stability.¹⁷ In this paper, we also show such examples of the CF₃ substituted 7-azaindoles (Table 3, entries 9–16). We found that the reaction conditions were adaptable with phenyl (3i) and electron-donating substituents at the aryl alkynes, such as methoxy group (3k). In addition to electron-donating substituent, halogens such as F and Cl were also tolerated (3j, 3m and 3n). However, aryl groups with an electron deficient substituent led to a lower yield (3l). Additionally, aliphatic groups afforded the corresponding products with reduced yields (3o and 3p).

Finally, this procedure was applied to the preparation of 1,2disubstituted 7-azaindoles 5. As shown in Table 4, the Sonogashira and consequently cyclization could be used for preparation the N-arylation of azaindoles. The electron-donating groups on the phenyl acetylene, such as OCH₃ (5c and 5m) were obtained the much higher yield of 78% than the electronwithdrawing group F (5b and 5i), Cl (5d, 5k and 5l) and CF₃ (**5e** and **5j**). 3,3,-Dimethyl-but-1-yne led to 2-*tert*-butyl-1-(3,4,5trimethoxy-phenyl)-1*H*-pyrrolo[2,3-*b*]pyridine 5f (42%), the decrease in yield may be due to decomposition of compound 2 under the MW conditions. Interestingly, the ethyl ester substituted group 5g was also separated with 33% yield under our MW protocol. In addition to substituted anyl group of R³, the alkyl-substituted compound 4 were also compatible with this process, furnishing the desired product 5n and 5o in 34% and 38% yield.

To investigate the mechanism of this transformation, experiments were carried out. The desired product 3a was

Table 3 Preparation of 7-azaindoles 3a-p^a

$$R^{1}$$
 $=$ R^{2} R^{2} R^{1} $=$ R^{2} R^{1} $=$ R^{2} R^{1} $=$ R^{2} R^{2} $=$ R^{2} $=$

Entry R_1	R_2	Product	Yield ^b (%)
1 H	Ph	3a	72
2 H	4-OCH ₃ -Ph	3 b	73
3 Me	Ph	3c	73
4 Me	4-F-Ph	3d	68
5 CN	Ph	3e	65
6 CN	4-F-Ph	3f	61
7 CN	4-OCH ₃ -Ph	3g	66
8 CN	Me	3h	43
9 CF ₃	Ph	3i	62
10 CF ₃	4-F-Ph	3j	56
11 CF ₃	4-OCH ₃ -Ph	3k	79
12 CF ₃	4-CF ₃ -Ph	31	45
13 CF ₃	4-Cl-Ph	3m	58
14 CF ₃	2-Cl-Ph	3n	32
15 CF ₃	Ме	30	33
16 CF ₃	t-Butyl	3 p	48

 $[^]a$ Reagents and conditions: 1 (1 mmol), 2 (3 mmol), Fe(acac) $_3$ (0.1 mmol), CuI (0.1 mmol), KO t Bu (1.5 mmol), NMP (2 mL), 60 min MW, 130 $^\circ$ C. b Isolated yields.

obtained in 70% yield under standardized reaction conditions (Scheme 1). We also tested the absence of Fe(acac)₃ and CuI, it was 22% yield when **3aa** and KO⁶Bu were used in NMP under MW conditions for 60 min. Base on the observed experimental results and pioneering reports, ^{15,18} we have described a plausible mechanistic pathway in Scheme 2. Oxidation addition of Fe(acac)₃ by the amine and pyridine functional units resulting in an organoiron complex **A**. At this stage, trans-metalation by copper complex **B** provides an intermediate Fe-species **C**, which could then undergo reductive elimination to forge the new carbon–carbon triple bond, thus yielding the compound **3aa**. After furnished the typical Sonogashira coupling, the

Table 4 Preparation of 1,2-disubstituted 7-azaindoles 5a-m^a

 a Reagents and conditions: 4 (1 mmol), 2 (3 mmol), Fe(acac) $_3$ (0.1 mmol), CuI (0.1 mmol), KO t Bu (1.5 mmol), NMP (2 mL), 60 min MW, 130 $^\circ$ C.

Scheme 1 Complementary experiments.

Scheme 2 Proposed mechanism.

desired 7-azaindole was generated under the microwave condition.

Conclusions

In summary, we have described the synthesis of biologically relevant 7-azaindole ring systems using an iron-catalyzed cyclic reaction under microwave irradiation. This approach provides a rapid and economical access to a diverse range of 7-azaindoles. The method employs readily available reagents and possesses broad scope and good tolerance of functional group, iron catalysis could be an interesting. Further efforts to utilize these compounds as versatile building blocks for assembling interesting heterocyclic molecules which can be applied in medicinal chemistry research are currently underway in our laboratories.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This project is supported by the National Natural Science Foundation of China (21867004), State Key Laboratory of Functions and Applications of Medicinal Plants (No. FAMP201707K) and (No. FAMP201801K).

RSC Advances Paper

Notes and references

- (a) D. Dallinger and C. O. Kappe, Chem. Rev., 2007, 107, 2563;
 (b) P. Appukkuttan, V. P. Mehta and E. V.-d. Eycken, Chem. Soc. Rev., 2010, 39, 1467;
 (c) M. Driowya, A. Saber, H. Marzaq, L. Demange, K. Bougrin and R. Benhida, Molecules, 2016, 21, 1032.
- 2 Z. Yang, J. Wang, L. Peng, X. You and H. Cui, *Tetrahedron Lett.*, 2016, 57, 5219.
- 3 D. Roy and Y. Uozumi, Adv. Synth. Catal., 2018, 360, 602.
- 4 M. B. Schuetz, L. Xiao, T. Lehnen, T. Fischer and S. Mathur, *Int. Mater. Rev.*, 2018, **63**, 341.
- 5 S. R. Walker, E. J. Carter, B. C. Huff and J. C. Morris, *Chem. Rev.*, 2009, **109**, 3080.
- 6 E. Spangenberg, P. L. Severson, L. A. Hohsfield, J. Crapser, J. Zhang, E. A. Burton, Y. Zhang, W. Spevak, J. Lin, N. Y. Phan, G. Habets, A. Rymar, G. Tsang, J. Walters, M. Nespi, P. Singh, S. Broome, P. Ibrahim, C. Zhang, G. Bollag, B. L. West and K. N. Green, *Nat. Commun.*, 2019, 10, 3758.
- 7 R. F. Goundry, K. Dai, M. Gonzalez, D. Legg, A. Kearney-McMullan, J. Morrison, A. Stark, P. Siedlecki, P. Tomlin and J. Yang, *Org. Process Res. Dev.*, 2019, 23, 1333.
- 8 N. D. Adams, J. L. Adams, J. L. Burgess, A. M. Chaudhari, R. A. Copeland, C. A. Donatelli, D. H. Drewry, K. E. Fisher, T. Hamajima, M. A. Hardwicke, W. F. Huffman, K. K. Koretke-Brown, Z. V. Lai, O. B. McDonald, H. Nakamura, K. A. Newlander, C. A. Oleykowski, C. A. Parrish, D. R. Patrick, R. Plant, M. A. Sarpong, K. Sasaki, S. J. Schmidt, D. J. Silva, D. Sutton, J. Tang, C. S. Thompson, P. J. Tummino, J. Wang, H. Xiang, J. Yang and D. Dhanak, J. Med. Chem., 2010, 53, 3973.
- 9 (a) S. Zhao and S. Wang, Chem. Soc. Rev., 2010, 39, 3142; (b)
 W. D. Tap, Z. A. Wainberg, S. P. Anthony, P. N. Ibrahim,
 C. Zhang, J. H. Healey, B. Chmielowski, A. P. Staddon,

- A. L. Cohn, G. I. Shapiro, V. L. Keedy, A. S. Singh, I. Puzanov, E. L. Kwak, A. J. Wagner, D. D. von-Hoff, G. J. Weiss, R. K. Ramanathan, J. Zhang, G. Habets, Y. Zhang, E. A. Burton, G. Visor, L. Sanftner, P. Severson, H. Nguyen, M. J. Kim, A. Marimuthu, G. Tsang, R. Shellooe, C. Gee, B. L. West, P. Hirth, K. Nolop, M. vande-Rijn, H. H. Hsu, C. Peterfy, P. S. Lin, S. Tong-Starksen and G. Bollag, N. Engl. J. Med., 2015, 373, 428; (c) T. Irie and M. Sawa, Chem. Pharm. Bull., 2018, 66, 29; (d) W. Li and L. Dong, Adv. Synth. Catal., 2018, 360, 1104; (e) N. Sharma and Anurag, Mini-Rev. Med. Chem., 2019, 19, 727.
- 10 (a) L. Yan, H. Wang, Y. Chen, Z. Li and Y. Pei, Mol. Diversity, 2018, 22, 323; (b) L. Yan, M. Deng, A. Chen, Y. Li, W. Zhang, Z. Du, C. Dong, B. Meunier and H. Chen, Tetrahedron Lett., 2019, 60, 1359.
- 11 F. Popowycz, J. Y. Merour and B. Joseph, *Tetrahedron*, 2007, 63, 8689.
- 12 (a) T. Lessing, F. Sterzenbach and T. J. J. Mueller, *Synlett*,
 2015, 26, 1217; (b) S. I. Purificacao, M. J. D. Pires,
 R. Rippel, A. S. Santos and M. M. B. Marques, *Org. Lett.*,
 2017, 19, 5118; (c) E. K. Yum and K. B. Hong, *Tetrahedron*,
 2017, 73, 6581.
- 13 T. Cernak, K. D. Dykstra, S. Tyaqarajan, P. Vachal and S. W. Krska, Chem. Soc. Rev., 2016, 45, 546.
- 14 A. Piontek, E. Bisz and M. Szostak, *Angew. Chem., Int. Ed. Engl.*, 2018, 57, 11116.
- 15 J. Mao, G. Xie, M. Wu, J. Guo and S. Ji, *Adv. Synth. Catal.*, 2008, **350**, 2477.
- 16 A. Carpita, A. Ribecai and P. Stabile, *Tetrahedron*, 2010, 66, 7169.
- 17 T. Furuya, A. S. Kamlet and T. Ritter, Nature, 2011, 473, 470.
- 18 (a) S. Handa, B. Jin, P. P. Bora, Y. Wang, X. Zhang, F. Gallou,
 J. Reilly and B. H. Lipshutz, ACS Catal., 2019, 9, 2423; (b)
 K. S. Sindhu, A. P. Thankachan, A. M. Thomas and
 G. Anilkumar, ChemistrySelect, 2016, 1, 556.