RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2022, 12, 21022

Metal-free synthesis of C2-quaternary indolinones by $(NH_4)_2S_2O_8$ mediated oxidative dearomatization of indoles†

Yu Dong, (1) ** Jun-Hu Qian, ** Xiang-Long Chen, ** Hui Jiang, ** Xue Li, ** Qiang Zhou, ** Ting Mei, ** Zhi-Chuan Shi, ** Zhong-Hui Li** and Bing He**

An efficient metal-free, $(NH_4)_2S_2O_8$ mediated oxidative dearomatization of indoles for the construction of C2-quaternary indolinones was disclosed. A series of C2-quaternary indolinones derivatives with good functional group tolerance were obtained in moderate to excellent yields. This methodology provides an alternative approach for the direct generation of all-carbon quaternary centers at the C2 position of indoles. This catalytic approach represents a step-economic and convenient strategy for the oxidative dearomatization of indoles.

Received 7th July 2022 Accepted 15th July 2022

DOI: 10.1039/d2ra04191j

rsc.li/rsc-advances

Introduction

N-Bonded heterocycles are privileged moieties in the molecular skeleton of various natural products and pharmaceuticals.1 Indole and its derivatives are important compounds, a subset of N-bonded heterocycles, are widespread in nature and exhibit significant biological activity² and oxindole derivatives (Fig. 1) are known to possess a variety of biological activities.3 Oxidative dearomatization reaction of indole leads to the formation of a diverse class of products such as 2/3-oxoindoles, indirubin, indigo, isatin and indoline derivatives.4 Among them, the transformation of indoles to C2/C3-quaternary indolinone derivatives is synthetically quite valuable since it converts structurally simpler, planar indole skeleton to a complex threedimensional architecture. For example, 2-(1H-indol-3-yl)-2,3'biindolin-3-one 1 was isolated as the product of indole oxidation by a strain of Claviceps purpurea.5 This compound has also been characterised from natural (bacterial) sources such as Vibrio parahaemolyticus⁶ and Haemophilus influenzae.⁷ In addition, isatisine A 2, an oxindole system having indole 2-substituents, is present in the roots and leaves of Isatis indigotica Fort (Cruciferae). This biennial herbaceous plant is widely cultivated in China and East Asia for the prevention and treatment of viral diseases such as influenza, viral pneumonia, mumps, and hepatitis.8 Therefore, we believe that synthesis of 2,2-disubstituted indolin-3-ones derivatives is valuable.




Fig. 1 Some important bioactive synthetic derivatives with 2,2-disubstituted indolin-3-one structural unit.

In 2012, Liu and coworkers reported the TEMPO [(2,2,6,6tetramethylpiperidin-1-yl)oxyl] mediated synthesis of C2quaternary indolinones from free (NH)-indoles (Scheme 1a).9 However, the reaction required three days and the high loading of TEMPO for completion. Subsequently, in 2013, Liu and coworkers reported a tandem oxidative homocoupling reaction for the generation of all-carbon quaternary centers at the C2 position of indoles by using NaNO2 with CH3SO3H in pyridine (Scheme 1b).10 In 2018, Ganesan and coworkers reported one pot oxidative dearomatization reaction of indole leading to the formation of the corresponding C2/C3-quaternary indolinones (Scheme 1c).11 Free (NH)-indoles gave C2-quaternary indolinone derivatives whilst (NR)-indoles yielded C3-quaternary indolinones as the major product. In 2020, Thakur and coworkers reported 'on-water' synthesis of 2,2-bis(indoly-3-yl)indoline-3ones via N2-selective dearomatization of '(N-H) protectionfree' indole derivatives (Scheme 1d).12 To the best of our knowledge, there is only limited precedent are available for the one-pot transformation of indole to the corresponding C2 quaternary indolinone derivatives, with no reported study for (NH₄)₂S₂O₈-catalysed methodologies. One relevant piece of work is by Ganesan et al. (Scheme 1), who described the TBHP

^aCollege of Chemistry and Life Science, Sichuan Provincial Key Laboratory for Structural Optimization and Application of Functional Molecules, Chengdu Normal University, Chengdu, 611130, P. R. China

^bSouthwest Minzu University, Chengdu 610041, P. R. China

Paper

Scheme 1 Previous and present approaches for the synthesis of 2,2-diaryloxindole (R = alkyl or halide group).

reagent system to convert free indoles to the corresponding C2-quaternary indolinone derivatives. Therefore, development of methods to meet the challenges of optimum reaction conditions would significantly enhance the utility of 2,2-diary-loxindoles. Herein, we describe successful implementation of the strategy shown in Scheme 1, which affords 2,2-bis(indoly-3-yl)indoline-3-ones from indoles using $(NH_4)_2S_2O_8$ in DMF as oxidant.

Results and discussion

The reaction conditions were firstly optimized by 1-methyl-1H-indole **1a** as the template substrates. As shown in Table 1, the model reaction employed (NH₄)₂S₂O₈ as oxidant in DMF (2 ml) as the solvent at 60 °C for 3 h, giving the desired product of 2,2-bis(indoly-3-yl)indoline-3-ones **2a** in a yield of 89% (entry 1). Subsequently, the effects of different oxidants on the reaction were investigated. For instance, K₂S₂O₈, oxone, H₂O₂, and TBHP were applied as oxidants for the one-pot reaction, respectively (entries 2–5). Among those, (NH₄)₂S₂O₈ displayed the highest activity, and the yield of **2a** was 89% (entry 1). In addition, **4a** was not observed in the absence of the oxidant, suggesting the importance of (NH₄)₂S₂O₈ as oxidant (entry 6). Increasing or

Table 1 Optimization of the reaction conditions^a

Entry	[Cat.]	Solvent	Temp. (°C)	Yield ^b (%)
1	$(NH_4)_2S_2O_8(2)$	DMF (2)	60	89
2	$K_2S_2O_8(2)$	DMF (2)	60	72
3	Oxone (2)	DMF (2)	60	38
4	$H_2O_2(2)$	DMF (2)	60	ND^c
5	TBHP (2)	DMF (2)	60	NR^d
6	No	DMF (2)	60	NR
7	$(NH_4)_2S_2O_8(1)$	DMF (2)	60	46
8	$(NH_4)_2S_2O_8(3)$	DMF (2)	60	52
9	$(NH_4)_2S_2O_8(2)$	DCM (2)	60	Trace
10	$(NH_4)_2S_2O_8(2)$	DCE (2)	60	10
11	$(NH_4)_2S_2O_8(2)$	$PhCH_3(2)$	60	22
12	$(NH_4)_2S_2O_8(2)$	MeCN (2)	60	ND
13	$(NH_4)_2S_2O_8(2)$	DMSO (2)	60	NR
14	$(NH_4)_2S_2O_8(2)$	EtOH (2)	60	71
15	$(NH_4)_2S_2O_8(2)$	DMF (1)	60	60
16	$(NH_4)_2S_2O_8(2)$	DMF (3)	60	32
17	$(NH_4)_2S_2O_8(2)$	DMF (2)	40	42
18	$(NH_4)_2S_2O_8(2)$	DMF (2)	80	45
19^e	$(NH_4)_2S_2O_8(2)$	DMF (2)	60	28
20^f	$(NH_4)_2S_2O_8(2)$	DMF(2)	60	50

^a Reaction conditions: 1-methyl-1*H*-indole **1a** (0.3 mmol), [oxidants] (equiv.), solvent (mL), T °C, air tube. ^b Isolated yields. ^c N.D. = no detected. ^d NR means not reaction. ^e Reaction for 1 h. ^f Reaction for 5 h. H₂O₂: 30% aqueous solution. TBHP: 70% aqueous solution.

decreasing the amount of $(NH_4)_2S_2O_8$ resulted in the decrease of yield (entry 7 and 8). Next, the solvent was screened. DMF was proved to be more efficient than others, such as DCM, DCE, PhCH₃, MeCN, DMSO or EtOH (entries 9–14). Changing the amount of DMF led to the reduced of the product yield (entries 15 and 16). The heating effect was also evaluated for this reaction and the reaction led to maximum yield at 60 °C while increasing or decreasing the temperature was not fruitful for this process (entry 17 and 18). Further investigation of the reaction for a shorter or longer time period (1 or 5 h) did not give a higher yield of 2a (entries 19 and 20). Finally, the approach employed $(NH_4)_2S_2O_8$ as oxidant in DMF (2 ml) as the solvent at 60 °C for 3 h, giving the desired product of 2,2-bis(indoly-3-yl)indoline-3-ones 2a in a yield of 89% (entry 1).

After achieving the optimized conditions, a series of substituted indole derivatives were tested for the oxidative dearomatization (Table 2). As summarized in Table 2, the reaction was compatible with a variety of indole moieties (1a–1u) bearing electron-donating and electron-withdrawing substituents to produce the desired C2-quaternary indolinones products (2a–2u) in moderate to good yields (73–89%). The structure of the desired C2-quaternary indolinones 2a was

Table 2 Scope for indoles^a

$$\begin{array}{c|c}
 & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \circ & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \bullet & \circ & \circ \\
 & \bullet & \bullet & \circ & \circ & \circ \\
 & \bullet & \bullet & \bullet & \circ & \circ \\
 & \bullet & \bullet & \bullet & \circ & \circ \\
 & \bullet & \bullet & \bullet & \circ & \circ \\
 & \bullet$$

Scheme 2 Plausible reaction mechanism

confirmed by X-ray analysis. Notably, the gram-scale synthesis afforded 1.05 g of 2a in 86% yield. To our delight, the reaction also showed good compatibility with a wide range of valuable functional groups such as chloro (2n and 2r) and bromo (2o and 2u). Tolerance to the halogen atoms was noteworthy since they have been frequently used for further modification. N-Alkylindoles can smoothly proceed to give the corresponding products in good yields (2a-2h). The 2-methylindoles and 2-phenylindoles gave the corresponding unsymmetrically substituted C2-quaternary indolinone product 2i-2l in 74-80% yield. Moreover, we were pleased to find that the position of the substituent on the indole moiety showed no obvious influence on the reaction outcome, and substitutions at the C5-(2m-2q), C6-(2r and 2s), or C7-(2t and 2u) were all well tolerated in the reaction.

On the basis of substrate diversity and previous reports, 13 the attack of the $(NH_4)_2S_2O_8$ onto the nucleophilic centre C-3 of indole (1a) generates intermediate (B) *in situ* through the various intermediate compounds (Scheme 2). As depicted in Scheme 2, the electrophilic intermediate (B) easily facilitates nucleophilic attack by the indole molecule (1a) to form intermediate compound C on oxidation condition. Subsequently, the electrophilic intermediate (C) easily facilitates nucleophilic attack by the indole molecule (1a) to form the C2-quaternary indolinone (1a)

Conclusions

In conclusion, we have successfully demonstrated an efficient metal-free, $(NH_4)_2S_2O_8$ mediated oxidative dearomatization of indoles for the construction of C2-quaternary indolinones derivatives in moderate to good yields. The significant aspects of our work allows modest functional group tolerance, were compatible under the current methodology. This methodology provides an alternative approach for the direct generation of all-carbon quaternary centers at the C2 position of indoles. This catalytic approach represents a step-economic and convenient strategy for the oxidative dearomatization of indoles.

Conflicts of interest

There are no conflicts to declare.

^a Reaction conditions: 1 (0.3 mmol), $(NH_4)_2S_2O_8$ (2 equiv.), DMF (2 mL), 60 °C, air tube for 3 h. Isolated yields. ^b In a 9 mmol scale.

Acknowledgements

We are grateful for the financial support from the project supported by Chengdu Normal University Project (2021CS21ZCY02) and National undergraduate training program for innovation and entrepreneurship (Grant No. 202114389016 and 202114389005), the Foundation of Chengdu Normal University Talent Introduction Research Funding (2021YJRC202020).

Notes and references

- 1 (*a*) E. Vitaku, D. T. Smith and J. T. Njardarson, Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among US FDA approved pharmaceuticals: miniperspective, *J. Med. Chem.*, 2014, 57, 10257–10274; (*b*) R. D. Taylor, M. MacCoss and A. D. Lawson, Rings in drugs: Miniperspective, *J. Med. Chem.*, 2014, 57, 5845–5859.
- 2 (a) R. Sundberg, *Indoles*, Academic Press, San Diego, 1996, p. 113; (b) S. Cacchi and G. Fabrizi, Synthesis and functionalization of indoles through palladium-catalyzed reactions, *Chem. Rev.*, 2005, **105**, 2873–2920; (c) L. Joucla and L. Djakovitch, Transition Metal-Catalysed, Direct and Site-Selective *N*1-, C2-or C3-Arylation of the Indole Nucleus: 20 Years of Improvements, *Adv. Synth. Catal.*, 2009, **351**, 673–714; (d) M. Bandini and A. Eichholzer, Catalytic functionalization of indoles in a new dimension, *Angew. Chem., Int. Ed.*, 2009, **48**, 9608–9644; (e) G. Bartoli, G. Bencivenni and R. Dalpozzo, Organocatalytic strategies for the asymmetric functionalization of indoles, *Chem. Soc. Rev.*, 2010, **39**, 4449–4465.
- 3 (a) A. Gazit, N. Osherov, I. Posner, P. Yaish, E. Poradosu, C. Gilon and A. Levitzki, Tyrphostins. II. Heterocyclic and. alpha.-substituted benzylidenemalononitrile tyrphostins as potent inhibitors of EGF receptor and ErbB2/neu tyrosine kinases, *J. Med. Chem.*, 1991, 34, 1896–1907; (b) S. Dilber, M. Saban, J. Jelaca, A. Gelineo, L. Arsenijević and M. Bogavac, Investigation of antimicrobial activity of some isatin derivatives, *Pharmazie*, 1989, 44, 649–650; (c) A. Wetzel and F. Gagosz, Gold-Catalyzed Transformation of 2-Alkynyl Arylazides: Efficient Access to the Valuable Pseudoindoxyl and Indolyl Frameworks, *Angew. Chem., Int. Ed.*, 2011, 123, 7492–7496; (d) C. V. S. Kumar, V. G. Puranik and C. V. Ramana, InCl₃-Mediated Addition of Indole to Isatogens: An Expeditious Synthesis of 13-deoxy-Isatisine A, *Chem.-Eur. J.*, 2012, 18, 9601–9611.
- 4 M. Linhares, S. L. Rebelo, M. M. Simoes, A. M. Silva, M. G. P. Neves, J. A. Cavaleiro and C. Freire, Biomimetic oxidation of indole by Mn (III) porphyrins, *Appl. Catal.*, 2014, 470, 427-433.
- 5 R. Bell, S. Carmeli and N. Sar, Vibrindole A, a metabolite of the marine bacterium, Vibrio parahaemolyticus, isolated

- from the toxic mucus of the boxfish Ostracion cubicus, *J. Nat. Prod.*, 1994, 57, 1587–1590.
- 6 A. Gazit, N. Osherov, I. Posner, P. Yaish, E. Poradosu, C. Gilon and A. Levitzki, Tyrphostins. II. Heterocyclic and. alpha.-substituted benzylidenemalononitrile tyrphostins as potent inhibitors of EGF receptor and ErbB2/neu tyrosine kinases, *J. Med. Chem.*, 1991, 34, 1896–1907.
- 7 T. L. Stull, L. Hyun, C. Sharetzsky, J. Wooten, J. P. McCauley and A. B. Smith, Production and Oxidation of Indole by Haemophilus influenzae, *J. Biol. Chem.*, 1995, 270, 5–8.
- 8 (a) J.-F. Liu, Z.-Y. Jiang, R.-R. Wang, Y.-T. Zheng, J.-J. Chen, X.-M. Zhang and Y.-B. Ma, Isatisine A, a novel alkaloid with an unprecedented skeleton from leaves of Isatis indigotica, *Org. Lett.*, 2007, **9**, 4127–4129; (b) J. Lee and J. S. Panek, Total synthesis of (+)-isatisine A, *Org. Lett.*, 2011, **13**, 502–505; (c) H.-Z. Zheng, Z.-H. Dong and Q. Yu, *Modern Study of Traditional Chinese Medicine*, Xueyaun Press, Beijing China, 1997, vol. 1, pp. 328–334.
- 9 (a) W.-B. Qin, Q. Chang, Y.-H. Bao, N. Wang, Z.-W. Chen and L.-X. Liu, Metal-free catalyzed oxidative trimerization of indoles by using TEMPO in air: a biomimetic approach to 2-(1H-indol-3-yl)-2, 3'-biindolin-3-ones, Org. Biomol. Chem., 2012, 10, 8814–8821; (b) Y.-B. Kong, J.-Y. Zhu, Z.-W. Chen and L.-X. Liu, Copper-catalyzed oxidative trimerization of indoles by using TEMPO to construct quaternary carbon centers: the synthesis of 2-(1H-indol-3-yl)-2, 3'-biindolin-3-ones, Can. J. Chem., 2014, 92, 269–273.
- 10 J. Xue, Y. Bao, W. Qin, J. Zhu, Y. Kong, H. Qu, Z. Chen and L. Liu, Metal-Free–Catalyzed Oxidative Trimerization of Indoles Using NaNO₂ to Construct Quaternary Carbon Centers: Synthesis of 2-(1*H*-Indol-3-yl)-2, 3'-biindolin-3ones, Synth. Commun., 2014, 44, 2215–2221.
- 11 J. Kothandapani, S. M. K. Reddy, S. Thamotharan, S. M. Kumar, K. Byrappa and S. S. Ganesan, TBHP Mediated Substrate Controlled Oxidative Dearomatization of Indoles to C2/C3-Quaternary Indolinones, *Eur. J. Org. Chem.*, 2018, 2018, 2762–2767.
- 12 S. B. Gohain, M. Basumatary, P. K. Boruah, M. R. Das and A. J. Thakur, Nano Au/Pd-catalysed 'on-water'synthesis of C3–C3' diaryl-oxindole scaffolds via N 2-selective dearomatization of indole, *Green Chem.*, 2020, 22, 170–179.
- 13 (a) S. Meenakshisundaram and N. Sarathi, Kinetics and mechanism of oxidation of indole by HSO, *Int. J. Chem. Kinet.*, 2007, **39**, 46–51; (b) See ref. 11.
- 14 A. Bahuguna, A. Kumar, S. Kumar, T. Chhabra and V. Krishnan, 2D-2D nanocomposite of MoS₂-graphitic carbon nitride as multifunctional catalyst for sustainable synthesis of C3-functionalized indoles, *ChemCatChem*, 2018, **10**, 3121–3132.