

## REVIEW

[View Article Online](#)  
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2022, 13, 6457

## Functional amyloids from bacterial biofilms – structural properties and interaction partners

Ümit Akbey<sup>\*a</sup> and Maria Andreassen <sup>\*b</sup>

Protein aggregation and amyloid formation have historically been linked with various diseases such as Alzheimer's and Parkinson's disease, but recently functional amyloids have gained a great deal of interest in not causing a disease and having a distinct function *in vivo*. Functional bacterial amyloids form the structural scaffold in bacterial biofilms and provide a survival strategy for the bacteria along with antibiotic resistance. The formation of functional amyloids happens extracellularly which differs from most disease related amyloids. Studies of functional amyloids have revealed several distinctions compared to disease related amyloids including primary structures designed to optimize amyloid formation while still retaining a controlled assembly of the individual subunits into classical cross- $\beta$ -sheet structures, along with a unique cross- $\alpha$ -sheet amyloid fold. Studies have revealed that functional amyloids interact with components found in the extracellular matrix space such as lipids from membranes and polymers from the biofilm. Intriguingly, a level of complexity is added as functional amyloids also interact with several disease related amyloids and a causative link has even been established between functional amyloids and neurodegenerative diseases. It is hence becoming increasingly clear that functional amyloids are not inert protein structures found in bacterial biofilms but interact with many different components including human proteins related to pathology. Gaining a clear understanding of the factors governing the interactions will lead to improved strategies to combat biofilm associated infections and the correlated antibiotic resistance. In the current review we summarize the current state of the art knowledge on this exciting and fast growing research field of biofilm forming bacterial functional amyloids, their structural features and interaction partners.

Received 1st February 2022  
Accepted 5th May 2022

DOI: 10.1039/d2sc00645f

[rsc.li/chemical-science](https://rsc.li/chemical-science)

## Introduction

Historically the formation of protein aggregates or amyloids has been associated with various diseases such as neurodegenerative diseases.<sup>1</sup> More recently it has been found that amyloids can have biological functions in the organism producing them, hence differing from the pathological amyloids and coining the term functional amyloids.<sup>2</sup> Amyloid fibrils and functional amyloids are a topic of great interest and research, and the exponential growth of the field, research output and citations are shown in Fig. 1. Functional amyloids are found in various organisms ranging from bacteria to humans and currently a whole range is known, Fig. 2.<sup>2</sup> They provide different types of functionality ranging from virulence during infection<sup>3</sup> to structural scaffolds in biofilms,<sup>4</sup> to storage of peptide hormones,<sup>5</sup> to formation of melanin granula in melanocytes,<sup>6</sup> to memory storage<sup>7</sup> and during cell death,<sup>8</sup> and many others.

Many bacteria form biofilms to survive under stress-inducing conditions to achieve communal living. Biofilms are microbial communities where the microorganisms live in a matrix of hydrated extracellular polymeric substances (EPS) that forms the scaffold of the biofilm.<sup>9</sup> The EPS is produced and secreted by the bacterial cells living in the biofilm, and it is composed of biopolymers such as polysaccharides, proteins and nucleic acids in the form of extracellular DNA (eDNA), Fig. 3. The formation of biofilm by bacteria enables adhesion to surfaces and other bacteria cells, retention of nutrients and water, and additionally acts as a protective barrier. Biofilm-associated pathogenic microbes are protected from antimicrobial agents and host immune system attacks, as a result they are more infectious and difficult to treat. Eighty percent of all chronic infections are related to bacterial biofilms.<sup>10,11</sup>

Aggregated proteins in the form of amyloid fibrils play a key role in maintaining the structural integrity of biofilms and when these proteins are mutated, the biofilms are disrupted and bacteria become accessible to, *e.g.*, antibiotic treatments. Functional amyloids strengthen biofilms and are a major threat to human health, since the (chronic) infections they cause are difficult to treat due to the biofilm structural integrity and

<sup>a</sup>Department of Structural Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA 15261, USA. E-mail: [umitakbey@pitt.edu](mailto:umitakbey@pitt.edu)<sup>b</sup>Department of Biomedicine, Aarhus University, Wilhelm Meyers Allé 3, 8000 Aarhus, Denmark. E-mail: [marij@biomed.au.dk](mailto:marij@biomed.au.dk)



**Fig. 1** The number of publications per year (left y-axis) and total citations (right y-axis) in the Web of Science (WoS) in the research fields of (A) "amyloid fibrils" and (B) "functional amyloid", by the end of 2021. Remarkable progress in these fields has been made in the recent years as evident from the increase in the research output in those fields.

insufficient penetration of drugs, thus promoting antibiotic resistance (antimicrobial resistance, AMR).<sup>12</sup>

Targeting biofilms and their amyloid components could be a novel approach to fight AMR, which results in ~million casualties per year, and is estimated to cause more death than cancer by 2050.<sup>13</sup> However, very little structural information exists about biofilms and their fibrillar components and how these components interact with other proteins. Determining these unknown structures and unraveling the structure–activity relationship of such fibrils will help to develop better therapies.

In this current review we focus on functional amyloids related to bacterial biofilm formation or functional bacterial amyloids (FuBAs), more specifically curli from *E. coli*, FapC from *Pseudomonas*, TasA from *B. subtilis* and phenol-soluble modulins (PSMs) from *S. aureus*, Fig. 2. For an extensive list of

various functional amyloids along with coverage of the broader theme of amyloids, functional amyloids, prions and functional prions we refer to other recent excellent reviews.<sup>14–22</sup>

Other functional amyloids related to bacteria but not directly involved in the biofilm formation have also been observed. These include but are not limited to *S. coelicolor* chaplins which facilitates the rising of aerial hyphae,<sup>23,24</sup> *M. tuberculosis* amyloid pili (MTP) which mediates host interactions during pathogenesis,<sup>25</sup> and *X. axonopodis* harpins which act as virulence factors.<sup>3,26</sup> Recently other functional amyloids from bacterial biofilms such as the Esp from *E. faecalis*<sup>27</sup> have also been discovered but due to the recent discovery these are still poorly understood and will not be described in this paper. Moreover, due to the homology between the *Salmonella* Tafi system and *E. coli* curli,<sup>28</sup> the Tafi system will also not be addressed here.



**Fig. 2** (A) Schematic representation of different bacterial biofilm forming functional amyloids: PSMs from *S. aureus*, CsgA from *E. coli*, FapC from *P. aeruginosa* and TasA from *B. subtilis*. The signal sequences are shown in red, the amyloid regions (AR) in blue and loop/linker and C-terminus regions in grey. The number of amino acids forming the amyloid region is given for each protein in blue. The lengths of the full-length proteins are given in parenthesis in black. The PSMs are short proteins and AR themselves. The CsgA is composed of 5 AR repeats, FapC from 5 AR repeats, whereas, TasA is considered to be composed of a large AR without repeats (note that other hypotheses exist on the nature of TasA amyloid region). (B) The amino acid composition of the five functional amyloids. The Uniprot identification numbers for each protein are given in parenthesis. The consecutive AR repeats are indicated as bold versus normal font for CsgA. (C) Illustration of amyloid fibril construction mechanisms by using different building blocks to the similar final amyloid fibril fold.

### Curli from *E. coli*

Curli was the first discovered biofilm forming functional amyloid fibril and hence curli fibrils from *E. coli* are the most extensively characterized FuBAs.<sup>4</sup> Curli fibrils have been found to be essential for biofilm formation and attachment to surfaces including plant cells, stainless steel, glass and plastics.<sup>29–33</sup> They also play key roles in interaction with host proteins and invasion of host cells.<sup>34–36</sup> The curli biogenesis involves the expression of two divergently evolved operons, *csgBA* and *csgDEF*.<sup>35</sup> CsgA is the major biofilm forming amyloid protein component of curli and CsgB is the minor amyloid protein. CsgA and CsgB are secreted as unstructured monomeric proteins to the cell surface to self-assemble into the final amyloid fibrils. CsgB facilitates CsgA fibrillation as a nucleator and is required for the complete assembly of curli fibrils.<sup>35,37</sup> CsgC is a chaperon-like accessory protein that prevents fibrillation in the periplasm to ensure that CsgA and CsgB are secreted as monomeric units to the extracellular compartment.<sup>38</sup> CsgE is another chaperon-like accessory protein that reversibly interacts with CsgA (in a 1 : 1 ratio) while the pore-forming protein CsgG promotes translocation, thereby achieving specificity for secretion.<sup>39</sup> CsgF and also CsgB help localize CsgA to the cell surface<sup>40</sup> while CsgD regulates the expression of the *csgABC* operon.<sup>19</sup> This amazingly regulated protein system controls and directs curli amyloid

formation and maturation, and prevents the formation of toxic oligomers or fibrils inside the cells that may otherwise harm the bacteria itself. CsgA and CsgB are homologs and share ~50% sequence identity. They both consist of five imperfect repeats of ~20 amino acids and the imperfect repeats of CsgA contain highly conserved glutamine and asparagine residues, which are important for the amyloid formation,<sup>37,41</sup> Fig. 2. Repeating the amyloidogenic region several times likely comprises an effective way of fibrillation compared to non-repeat containing proteins.<sup>42</sup> This may explain the fast fibril formation without a lag-phase compared to other amyloids. The repeats R1, R3 and R5 of CsgA are amyloidogenic on their own and can form fibrils similar to wild type (WT) CsgA, whereas R2 and R4 do not form fibrils on their own and are dispensable for WT fibril formation.<sup>40</sup> Interestingly, the amyloidogenic R1 and R5 contain sequences that contribute significantly to the ability of CsgA to bind human proteins such as fibronectin, plasminogen, tissue plasminogen activator, and  $\beta$ 2-microglobulin.<sup>43</sup> *In vitro* in the absence of the other curli components CsgA aggregates in a nucleation dependent manner.<sup>44</sup>

### FapC from *Pseudomonas*

Functional amyloids in *Pseudomonas* (Fap) have more recently been discovered and are encoded by a single operon with six









**Fig. 3** (A) Representation of bacteria mediated biofilm formation, initiated by the motile species gathering together into the mature solid biofilm matrix. The molecular details of the biofilm composed of bacteria themselves, polysaccharides, extracellular DNA and protein amyloid fibrils are additionally shown. This figure is reproduced with permission.<sup>9</sup> (B) The wild-type (WT) *versus* the deletion-mutants ( $\Delta$ ) of *B. subtilis* bacteria and the biofilm formed by them. The WT,  $\Delta$ eps,  $\Delta$ tasA and  $\Delta$ eps-tasA. The deletion mutants form bacterial communities that are healthy but with a weakened/disrupted biofilm. This figure is reproduced with permission.<sup>63</sup> (C) Reconstruction of a native-like *B. subtilis* biofilm from a  $\Delta$ tasA deletion mutant that cannot form biofilm by itself. The supplementation of recombinantly produced TasA protein to the growth medium of the  $\Delta$ tasA mutant *B. subtilis* strain reconstitutes the native-like biofilm. This allows incorporation of isotope-labelled TasA into the biofilm for *in vivo* NMR spectroscopy without any background signal. (D) 2D  $^1\text{H}$ - $^{15}\text{N}$  NMR spectra of different TasA samples prepared *in vitro*, soluble monomer, oligomer and fibril, as well as *in vivo* TasA in biofilm. Spectral differences indicate structural differences and shed light on the biofilm TasA form. For more details on the figure and explanation of the additional indications, the reader is referred to the original article. This figure is reproduced with permission.<sup>81</sup>




are structures available from small peptide fragments from functional amyloids, monomeric protein forms as well as some structural models. This lack of structural information limits our understanding of functional amyloids and the formation of biofilms to a large extent, and hampers the strategic fight against, *e.g.*, biofilm related infections, pathogenesis and antimicrobial resistance (AMR).<sup>73,74</sup>

## General structural features of (functional) amyloid fibrils

The function of a protein in living organisms is directed by its three-dimensional (3D) structure, entailing a structure–function relationship. Every protein has to adopt its correct fold to function, which is uniquely determined by the amino acid sequence. However, one protein can adopt multiple structures if the conditions (buffer, pH, salt, concentration *etc.*) are manipulated towards different local/globular minima in the protein folding energy landscape. The amyloid fold derived by intermolecular contacts, rather than intramolecular contacts, is one of these possible structures and with its high stability and lower total energy, once formed they are usually trapped in this conformation.<sup>75</sup>

Amyloid fibrils can be obtained for many protein sequences, *e.g. via* misfolding along the folding pathway towards a native conformation. Protein misfolding and/or fibrillation is usually associated with biological anomalies and diseases.<sup>1,76</sup> Neurodegenerative diseases such as Alzheimer's disease (AD) and Parkinson's disease (PD) are known examples that are caused by pathological amyloid proteins. The loss of normal protein function and subsequent gain of toxicity are key phenotypes for this type of pathology. Functional amyloids, in contrast, have a distinct biological function *in vivo*.<sup>1,77,78</sup> By applying harsh conditions such as extreme pH or denaturing conditions, many globular or unfolded proteins can be directed to amyloid fibrils.<sup>79</sup> Starting from a natively unfolded protein state (intrinsically disordered protein, IDP), rather than a well-defined native fold, highly stable amyloid structures can be formed as seen in AD and PD or from functional amyloids like CsgA and FapC.<sup>20,42</sup> Amyloid formation has been observed from proteins with all types of native structures, *e.g.* in an  $\alpha$ -helical structure from serum amyloid,  $\beta$ -sheet rich structure from transthyretin, or  $\alpha$ - and  $\beta$ -rich structure from lysozyme,<sup>80</sup> as well as in a more controlled manner from a globular protein fold.<sup>63,81</sup> When amyloid fibril formation occurs from an initial folded



**Fig. 4** (A) Representation of the cross- $\beta$ -sheet amyloid fibril fold. The interstrand and intersheet distances of  $\sim 4.7$  and  $\sim 10$  Å are depicted. Here the  $\beta$ -strands form intermolecular  $\beta$ -sheets and are orthogonal to the fiber axis. (B) The negative-stain electron microscopy (EM) micrograph of four different biofilm forming functional amyloids. The white scale bar represents 200 nm at each micrograph. The CsgA EM is reproduced with permission<sup>148</sup>

protein extensive conformational rearrangements are required, which may require unfolding to an intermediate that then forms the final amyloid fibril structure.<sup>82,83</sup> Despite many initial forms of fibril forming proteins, the final structure consists of cross- $\beta$ -sheets as explained below.<sup>84</sup>

Amyloid fibrils are insoluble protein aggregates and are rich in  $\beta$ -sheet content if not composed predominantly by them. For a globular protein the  $\beta$ -strands form intramolecular  $\beta$ -sheets, however, for amyloid fibrils the  $\beta$ -strands form intermolecular  $\beta$ -sheets which are arranged orthogonal to the fiber axis. The longer  $\beta$ -strands can be bent into a  $\beta$ -hairpin motif connecting two  $\beta$ -strands. This cross- $\beta$ -sheet structure is stabilized by hydrogen bonds running parallel to the fiber axis, Fig. 4A.

Steric-zippers formed from tightly packed protein sidechains between the  $\beta$ -sheets increases the amyloid fibril stability particularly for the small-peptide amyloid examples. One or more cross- $\beta$ -sheets then form one or more protofilaments. Due to this protein architecture a typical X-ray diffraction pattern is observed, with two typical reflections at  $\sim 4.7$  Å (vertical axis) corresponding to the inter-strand spacing and at  $\sim 10$  Å (horizontal axis) corresponding to the inter-sheet spacing due to the sidechains,<sup>85–87</sup> Fig. 4. The kinetics of amyloid formation is typically studied using the amyloid binding dye Thioflavin T (ThT). The fluorescence intensity of the dye increases due to binding of ThT in grooves at the surface of the amyloid fibrils perpendicular to the fibril axis. This results in rotational



Fig. 5 (A) Schematic representation of amyloid fibril formation. The soluble protein fold (in red) can form oligomers (in blue) or amyloid fibrils (in green) depending on the system and condition. The oligomers then form amyloid fibrils. The amyloid fibril maturation can happen with different steps/mechanisms as depicted in the figure; 1° nucleation, elongation, 2° nucleation/self-replication and a final maturation. (B) Amyloid binding Thioflavin-T dye results monitoring the formation of amyloid fibrils from soluble monomers for PSM $\alpha$ 1 and PSM $\alpha$ 3 proteins from *S. aureus* (repeated three times for each protein, please note the different time scales).

A quantitative comparison of the amino acid composition of functional *versus* pathological amyloids reveals more details on the evolutionary optimization of amino acids and their selection accordingly.<sup>17</sup> A comparison of four pathological and six functional amyloid sequences indicated that despite high-frequency occurrence of nonpolar residues in both, the polar residues were more dominant for functional amyloids. The polar residues can form ladders of hydrogen bonds along the fibril, with higher frequency in functional amyloids. Positively charged residues such as lysine, arginine and histidine are observed more frequently in pathological amyloids, whereas, negatively charged residues are observed for both types to a similar degree (except glutamate showing higher frequency for pathological amyloids). This phenomenon has alternatively been described as gatekeeper residues composed of arginine, aspartate, glycine, lysine or proline.<sup>42</sup> Overall, more frequent occurrence of glycine, serine and glutamine in functional amyloids correlates with them being more reversible with more hydrophilic and flexible interfaces, in contrast to the hydrophobic, dry and tight interfaces observed for pathological amyloids.<sup>16</sup> The aliphatic residues constituting highly stable steric-zipper conformations are observed often in small-peptide

Solution NMR spectroscopy has been applied to study functional amyloids in their monomeric soluble form.<sup>81,118–120</sup> Since most of the known functional amyloids are unfolded proteins, the use of this technique is limited to understanding



ssNMR spectroscopy historically relied on 2D/3D carbon-detection experiments for chemical shift dispersion and sequential assignment,<sup>99,127–129</sup> requiring ~10 mg of fully/selective/sparse isotope labeled protein. More recently, proton-detection experiments relying on protein deuteration and/or ultra-fast sample spinning (MAS) have expanded the scope of ssNMR by improving the sensitivity significantly and reducing the requirement for the protein amount needed to <1 mg.<sup>130–134</sup> Nowadays, these proton-detecting ssNMR experiments at high magnetic fields above 1 GHz combined with MAS frequencies above 100 kHz allow a fast and sensitive structure determination of amyloids or other protein forms.<sup>135–138</sup> It is important to note that combining long-range structural information from cryo-EM and atomic resolution of a single unit from ssNMR can be a very powerful experimental approach. In addition, dynamic nuclear polarization based (DNP) ssNMR increases the sensitivity by several orders of magnitude, and opens up the possibility to study systems otherwise very difficult or impossible, *e.g.* proteins in complex environments at low concentrations.<sup>139–143</sup>

**PSMα1**  
monomer / fragment  
(NMR - 5KHB)

**PSMα3**  
monomer / fibril  
(NMR - 5KGY)

**TasA**  
monomer  
XRay - 5OF1

**FapC**  
model

**CsgA**  
fragments  
45LNIQY50 (XRay - 6G8D)  
47IYQYGG52 (XRay - 6G8C)  
137VTQGVF142 (XRay - 6G8E)  
129TASNSS134 (XRay - 6G9G)

**CsgA**  
model

**FapC**  
model

**CsgA**  
AlphaFold model

Chem. Sci., 2022, 13, 6457-6477 | 6465

## Structural studies on FuBAs

**PSMs.** PSMs are characterized as amphipathic  $\alpha$ -helices in solution, Fig. 6.<sup>144,145</sup> They can adopt either a  $\beta$ -sheet or  $\alpha$ -helical structure after aggregation *in vitro* and in biofilms, as determined from ThT-binding and other biophysical assays.<sup>55</sup> The atomic resolution structure of full-length PSM $\alpha$ 3 amyloid-like fibrils (22 amino acids) was determined by X-ray crystallography in 2017,<sup>59</sup> Fig. 6. This unique structure is composed of a cross- $\alpha$ -architecture, in contrast to the “canonical” cross- $\beta$  structural features of other known amyloids. This was the first determined prokaryotic cross- $\alpha$  amyloid fibril structure. The X-ray fibril diffraction pattern indicates reflections at  $\sim$ 4.6, 9.2 and 11.5 Å. This non-canonical structure is made of amphipathic  $\alpha$ -helices stacking perpendicular to the fiber axis forming sheets, which are tightly packed with their hydrophobic face towards each other between sheets. PSM $\alpha$ 3 was determined to be  $\alpha$ -helical both in solution as a monomer and in the fibrillar form, Fig. 6.<sup>59,118</sup> The buried surface area, aromatic residues, and cavities inducing ThT binding are properties similar to the cross- $\beta$  fibrils. It was shown that the toxicity of this protein is strongly coupled to its ability to form fibrils in the cross- $\alpha$  form.<sup>145</sup>

Truncation mutants of PSM $\alpha$ 3 were studied for structure determination.<sup>61</sup> In contrast to the cross- $\alpha$  structure of the full length PSM $\alpha$ 3, the  $\gamma$ LFKFFK<sub>12</sub> fragment forms two different atypical cross- $\beta$  amyloid fibrils composed of anti-parallel strands, Fig. 6. One polymorph is composed of hexameric architecture with cylindrical cavities along the fiber axis, and the other one is composed of out-of-register  $\beta$ -sheets that are  $-50^\circ$  off from the fibril axis in contrast to the  $90^\circ$  found for in-register sheets. These structures were found to be less stable compared to the classic cross- $\beta$  structures containing steric-zipper elements. In contrast to this picture, the structures of truncation spine elements of PSM $\alpha$ 1  $\gamma$ IHKVIK<sub>12</sub> and PSM $\alpha$ 4  $\gamma$ -IHKIK<sub>12</sub> form cross- $\beta$  fibrils with steric-zippers and have classical X-ray reflections at 4.6 and 10 Å. Full length PSM $\alpha$ 1 and  $\alpha$ 4 undergo a structural transition from helical to  $\beta$ -sheet in solution during aggregation within several days, in contrast to PSM $\alpha$ 3.<sup>146,147</sup> These transitions or non-transitions could be related to the functions of these proteins, which are strikingly colorful. More structural studies on full length PSMs are needed to unravel the full picture of these systems.

**CsgA.** The secondary structure prediction and ssNMR data suggest the formation of strand-loop-strand type of structures for the repeat units of the CsgA amyloid fibril.<sup>35,148</sup> Detailed structural work on CsgA was performed by Tycko and later by Ritter and coworkers, Fig. 7.<sup>148</sup> The earlier work comprises ssNMR characterization that was supported with X-ray diffraction, EM, MPL measurements and biophysical methods. The ssNMR results give hints about the CsgA fibril structure prepared without signal-peptides. The lyophilized and rehydrated samples showed low spectral quality in contrast to the functional amyloid HET-s with superb resolution.<sup>99</sup> Nevertheless, superior NMR resolution was obtained from a recombinant prepared CsgA sample.<sup>149</sup>

Typical X-ray diffraction patterns were observed for both CsgA and CsgB indicating cross- $\beta$ -sheet spacings of  $\sim$ 4.7 and  $\sim$ 9 Å. Secondary structure contributions were determined by circular dichroism (CD) to be  $\sim$ 16%  $\alpha$ -helix,  $\sim$ 40%  $\beta$ -sheet,  $\sim$ 13%  $\beta$ -turn and  $\sim$ 31% others. These *in vitro* prepared fibrils of CsgA were more resistant to proteinase-K as compared to the soluble form. A comparison of the observed chemical shifts with the random-coil values<sup>150,151</sup> indicates the presence of major and minor contributions for many of the selectively labeled sites. Quite surprisingly, these chemical shifts indicate both  $\beta$ -sheet and non- $\beta$ -sheet contributions, which is a major difference between CsgA and other pathological amyloids that are composed of in-register  $\beta$ -sheet fibrils. This situation was similar to that of the HET-s structure which was found to adopt a unique  $\beta$ -solenoid structure.<sup>99</sup> The distance measurements based on dipolar-dephasing experiments were not definitive but indicated a non in-register  $\beta$ -sheet structure. This was supported by the MPL measurements and EM images showing the fibrils having a narrow ( $\sim$ 3–4 nm) width. Proposed flexible loops that may be connecting the rigid repeat units were not observed by NMR.

A CsgA structural model based on amino acid contacts from a multiple sequence alignment of homologous proteins suggests that the hairpins stack on top of each other to form a  $\beta$ -helical structure, Fig. 6,<sup>48,152</sup> similar to HET-s and the recently determined structure of a HET-s homolog HELLF functional amyloid fibrils.<sup>99,153</sup> The model is consistent with the repeats identified for CsgA, Fig. 2. The Alphafold structural model is overall similar to this model, and both are consistent with the NMR data described above.<sup>154</sup>

A more recent ssNMR study revealed more insights on the structural fold of CsgA and supports the proposed models, Fig. 7.<sup>149</sup> This work presents evidence *via* ssNMR that the biofilm-extracted and recombinant produced proteins adopt a similar homogeneous structure. The spectral resolution ( $\sim$ 0.35 ppm) of these preparations is superior compared to that of the lyophilized samples,<sup>148</sup> and similar to that of HET-s or  $\alpha$ Syn.<sup>99,129,155</sup> Sample preparation of amyloid fibrils remains a challenge and needs to be optimized carefully for each system. The segmental labeling strategy of a single repeat unit R1 (21–69 amino acids) greatly simplifies the NMR spectrum and allows sequential assignment and secondary structure propensity determination.<sup>156</sup> These results support the strand-loop-strand structural motif with two  $\beta$ -strands from S43–Y50 and S55–T61 separated by a short rigid glycine loop. Moreover, the R1 repeat unit and the full CsgA adopts a similar  $\beta$ -solenoid structure, and most probably all the repeats are structurally equivalent. Observation of long-range distance restraints defines a tight turn between the individual  $\beta$ -strands of the repeats. Finally, ssNMR spectra monitoring only mobile segments contain signals from the N-terminus and the loop that connects repeats.

**FapC.** A detailed structural and biophysical study depicts our current understanding of the FapC functional amyloid, Fig. 7.<sup>157</sup> In this work, full-length (FL: NR1L1R2L2R3C) and three truncation mutants (NL1L2R3C, L2R3C and R3C) were studied in detail. The effect of loop regions on the fibrillation and fibril structure was studied using these different constructs. The X-





**Fig. 7** NMR is a powerful technique to study protein structures of soluble and insoluble forms. (A) 2D  $^1\text{H}$ – $^{15}\text{N}$  HSQC NMR spectra of soluble monomeric CsgA (in DMSO), FapC (in DMSO) and TasA (in  $\text{H}_2\text{O}$ ) functional amyloids. The unfolded nature of the CsgA and FapC can be seen directly from the narrow chemical shift dispersion in the proton dimension in the HSQC spectra. In contrast, the TasA has a well-dispersed HSQC spectrum, which is an indication of its folded globular protein fold. (B) The 2D  $^{13}\text{C}$ – $^{13}\text{C}$  MAS NMR spectra of CsgA, FapC (R3C construct) and TasA are shown. Solid-state NMR spectroscopy is used to study structures of functional amyloids in their insoluble fibrillar form. The NMR spectra show high quality that is suitable for structure determination. Sparse/segmental labeling combined with proton-detection experiments will additionally increase the full structure determination possibilities for these kinds of difficult proteins. Such segmental labeling of only a single repeat for CsgA results in superior spectral resolution, isotope labeling of full-length is shown in black and single-repeat is shown in blue with several signal assignments. Frans Mulder kindly provided the FapC HSQC spectrum. The 2D  $^{13}\text{C}$ – $^{13}\text{C}$  ssNMR spectrum of CsgA is reproduced with permission.<sup>149</sup>

ray diffraction analysis resulted in the typical reflections at  $\sim 4.6$  and  $\sim 10$  Å, as well as an additional  $\sim 3.9$  Å reflection for all preparations. The smaller distance reflection was assumed to arise from the twisting of the fibrils. This indicates that truncation mutants comprise similar structures to the FL FapC. EM and infrared spectroscopy (IR) provided additional proof that the truncation preparations are structurally similar to the FL FapC. Initial sequence analyses predicted three imperfect repeats separated by linker regions that are assumed to be flexible loops.<sup>48,158</sup> The repeats contain mostly hydrophobic amino acids, whereas the linker and N-terminal region contain charged amino acids that were the origin of the initial nomenclature. This picture is modified now by the determination of four amyloidogenic hot spots in FapC (between 37–43, 102–111, 157–163 and 215–219), present in the linkers, termini and R3 regions. The aggregation of the FL has a lag phase, which is further slowed down by the removal of R1 and R2. The shortest truncation unit R3C aggregates *via* secondary

nucleation, whereas, FL aggregates *via* primary nucleation and elongation. Surprisingly, the predicted loop regions increase the aggregation propensity, highlighting the importance of the non-repeat regions for FapC. The removal of all three repeats still results in amyloid fibril formation,<sup>53</sup> indicating the presence of an amyloidogenic hot spot in the linker regions.

A recent study by Pedersen and coworkers expands the macromolecular structural details of FapC amyloid fibrils.<sup>159</sup> In this work, an MPL value of  $\sim 33$  kDa  $\text{nm}^{-1}$  was determined by EM and small angle X-ray scattering (SAXS), indicating that the mature fibrils consisted of two-protofilaments per fibril. The protofilaments were formed without any intermediate species besides monomers and oligomers, which then mature into more compact fibrils.

2D ssNMR spectra revealed further insights about FapC.<sup>157</sup> First of all, the spectral quality is decent compared to the CsgA, however, it was significantly lower compared to HET-s. Nevertheless, the R3C construct gives an opportunity to extract



Freshly prepared TasA<sub>28–261</sub> and TasA<sub>28–239</sub> samples at pH 7 are mostly monomeric with a small oligomeric fraction, whereas TasA<sub>261</sub> favors oligomer formation to a greater extent. The C-terminus of TasA (~240–261 aa) is unstructured as determined by NMR spectroscopy. The crystal structure was obtained at 1.56 Å for the monomeric TasA<sub>239</sub> at pH 4.6. The jellyroll-fold consists of antiparallel β-sheets separated by short helices and longer loops, with surprisingly differential rigidity observed among the proteins.

In contrast to the three different TasA forms obtained for the recombinant protein *in vitro*, a pure fibrillar form is obtained *in vivo*.<sup>149</sup> The model *B. subtilis* system that cannot produce TasA ( $\Delta$ tasA) was used for understanding insights into the native biofilm. This strain can grow healthy, but cannot produce biofilm. Externally supplied recombinant TasA protein reconstitutes the native-like biofilm, Fig. 3C.<sup>63,81</sup> A DCN-labeled TasA<sub>261</sub> was used in this model system and a fingerprint HSQC-type ssNMR spectrum was obtained, Fig. 3D. This experimental setup is the first demonstration of *in vivo* (in the biofilm) ssNMR spectroscopy, to the best of our knowledge. The resulting NMR spectrum with decent resolution contains all the features from the spectrum of the fibrillar sample. *In vivo* biofilm and *in vitro* fibrillar spectra overlap to a large extent, however they still have small differences which indicate that additional factors within the biofilm can modulate the protein structure. This



remarkable observation indicates that the biofilm contains homogeneous TasA fibrils that are resistant to proteases.

In summary, biofilm forming functional amyloids and functional amyloids in general have many properties similar to pathological amyloids. These include, *e.g.*, a common cross- $\beta$  structure, X-ray diffraction pattern and limited proteolysis results. The details of how these two classes of amyloids form structures are still surprisingly unique. As unique features, functional amyloids have a control over the mechanism that regulates their *in vivo* properties and fibrillation, whereas pathological ones has not been evolutionary optimized in the same manner due to their lack of function. Structurally we have limited information and except from a special crystalline case of PSM $\alpha$ 3, no high-resolution structure is available. Nevertheless, studies unraveling structural features exist as we have summarized above and comprise a solid basis for the future full structure determination studies. With the advancements in structural biology techniques, we expect a bright future with many high-resolution structures enlightening the molecular mechanisms of functional amyloids and biofilm formation *in vitro* and *in vivo*.

## Interactions between functional amyloids and components present in the biofilm matrix

As mentioned earlier the functional amyloids in bacterial biofilms are formed extracellularly. The assembly process therefore happens in close proximity to both cell membranes and the biopolymers constituting the EPS of the biofilm matrix. The interactions between functional amyloids and other biofilm components during the formation of functional amyloids and maturation of biofilms are therefore crucial to understand the biogenesis of biofilms and have thus been a topic of interest.

### Interactions between functional amyloids and lipids

During a study on how the cell membrane of the bacterial cell affects the formation of curli amyloids in *E. coli* it was found that in the presence of lipopolysaccharides (LPS), which are found in the outer membrane of Gram-negative bacteria, the ability of CsgB to nucleate and seed the formation of CsgA is enhanced.<sup>165</sup> This was found to be due to electrostatic interactions between the lipid bilayer and the CsgB protein. Hence the bacterial cell membrane adds a layer of cooperativity to the formation of curli amyloids ensuring that the formation of the functional amyloids occurs at the cell membrane to tether the amyloid structures to the cells and hence keeping the amyloid structures in place in the biofilm.

LPS have also been found to accelerate the aggregation of FapC by bypassing the nucleation lag-phase.<sup>164</sup> This is achieved through reduction of the population of intermediates during aggregation in addition to rearrangement of early aggregates. A similar effect on aggregation is observed for the biosurfactant rhamnolipid which is produced in *Pseudomonas* species alongside the FapC protein. At high concentrations of the rhamnolipid micelles, they were found to be decorated with FapC fibrils

indicating a more long-term interaction between FapC and rhamnolipids.

TasA also interacts differently with bacterial cell membranes compared to eukaryotic cell membranes.<sup>166</sup> In the presence of bacterial cell membranes the fibers formed by TasA are disordered and have a different  $\beta$ -sheet signature than fibers formed in the absence of membranes. In the presence of eukaryotic cell membranes TasA fibers display the same morphology and  $\beta$ -sheet signature as in the absence of membranes indicating that the presence of eukaryotic cell membranes does not affect the TasA fiber formation. Similarly the bacterial membranes were found to deform significantly upon interaction with TasA. Again this effect was not observed for eukaryotic cell membranes suggesting that TasA penetrates the bacterial cell membranes but not eukaryotic cell membranes.<sup>166</sup>

Other functional amyloids have also been found to interact with lipid cell membranes. *S. aureus* PSM peptides are able to cause cell lysis and hence interact with the cell membrane in a variety of eukaryotic cells, including monocytes, leukocytes, erythrocytes, epithelial cells, endothelial cells, and osteoblasts.<sup>58,167,168</sup> The lytic activity of the PSM peptides is closely linked to the degree of  $\alpha$ -helicity of the peptides in the monomeric form<sup>169</sup> and can furthermore be counteracted by serum lipoproteins.<sup>170</sup> Besides the  $\alpha$ -helicity of the peptides the presence of cholesterol also increases the susceptibility to lysis in vesicles containing between 10 and 30 mol% cholesterol.<sup>169</sup> However, when a shift from liquid-disordered to liquid-ordered state occurs in the lipid bilayer upon addition of higher amounts of cholesterol, a decrease in lytic activity from the PSM peptides occurs. Based on these results it was speculated that differences in cholesterol concentration in phagosomes from different cell types and different cell species could impact PSM-mediated phagosome escape and subsequent survival of *S. aureus* during an infection.

The ability of lipid bilayers to alter the aggregation of proteins is not limited to functional amyloids as lipids have also been found to play a key role in the aggregation  $\alpha$ Syn. Recently, Galvagnion and coworkers showed that lipid bilayers enhance the rate of heterogeneous primary nucleation of  $\alpha$ Syn by three orders of magnitude.<sup>171</sup>

### Interactions between functional amyloids and extracellular polymeric substances

Since the formation of functional amyloids occurs extracellularly, other biomolecules of the biofilm matrix come in close proximity with functional amyloids. Interactions with the other matrix components could therefore play a role during the formation of the functional amyloids. The EPS of the biofilm shows high biodiversity with distinct biomolecules being expressed in different bacterial species however eDNA is emerging as a nearly universal component of the biofilm across species even including fungi.<sup>172</sup> eDNA is found to play a role in facilitating adhesion<sup>173</sup> and promoting intercellular aggregation<sup>174</sup> along with providing structural integrity to the biofilm architecture.<sup>175–177</sup>



fibers and also nucleate endogenously added recombinant purified CsgA.<sup>4,40,185</sup> Substoichiometric concentrations of CsgB have been found to alter the mechanism of aggregation of CsgA. In the absence of CsgB, CsgA was found to aggregate through the formation of amorphous aggregates but in the presence of CsgB ordered oligomers with a  $\beta$ -sheet structure were seen, thus resembling the mechanism seen for CsgB in the absence of CsgA.<sup>187</sup>

Through investigation of the individual domains in the proteins the C-terminal parts of both CsgA and CsgB are found to play a crucial role in the nucleation mechanism. Besides being amyloidogenic, the C-terminal repeat 5 (R5) in CsgA has been found to be critical for *in vitro* nucleation of CsgA.<sup>40</sup> The R4 and R5 of CsgB are required to tether the CsgA aggregates to the lipid membrane, and mutants lacking these domains are no longer localized to the membrane but are secreted into the extracellular space. The C-terminal repeat domain, R5, is required to anchor the protein to the cell membrane although this domain in CsgB is not amyloidogenic on its own.<sup>188</sup> In addition, CsgA nucleation by a CsgB truncation mutant lacking the C-terminal 19 amino acids is less efficient than full-length CsgB *in vivo*.<sup>37</sup>

While TasA constitutes the major components of the fibers present in *B. subtilis* biofilm, TapA has been found to co-purify as a minor component of the fibers in a 1 : 100 ratio with TasA.<sup>189</sup> *In vitro* TapA contributes to the polymerization of TasA into fibers. More specifically, a sequence of 8 amino acids in the N-terminal of TapA has been identified as being crucial in the initiation of the self-assembly of TasA.<sup>190</sup> Additionally, the 5 conserved cysteine residues in TapA play a minor role in the formation of a robust biofilm. The presence of TapA has furthermore been shown to be required in order to anchor TasA fibers to the cell surface.<sup>63,189</sup> Incorporation of D-amino acids into the peptidoglycan results in release of TasA fibers from the cell surface and disassembly of the biofilm.<sup>191</sup> Besides TapA, TasA has also been found to interact with BslA during fiber formation in the biofilm. BslA acts synergistically with other matrix components to facilitate the assembly of the biofilm matrix.<sup>70</sup> *In vivo*, overexpression of TasA was not sufficient to overcome the effects of deletion of BslA. Besides aiding in the self-assembly of TasA, BslA also self-assembles on its own at interfaces to form a hydrophobic surface layer around the biofilm.<sup>72</sup>

Although no specific nucleator protein has been identified for the PSM peptides from *S. aureus*, interactions between the individual peptides have been investigated through cross-seeding experiments. No specific pattern in cross-seeding was seen between  $\alpha$ PSMs and  $\beta$ PSMs but PSM $\alpha$ 1 was found to cross-seed all the other PSM peptides.<sup>192</sup> Based on the kinetics of the aggregation of PSM peptides along with the cross-seeding it was speculated that PSM $\alpha$ 3 that aggregates extremely fast is able to kick-start the functional amyloid formation in biofilm. The PSM $\alpha$ 3 aggregates are then able to cross-seed PSM $\alpha$ 1 that normally aggregates much slower in the absence of PSM $\alpha$ 3 aggregates. The PSM $\alpha$ 1 aggregates can then proceed to cross-seed all the other remaining PSM peptides to ensure that even the more slowly aggregating peptides aggregate on a timescale relevant

## Cross-seeding between different functional amyloids

6470 | Chem. Sci., 2022, 13, 6457–6477



for the bacteria during biofilm formation. This ensures that  $\beta$ PSM peptides which have been found to have the most pronounced impact on *S. aureus* biofilm structuring<sup>60</sup> are able to aggregate in a time scale relevant for the bacteria.

This mechanism of less amyloidogenic peptides/proteins being seeded by more highly amyloidogenic peptides/proteins has also been observed for other functional amyloids not involved in biofilm formation. Different members of the CRES-family which comprise part of the complex extracellular amyloid matrix in mouse spermatozoa have been found to cross-seed each other in a manner similar to the PSM peptides and these are also seeding of CsgA by CsgB.<sup>193</sup> The cross-seeding between the CRES subgroup members suggests that it may be an evolutionarily conserved mechanism to control the assembly of some functional amyloids.

### Cross-seeding between functional amyloids and pathological amyloids

Cross-seeding between functional amyloids and pathological amyloids is an emerging interesting research area due to recent discoveries of a link between functional amyloids in the gut and progression in neurodegenerative diseases. During the progression of PD it has been observed that  $\alpha$ Syn aggregates in the form of an  $\alpha$ Syn-immunoreactive Lewy body and Lewy neurite pathology spreads through the brain in a progressive manner from the brainstem to the telencephalon.<sup>194</sup> In the earliest stages aggregates are detected in the olfactory bulb, as well as in the dorsal motor nucleus of the vagus nerve in the medulla oblongata. In later stages  $\alpha$ Syn aggregate pathology is found more widespread in the brainstem *via* the pons and midbrain, in the basal forebrain and, ultimately, in the neocortex. The spreading of aggregated species of  $\alpha$ Syn in mice has been demonstrated to happen through cell-to-cell transmission of pathologic  $\alpha$ Syn aggregates in anatomically interconnected regions upon injections of synthetic  $\alpha$ Syn fibers leading to nigrostriatal degeneration.<sup>195</sup> Furthermore, the injections of  $\alpha$ Syn aggregates in rats and the subsequent pattern of  $\alpha$ Syn accumulation observed suggests that the cell-to-cell spreading occurs only through retrograde transmission.<sup>196</sup> In rodent models the cell-to-cell spreading of  $\alpha$ Syn aggregates has also been used to demonstrate a spreading and propagation from neurons in the gut to the brain through the vagus nerve and/or the spinal cord.<sup>197–199</sup> Furthermore, the bacterial composition of the gut microbiome has been observed to lead to increased neuroinflammation and  $\alpha$ Syn aggregate deposits in the brain of  $\alpha$ Syn expressing mice. As a link emerges between neurons in the gut and the brain and the possible spread of pathological amyloid species, the interactions between functional amyloids formed from bacteria and pathological amyloids prove to be an interesting topic highly related to bacterial biofilm forming functional amyloids.<sup>200</sup>

In mice it was found that monomeric CsgA was able to accelerate the aggregation of  $\alpha$ Syn even under substoichiometric conditions.<sup>200</sup> The accelerated aggregation occurs through transient interactions between CsgA and  $\alpha$ Syn. In the same mouse model Tau aggregation was not found to be

accelerated by CsgA monomers. When  $\alpha$ Syn is overexpressed in mice it was seen that simultaneous colonization with curli-producing *E. coli* in the gut promotes  $\alpha$ Syn pathology in both the gut and the brain. Furthermore, curli expression is required for *E. coli* to intensify  $\alpha$ Syn-induced behavioral deficits, including intestinal and motor impairments in the mice. Interestingly CsgA is not the only component of the curli apparatus found to interact with  $\alpha$ Syn. CsgC and CsgE, both of which act as periplasmic chaperones and prevent aggregation of CsgA intracellularly,<sup>4,38</sup> also interact with  $\alpha$ Syn although in different ways. CsgC inhibits the aggregation of  $\alpha$ Syn while CsgE surprisingly accelerates the aggregation of  $\alpha$ Syn.<sup>201</sup>

PD is not the only neurodegenerative disease to be linked to functional bacterial amyloids, as CsgA pre-formed seeds have been found to also accelerate the aggregation of A $\beta$ , and fibrillation inhibitors are shared between CsgA and A $\beta$ .<sup>202</sup> Curli cross-seeding exerted a complicated concentration dependent effect on A $\beta$ 1–40 fibrillogenesis kinetics.<sup>203</sup>

In human semen fragments of prostatic acid phosphatase (PAP248–286) increase HIV infection efficiency through increased virus adhesion to target cells. This enhancement of infection only occurs when PAP248–286 is in the form of amyloid aggregates termed SEVI (Semen Enhancer of Viral Infection), however monomeric PAP248–286 aggregates very slowly and exogenous factors have been suggested as a promoter of SEVI fiber formation *in vivo*.<sup>203</sup> Indeed it has been demonstrated that curli amyloids of both CsgA and CsgB can act as a catalyst for SEVI formation from PAP248–286. At low concentrations *in vitro*, cross-seeding with curli fibers results in fibers of PAP248–286 that retain the ability to enhance HIV infection. Kinetic analysis of the cross-seeding of PAP248–286 with curli moderately affects the nucleation rate while significantly enhancing the growth of fibers from existing nuclei and hence the elongation rate. During the study it was also shown that curli amyloids of both CsgA and CsgB were able to cross-seed IAPP leading to a decreased lag-time but simultaneously a strong inhibition is observed in IAPP elongation.<sup>203</sup>

FapC has also been found to interact with proteins associated with neurodegenerative diseases, specifically  $\alpha$ Syn and A $\beta$ . *In vitro* FapC can seed A $\beta$  while templating the structure of the resulting A $\beta$  amyloid fibers to resemble that of FapC fibers.<sup>204</sup> *In vivo* in zebrafish larvae FapC was able to induce accelerated A $\beta$  aggregation and A $\beta$  pathology. In adult zebrafish FapC was found to accelerate A $\beta$ -induced cognitive pathology. To extend the results to the bacterial biofilm it was found that co-injection of A $\beta$  and the protein part of *P. aeruginosa* biofilm samples into adult zebrafish elevated A $\beta$  pathology. This elevated pathology was not observed in zebrafish injected with only the biofilm protein sample. Based on these results A $\beta$  can also interact with the fibers when they are present in a complex setting such as a biofilm despite the morphology of the biofilm protein sample being different from that of FapC. The changes in morphology were attributed to the presence of other structural components of the biofilm in the sample.<sup>204</sup>

FapC is also capable of interacting with  $\alpha$ Syn. FapC binds *in vitro* to  $\alpha$ Syn monomers, oligomers, and aggregates.<sup>205</sup> Full-length FapC protein slightly accelerates  $\alpha$ Syn aggregation



while the FapC deletion mutant where all three repeat regions have been deleted and hence only the loop regions remain ( $\Delta R1R2R3$ ) increases the lag-time of  $\alpha$ Syn aggregation. The fibrillation inhibition was found to be due to the formation of hetero-oligomers possibly enhanced by disulfide bond formation in FapC  $\Delta R1R2R3$ . Other deletion mutants where only one or two repeat regions had been deleted did not increase the lag-time of  $\alpha$ Syn. Interestingly, only the monomeric protein is responsible for the interactions between FapC and  $\alpha$ Syn as no seeding effects were seen for FapC aggregates on  $\alpha$ Syn aggregation.

Although several links have been found between functional bacterial amyloids and neurodegenerative diseases, it cannot be concluded that biofilm formation is involved in the causative effects of such neurodegenerative diseases. For example, it has also been shown that biofilm can actually protect against protein aggregation. In *C. elegans* the probiotic *B. subtilis* has been found to inhibit  $\alpha$ Syn aggregation by altering the host sphingolipid metabolism and thereby reduce  $\alpha$ Syn aggregation through biofilm formation in the gut.<sup>206</sup>

In summary, functional amyloids are found to interact with a wide variety of other proteins both from bacteria and from humans. They are even emerging as contributing factors involved in the development of the pathology of various human diseases with pathology linked to protein aggregation.

## Concluding remarks

It is becoming increasingly clear that functional bacterial amyloids involved in biofilm formation are not solitary inert structural components in the biofilm. Cross-talk between different aggregated species is emerging as a common theme and not only a phenomenon observed between various functional amyloids and pathological amyloids. Self-assembled structures of various metabolites have also been found to induce aggregation of pathological amyloid species. Fibrillar aggregates of homocysteine have been found to display cross-talk with A $\beta$ ,<sup>207</sup> and amyloid-like aggregates of quinolinic acid have been observed to cross-seed  $\alpha$ Syn aggregation.<sup>208</sup> Despite the recent focus and many new insights gained in the field, the molecular mechanisms underlining the interactions between different amyloid species are very limited. This lack of molecular insight limits possibilities to develop strategies to better combat biofilm related bacterial infections. We hope that this review will make a contribution towards a more complete understanding.

Our fight against AMR and biofilm related infections relies strongly on our knowledge of the molecular details of biofilm forming functional amyloids. With the recent fast pace and exponential growth in biofilm, amyloid fibril and functional amyloid related research we believe that the future is very bright. As of today, structural and mechanistic models are predominantly still missing, along with high-resolution structures of the key elements of this complex network in biofilms. Recent developments in the field of structural biology by cryo-EM and solid-state NMR will pave the way towards our atomic level understanding of proteins *in vitro* and *in vivo*. All in all, we

will be able to understand, manipulate and fight persistent infections, AMR, biofilms and amyloids forming them.

## Author contributions

The manuscript was conceived, designed and written by UA and MA with equal contributions.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We acknowledge Roberto Kolter for the permission to reuse the *B. subtilis* biofilm pictures, and the inspiration for our biofilm illustration. We thank Daniel Lopez for additional biofilm pictures. Frans Mulder and Daniel Otzen are acknowledged for supplying the solution NMR spectrum of monomeric FapC and helpful discussion. Robert Tycko is acknowledged for supplying the solid-state NMR spectrum of CsgA. We are thankful to Kresten Lindorff-Larsen for the PDB-like structure file of CsgA and to Morten S. Dueholm for discussions on FapC structure model. Meytal Landau is acknowledged for providing the assembly crystal structure file for CsgA 6G9G fragment. UA acknowledges Anne Diehl, Madhu Nagaraj, Kürsat Turgay, Matthias Hiller and Hartmut Oshkinat for making the initial stages of functional amyloid work possible. UA acknowledges financial support from the DFF Mobilex and Aarhus University AIAS Cofund grants. MA gratefully acknowledges financial support from Aarhus University Research Foundation AUFF-E-2017-7-16 and L'oreal and Unesco For Women in Science National Award (Denmark).

## References

- 1 F. Chiti and C. M. Dobson, *Annu. Rev. Biochem.*, 2006, **75**, 333–366.
- 2 F. Chiti and C. M. Dobson, *Annu. Rev. Biochem.*, 2017, **86**, 27–68.
- 3 J. Oh, J. G. Kim, E. Jeon, C. H. Yoo, J. S. Moon, S. Rhee and I. Hwang, *J. Biol. Chem.*, 2007, **282**, 13601–13609.
- 4 M. R. Chapman, L. S. Robinson, J. S. Pinkner, R. Roth, J. Heuser, M. Hammar, S. Normark and S. J. Hultgren, *Science*, 2002, **295**, 851–855.
- 5 S. K. Maji, M. H. Perrin, M. R. Sawaya, S. Jessberger, K. Vadodaria, R. A. Rissman, P. S. Singru, K. P. Nilsson, R. Simon, D. Schubert, D. Eisenberg, J. Rivier, P. Sawchenko, W. Vale and R. Riek, *Science*, 2009, **325**, 328–332.
- 6 J. F. Berson, A. C. Theos, D. C. Harper, D. Tenza, G. Raposo and M. S. Marks, *J. Cell Biol.*, 2003, **161**, 521–533.
- 7 K. Si, S. Lindquist and E. R. Kandel, *Cell*, 2003, **115**, 879–891.
- 8 V. Coustou, C. Deleu, S. Saupe and J. Begueret, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 9773–9778.





- 9 H. C. Flemming and J. Wingender, *Nat. Rev. Microbiol.*, 2010, **8**, 623–633.
- 10 M. Malone, T. Bjarnsholt, A. J. McBain, G. A. James, P. Stoodley, D. Leaper, M. Tachi, G. Schultz, T. Swanson and R. D. Wolcott, *J. Wound Care*, 2017, **26**, 20–25.
- 11 B. C. Wu, E. F. Haney, N. Akhoundsadegh, D. Pletzer, M. J. Trimble, A. E. Adriaans, P. H. Nibbering and R. E. Hancock, *npj Biofilms Microbiomes*, 2021, **7**, 1–13.
- 12 T. F. Mah and G. A. O'Toole, *Trends Microbiol.*, 2001, **9**, 34–39.
- 13 C. J. Murray, K. S. Ikuta, F. Sharara, L. Swetschinski, G. R. Aguilar, A. Gray, C. Han, C. Bisignano, P. Rao and E. Wool, *Lancet*, 2022, **399**, 629–655.
- 14 R. Gallardo, N. A. Ranson and S. E. Radford, *Curr. Opin. Struct. Biol.*, 2020, **60**, 7–16.
- 15 S. M. Ulamec, D. J. Brockwell and S. E. Radford, *Front. Neurosci.*, 2020, **14**, 611285.
- 16 M. R. Sawaya, M. P. Hughes, J. A. Rodriguez, R. Riek and D. S. Eisenberg, *Cell*, 2021, **184**, 4857–4873.
- 17 N. Salinas, T. L. Povolotsky, M. Landau and I. Kolodkin-Gal, *Microbiol. Mol. Biol. Rev.*, 2020, **85**, e00062–00020.
- 18 P. Ragonis-Bachar and M. Landau, *Curr. Opin. Struct. Biol.*, 2021, **68**, 184–193.
- 19 S. A. Levkovich, E. Gazit and D. Laor Bar-Yosef, *Trends Microbiol.*, 2021, **29**, 251–265.
- 20 P. C. Ke, R. Zhou, L. C. Serpell, R. Riek, T. P. Knowles, H. A. Lashuel, E. Gazit, I. W. Hamley, T. P. Davis and M. Fändrich, *Chem. Soc. Rev.*, 2020, **49**, 5473–5509.
- 21 V. A. Sergeeva, N. V. Zakharova, A. E. Bugrova, N. L. Starodubtseva, M. I. Indeykina, A. S. Kononikhin, V. E. Frankevich and E. N. Nikolaev, *Eur. J. Mass Spectrom.*, 2020, **26**, 158–161.
- 22 M. S. Rubel, S. A. Fedotov, A. V. Grizel, J. V. Sopova, O. A. Malikova, Y. O. Chernoff and A. A. Rubel, *Life*, 2020, **10**, 156.
- 23 M. Bokhove, D. Claessen, W. de Jong, L. Dijkhuizen, E. J. Boekema and G. T. Oostergetel, *J. Struct. Biol.*, 2013, **184**, 301–309.
- 24 W. De Jong, H. A. Wösten, L. Dijkhuizen and D. Claessen, *Mol. Microbiol.*, 2009, **73**, 1128–1140.
- 25 C. J. Alteri, J. Xicohtencatl-Cortes, S. Hess, G. Caballero-Olin, J. A. Giron and R. L. Friedman, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 5145–5150.
- 26 G. G. Sgro, F. A. Ficarra, G. Dunger, T. E. Scarpeci, E. M. Valle, A. Cortadi, E. G. Orellano, N. Gottig and J. Ottado, *Mol. Plant Pathol.*, 2012, **13**, 1047–1059.
- 27 A. Taglialegna, L. Matilla-Cuenca, P. Dorado-Morales, S. Navarro, S. Ventura, J. A. Garnett, I. Lasa and J. Valle, *npj Biofilms Microbiomes*, 2020, **6**, 15.
- 28 D. Romero and R. Kolter, *Int. Microbiol.*, 2014, **17**, 65–73.
- 29 C. Jeter and A. G. Matthysse, *Mol. Plant-Microbe Interact.*, 2005, **18**, 1235–1242.
- 30 J. H. Ryu and L. R. Beuchat, *Appl. Environ. Microbiol.*, 2005, **71**, 247–254.
- 31 J. H. Ryu, H. Kim, J. F. Frank and L. R. Beuchat, *Lett. Appl. Microbiol.*, 2004, **39**, 359–362.
- 32 C. Prigent-Combaret, G. Prensier, T. T. Le Thi, O. Vidal, P. Lejeune and C. Dorel, *Environ. Microbiol.*, 2000, **2**, 450–464.
- 33 D. Otzen and P. H. Nielsen, *Cell. Mol. Life Sci.*, 2008, **65**, 910–927.
- 34 M. F. Gebbink, D. Claessen, B. Bouma, L. Dijkhuizen and H. A. Wosten, *Nat. Rev. Microbiol.*, 2005, **3**, 333–341.
- 35 M. M. Barnhart and M. R. Chapman, *Annu. Rev. Microbiol.*, 2006, **60**, 131–147.
- 36 S. Kanamaru, H. Kurazono, A. Terai, K. Monden, H. Kumon, Y. Mizunoe, O. Ogawa and S. Yamamoto, *Int. J. Antimicrob. Agents*, 2006, **28**(Suppl 1), S21–S25.
- 37 N. D. Hammer, J. C. Schmidt and M. R. Chapman, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 12494–12499.
- 38 M. L. Evans, E. Chorell, J. D. Taylor, J. Aden, A. Gotheson, F. Li, M. Koch, L. Sefer, S. J. Matthews, P. Wittung-Stafshede, F. Almqvist and M. R. Chapman, *Mol. Cell*, 2015, **57**, 445–455.
- 39 A. A. Nenninger, L. S. Robinson, N. D. Hammer, E. A. Epstein, M. P. Badtke, S. J. Hultgren and M. R. Chapman, *Mol. Microbiol.*, 2011, **81**, 486–499.
- 40 X. Wang, N. D. Hammer and M. R. Chapman, *J. Biol. Chem.*, 2008, **283**, 21530–21539.
- 41 X. Wang and M. R. Chapman, *J. Mol. Biol.*, 2008, **380**, 570–580.
- 42 D. Otzen and R. Riek, *Cold Spring Harbor Perspect. Biol.*, 2019, **11**, a033860.
- 43 A. Olsen, H. Herwald, M. Wikstrom, K. Persson, E. Mattsson and L. Bjorck, *J. Biol. Chem.*, 2002, **277**, 34568–34572.
- 44 X. Wang, D. R. Smith, J. W. Jones and M. R. Chapman, *J. Biol. Chem.*, 2007, **282**, 3713–3719.
- 45 M. S. Dueholm, S. V. Petersen, M. Sonderkaer, P. Larsen, G. Christiansen, K. L. Hein, J. J. Enghild, J. L. Nielsen, K. L. Nielsen, P. H. Nielsen and D. E. Otzen, *Mol. Microbiol.*, 2010, **77**, 1009–1020.
- 46 M. S. Dueholm, M. T. Sondergaard, M. Nilsson, G. Christiansen, A. Stensballe, M. T. Overgaard, M. Givskov, T. Tolker-Nielsen, D. E. Otzen and P. H. Nielsen, *MicrobiologyOpen*, 2013, **2**, 365–382.
- 47 K. L. Palmer, L. M. Mashburn, P. K. Singh and M. Whiteley, *J. Bacteriol.*, 2005, **187**, 5267–5277.
- 48 M. S. Dueholm, D. Otzen and P. H. Nielsen, *PLoS One*, 2013, **8**, e76630.
- 49 P. H. Nielsen, M. S. Dueholm, T. R. Thomsen, J. L. Nielsen and D. Otzen, in *Biofilm highlights*, Springer, 2011, pp. 41–62.
- 50 S. L. Rouse, W. J. Hawthorne, J. L. Berry, D. S. Chorev, S. A. Ionescu, S. Lambert, F. Stylianou, W. Ewert, U. Mackie, R. M. L. Morgan, D. Otzen, F. A. Herbst, P. H. Nielsen, M. Dueholm, H. Bayley, C. V. Robinson, S. Hare and S. Matthews, *Nat. Commun.*, 2017, **8**, 263.
- 51 S. L. Rouse, W. J. Hawthorne, S. Lambert, M. L. Morgan, S. A. Hare and S. Matthews, *Acta Crystallogr., Sect. F: Struct. Biol. Commun.*, 2016, **72**, 892–896.
- 52 L. F. B. Christensen, J. S. Nowak, T. V. Sonderby, S. A. Frank and D. E. Otzen, *J. Biol. Chem.*, 2020, **295**, 13031–13046.



- 53 C. B. Rasmussen, G. Christiansen, B. S. Vad, C. Lynggaard, J. J. Enghild, M. Andreassen and D. Otzen, *Protein Sci.*, 2019, **28**, 633–642.
- 54 T. V. Sonderby, H. O. Rasmussen, S. A. Frank, J. Skov Pedersen and D. E. Otzen, *J. Mol. Biol.*, 2021, **434**, 167337.
- 55 K. Schwartz, A. K. Syed, R. E. Stephenson, A. H. Rickard and B. R. Boles, *PLoS Pathog.*, 2012, **8**, e1002744.
- 56 A. Peschel and M. Otto, *Nat. Rev. Microbiol.*, 2013, **11**, 667–673.
- 57 A. Bleem, R. Francisco, J. D. Bryers and V. Daggett, *npj Biofilms Microbiomes*, 2017, **3**, 16.
- 58 R. Wang, K. R. Braughton, D. Kretschmer, T. H. Bach, S. Y. Queck, M. Li, A. D. Kennedy, D. W. Dorward, S. J. Klebanoff, A. Peschel, F. R. DeLeo and M. Otto, *Nat. Med.*, 2007, **13**, 1510–1514.
- 59 E. Tayeb-Fligelman, O. Tabachnikov, A. Moshe, O. Goldshmidt-Tran, M. R. Sawaya, N. Coquelle, J. P. Colletier and M. Landau, *Science*, 2017, **355**, 831–833.
- 60 S. Periasamy, H. S. Joo, A. C. Duong, T. H. Bach, V. Y. Tan, S. S. Chatterjee, G. Y. Cheung and M. Otto, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 1281–1286.
- 61 N. Salinas, J. P. Colletier, A. Moshe and M. Landau, *Nat. Commun.*, 2018, **9**, 3512.
- 62 S. S. Branda, F. Chu, D. B. Kearns, R. Losick and R. Kolter, *Mol. Microbiol.*, 2006, **59**, 1229–1238.
- 63 D. Romero, C. Aguilar, R. Losick and R. Kolter, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 2230–2234.
- 64 A. Driks, *Mol. Microbiol.*, 2011, **80**, 1133–1136.
- 65 N. R. Stanley and B. A. Lazazzera, *Mol. Microbiol.*, 2004, **52**, 917–924.
- 66 F. Chu, D. B. Kearns, S. S. Branda, R. Kolter and R. Losick, *Mol. Microbiol.*, 2006, **59**, 1216–1228.
- 67 A. G. Stover and A. Driks, *J. Bacteriol.*, 1999, **181**, 1664–1672.
- 68 A. G. Stover and A. Driks, *J. Bacteriol.*, 1999, **181**, 7065–7069.
- 69 R. Terra, N. R. Stanley-Wall, G. Cao and B. A. Lazazzera, *J. Bacteriol.*, 2012, **194**, 2781–2790.
- 70 A. Ostrowski, A. Mehert, A. Prescott, T. B. Kiley and N. R. Stanley-Wall, *J. Bacteriol.*, 2011, **193**, 4821–4831.
- 71 K. Kobayashi and M. Iwano, *Mol. Microbiol.*, 2012, **85**, 51–66.
- 72 L. Hobley, A. Ostrowski, F. V. Rao, K. M. Bromley, M. Porter, A. R. Prescott, C. E. MacPhee, D. M. van Aalten and N. R. Stanley-Wall, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 13600–13605.
- 73 J. W. Costerton, P. S. Stewart and E. P. Greenberg, *Science*, 1999, **284**, 1318–1322.
- 74 H. C. Flemming, J. Wingender, U. Szewzyk, P. Steinberg, S. A. Rice and S. Kjelleberg, *Nat. Rev. Microbiol.*, 2016, **14**, 563–575.
- 75 C. M. Dobson, *Trends Biochem. Sci.*, 1999, **24**, 329–332.
- 76 A. Berry and S. E. Radford, *Biochem. Soc. Symp.*, 2001, **68**, 1–26.
- 77 D. M. Fowler, A. V. Koulov, C. Alory-Jost, M. S. Marks, W. E. Balch and J. W. Kelly, *PLoS Biol.*, 2006, **4**, e6.
- 78 D. M. Fowler, A. V. Koulov, W. E. Balch and J. W. Kelly, *Trends Biochem. Sci.*, 2007, **32**, 217–224.
- 79 L. Giehm, N. Lorenzen and D. E. Otzen, *Methods*, 2011, **53**, 295–305.
- 80 J. D. Sipe, M. D. Benson, J. N. Buxbaum, S. Ikeda, G. Merlini, M. J. M. Saraiva and P. Westermark, *Amyloid*, 2014, **21**, 221–224.
- 81 A. Diehl, Y. Roske, L. Ball, A. Chowdhury, M. Hiller, N. Moliere, R. Kramer, D. Stoppler, C. L. Worth, B. Schlegel, M. Leidert, N. Cremer, N. Erdmann, D. Lopez, H. Stephanowitz, E. Krause, B. J. van Rossum, P. Schmieder, U. Heinemann, K. Turgay, U. Akbey and H. Oschkinat, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, 3237–3242.
- 82 M. Torok, S. Milton, R. Kaye, P. Wu, T. McIntire, C. G. Glabe and R. Langen, *J. Biol. Chem.*, 2002, **277**, 40810–40815.
- 83 V. N. Uversky and A. L. Fink, *Biochim. Biophys. Acta, Proteins Proteomics*, 2004, **1698**, 131–153.
- 84 R. Nelson, M. R. Sawaya, M. Balbirnie, A. O. Madsen, C. Riekel, R. Grothe and D. Eisenberg, *Nature*, 2005, **435**, 773–778.
- 85 W. T. Astbury, S. Dickinson and K. Bailey, *Biochem. J.*, 1935, **29**, 2351–2360.
- 86 A. J. Geddes, K. D. Parker, E. D. T. Atkins and E. Beighton, *J. Mol. Biol.*, 1968, **32**, 343–344.
- 87 M. Fandrich and C. M. Dobson, *EMBO J.*, 2002, **21**, 5682–5690.
- 88 M. Biancalana and S. Koide, *Biochim. Biophys. Acta*, 2010, **1804**, 1405–1412.
- 89 M. Groenning, *J. Chem. Biol.*, 2010, **3**, 1–18.
- 90 H. LeVine, *Amyloid*, 1995, **2**, 1–6.
- 91 A. M. Morris, M. A. Watzky and R. G. Finke, *Biochim. Biophys. Acta*, 2009, **1794**, 375–397.
- 92 C. C. F. Blake, L. C. Serpell, M. Sunde, O. Sandgren and E. Lundgren, *Nature and Origin of Amyloid Fibrils*, 1996, vol. 199, pp. 6–21.
- 93 M. Sunde and C. Blake, in *Advances in Protein Chemistry*, ed. R. Wetzel, 1997, vol. 50, Protein Misassembly, pp. 123–159.
- 94 L. C. Serpell, C. C. F. Blake and P. E. Fraser, *Biochemistry*, 2000, **39**, 13269–13275.
- 95 N. P. Reynolds, J. Adamcik, J. T. Berryman, S. Handschin, A. A. H. Zanjani, W. Li, K. Liu, A. Zhang and R. Mezzenga, *Nat. Commun.*, 2017, **8**, 1338.
- 96 C. P. Jaronec, C. E. MacPhee, V. S. Bajaj, M. T. McMahon, C. M. Dobson and R. G. Griffin, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 711–716.
- 97 R. Tycko and R. B. Wickner, *Acc. Chem. Res.*, 2013, **46**, 1487–1496.
- 98 F. Shewmaker, R. P. McGlinchey and R. B. Wickner, *J. Biol. Chem.*, 2011, **286**, 16533–16540.
- 99 C. Wasmer, A. Lange, H. Van Melckebeke, A. B. Siemer, R. Riek and B. H. Meier, *Science*, 2008, **319**, 1523–1526.
- 100 R. Guerrero-Ferreira, N. M. Taylor, D. Mona, P. Ringler, M. E. Lauer, R. Riek, M. Britschgi and H. Stahlberg, *eLife*, 2018, **7**, e36402.
- 101 M. A. Walti, F. Ravotti, H. Arai, C. G. Glabe, J. S. Wall, A. Bockmann, P. Guntert, B. H. Meier and R. Riek, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, E4976–E4984.



- 102 S. Campioni, B. Mannini, M. Zampagni, A. Pensalfini, C. Parrini, E. Evangelisti, A. Relini, M. Stefani, C. M. Dobson, C. Cecchi and F. Chiti, *Nat. Chem. Biol.*, 2010, **6**, 140–147.
- 103 P. G. Wolynes, *Biochimie*, 2015, **119**, 218–230.
- 104 H. Van Melckebeke, C. Wasmer, A. Lange, E. Ab, A. Loquet, A. Bockmann and B. H. Meier, *J. Am. Chem. Soc.*, 2010, **132**, 13765–13775.
- 105 D. Li and C. Liu, *Nat. Chem. Biol.*, 2021, **17**, 237–245.
- 106 M. Fandrich, S. Nystrom, K. P. R. Nilsson, A. Bockmann, H. LeVine and P. Hammarstrom, *J. Intern. Med.*, 2018, **283**, 218–237.
- 107 C. Seuring, J. Verasdonck, P. Ringler, R. Cadalbert, H. Stahlberg, A. Bockmann, B. H. Meier and R. Riek, *J. Phys. Chem. B*, 2021, **125**, 484.
- 108 D. T. Murray, M. Kato, Y. Lin, K. R. Thurber, I. Hung, S. L. McKnight and R. Tycko, *Cell*, 2017, **171**, 615–627.
- 109 M. P. Hughes, M. R. Sawaya, D. R. Boyer, L. Goldschmidt, J. A. Rodriguez, D. Cascio, L. Chong, T. Gonen and D. S. Eisenberg, *Science*, 2018, **359**, 698–701.
- 110 R. Hervas, M. J. Rau, Y. Park, W. J. Zhang, A. G. Murzin, J. A. J. Fitzpatrick, S. H. W. Scheres and K. Si, *Science*, 2020, **367**, 1230.
- 111 M. Andreasen, G. Meisl, J. D. Taylor, T. C. T. Michaels, A. Levin, D. E. Otzen, M. R. Chapman, C. M. Dobson, S. J. Matthews and T. P. J. Knowles, *mBio*, 2019, **10**, e02279-18.
- 112 S. H. Alijanvand, A. Peduzzo and A. K. Buell, *Front. Mol. Biosci.*, 2021, **8**, 669994.
- 113 M. R. Sawaya, S. Sambashivan, R. Nelson, M. I. Ivanova, S. A. Sievers, M. I. Apostol, M. J. Thompson, M. Balbirnie, J. J. W. Wiltzius, H. T. McFarlane, A. O. Madsen, C. Riekel and D. Eisenberg, *Nature*, 2007, **447**, 453–457.
- 114 D. Willbold, B. Strodel, G. F. Schroder, W. Hoyer and H. Heise, *Chem. Rev.*, 2021, **121**, 8285–8307.
- 115 E. Nogales and S. H. W. Scheres, *Mol. Cell*, 2015, **58**, 677–689.
- 116 E. Callaway, *Nature*, 2015, **525**, 172–174.
- 117 W. Kuhlbrandt, *Science*, 2014, **343**, 1443–1444.
- 118 K. M. Towle, C. T. Lohans, M. Miskolzie, J. Z. Acedo, M. J. van Belkum and J. C. Vederas, *Biochemistry*, 2016, **55**, 4798–4806.
- 119 N. E. Mammeri, J. Hierrezuelo, J. Tolchard, J. Cámara-Almirón, J. Caro-Astorga, A. Álvarez-Mena, A. Dutour, M. Berbon, J. Shenoy and E. Morvan, *FASEB J.*, 2019, **33**, 12146–12163.
- 120 L. Sewell, F. Stylianou, Y. Xu, J. Taylor, L. Sefer and S. Matthews, *Sci. Rep.*, 2020, **10**, 7896.
- 121 T. Konuma, E. Chatani, M. Yagi, K. Sakurai, T. Ikegami, H. Naiki and Y. Goto, *J. Mol. Biol.*, 2011, **405**, 851–862.
- 122 N. A. Whittemore, R. Mishra, I. Kheterpal, A. D. Williams, R. Wetzler and E. H. Serpersu, *Biochemistry*, 2005, **44**, 4434–4441.
- 123 R. Tycko, in *Annual Review of Physical Chemistry*, ed. S. R. Leone, P. S. Cremer, J. T. Groves and M. A. Johnson, 2011, vol. 62, pp. 279–299.
- 124 B. H. Meier, R. Riek and A. Bockmann, *Trends Biochem. Sci.*, 2017, **42**, 777–787.
- 125 L. Bousset, L. Pieri, G. Ruiz-Arlandis, J. Gath, P. H. Jensen, B. Habenstein, K. Madiona, V. Olieric, A. Bockmann, B. H. Meier and R. Melki, *Nat. Commun.*, 2013, **4**, 2575.
- 126 R. Guerrero-Ferreira, N. M. I. Taylor, A. A. Arteni, P. Kumari, D. Mona, P. Ringler, M. Britschgi, M. E. Lauer, A. Makky, J. Verasdonck, R. Riek, R. Melki, B. H. Meier, A. Bockmann, L. Bousset and H. Stahlberg, *Elife*, 2019, **8**, e48907.
- 127 F. Castellani, B. van Rossum, A. Diehl, M. Schubert, K. Rehbein and H. Oschkinat, *Nature*, 2002, **420**, 98–102.
- 128 A. K. Schutz, T. Vagt, M. Huber, O. Y. Ovchinnikova, R. Cadalbert, J. Wall, P. Guntert, A. Bockmann, R. Glockshuber and B. H. Meier, *Angew. Chem., Int. Ed.*, 2015, **54**, 331–335.
- 129 M. D. Tuttle, G. Comellas, A. J. Nieuwkoop, D. J. Covell, D. A. Berthold, K. D. Kloepper, J. M. Courtney, J. K. Kim, A. M. Barclay, A. Kendall, W. Wan, G. Stubbs, C. D. Schwieters, V. M. Y. Lee, J. M. George and C. M. Rienstra, *Nat. Struct. Mol. Biol.*, 2016, **23**, 409–415.
- 130 V. Chevelkov, K. Rehbein, A. Diehl and B. Reif, *Angew. Chem., Int. Ed.*, 2006, **45**, 3878–3881.
- 131 U. Akbey, S. Lange, W. T. Franks, R. Linser, K. Rehbein, A. Diehl, B. J. van Rossum, B. Reif and H. Oschkinat, *J. Biomol. NMR*, 2010, **46**, 67–73.
- 132 J. R. Lewandowski, J. N. Dumez, U. Akbey, S. Lange, L. Emsley and H. Oschkinat, *J. Phys. Chem. Lett.*, 2011, **2**, 2205–2211.
- 133 V. Agarwal, S. Penzel, K. Szekely, R. Cadalbert, E. Testori, A. Oss, J. Past, A. Samoson, M. Ernst, A. Bockmann and B. H. Meier, *Angew. Chem., Int. Ed.*, 2014, **53**, 12253–12256.
- 134 S. K. Vasa, P. Rovo and R. Linser, *Acc. Chem. Res.*, 2018, **51**, 1386–1395.
- 135 E. Barbet-Massin, A. J. Pell, J. S. Retel, L. B. Andreas, K. Jaudzems, W. T. Franks, A. J. Nieuwkoop, M. Hiller, V. Higman, P. Guerry, A. Bertarello, M. J. Knight, M. Felletti, T. Le Marchand, S. Kotelovica, I. Akopjana, K. Tars, M. Stoppini, V. Bellotti, M. Bolognesi, S. Ricagno, J. J. Chou, R. G. Griffin, H. Oschkinat, A. Lesage, L. Emsley, T. Herrmann and G. Pintacuda, *J. Am. Chem. Soc.*, 2014, **136**, 12489–12497.
- 136 K. Xue, R. Sarkar, C. Motz, S. Asami, D. C. R. Camargo, V. Decker, S. Wegner, Z. Tosner and B. Reif, *Sci. Rep.*, 2017, **7**, 7444.
- 137 C. M. Quinn, M. Z. Wang and T. Polenova, in *Protein NMR: Methods and Protocols*, ed. R. Ghose, 2018, vol. 1688, pp. 1–35.
- 138 B. Reif, S. E. Ashbrook, L. Emsley and M. Hong, *Nat. Rev. Methods Primers*, 2021, **1**, 2.
- 139 A. Konig, D. Scholzel, B. Uluca, T. Viennet, U. Akbey and H. Heise, *Solid State Nucl. Magn. Reson.*, 2019, **98**, 1–11.
- 140 U. Akbey and H. Oschkinat, *J. Magn. Reson.*, 2016, **269**, 213–224.
- 141 K. K. Frederick, *Biophys. J.*, 2019, **116**, 203A.
- 142 K. Jaudzems, T. Polenova, G. Pintacuda, H. Oschkinat and A. Lesage, *J. Struct. Biol.*, 2019, **206**, 90–98.



- 143 U. Akbey, W. T. Franks, A. Linden, S. Lange, R. G. Griffin, B. J. van Rossum and H. Oschkinat, *Angew. Chem., Int. Ed.*, 2010, **49**, 7803–7806.
- 144 A. L. Cogen, K. Yamasaki, J. Muto, K. M. Sanchez, L. C. Alexander, J. Tanios, Y. P. Lai, J. E. Kim, V. Nizet and R. L. Gallo, *PLoS One*, 2010, **5**, e8557.
- 145 E. Tayeb-Fligelman, N. Salinas, O. Tabachnikov and M. Landau, *Structure*, 2020, **28**, 301–313.
- 146 P. Marinelli, I. Pallares, S. Navarro and S. Ventura, *Sci. Rep.*, 2016, **6**, 34552.
- 147 M. Zaman and M. Andreassen, *Microorganisms*, 2021, **9**, 117.
- 148 F. Shewmaker, R. P. McGlinchey, K. R. Thurber, P. McPhie, F. Dyda, R. Tycko and R. B. Wickner, *J. Biol. Chem.*, 2009, **284**, 25065–25076.
- 149 T. Schubeis, P. Yuan, M. Ahmed, M. Nagaraj, B. J. van Rossum and C. Ritter, *Angew. Chem., Int. Ed. Engl.*, 2015, **54**, 14669–14672.
- 150 D. S. Wishart, B. D. Sykes and F. M. Richards, *J. Mol. Biol.*, 1991, **222**, 311–333.
- 151 D. S. Wishart, C. G. Bigam, A. Holm, R. S. Hodges and B. D. Sykes, *J. Biomol. NMR*, 1995, **5**, 67–81.
- 152 P. Tian, W. Boomsma, Y. Wang, D. E. Otzen, M. H. Jensen and K. Lindorff-Larsen, *J. Am. Chem. Soc.*, 2015, **137**, 22–25.
- 153 A. Daskalov, N. El Mammeri, A. Lends, J. Shenoy, G. Lamon, Y. Fichou, A. Saad, D. Martinez, E. Morvan, M. Berbon, A. Grelard, B. Kauffmann, M. Ferber, B. Bardiaux, B. Habenstein, S. J. Saupe and A. Loquet, *Front. Mol. Neurosci.*, 2021, **14**, 670513.
- 154 J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli and D. Hassabis, *Nature*, 2021, **596**, 583–589.
- 155 N. Ferreira, H. Gram, Z. A. Sorrentino, E. Gregersen, S. I. Schmidt, L. Reimer, C. Betzer, C. Perez-Gozalbo, M. Beltoja, M. Nagaraj, J. Wang, J. S. Nowak, M. D. Dong, K. Willen, E. Cholak, K. Bjerregaard-Andersen, N. Mendez, P. Rabadia, M. Shah Nawaz, C. Soto, D. E. Otzen, U. Akbey, M. Meyer, B. I. Giasson, M. Romero-Ramos and P. H. Jensen, *Acta Neuropathol.*, 2021, **142**, 87–115.
- 156 T. Schubeis, M. Nagaraj and C. Ritter, in *Split Inteins: Methods and Protocols*, ed. H. D. Mootz, 2017, vol. 1495, pp. 147–160.
- 157 M. Nagaraj, M. Ahmed, J. Lyngso, B. S. Vad, A. Boggild, A. Fillipsen, J. S. Pedersen, D. E. Otzen and U. Akbey, *J. Mol. Biol.*, 2020, **432**, 2232–2252.
- 158 S. L. Rouse, S. J. Matthews and M. S. Dueholm, *J. Mol. Biol.*, 2018, **430**, 3685–3695.
- 159 H. O. Rasmussen, D. E. Otzen and J. S. Pedersen, *Biophys. J.*, 2021, **120**, 2262–2275.
- 160 N. El Mammeri, J. Hierrezuelo, J. Tolchard, J. Camara-Almiron, J. Caro-Astorga, A. Alvarez-Mena, A. Dutour, M. Berbon, J. Shenoy, E. Morvan, A. Grelard, B. Kauffmann, S. Lecomte, A. de Vicente, B. Habenstein, D. Romero and A. Loquet, *FASEB J.*, 2019, **33**, 12146–12163.
- 161 M. Ghayeb, S. Hayet, N. Lester-Zer, Y. Levi-Kalishman and L. Chai, *Microorganisms*, 2021, **9**, 529.
- 162 N. Lorenzen, S. B. Nielsen, A. K. Buell, J. D. Kaspersen, P. Arosio, B. S. Vad, W. Paslawski, G. Christiansen, Z. Valnickova-Hansen, M. Andreassen, J. J. Enghild, J. S. Pedersen, C. M. Dobson, T. P. Knowles and D. E. Otzen, *J. Am. Chem. Soc.*, 2014, **136**, 3859–3868.
- 163 Z. Najarzadeh, H. Mohammad-Beigi, J. Nedergaard Pedersen, G. Christiansen, T. V. Sonderby, S. A. Shojaosadati, D. Morshedi, K. Stromgaard, G. Meisl, D. Sutherland, J. Skov Pedersen and D. E. Otzen, *Biomolecules*, 2019, **9**, 659.
- 164 Z. Najarzadeh, J. N. Pedersen, G. Christiansen, S. A. Shojaosadati, J. S. Pedersen and D. E. Otzen, *Biochim. Biophys. Acta, Proteins Proteomics*, 2019, **1867**, 140263.
- 165 H. M. Swasthi and S. Mukhopadhyay, *J. Biol. Chem.*, 2017, **292**, 19861–19872.
- 166 R. Malishev, R. Abbasi, R. Jelinek and L. Chai, *Biochemistry*, 2018, **57**, 5230–5238.
- 167 G. Y. Cheung, K. Rigby, R. Wang, S. Y. Queck, K. R. Braughton, A. R. Whitney, M. Teintze, F. R. DeLeo and M. Otto, *PLoS Pathog.*, 2010, **6**, e1001133.
- 168 J. P. Rasigade, S. Trouillet-Assant, T. Ferry, B. A. Diep, A. Sapin, Y. Lhoste, J. Ranfaing, C. Badiou, Y. Benito, M. Bes, F. Couzon, S. Tigaud, G. Lina, J. Etienne, F. Vandenesch and F. Laurent, *PLoS One*, 2013, **8**, e63176.
- 169 M. Laabei, W. D. Jamieson, Y. Yang, J. van den Elsen and A. T. Jenkins, *Biochim. Biophys. Acta*, 2014, **1838**, 3153–3161.
- 170 B. G. Surewaard, R. Nijland, A. N. Spaan, J. A. Kruijtz, C. J. de Haas and J. A. van Strijp, *PLoS Pathog.*, 2012, **8**, e1002606.
- 171 C. Galvagnion, A. K. Buell, G. Meisl, T. C. Michaels, M. Vendruscolo, T. P. Knowles and C. M. Dobson, *Nat. Chem. Biol.*, 2015, **11**, 229–234.
- 172 D. Campoccia, L. Montanaro and C. R. Arciola, *Int. J. Mol. Sci.*, 2021, **22**, 9100.
- 173 J. T. Blakeman, A. L. Morales-García, J. Mukherjee, K. Gori, A. S. Hayward, N. J. Lant and M. Geoghegan, *Langmuir*, 2019, **35**, 6468–6475.
- 174 J. L. Lister and A. R. Horswill, *Front. Cell. Infect. Microbiol.*, 2014, **4**, 178.
- 175 J. A. Jurcisek and L. O. Bakaletz, *J. Bacteriol.*, 2007, **189**, 3868–3875.
- 176 E. E. Mann, K. C. Rice, B. R. Boles, J. L. Endres, D. Ranjit, L. Chandramohan, L. H. Tsang, M. S. Smeltzer, A. R. Horswill and K. W. Bayles, *PLoS One*, 2009, **4**, e5822.
- 177 M. R. Kiedrowski, J. S. Kavanaugh, C. L. Malone, J. M. Mootz, J. M. Voyich, M. S. Smeltzer, K. W. Bayles and A. R. Horswill, *PLoS One*, 2011, **6**, e26714.
- 178 P. M. Gallo, G. J. Rapsinski, R. P. Wilson, G. O. Oppong, U. Sriram, M. Goulian, B. Buttarro, R. Caricchio, S. Gallucci and Ç. Tükel, *Immunity*, 2015, **42**, 1171–1184.





- 179 K. Schwartz, M. Ganesan, D. E. Payne, M. J. Solomon and B. R. Boles, *Mol. Microbiol.*, 2016, **99**, 123–134.
- 180 R. M. Shanks, N. P. Donegan, M. L. Graber, S. E. Buckingham, M. E. Zegans, A. L. Cheung and G. A. O'Toole, *Infect. Immun.*, 2005, **73**, 4596–4606.
- 181 C. B. Ibberson, C. P. Parlet, J. Kwiecinski, H. A. Crosby, D. K. Meyerholz and A. R. Horswill, *Infect. Immun.*, 2016, **84**, 1917–1929.
- 182 S. Mishra and A. R. Horswill, *mSphere*, 2017, **2**, e00135-17.
- 183 Z. Najarzadeh, M. Zaman, V. Sereikaite, K. Stromgaard, M. Andreassen and D. E. Otzen, *J. Biol. Chem.*, 2021, **297**, 100953.
- 184 M. Hammar, Z. Bian and S. Normark, *Proc. Natl. Acad. Sci. U. S. A.*, 1996, **93**, 6562–6566.
- 185 M. Hammar, A. Arnqvist, Z. Bian, A. Olsen and S. Normark, *Mol. Microbiol.*, 1995, **18**, 661–670.
- 186 Z. Bian and S. Normark, *EMBO J.*, 1997, **16**, 5827–5836.
- 187 Q. Shu, S. L. Crick, J. S. Pinkner, B. Ford, S. J. Hultgren and C. Frieden, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 6502–6507.
- 188 N. D. Hammer, B. A. McGuffie, Y. Zhou, M. P. Badtke, A. A. Reinke, K. Brannstrom, J. E. Gestwicki, A. Olofsson, F. Almqvist and M. R. Chapman, *J. Mol. Biol.*, 2012, **422**, 376–389.
- 189 D. Romero, H. Vlamakis, R. Losick and R. Kolter, *Mol. Microbiol.*, 2011, **80**, 1155–1168.
- 190 D. Romero, H. Vlamakis, R. Losick and R. Kolter, *J. Bacteriol.*, 2014, **196**, 1505–1513.
- 191 I. Kolodkin-Gal, D. Romero, S. Cao, J. Clardy, R. Kolter and R. Losick, *Science*, 2010, **328**, 627–629.
- 192 M. Zaman and M. Andreassen, *eLife*, 2020, **9**, e59776.
- 193 H. Q. Do, A. Hewetson, C. G. Borcik, M. C. Hastert, S. Whelly, B. J. Wylie, R. B. Sutton and G. A. Cornwall, *J. Biol. Chem.*, 2021, **296**, 100250.
- 194 H. Braak, K. Del Tredici, U. Rub, R. A. de Vos, E. N. Jansen Steur and E. Braak, *Neurobiol. Aging*, 2003, **24**, 197–211.
- 195 K. C. Luk, V. Kehm, J. Carroll, B. Zhang, P. O'Brien, J. Q. Trojanowski and V. M. Lee, *Science*, 2012, **338**, 949–953.
- 196 K. L. Paumier, K. C. Luk, F. P. Manfredsson, N. M. Kanaan, J. W. Lipton, T. J. Collier, K. Steece-Collier, C. J. Kemp, S. Celano, E. Schulz, I. M. Sandoval, S. Fleming, E. Dirr, N. K. Polinski, J. Q. Trojanowski, V. M. Lee and C. E. Sortwell, *Neurobiol. Dis.*, 2015, **82**, 185–199.
- 197 S. Holmqvist, O. Chutna, L. Bousset, P. Aldrin-Kirk, W. Li, T. Bjorklund, Z. Y. Wang, L. Roybon, R. Melki and J. Y. Li, *Acta Neuropathol.*, 2014, **128**, 805–820.
- 198 N. Uemura, H. Yagi, M. T. Uemura, Y. Hatanaka, H. Yamakado and R. Takahashi, *Mol. Neurodegener.*, 2018, **13**, 21.
- 199 N. Van Den Berge, N. Ferreira, H. Gram, T. W. Mikkelsen, A. K. O. Alstrup, N. Casadei, P. Tsung-Pin, O. Riess, J. R. Nyengaard, G. Tamguney, P. H. Jensen and P. Borghammer, *Acta Neuropathol.*, 2019, **138**, 535–550.
- 200 T. R. Sampson, C. Challis, N. Jain, A. Moiseyenko, M. S. Ladinsky, G. G. Shastri, T. Thron, B. D. Needham, I. Horvath, J. W. Debelius, S. Janssen, R. Knight, P. Wittung-Stafshede, V. Gradinaru, M. Chapman and S. K. Mazmanian, *eLife*, 2020, **9**, e53111.
- 201 E. Chorell, E. Andersson, M. L. Evans, N. Jain, A. Gotheson, J. Aden, M. R. Chapman, F. Almqvist and P. Wittung-Stafshede, *PLoS One*, 2015, **10**, e0140194.
- 202 S. Perov, O. Lidor, N. Salinas, N. Golan, E. Tayeb-Fligelman, M. Deshmukh, D. Willbold and M. Landau, *PLoS Pathog.*, 2019, **15**, e1007978.
- 203 K. Hartman, J. R. Brender, K. Monde, A. Ono, M. L. Evans, N. Popovych, M. R. Chapman and A. Ramamoorthy, *PeerJ*, 2013, **1**, e5.
- 204 I. Javed, Z. Zhang, J. Adamcik, N. Andrikopoulos, Y. Li, D. E. Otzen, S. Lin, R. Mezzenga, T. P. Davis, F. Ding and P. C. Ke, *Adv. Sci.*, 2020, **7**, 2001299.
- 205 L. F. B. Christensen, K. F. Jensen, J. Nielsen, B. S. Vad, G. Christiansen and D. E. Otzen, *ACS Omega*, 2019, **4**, 4029–4039.
- 206 M. E. Goya, F. Xue, C. Sampedro-Torres-Quevedo, S. Arnaouteli, L. Riquelme-Dominguez, A. Romanowski, J. Brydon, K. L. Ball, N. R. Stanley-Wall and M. Doitsidou, *Cell Rep.*, 2020, **30**, 367–380 e367.
- 207 D. Sade Yazdi, D. Laor Bar-Yosef, H. Adsi, T. Kreiser, S. Sigal, S. Bera, D. Zaguri, S. Shaham-Niv, D. S. Oluwatoba, D. Levy, M. Gartner, T. D. Do, D. Frenkel and E. Gazit, *Proc. Natl. Acad. Sci. U. S. A.*, 2021, **118**, e2017575118.
- 208 O. Tavassoly, D. Sade, S. Bera, S. Shaham-Niv, D. J. Voadlo and E. Gazit, *J. Mol. Biol.*, 2018, **430**, 3847–3862.

