

RESEARCH ARTICLE

[View Article Online](#)[View Journal](#) | [View Issue](#)

Check for updates

Cite this: *Org. Chem. Front.*, 2024, **11**, 690

Received 23rd October 2023,
Accepted 4th December 2023

DOI: 10.1039/d3qo01749d

rsc.li/frontiers-organic

Asymmetric copper-catalyzed alkynylallylic dimethylamination†

Si-Yuan Luo,^{a,b,c} Guo-Qiang Lin^{a,b} and Zhi-Tao He *^{c,d}

A feasible protocol for Cu-catalyzed asymmetric alkynylallylic dimethylamination is developed with the discovery of tetramethyldiaminomethane as a new, stable and convenient surrogate of dimethylamine. A series of enantioenriched 1,4-enynes are constructed in reasonable yields, high regioselectivities and moderate to good enantioselectivities. Mechanistic experiments show that the tertiary amine works as a nucleophile directly followed by the release of the expected dialkylamine unit, different from the conventional primary and secondary amines with the cleavage of a proton as the nucleophile. DFT calculations elucidate the origin of regio- and enantioselectivity for the present transformation.

Introduction

Transition metal-catalyzed η^3 -substitution reactions have been widely used for the construction of different stereogenic centers.^{1–6} In particular, since the pioneering work of van Maarseveen⁷ and Nishibayashi⁸ groups, Cu-catalyzed propargylic substitution has emerged as a reliable strategy to introduce C–N,^{9–16} C–O,^{17–20} C–C^{21–29} and C–S^{30,31} bonds vicinal to an alkyne unit stereoselectively. In this case, a propargylic leaving group is required to be preprepared for the following generation of a critical Cu-allenylidene intermediate. However, when a leaving group is located at a remote position to the alkyne unit, such a motif is generally not considered a potential substrate for propargylic substitution. Recently, Fang,³² Xu³³ and our group^{34,35} independently demonstrated the concept of Cu-catalyzed remote propargylic substitution reactions. In these studies, an olefin^{32–34} or aryl unit³⁵ was used to link the alkyne unit and the γ -leaving group, and the reactive alkene-tethered or dearomatization-induced Cu-allenylidene intermediate was smoothly formed to induce the expected substitution. In particular, our group³⁴ and Xu³³ sequentially described the first asymmetric alkynylallylic substitution processes, in which both S_N2' and *in situ* substitution models were shown to be feasible

to form chiral 1,4-enyne products and spirocycles, respectively. However, the substrate scope and reaction models are heavily limited to the use of common nucleophiles. Thus, the discovery of new applications and the establishment of efficient catalytic systems suitable for the less developed remote propargylic substitution reactions are highly desired.

The chiral dimethylamino unit exists widely in bioactive molecules, such as rivastigmine,³⁶ which is used to treat Alzheimer's disease, and the tetracycline family including tetracycline, doxycycline and tigecycline, a series of broad-spectrum antibiotics³⁷ (Scheme 1a). Based on our previous work,³⁴ we envisioned that with dimethylamine as the nucleophile, Cu-catalyzed asymmetric alkynylallylic substitution might provide a novel and valuable route to construct chiral 1,4-enynes bearing a dimethylamino stereocenter. The vicinal olefin and alkyne units could further be transformed into a series of functional groups, thus largely broadening the application potential of this methodology. However, dimethylamine is gaseous at room temperature and usually stored in special solvents, which limits the convenient use of this reagent and also influences the efficiency of the corresponding transformation (Scheme 1b). Therefore, the identification of different and efficient surrogates of the dimethylamine nucleophile is critical to realize the expected alkynylallylic dimethylation reaction. Here, we discovered that tetramethyldiaminomethane³⁸ worked as a novel and reliable precursor of dimethylamine and enabled enantioselective Cu-catalyzed alkynylallylic dimethylaminations to construct valuable chiral 1,4-enynes³⁹ (Scheme 1b). A variety of 1,4-enynes bearing a dimethylamino stereocenter were constructed in good yields and moderate to good enantioselectivities. Mechanistic experiments and DFT calculations elucidated the origin of regio- and enantioselectivity and revealed that tetramethyldiaminomethane first participated in the reaction as a neutral

^aState Key Laboratory of Chemical Biology, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Shanghai, 200032, China.



E-mail: hezt@sioc.ac.cn

^b*School of Physical Science and Technology, ShanghaiTech University, Shanghai, 201210, China*

^cState Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Shanghai, 200032, China

^dSchool of Chemistry and Materials Science, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China



† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3qo01749d>



Scheme 2 The scope of electrophiles. The reactions were carried out on a 0.1 mmol scale. Isolated yields. The rr values were determined by ^1H NMR. The er values were determined by HPLC analysis.

methylamination products in 64–93% yields with generally >20:1 rr and >90:10 er (3b–3j and 3m). Notably, substrates that are prone to cross-coupling reactions also yielded amination products smoothly (3f and 3g). Multi-substituted aryl enyne was also a suitable electrophile for the transformation (3k). In addition, 1,3-enyne bearing an *ortho*-methoxy unit afforded 3l in 86% yield, >20:1 rr and 90:10 er. However, alkyl-derived enyne as the electrophile failed to undergo the alkynylallylic dimethylamination reaction (3n).

After demonstrating the feasibility of the remote propargylic dimethylamination with this novel dimethylamine source, we further explored the application potential of this protocol with other amine nucleophiles and the results are summarized in Scheme 3. A set of secondary amine sources as the surrogates




Scheme 3 The scope of amine precursors. The reactions were carried out on a 0.1 mmol scale. Isolated yields. The rr values were determined by ^1H NMR. The er values were determined by HPLC analysis.


of dimethylamine, pyrrolidine and morpholine were suitable for the alkynylallylic substitution, furnishing the corresponding products 4a–4c in high yields, >20:1 regioselectivity and up to 94:6 er. This strategy was also applicable to the introduction of different diarylmethyl amines, and the corresponding products 4d and 4e were obtained in 95:5 and 96:4 er, respectively. The absolute configuration of 4d was determined to be (*R*)-4d based on the same reported compound.^{34a} However, the introduction of the diarylamine moiety *via* this route did not work (4f).

To figure out the potential reaction mechanism, a set of mechanistic studies were conducted (Scheme 4). As it was unclear whether the nucleophile 2a underwent the substitution *via* the *in situ* release of dimethylamine or serving as the tertiary amine nucleophile first and then releasing the dimethylamine unit, several control experiments were carried out with 2e as the substrate (Scheme 4a). When the reaction was conducted under standard conditions without a Cu catalyst or ligand or Cu/ligand, no product 4d was formed in these cases. In particular, dibenzylamine 5 was not detected. When electrophile 1a was not used under the standard conditions, no dibenzylamine 5 was observed similarly. These facts indicated that the *in situ* generation of 5 from 2e might be unfavourable under the present reaction system, and tertiary amine 2e might directly work as the nucleophile.³⁸


Next, a nonlinear relationship experiment showed a linear curve between the enantiopurity of chiral ligand L6 and that of the product 3a (Scheme 4b). This result suggested that mono-copper/L6 as the catalyst might be involved in the enantio-determining step. Based on this fact, DFT calculations were conducted to uncover the origin of regio- and enantio-selectivity for the alkynylallylic amination reaction. Both the condensed dual descriptor (CDD) and electrophilicity index (EI) were determined for the optimized structure of **int-1**




a. Control experiments




b. Nonlinear relationship experiments



c. Origin of regioselectivity




d. DFT calculations for the origin of enantioselectivity



Scheme 4 Preliminary mechanistic studies. BP86/def2tzvp/SMD(methanol)//B3LYP-D3(BJ)/6-31g*/SDD was used for DFT calculations.

(Scheme 4c).^{40,41} The relatively larger CDD and EI data for the α -carbon center than those for the γ -carbon center showed that the α -carbon center is more electrophilic than the γ -carbon center. This is consistent with the observed preferred nucleophilic attack at the α -carbon position. Finally, calculations were conducted to uncover the energy profile of enantioselective substitution processes (Scheme 4d). It was obvious that the *Re*-face nucleophilic attack occurred via **TS1** with only 17.9 kcal mol⁻¹ while the *Si*-face nucleophilic attack via **TS2** was 20.5 kcal mol⁻¹. Therefore, the (*R*)-configuration for the formed stereocenter was observed in the present transformation.

Based on the above experimental and computational results, the proposed catalytic cycle is described in Scheme 5. The enyne substrate **1a** first reacted with the copper catalyst in the presence of a base to give alkynyl copper intermediate **int-4**, which sequentially generated the Cu-allenylidene intermediate **int-1** and tautomers **int-5** and **int-6** via the cleavage of the leaving group. Then an outer-sphere nucleophilic attack by tertiary amine **2a** occurred to provide ammonium salt **int-7**, which then released product **3a** and regenerated the Cu cata-



Scheme 5 Proposed mechanism.

lyst. The released dimethylaminium might be further captured by potential nucleophiles in the reaction systems, such as methoxide.^{38,42}



Conclusions

In summary, we have established a feasible protocol for an asymmetric alkynylallylic dimethylamination reaction with the use of tetramethylmethanediamine as a new dimethylamine source. The novel protocol features reasonable functional group compatibility, high regioselectivity and moderate to good enantioselectivity. Different from the commonly used primary and secondary amines as nucleophiles with the cleavage of a proton, mechanistic experiments support the involvement of the tertiary amine as a neutral nucleophile directly followed by the release of dialkylamine. DFT calculations further elucidate the origin of regio- and enantioselectivity for the alkynylallylic substitution reaction.

Author contributions

Z.-T. H. conceived the project. S.-Y. L. conducted the experiments and calculations. Z.-T. H. and G.-Q. L. supervised the project. Z.-T. H. and S.-Y. L. co-wrote the manuscript with feedback from all authors. Yuan-Xiang Yang is acknowledged for checking some results.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We acknowledge the National Natural Science Foundation of China (22371292 and 22071262), the Natural Science Foundation of Ningbo (2023J036), the Shanghai Municipal Committee of Science and Technology (22ZR1475200), and the Shanghai Institute of Organic Chemistry for financial support.

References

- 1 L. Süssé and B. M. Stoltz, Enantioselective formation of quaternary centers by allylic alkylation with first-row transition-metal catalysts, *Chem. Rev.*, 2021, **121**, 4084–4099.
- 2 G. Li, X. Huo, X. Jiang and W. Zhang, Asymmetric synthesis of allylic compounds via hydrofunctionalisation and difunctionalisation of dienes, allenes, and alkynes, *Chem. Soc. Rev.*, 2020, **49**, 2060–2118.
- 3 R. J. Detz, H. Hiemstra and J. H. Van Maarseveen, Catalyzed propargylic substitution, *Eur. J. Org. Chem.*, 2009, 6263–6376.
- 4 Y. Miyake, S. Uemura and Y. Nishibayashi, Catalytic propargylic substitution reactions, *ChemCatChem*, 2009, **1**, 342–356.
- 5 D.-Y. Zhang and X.-P. Hu, Recent advances in copper-catalyzed propargylic substitution, *Tetrahedron Lett.*, 2015, **56**, 283–295.
- 6 K. Sakata and Y. Nishibayashi, Mechanism and reactivity of catalytic propargylic substitution reactions via metal-allynylidene intermediates: a theoretical perspective, *Catal. Sci. Technol.*, 2018, **8**, 12–25.
- 7 R. J. Detz, M. M. Delville, H. Hiemstra and J. H. Van Maarseveen, Enantioselective copper-catalyzed propargylic amination, *Angew. Chem., Int. Ed.*, 2008, **47**, 3777–3780.
- 8 G. Hattori, H. Matsuzawa, Y. Miyake and Y. Nishibayashi, Copper-catalyzed asymmetric propargylic substitution reactions of propargylic acetates with amines, *Angew. Chem., Int. Ed.*, 2008, **47**, 3781–3783.
- 9 C. Zhang, Y.-H. Wang, X.-H. Hu, Z. Zheng, J. Xu and X.-P. Hu, Chiral tridentate *P,N,N* ligands for highly enantioselective copper-catalyzed propargylic amination with both primary and secondary amines as nucleophiles, *Adv. Synth. Catal.*, 2012, **354**, 2854–2858.
- 10 G. Hattori, K. Sakata, H. Matsuzawa, Y. Tanabe, Y. Miyake and Y. Nishibayashi, Copper-catalyzed enantioselective propargylic amination of propargylic esters with amines: copper-allynylidene complexes as key intermediates, *J. Am. Chem. Soc.*, 2010, **132**, 10592–10608.
- 11 R. J. Detz, Z. Abiri, R. I. Griel, H. Hiemstra and J. H. Van Maarseveen, Enantioselective copper-catalysed propargylic substitution: synthetic scope study and application in formal total syntheses of (+)-anisomycin and (–)-cytoxazone, *Chem. – Eur. J.*, 2011, **17**, 5921–5930.
- 12 T. Mino, H. Taguchi, M. Hashimoto and M. Sakamoto, Copper-catalyzed asymmetric propargylic amination of propargylic acetates with amines using BICMAP, *Tetrahedron: Asymmetry*, 2013, **24**, 1520–1523.
- 13 M. Shibata, K. Nakajima and Y. Nishibayashi, Enantioselective intramolecular propargylic amination using chiral copper-pybox complexes as catalysts, *Chem. Commun.*, 2014, **50**, 7874–7877.
- 14 D.-Y. Zhang, L. Shao, J. Xu and X.-P. Hu, Copper-catalyzed asymmetric formal [3 + 2] cycloaddition of propargylic acetates with hydrazines: enantioselective synthesis of optically active 2-pyrazolines, *ACS Catal.*, 2015, **5**, 5026–5030.
- 15 Z.-T. Liu, Y.-H. Wang, F.-L. Zhu and X.-P. Hu, Enantioselective copper-catalyzed formal [4 + 2] cycloaddition of o-aminophenol derivatives with propargylic esters for synthesis of optically active 3,4-dihydro-2*H*-1,4-benzoxazines, *Org. Lett.*, 2016, **18**, 1190–1193.
- 16 W. Guo, L. Zuo, M. Cui, B. Yan and S. Ni, Propargylic amination enabled the access to enantioenriched acyclic α -quaternary α -amino ketones, *J. Am. Chem. Soc.*, 2021, **143**, 7629–7634.
- 17 K. Nakajima, M. Shibata and Y. Nishibayashi, Copper-catalyzed enantioselective propargylic etherification of propargylic esters with alcohols, *J. Am. Chem. Soc.*, 2015, **137**, 2472–2475.
- 18 R.-Z. Li, H. Tang, K.-R. Yang, L.-Q. Wan, X. Zhang, J. Liu, Z. Fu and D.-W. Niu, Enantioselective propargylation of polyols and desymmetrization of meso 1,2-diols by copper/boronic acid dual catalysis, *Angew. Chem., Int. Ed.*, 2017, **56**, 7213–7217.



- 19 R.-Z. Li, D.-Q. Liu and D.-W. Niu, Asymmetric O-propargylation of secondary aliphatic alcohols, *Nat. Catal.*, 2020, **3**, 672–680.
- 20 Y. Hou, Z. Zhang, X. Sun, Z. Yang, Y.-X. Luan and P. Tang, Copper-catalyzed enantioselective trifluoromethoxylation of propargyl sulfonates, *Angew. Chem., Int. Ed.*, 2023, **62**, e202218919.
- 21 P. Fang and X.-L. Hou, Asymmetric copper-catalyzed propargylic substitution reaction of propargylic acetates with enamines, *Org. Lett.*, 2009, **11**, 4612–4615.
- 22 F.-L. Zhu, Y. Zou, D.-Y. Zhang, Y.-H. Wang, X.-H. Hu, S. Chen, J. Xu and X.-P. Hu, Enantioselective copper-catalyzed decarboxylative propargylic alkylation of propargyl beta-ketoesters with a chiral ketimine *P,N,N*-ligand, *Angew. Chem., Int. Ed.*, 2014, **53**, 1410–1414.
- 23 W. Shao, H. Li, C. Liu, C. J. Liu and S.-L. You, Copper-catalyzed intermolecular asymmetric propargylic dearomatization of indoles, *Angew. Chem., Int. Ed.*, 2015, **54**, 7684–7687.
- 24 K. Tsuchida, Y. Senda, K. Nakajima and Y. Nishibayashi, Construction of chiral tri- and tetra-arylmethanes bearing quaternary carbon centers: copper-catalyzed enantioselective propargylation of indoles with propargylic esters, *Angew. Chem., Int. Ed.*, 2016, **55**, 9728–9732.
- 25 Q. Wang, T.-R. Li, L.-Q. Lu, M.-M. Li, K. Zhang and W.-J. Xiao, Catalytic asymmetric [4 + 1] annulation of sulfur ylides with copper-allenylidene intermediates, *J. Am. Chem. Soc.*, 2016, **138**, 8360–8363.
- 26 K. Zhang, L.-Q. Lu, S. Yao, J.-R. Chen, D.-Q. Shi and W.-J. Xiao, Enantioconvergent copper catalysis: in situ generation of the chiral phosphorus ylide and its Wittig reactions, *J. Am. Chem. Soc.*, 2017, **139**, 12847–12854.
- 27 J. Song, Z.-J. Zhang and L.-Z. Gong, Asymmetric [4 + 2] annulation of C1 ammonium enolates with copper-allenylidenes, *Angew. Chem., Int. Ed.*, 2017, **56**, 5212–5216.
- 28 H.-D. Qian, Z.-H. Li, S. Deng, C. Yao, H.-M. Xiang, G. Xu, Z.-Q. Geng, Z. Wang, L. Chen, C. Liu, C. Zhu, X. Qi and H. Xu, Catalytic asymmetric vinylogous and bisvinylogous propargylic substitution, *J. Am. Chem. Soc.*, 2022, **144**, 15779–15785.
- 29 F. Gong, X. Meng, S. Lan, J. Liu, S. Yang and X. Fang, Asymmetric semipinacol rearrangement enabled by copper-catalyzed propargylic alkylation, *ACS Catal.*, 2022, **12**, 12036–12044.
- 30 J. E. Gomez, A. Cristofol and A. W. Kleij, Copper-catalyzed enantioselective construction of tertiary propargylic sulfones, *Angew. Chem., Int. Ed.*, 2019, **58**, 3903–3907.
- 31 X. Gao, Y.-L. Xiao, S. Zhang, J. Wu and X.-G. Zhang, Copper-catalyzed enantioselective trifluoromethylthiolation of secondary propargyl sulfonates, *CCS Chem.*, 2021, **3**, 1463–1471.
- 32 S. Niu, Y. Luo, C. Xu, J. Liu, S. Yang and X. Fang, Copper-catalyzed yne-allylic substitutions using stabilized nucleophiles, *ACS Catal.*, 2022, **12**, 6840–6850.
- 33 H.-H. Kong, C. Zhu, S. Deng, G. Xu, R. Zhao, C. Yao, H.-M. Xiang, C. Zhao, X. Qi and H. Xu, Remote enantioselective [4 + 1] annulation with copper-vinylvinylidene intermediates, *J. Am. Chem. Soc.*, 2022, **144**, 21347–21355.
- 34 (a) J.-S. Ma, H.-Y. Lu, Y.-W. Chen, W.-C. Zhao, Y.-Z. Sun, R.-P. Li, H.-X. Wang, G.-Q. Lin and Z.-T. He, Copper-catalysed convergent regio- and enantioselective alkynylallylic substitution, *Nat. Synth.*, 2022, **2**, 37–48; (b) J. H. van Maarseveen, Copper for alkynylallylic substitution, *Nat. Synth.*, 2023, **2**, 11–12.
- 35 Y.-Z. Sun, Z.-Y. Ren, Y.-X. Yang, Y. Liu, G.-Q. Lin and Z.-T. He, Asymmetric Substitution by Alkynyl Copper Driven Dearomatization and Rearomatization, *Angew. Chem., Int. Ed.*, 2023, **62**, e202314517.
- 36 M. Rosler, R. Anand, A. C. Sain, S. Gauthier, Y. Agid, P. D. Bianco, H. B. Stahelin, R. Hartman and M. Gharabawi, Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomized controlled trial, *Br. Med. J.*, 1999, **318**, 633–638.
- 37 (a) D. Schnappinger and W. Hillen, Tetracyclines: antibiotic action, uptake, and resistance mechanisms, *Arch. Microbiol.*, 1996, **165**, 359–369; (b) A. N. Sapadin and R. Fleischmajer, Tetracyclines: nonantibiotic properties and their clinical implications, *J. Am. Acad. Dermatol.*, 2006, **54**, 258–265; (c) G. A. Pankey, Tigecycline, *J. Antimicrob. Chemother.*, 2005, **56**, 470–480.
- 38 Y. Zhou, Y. Xie, L. Yang, P. Xie and H.-M. Huang, Copper-catalyzed aerobic oxidative amination of arylboronic acid with amination under base-free conditions, *Tetrahedron Lett.*, 2013, **54**, 2713–2716.
- 39 H.-Y. Lu and Z.-T. He, Catalytic asymmetric synthesis of 1,4- enynes, *Chin. Chem. Lett.*, 2023, **34**, 108105–108115.
- 40 T. Lu and F. Chen, Multiwfn: A Multifunctional Wavefunction Analyzer, *J. Comput. Chem.*, 2012, **33**, 580–592.
- 41 T. Lu and Q. Chen, Realization of conceptual density functional theory and information-theoretic approach in Multiwfn program, in *Conceptual Density Functional Theory*, ed. S. Liu, Wiley-VCH, 2022, pp. 631–647.
- 42 (a) B. E. Love, Facile synthesis of *N*-dialkylaminomethylsubstituted heterocycles, *J. Org. Chem.*, 2007, **72**, 630–632; (b) R. R. Khairullina, B. F. Akmanov, R. V. Kunakova and A. G. Ibragimov, Synthesis of amino sulfides in the presence of rare-earth and transition metal catalysts, *Russ. J. Org. Chem.*, 2012, **48**, 902–907.