



Cite this: *Chem. Commun.*, 2025, 61, 4192

Received 11th December 2024,
Accepted 11th February 2025

DOI: 10.1039/d4cc06483f

rsc.li/chemcomm

We report a palladium catalyzed, multicomponent synthesis of pyrido[2,1-*a*]isoindoles via carbonylative coupling of imines and 2-bromopyridines to form mesoionic dipoles, followed by cycloaddition with *in situ* generated arynes. This one-pot method allows efficient, modular assembly of polysubstituted products with independent control of each building block.

The development of modular and efficient approaches to assemble heterocyclic products has become an important thrust in synthetic chemistry. One common approach to this goal is *via* multicomponent coupling reactions.¹ Unlike more classical multistep synthetic procedures, multicomponent reactions provide an avenue to couple three or more starting materials in a single reaction. This can not only create a more streamlined synthesis, but, due to their modularity, can be readily diversified to change product structures, making them attractive as well for product design. Nevertheless, a challenge in the design of multicomponent coupling reactions is accessing the reactive building blocks needed to drive the consecutive formation of multiple bonds in a single reaction. These can themselves often require a multistep synthesis and thus detracts from the efficiency benefits of employing multicomponent reactions.

A useful approach to address this challenge is to exploit metal catalysis with energetic building blocks such as carbon monoxide. Carbon monoxide is a broadly available feedstock chemical, which has helped make metal catalyzed carbonylation reactions among the most heavily employed transformations in chemical synthesis.² In addition, an underexploited feature of carbon monoxide is its energetics. The conversion of carbon monoxide to a carboxylic acid derivative is often highly

Design of a modular, palladium-catalyzed carbonylative synthesis of pyrido[2,1-*a*]isoindoles *via* benzyne 1,3-dipolar cycloaddition†

Karina S. Wong,^a Mohammad Ghanbari^{b,*} and Bruce A. Arndtsen^{a*}

exergonic. We and others have shown that this feature can be exploited to drive the catalytic build-up of a range of reactive products,^{3,4} such as the modular assembly of 1,3-dipoles as a route into multicomponent heterocycle synthesis.⁵ An example of this chemistry is the formation of pyridine-based 1,3-dipole **1**, which, following cycloaddition with electron deficient alkynes offers a multicomponent synthesis of indolizines (Fig. 1a).⁶

In considering the reactivity of **1**, we questioned if this 1,3-dipolar cycloaddition manifold might be directed toward other products. For example, benzyne have attracted significant research attention due to the reactivity of their strained triple bonds, including their use in nucleophilic additions, pericyclic chemistry, and related transformations.⁷ This synthetic utility is facilitated in part by the low lying LUMO energy of the triple




Fig. 1 Carbonylative formation of 1,3-dipoles and benzyne cycloaddition routes to pyrido[2,1-*a*]isoindole synthesis.

^a Department of Chemistry, McGill University, 801 Sherbrooke Street West, Montreal, QC, H3A 0B8, Canada. E-mail: bruce.arndtsen@mcgill.ca

^b Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan 8731753153, Iran. E-mail: ghanbari-m@kashanu.ac.ir

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data, NMR spectra, and X-ray crystal data. CCDC 2386038. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4cc06483f>



bond in benzyne, which imparts similar reactivity to electron deficient alkynes.⁸ The latter suggests that benzyne cycloaddition might be employed in this sequence with **1** to access, in this case, fused ring pyrido[2,1- α]isoindoles. Although less extensively studied as indolizines, pyrido[2,1- α]isoindoles have found use in pharmaceutically relevant products, and their delocalized 14 π -electron aromatic system has led to their application in electronic materials and dyes (Fig. 1b).⁹ Additionally, reduced forms of pyrido[2,1- α]isoindoles are found in various alkaloids and therapeutic agents.¹⁰ Pyrido[2,1- α]isoindoles are most commonly prepared *via* cyclization reactions. While effective, these typically require the multistep build-up of the appropriately substituted precursor.¹¹ Benzyne cycloaddition to pyridinium ylides can also access these structures (Fig. 1c), although these are limited to forming products with strong electron withdrawing *C*-acyl units to access the dipole itself.¹² As an alternative, we describe here how pyrido[2,1- α]isoindoles can be generated in a modular fashion *via* palladium-catalyzed carbonylation reactions. This involves the cycloaddition of benzyne derivatives to *in situ* generated **1** and provides a method to assemble these structures in one pot, with minimal waste, and where each of the imine, bromopyridine, and benzyne units can be systematically modified to access a range of new variants of these structures (Fig. 1d).

There are a number of methods available to generate arynes.¹³ One of the most versatile is the fluoride induced 1,2-elimination from *o*-trimethylsilylphenyl triflates developed by Kobayashi.¹⁴ Considering the mild conditions of this reaction and the stability of *o*-trimethylsilylphenyl triflates, our initial studies explored if this approach could allow benzyne generation to be compatible with the formation of 1,3-dipole **1**. The palladium-catalyzed carbonylative coupling of the imine *p*-tolyl(H)C=Nbn, 2-bromopyridine, and *o*-trimethylsilylphenyl triflate in the presence of CsF as an activating agent in acetonitrile does lead to the formation of 6-(*p*-tolyl)pyrido[2,1- α]isoindole **2a**, but does so in very low yield (9%, Table 1, entry 1). Examination of the reaction mixture by ¹H NMR analysis shows significant amounts of unreacted imine and 2-bromopyridine. Increasing the amount of benzyne precursor completely inhibits the reaction (entry 2). Similar results were observed with other palladium catalysts (entries 3–7, see Table S1 for full development, ESI†).

Our previous results have shown that 1,3-dipole **1a** is generated in high yield with these substrates,⁶ suggesting the presence of benzyne inhibiting catalysis. As such, a straightforward solution is to add *o*-trimethylsilylphenyl triflate and CsF to the catalytically generated 1,3-dipole, which results in cycloaddition to afford **2a** in 53% yield (entry 8). We observe the formation of side products in this reaction that may arise from the cycloaddition of benzyne to the product **2a**.¹⁵ This can be minimized by employing benzyne as the limiting reagent and leads to the formation of **2a** in slightly enhanced yield of 65% (entry 9).¹⁶ Under these conditions in benzene solvent, benzyne generation can be carried out during the palladium catalyzed reaction in somewhat lower yield (57%, entry 10).

A useful feature of this transformation is its modularity, where systematic changes to the imine, bromopyridine, and

Table 1 Development of a palladium-catalyzed synthesis of pyrido[2,1- α]isoindoles

2.5% Pd(dba)₃·CHCl₃, 5% ligand, *i*-Pr₂NEt, Bu₄NCl, 80 °C

CSF

2a

Procedures

A: Reaction of all reagents, 24 hr

B: Benzyne precursor / CsF in CH₃CN added after catalysis, RT, 24 hr.

Entry	Procedure	Ligand	Solvent	% 2a ^d
1	A	Xantphos	CH ₃ CN	9%
2 ^a	A	Xantphos	CH ₃ CN	0%
3 ^{b,f}	A	DPEphos	C ₆ H ₆	5%
4 ^{b,f}	A	dppe	C ₆ H ₆	0%
5 ^{b,f}	A	P(<i>t</i> -Bu) ₃	C ₆ H ₆	6%
6 ^{b,f}	A	P(Ph) ₃	C ₆ H ₆	0%
7 ^{b,f}	A	JohnPhos	C ₆ H ₆	2%
8 ^c	A	Xantphos	C ₆ H ₆	53%
9 ^b	A	Xantphos	C ₆ H ₆	65% (60%) ^e
10 ^{b,f}	A	Xantphos	C ₆ H ₆	57%

Xantphos


DPEphos

JohnPhos

Procedure A: *p*-tolyl(H)C=Nbn (34 mg, 0.16 mmol); 2-bromopyridine (38 mg, 0.24 mmol); *o*-trimethylsilylphenyl triflate (48 mg, 0.16 mmol); CsF (109 mg, 0.72 mmol); *i*-Pr₂NEt (25 mg, 0.19 mmol); Pd(dba)₃·CHCl₃ (4.1 mg, 0.004 mmol); ligand (0.008 mmol bidentate, 0.016 mmol monodentate); Bu₄NCl (67 mg, 0.24 mmol); 3.2 mL solvent; 5 atm CO; 80 °C. Procedure B: all reagents except benzyne precursor/CsF in C₆H₆ 80 °C, 4 h, then *o*-trimethylsilylphenyl triflate (33 mg, 0.11 mmol); CsF (75 mg, 0.50 mmol), 3 mL CH₃CN, RT, 24 h. ^a 0.35 mmol. ^b 0.11 mmol. ^c 0.16 mmol *o*-trimethylsilylphenyl triflate. ^d ¹H NMR yield. ^e Isolated yield. ^f 48 h.

benzyne reagents can be employed to readily access a range of new variants of pyrido[2,1- α]isoindoles. For example, as shown in Table 2, a number of substituted *C*-aryl imines can be incorporated into the reaction. The reaction conditions for the build-up of the 1,3-dipole are influenced by the imine, with more nucleophilic imines accelerating the reaction (**2a–d**), while those with electron-withdrawing substituents (**2f**, **2h**) requiring more pressing conditions or longer reaction times (see ESI,† for details). Regardless of the reaction rate, imines with both electron donating and electron withdrawing *C*-aryl substituents form the corresponding pyrido[2,1- α]isoindole in good overall yield (**2a–e**, **2h**). Heterocycles such as pyridine (**2f**) and thiophene (**2g**) can also be incorporated into the product from the appropriate imine. The use of polyaromatic substituted imines lead to the corresponding pyrido[2,1- α]isoindoles (**2i–j**), as does the more sterically hindered *o*-tolyl imine (**2e**). *C*-Aliphatic imines were also viable substrates and form cycloaddition products in reasonable yields. In the case of **2k–l**, the formation of the 1,3-dipole must be performed in the more polar solvent, which presumably enhances the ability of this imine to trap an *in situ* generated acid chloride (*vide infra*).



Table 2 Modulation of imines in pyrido[2,1- α]isoindole synthesis

^a Imine (0.16 mmol); 2-bromopyridine (0.24 mmol); $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (4.1 mg, 0.004 mmol); xantphos (4.6 mg, 0.008 mmol); Bu_4NCl (67 mg, 0.24 mmol); $i\text{-Pr}_2\text{NEt}$ (25 mg, 0.19 mmol); CO (5 atm); C_6H_6 (3.2 mL), 80°C , 4 h, see ESI for reaction time variations. ^b *o*-Trimethylsilylaryl triflate (0.11 mmol); CsF (75 mg, 0.50 mmol); CH_3CN (3.2 mL). Step (a) 24 h. ^c 120°C , Bu_4NCl (0.48 mmol), 5% $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$, 10% xantphos. ^c In CH_3CN .

The 2-bromopyridine unit can also be modulated. As examples, electron donating and withdrawing groups on the pyridine component are well tolerated (**2m-o**), as are substituents closer to the pyrrole unit (e.g. **2p**). The 2-bromopyridine unit can be replaced with 2-bromoquinoline to form more extended conjugated products (**2q-r**). Finally, the benzyne precursor can also be tuned. In the case of 4,5-dimethyl substituted trimethylsilyl aryl triflate, the product **2s** is formed in good overall yield,




Fig. 2 Crystal structure of **2c**. Select bond lengths [Å]: N1–C13 1.385(3), N1–C2 1.396(3), N1–C6 1.415(3), C6–C7 1.403(3), C7–C8 1.418(3), C8–C13 1.395(3), C2–C3 1.354(4). Bond angles ($^\circ$): C13–N1–C6 110.08(18), N1–C13–C8 106.47(19), C13–C8–C7 109.65(18), C6–C7–C8 106.87(19) dihedral angles ($^\circ$): C5–C6–C7–C12 0.3, C2–N1–C13–C8 176.4, N1–C13–C8–C9 177.6.

as is the fused ring product **2t**. However, unsymmetrical arynes form a mixture of cycloaddition products (**2u**). The latter contrasts with results using substituted alkynes,⁶ and presumably arises from the high aryne reactivity and the moderate electronic influence of the remote 4-methoxy substituent on selectivity.¹⁷

Pyrido[2,1- α]isoindoles are often sensitive to oxidation.¹⁸ Nevertheless, we have found that the products here are indefinitely stable when stored under an inert atmosphere. Under these conditions, crystals of **2c** can be obtained from diethyl ether/acetonitrile, and its structure confirmed by X-ray crystallography. As shown in Fig. 2, **2c** has a relatively planar π -conjugated core (dihedral angle: C5–C6–C7–C12: 0.3°). The C–C bonds in the pyridine ring are shortened relative to those in the pyrrole (e.g. C2–C3: 1.354(4) Å vs. C8–C13: 1.395(3) Å), suggesting a higher degree of aromaticity in the pyridine. The aromatic substituent on C13 is twisted out of the plane of the pyrido[2,1- α]isoindole presumably due to steric clashes between the *ortho*-arene hydrogens and the heterocyclic core (N1–C13–C14–C19: 49.8°). The crystal does not exhibit a π -stacking system between pyrido[2,1- α]isoindoles; however, there are C–H π -interactions between C12–H and a second pyrido[2,1- α]isoindole (see ESI,[†] for details).

Our preliminary postulate for the mechanism of this multi-component reaction is shown in Fig. 3. The oxidative addition of aryl bromides to Pd(0) is well-established and would form here a Pd(II)–(2-pyridyl) complex **3** that can undergo CO insertion to form the Pd-acyl complex **4**. In the presence of chloride, anionic exchange can occur to form the chloride complex **5**. As we have previously noted,⁶ chloride significantly enhances the rate of 1,3-dipole **1** formation by allowing the reductive elimination of an electrophilic acid chloride, which offers a lower barrier pathway to incorporate the weakly nucleophilic imine than direct reaction with complex **4** for subsequent cyclization to 1,3-dipole **1**. The rigid, large bite-angle Xantphos ligand presumably also facilitates this step by creating a sterically encumbered palladium to drive the disfavored reductive elimination of acid chloride.⁶ In analogy to reports of benzyne addition to mesoionic 1,3-dipoles such as Münchnones,¹⁹ the




Fig. 3 Proposed reaction mechanism of pyrido[2,1- α]isoindole formation.

in situ generation of aryne after 1,3-dipole formation leads to its rapid cycloaddition followed by cycloreversion to release isocyanate to form pyrido[2,1- α]isoindole 2. The isocyanate can be clearly seen by *in situ* ^1H NMR analysis of the reaction mixture (see Fig. S1, ESI †). The inhibitory influence of benzyne generation during catalysis could be tied to the presence of fluoride, which may slow the generation of the 1,3-dipole 1, or the formation of benzyne before the 1,3-dipole is present, leading to its decomposition.

In conclusion, we have described a multicomponent route to form substituted pyrido[2,1- α]isoindoles. This couples the palladium-catalyzed carbonylative formation of pyridine-based 1,3-dipoles with their ability to undergo cycloaddition with *in situ* generated arynes. The systematic variation of any of the three substrates can allow a range of new variants of these structures to be generated in one pot, and from either commercially available or easily generated building blocks.

We would like to thank the Natural Science and Engineering Research Council of Canada (NSERC), McGill University and the FRQNT supported by the Centre for Green Chemistry and Catalysis for funding this research.

Data availability

The data supporting this article have been included as part of the ESI † .

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) D. M. D'Souza and T. J. J. Müller, *Chem. Soc. Rev.*, 2007, **36**, 1095–1108; (b) R. V. A. Orru and E. Ruijter, *Synthesis of Heterocycles via Multicomponent Reactions. I*, Springer-Verlag, Berlin, 2010; (c) B. H. Rotstein, S. Zaretsky, V. Rai and A. K. Yudin, *Chem. Rev.*, 2014, **114**, 8323–8359.
- B. Cornils, W. A. Herrmann, M. Beller and R. Paciello, *Applied Homogeneous Catalysis with Organometallic Compounds: A Comprehensive Handbook in Four Volumes*, Wiley, 2017, vol. 4.
- (a) J. S. Quesnel and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2013, **135**, 16841–16844; (b) G. Kinney, J. Tjutrins, G. M. Torres, N. J. Liu, O. Kulkarni and B. A. Arndtsen, *Nat. Chem.*, 2018, **10**, 193–199; (c) R. G. Kinney and B. A. Arndtsen, *Angew. Chem., Int. Ed.*, 2019, **58**, 5085–5089; (d) T. M. Levesque, R. G. Kinney and B. A. Arndtsen, *Chem. Sci.*, 2020, **11**, 3104–3109; (e) Y. Liu, A. M. Kaiser and B. A. Arndtsen, *Chem. Sci.*, 2020, **11**, 8610–8616.
- (a) R. Munday, J. R. Martinelli and S. L. Buchwald, *J. Am. Chem. Soc.*, 2008, **130**, 2754–2755; (b) A. Brennfürer, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2009, **48**, 4114–4133; (c) M. N. Burhardt, R. H. Taaning and T. Skrydstrup, *Org. Lett.*, 2013, **15**, 948–951.
- (a) B. A. Arndtsen, *Chem. – Eur. J.*, 2009, **15**, 302–313; (b) S. Bontemps, J. S. Quesnel, K. Worrall and B. A. Arndtsen, *Angew. Chem.*, 2011, **123**, 9110–9113; (c) D. C. Leitch, L. V. Kayser, Z.-Y. Han, A. R. Siamaki, E. N. Keyzer, A. Gefen and B. A. Arndtsen, *Nat. Commun.*, 2015, **6**, 7411; (d) J. Tjutrins and B. A. Arndtsen, *Chem. Sci.*, 2017, **8**, 1002–1007.
- S. A. Roy, J. Zgheib, C. Zhou and B. A. Arndtsen, *Chem. Sci.*, 2021, **12**, 2251–2256.
- (a) A. Biju, *Modern Aryne Chemistry*, Wiley-VCH, Weinheim, 2021; (b) H. Ne Pellissier and M. Santelli, *Tetrahedron*, 2003, **59**, 701–730.
- (a) N. G. Rondan, L. N. Domelsmith, K. N. Houk, A. T. Bownelb and R. H. Levin, *Tetrahedron Lett.*, 1979, 3237–5240; (b) W. F. Maier, G. C. Lau and A. B. McEwen, *J. Am. Chem. Soc.*, 1985, **107**, 4724.
- (a) J. S. A. Badaro, B. Koszarna, M. H. E. Bousquet, E. T. Ouellette, D. Jacquemin and D. T. Gryko, *Org. Chem. Front.*, 2022, **9**, 1861–1874; (b) T. Mitsumori, M. Bendikov, O. Dautel, F. Wudl, T. Shioya, H. Sato and Y. Sato, *J. Am. Chem. Soc.*, 2004, **126**, 16793–16803; (c) A. Pareek, M. Mehboob, M. Cieplac, M. Majdecki, H. Szabat, K. Noworyta, P. Polcynski, M. Morawiak, P. Sharma, C. Foroutan-Nejad and P. Gawel, *J. Am. Chem. Soc.*, 2025, DOI: [10.1021/jacs.4c16189](https://doi.org/10.1021/jacs.4c16189).
- (a) B. Song, X. Guo, L. Yang, H. Yu, X. Zong, X. Liu, H. Wang, Z. Xu, Z. Lin and W. Yang, *Angew. Chem., Int. Ed.*, 2023, **62**, e202218886; (b) K. Speck and T. Magauer, *Beilstein J. Org. Chem.*, 2013, **9**, 2048–2078; (c) A. M. Hamlin, J. K. Kisunzu and R. Sarpong, *Org. Biomol. Chem.*, 2014, **12**, 1846; (d) S. W. Pelletier, *Pure Appl. Chem.*, 1997, **69**, 119–124.
- (a) A. A. Pokhonenko, Z. V. Vojtenko and V. A. Kovtunencko, *Russ. Chem. Rev.*, 2004, **73**, 833–849; (b) B. Zhao, M. Yu, H. Liu, Y. Chen, Y. Yuan and X. Xie, *Adv. Synth. Catal.*, 2014, **356**, 3295–3301; (c) W. Li, S. Zhang, X. Feng, X. Yu, Y. Yamamoto and M. Bao, *Org. Lett.*, 2021, **23**, 2521–2526; (d) D. Shi, T. Zeng, X. Lei, X. Wu, M. Li and Y. Zhang, *Synthesis*, 2022, 5434–5444.
- (a) C. Xie, Y. Zhang and P. Xu, *Synlett*, 2008, 3115–3120; (b) X. Huang and T. Zhang, *Tetrahedron Lett.*, 2009, **50**, 208–211.
- (a) F. I. M. Idris and C. R. Jones, *Org. Biomol. Chem.*, 2017, **15**, 9044–9056; (b) H. Yoshida, *Comprehensive Aryne Synthetic Chemistry*, Elsevier Science, 2022.
- (a) Y. Himeshima, T. Sonoda and H. Kobayashi, *Chem. Lett.*, 1983, 1211–1214; (b) J. Shi, L. Li and Y. Li, *Chem. Rev.*, 2021, **121**, 3892–4044.
- H. Hennige, R. Kreher and J. Uhrig, *Synthesis*, 1982, 842–844.
- All benzyne is consumed under these conditions, and small amounts of products from its addition to 2a are formed, suggesting this is the highest efficiency accessible in the reaction.
- T. Ikawa, S. Masuda, A. Takagi and S. Akai, *Chem. Sci.*, 2016, **7**, 5206.
- S. Asako, T. Kobashi and K. Takai, *J. Am. Chem. Soc.*, 2018, **140**, 15425.
- (a) A. V. Dubrovskiy, N. A. Markina and R. C. Larock, *Org. Biomol. Chem.*, 2013, **11**, 191–218; (b) J. M. Lopchuk and G. W. Gribble, *Tetrahedron Lett.*, 2015, **56**, 3208–3211.

