

Food & Function

Linking the chemistry and physics of food with health and nutrition

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. Li, R. An, H. Wang, S. Yuan and X. Shi, *Food Funct.*, 2026, DOI: 10.1039/D6FO01315E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Plant-Derived Exosome-Like Nanoparticles Ameliorate Glycolipid Metabolism Diseases:**Molecular Mechanism, Advances and Bottlenecks**Xue Li ^{1,*}, Ran An ^{1,*}, Hairong Wang ¹, Shiya Yuan ², Xu Shi ¹

¹ Department of Laboratory Medicine, Lequn Branch, The First Hospital of Jilin University, Changchun, Jilin 130031, China

² College of Traditional Chinese Medicine, Tianjin University of Traditional Chinese Medicine, Tianjin 301617, China

*** These authors contributed equally to this work**

Correspondence: Xu Shi, Department of Laboratory Medicine, The First Hospital of Jilin University, No.3302 Jilin Road, Changchun, Jilin 130023, P.R. China. Email address: shixu@jlu.edu.cn.



14 **Abstract**View Article Online
DOI: 10.1039/D6FO01315E

15 Glycolipid Metabolism Diseases, including obesity, type 2 diabetes mellitus (T2DM), and
16 non-alcoholic fatty liver disease (NAFLD), are increasingly becoming a significant global public
17 health burden. Existing treatment approaches still face challenges in terms of long-term efficacy
18 and safety, highlighting an urgent need to develop innovative intervention strategies. Compared
19 to mammal-derived exosomes, exosome-like nanoparticles derived from natural plants exhibit
20 unique application prospects owing to their abundant sources, good biocompatibility and low
21 immunogenicity. This review systematically summarizes the recent progress of natural plant-
22 derived exosome-like nanoparticles (PELNs) in ameliorating disorders of glucolipid metabolism
23 through multi-target and multi-pathway synergistic effects, including enhancing insulin
24 sensitivity, alleviating oxidative stress, inhibiting inflammatory responses, and modulating gut
25 microbiota balance. We summarize the potential of PELNs as novel therapeutic agent and drug
26 delivery carriers, and analyze the the current issues and challenges faced in clinical applications.

27

28 **Key words:** natural plants; exosome-like nanoparticles; glycolipid metabolism; type II diabetes
29 mellitus; nonalcoholic fatty liver disease

30

31



32 Introduction

33 Disorders of glycolipid metabolism, such as type 2 diabetes mellitus (T2DM) and
34 nonalcoholic fatty liver disease (NAFLD), have become major global health burdens. The
35 pathogenesis of these diseases involves the interplay of multiple factors, including insulin
36 resistance, chronic inflammation, oxidative stress, dyslipidemia, and gut microbiota dysbiosis.¹⁻
37 ³ Traditional drug therapies acting on a single site often cause side effects. Consequently, the
38 development of novel intervention strategies with multiple targets and high safety profiles is a
39 research hotspot. Exosomes, key mediators of intercellular communication, have attracted
40 significant attention due to their intrinsic nanoscale vesicle structure and ability to encapsulate
41 bioactive molecules.^{4,5}

42 In recent years, studies have demonstrated that plants also produce nanoparticles similar to
43 mammalian-derived exosomes (MDEs), termed plant-derived exosome-like nanoparticles
44 (PELNs).⁶ Compared with MDEs, PELNs have attracted widespread attention due to their natural
45 origin, potent biological activity, high stability, and efficacy in drug encapsulation and targeted
46 delivery.

47 PELNs not only contain bioactive phytochemicals but also serve as natural nanocarriers,
48 thereby protecting the encapsulated cargo from degradation and facilitating their uptake by
49 mammalian cells.⁷⁻⁹ This provides a novel approach to utilizing abundant natural plant resources
50 to address metabolic diseases, thus potentially overcoming traditional drug delivery limitations
51 and enabling more precise treatments.¹⁰ This review systematically summarizes current research
52 on natural PELNs in glycolipid metabolism diseases. Furthermore, it explores the characteristics,
53 advantages, and molecular mechanisms of PELNs, as well as their therapeutic potential and
54 challenges as therapeutic agents and drug delivery carriers.

55 1. Overview of PELNs


56 1.1 Source and Biogenesis

57 Plant cells secrete various types of extracellular vesicles (EVs) that differ in structure and
58 function. The main categories include PELNs, shedding microvesicles (MVs), and vesicles
59 derived from exocyst-positive organelle (EXPO) (Figure 1). Among these EVs, PELNs have
60 attracted the most attention. The biogenesis of PELNs is similar to that of mammalian exosomes
61 and involves the formation of early endosome (EE) and multivesicular bodies (MVB), as well as



62 the fusion of MVB with the plasma membrane.

View Article Online
DOI: 10.1039/D6FO01315E



63

64 **Figure 1.** A schematic illustration of the source and biogenesis of plant extracellular vesicles
 65 (EVs). Plant EVs are mainly derived from three pathways: **(1)** Endosomal pathway: the plasma
 66 membrane invaginates to form the early endosome (EE) containing intracellular vesicles (ILVs),
 67 and then the EE membrane buds inward to generate multivesicular bodies (MVB). Upon fusion
 68 of the MVB membrane with the plasma membrane, PELNs are released into the extracellular
 69 space; **(2)** Budding or shedding pathway: forming microvesicle (MV) by direct budding from the
 70 plasma membrane or detaching; **(3)** Exotyst-positive organelle (EXPO) pathway: generating
 71 EXPO vesicle derived from specific organelles including endoplasmic reticulum (ER) and Golgi
 72 apparatus.

73

74 PELNs can be derived from different parts of plants, including fruits, vegetables, and leaves,
 75 ^{11,12} with diameters typically ranging from 30 to 150 nanometers. PELNs extracted from flowers
 76 and leaves often demonstrate particle sizes of less than 150 nm, ¹²⁻¹⁵ whereas those derived from



77 root organs such as ginseng usually exhibit sizes exceeding 200 nm.^{16,17} As mediators of
78 intercellular communication, PELNs encapsulate a rich array of bioactive molecules, including
79 proteins, lipids, miRNAs and plant-specific active components (such as curcumin and crocin).¹⁸⁻
80 ²¹ These substances collectively confer inherent multicomponent bioactivity to PELNs. The
81 unique physicochemical properties of PELNs, such as their small particle size, negative charge,
82 and lipid bilayer structure, enable them to penetrate the intestinal mucus layer, withstand both
83 extreme pH and enzymatic degradation, and adhere to intestinal epithelial cells through
84 electrostatic interactions, thereby offering advantages such as low immunogenicity, high stability,
85 and natural targeting.²²

86 The size of shedding MVs typically ranges from 100 to 1000 nanometers. Unlike PELNs
87 (which depend on the endosomal system for secretion), shedding MVs are primarily formed
88 through direct budding or shedding from the plasma membrane. Therefore, their membrane
89 composition more directly reflects the makeup of the source cell's plasma membrane, and their
90 cargo may include cell wall-modifying enzymes.²³ EXPO vesicles are derived from organelles
91 such as chloroplasts and the endoplasmic reticulum (ER).²⁴ The unique cargo composition of
92 these vesicles may confer special functions, thus representing a potential direction for future
93 research.²⁵ For example, some EXPO vesicles secreted by plants contain an abundant amount of
94 peroxisomes, thus suggesting their potential involvement in antioxidant defense or the transport
95 of lipid metabolites.²⁶

96 1.2 Isolation and purification

97 Currently, various techniques have been applied for the isolation and purification of PELNs.
98 Ultracentrifugation (particularly differential ultracentrifugation) is the traditional "gold standard"
99 method for isolating exosomes. This method progressively increases the centrifugation speed to
100 gradually remove cell debris and large vesicles, thereby ultimately precipitating the exosome
101 fraction at a high speed (typically at a speed of approximately 100,000×g). Although this method
102 can yield exosomes exhibiting relatively high purity, it is time-consuming and may cause vesicle
103 aggregation or damage.^{27,28}

104 The size exclusion chromatography (SEC) method utilizes a chromatographic column
105 formed by porous spherical fillers, thus allowing for molecules of different sizes to elute at
106 different rates and enabling the gentle separation of exosomes. SEC can better preserve the



107 biological activity and natural morphology of exosomes, along with effectively removing
108 copurified soluble proteins and other impurities.²⁹

109 The polymer precipitation method (including the use of polyethylene glycol) uses
110 hydrophilic polymers to “capture” exosomes, thereby altering their solubility to cause
111 precipitation. This method is simple and fast to perform; however, it may introduce polymer
112 impurities and lead to exosome aggregation.³⁰

113 Tangential flow filtration (TFF) and ultrafiltration (UF) are separation techniques based on
114 particle size. UF typically employs a series of membrane filters with pore sizes of 0.1, 0.22, and
115 0.45 micrometers to initially remove cells, debris, and larger particles, followed by the use of UF
116 membranes with appropriate pore sizes to separate soluble and aggregated proteins.³¹ TFF
117 achieves the efficient separation of exosomes from impurities through fluid shear forces parallel
118 to the membrane surface. Due to its high efficiency and scalability, this method has become a
119 core technology for PELNs extraction in recent years.^{32,33}

120 To obtain high-purity exosomes, researchers often utilize a combination of multiple methods,
121 such as preliminary enrichment through differential centrifugation, followed by fine purification
122 using SEC. Additionally, separation technologies based on microfluidics and novel materials
123 (such as titanium dioxide microspheres)^{34,35} are being continuously developed, with an aim of
124 improving separation efficiency, purity, and throughput. The features of the main isolation and
125 purification techniques are summarized in Table 1.

126 1.3 Characterization methods

127 The comprehensive characterization of isolated plant exosomes is necessary to confirm their
128 identity, assess their quality, and explore their functional properties. Nanoparticle tracking
129 analysis (NTA) and dynamic light scattering (DLS) are typically used to determine the particle
130 size distribution and concentration of exosomes.^{36,37} Zeta potential analysis is used to evaluate
131 the net charge on the particle surface, thereby reflecting its colloidal stability and its tendency to
132 interact with cells.^{37,38} Electron microscopy techniques, particularly transmission electron
133 microscopy (TEM) and cryo-electron microscopy (cryo-EM), enable the direct observation of the
134 morphology, size, and membrane structure of exosomes; moreover, these techniques are
135 considered to be the gold standard for confirming their vesicular morphology.³⁹

136 To thoroughly analyze the molecular composition and functional basis of exosomes, omics



137 analysis techniques are crucial. Proteomics can be used to identify the proteins carried by
138 exosomes through mass spectrometry, thus revealing their possible cellular origins, signaling
139 pathways, and potential uses as disease biomarkers.^{40,41} Lipidomics is used to systematically
140 analyze the types and contents of lipid molecules in the exosome membrane and interior;
141 moreover, these lipids not only form the structural basis of exosomes but also participate in cell
142 signaling and metabolic regulation.⁴² By integrating multiomics data such as proteomics,
143 lipidomics, and transcriptomics data, a systematic interpretation of the bioactive components of
144 PELNs and their potential mechanisms in regulating host cell metabolism can be achieved.

145 **2. Advantages of PELNs**

146 **2.1 Differences between PELNs and MDEs**

147 PELNs share structural and functional similarities with exosomes derived from animal cells
148 but also exhibit significant differences. Both types of exosomes are nanosized vesicles
149 demonstrating a lipid bilayer structure and are approximately 100 nm in size; additionally, they
150 serve as carriers for intercellular communication by transporting bioactive molecules such as
151 proteins, lipids, and nucleic acids. Furthermore, both types of exosomes exhibit good
152 biocompatibility and can cross biological barriers.¹¹ However, key distinctions have been
153 observed regarding their sources, acquisition methods, and specific characteristics. MDEs are
154 typically isolated from cell culture supernatants or bodily fluids and involve complex production
155 processes, high costs, and limited yields. In contrast, PELNs are derived from abundant plant
156 materials; moreover, they are easy to produce on a large scale, exhibit lower costs, and are
157 renewable.^{18,43} Furthermore, PELNs generally exhibit lower immunogenicity and cytotoxicity
158 compared to mammalian exosomes.⁴⁴ Functionally, research on MDEs often focuses on their
159 roles in disease progression (such as carcinoma) and as disease biomarkers, whereas PELNs
160 demonstrate unique pharmacological activities, such as anti-inflammatory, antioxidation, and
161 direct antitumor effects, due to the specific plant active components that they carry (such as
162 curcumin and berberine derivatives, among other components).⁷

163 **2.2 Engineering modification of PELNs**


164 In addition to their inherent unique advantages, such as intrinsic biological activity, high
165 stability, and ease of absorption, PELNs can also effectively encapsulate various chemicals, cross
166 biological barriers, and serve as stable, low-immunogenicity therapeutic platforms. As vectors,



167 engineering modifications for targeting and therapeutic efficacy have shown great potential. View Article Online
DOI: 10.1039/D6FO01315E

168 2.2.1 Surface modifications

169 Previous studies have combined exosomes with advanced delivery platforms, such as
170 hydrogel sustained-release platforms. For example, curcumin-derived ⁴⁵ and *Salvia miltiorrhiza*-
171 derived nanoparticles ⁴⁶ were mixed with thermosensitive gels and then cast into nanoparticle
172 array molds to prepare microneedle patches, thus enhancing the local sustained-release and long-
173 lasting effects of plant-derived nanoparticles (Figure 2). Grapefruit-derived exosomes can be
174 loaded into dissolvable hyaluronic acid microneedles to facilitate tendon repair, ⁴⁷ and *Lycium*
175 *barbarum*-derived nanosized vesicles can be encapsulated in fibrin gel to improve targeted
176 delivery efficiency and retention time. ⁴⁸



177 **Figure 2.** Fabrication and characterization of Tur-EVs-loaded microneedle (T-MN) in rat rotator
178 cuff repair. **(A)** Schematic of T-MN fabrication via micromolding. **(B)** Light microscopy images
179 of T-MN. **(C)** SEM analysis of T-MN. **(D)** Fluorescence shows that Tur-EVs are uniformly
180 distributed in microneedle. Reproduced with permission. ⁴⁵ Copyright 2025, Elsevier.

181 **Abbreviations:** Tur-EVs, turmeric-derived extracellular vesicles.

182

183 In addition to the aforementioned physical modifications, surface biological modifications
184 of nanoscale vesicles offer substantial therapeutic potential. Su et al. ⁴⁹ modified folate (FA) onto
185 the surface of ginger-derived extracellular vesicles (GDEVs) through folate-polyethylene glycol-
186 cholesterol (FA-PEG2000-Chol) modifications, thereby constructing engineered exosomes (FA-
187 GDEVs). These FA-GDEVs can specifically target pro-inflammatory M1 macrophages which
188 highly express folate receptors (FRs) in rheumatoid arthritis (RA)-affected joints, ⁵⁰⁻⁵² alleviating
189 both joint inflammation and cartilage destruction.



190 Chemical modification is also frequently employed. Chen et al.⁵³ incorporated low
191 concentration cholesterol into the membrane of nanovesicles from *Clematis filamentosa* Dunn
192 (CDNVs), thereby effectively inhibiting M1 macrophage polarization at lung injury sites.

193 Recently, Jin et al.⁵⁴ modified *Calendula officinalis* L.-derived extracellular vesicles
194 (COEVs) with phosphatidylserine (PS), enabling specific phagocytosis by M1 macrophages at
195 fracture sites. Subsequently, a ROS-responsive hydrogel was employed to encapsulate the PS-
196 modified COEVs to facilitate the on-demand release, thereby effectively alleviating inflammation
197 and promoting fracture healing. Therefore, the future surface modification of PELNs will no
198 longer be solely physical, chemical, or biological modifications, but rather a composite strategy
199 that integrates multiple modification approaches.

200 **2.2.2 Drug delivery**

201 Drug loading strategies for PELNs primarily involve introducing exogenous therapeutic
202 molecules into preformed natural vesicles. Common methods include passive loading and active
203 loading techniques. Passive loading strategies, such as coincubation, involve incubating drugs
204 with PELNs by utilizing the hydrophobic nature of drugs (e.g., ascorbic acid and doxorubicin) to
205 facilitate diffusion into the lipid bilayer of PELNs.⁵⁵⁻⁵⁷ In contrast, active loading strategies
206 temporarily disrupt the integrity of the vesicle lipid bilayer through physical methods to increase
207 permeability. Techniques such as electroporation, sonication, or freeze-thaw cycles transiently
208 disrupt the PELNs lipid membrane to permit the loading of hydrophilic macromolecules (e.g.,
209 proteins and nucleic acids).^{58,59} However, these methods may affect the structural integrity of
210 vesicles, and the loading process requires optimization to strike a balance between loading
211 efficiency and vesicle stability. In contrast, synthesized engineered nanoparticles often exhibit
212 superior drug loading capacity and encapsulation efficiency, which are issues that need to be
213 addressed in the future regarding PELN-loading strategies.

214 **2.2.3 Reconstruction of lipid components**

215 To overcome potential interference from endogenous components in natural PELNs and
216 achieve more uniform and controllable carrier production, some studies have adopted liquid-
217 liquid extraction methods to extract total lipids, followed by the self-assembly of lipids in the
218 aqueous phase to form nanoparticles through methods such as extrusion, sonication, or high-
219 pressure homogenization. For example, using the solvent-assisted vesicle hydration (SAVH)



220 method, the lipid fractions of natural nanosized vesicles derived from grapes and tomatoes were
221 extracted and reconstituted to construct grape-tomato hybrid nanosized vesicles.⁶⁰ This method
222 significantly enhances vesicle purity and stability through the selective extraction of lipid
223 fractions, along with achieving a notable increase in yield. Compared with the original vesicles,
224 the fused hybrid vesicles not only retain antioxidant substances from grape and tomato sources
225 (such as resveratrol and lycopene) but also exhibit a greater hydroxyl radical scavenging capacity,
226 thereby demonstrating potential synergistic antioxidant effects. The advantages of this strategy
227 are attributed to the precise control of carrier particle size and uniformity, as well as the
228 simultaneous loading of drugs or nucleic acids during the assembly process. This “biomimetic”
229 nanocarrier based on PELNs membrane lipids combines the safety of natural carriers with the
230 designability of synthetic carriers, thus offering new insights for the large-scale, low-cost
231 production of drug delivery systems.

232 **3. Mechanisms through which PELNs regulate glucolipid metabolism**

233 **3.1 Activation of the IP3K/AKT signaling pathway**

234 T2DM is a chronic metabolic disease characterized by persistent hyperglycemia and insulin
235 resistance, which often trigger a series of complications, such as NAFLD, cardiovascular diseases,
236 nephrosis, retinopathy, and neuropathy.^{61,62} Traditional treatments for T2DM include insulin,
237 biguanides, thiazolidinediones, dipeptidyl peptidase-4 inhibitors (DPP-4), and newer agents, such
238 as glucagon-like peptide-1 (GLP-1) receptor agonists and sodium glucose cotransporter protein 2
239 (SGLT-2) inhibitors.^{63,64} Although these drugs can effectively reduce blood glucose levels and
240 provide auxiliary benefits such as cardiovascular protection and weight loss, they are associated
241 with several limitations, including side effects (e.g., gastrointestinal disorders and liver injury),
242 low patient compliance, and the need for lifelong medication. PELNs are expected to become a
243 safer and more effective novel therapeutic strategy because of their high biocompatibility and low
244 toxicity.

245 Singhal et al.⁶⁵ systematically revealed the significant efficacy of ginger-derived exosome-
246 like nanoparticles (GELNs) in ameliorating insulin resistance in T2DM. In a T2DM mouse model,
247 GELNs significantly reduced fasting blood glucose levels, along with improving glucose
248 tolerance and insulin sensitivity, with efficacy being comparable to the metformin group.
249 Mechanistically, this study focused on the key PI3K/AKT signaling pathway in glucose



250 metabolism. After insulin binds to the insulin receptor, it recruits PI3K, activates PI3K/AKT
 251 signaling, and subsequently induces a series of changes in glucose metabolism-related proteins,
 252 such as promoting glucose uptake (via GLUT4 translocation),⁶⁶ inhibiting hepatic glucose output
 253 (via the phosphorylation of FoxO),⁶⁷ and promoting glycogen synthesis (via the inhibition of
 254 GSK3).⁶⁸ GELNs have been observed to increase the tyrosine phosphorylation of insulin receptor
 255 substrate-1 (IRS-1), thereby activating Akt-2 (via phosphorylation at Ser474), downregulating the
 256 expression of key gluconeogenic enzymes (such as PCK-1 and G6PC), and inhibiting excessive
 257 hepatic glucose production. Moreover, GELNs can upregulate the expression of glycogen
 258 synthase-2 (GYS-2), promote hepatic glycogen storage, and downregulate lipogenesis-related
 259 factors (such as SREBP-1c and FAS), thereby reducing hepatic ectopic fat deposition.⁶⁵
 260 Moreover, researchers have reported that GELNs contain 121 types of miRNAs, and the predicted
 261 target genes of these miRNAs were observed to also be significantly enriched in PI3K/Akt-related
 262 pathways. The transfection of a single-stranded mimic of synthesized GELNs miRNA (mtr-
 263 miR399q) into insulin-resistant HepG2 cells significantly downregulated the expression of the
 264 key intracellular gluconeogenic gene PCK-1.⁶⁵

265 Under normal physiological conditions, insulin phosphorylates Foxa2 via the PI3K/AKT
 266 pathway, thus resulting in its inactivation and translocation from the nucleus to the cytoplasm.
 267 However, in insulin-resistant states, chronic hyperinsulinemia causes Foxa2 to remain persistently
 268 localized in the cytoplasm and become inactive, thereby exacerbating hepatic lipid accumulation
 269 and insulin resistance.⁶⁹ Studies have demonstrated that phosphatidic acid (PA), which is
 270 abundant in ginger-derived nanoparticles (GDNPs), can directly bind to the Foxa2 protein,
 271 thereby covering its Thr156 phosphorylation site. This binding inhibits Akt-1-mediated Foxa2
 272 phosphorylation and prevents Foxa2 inactivation and nuclear export, thereby maintaining its
 273 transcriptional activity and driving lipolysis.⁷⁰ Importantly, GDNPs treatment alters the lipid
 274 composition of intestinal epithelial cell-derived exosomes by increasing the proportion of
 275 phosphatidic acid and reducing the level of phosphatidylcholine. These modified epithelial cell-
 276 derived exosomes are transported to the liver, where they also upregulate Foxa2 expression and
 277 inhibit its phosphorylation in hepatocytes. This mechanism, which systemically improves insulin
 278 sensitivity and glucolipid metabolism via the gut-liver axis, highlights GDNPs as a potential novel
 279 strategy for treating T2DM and its complications.



280 Mung bean sprouts, which are traditionally recognized as a hypoglycemic food, are rich in
281 various bioactive components, and multiple studies have suggested that their extracts have
282 hypoglycemic effects. ⁷¹ Tang et al. ⁷² isolated and purified exosome-like nanoparticles from
283 mung bean sprouts (MELNs) and evaluated their therapeutic effects in high-fat diet-induced
284 diabetic mice. They reported that MELNs could upregulate the expression of the glucose
285 transporter GLUT4 and promote its membrane translocation by activating the PI3K/Akt signaling
286 pathway. This activation enhances cellular glucose uptake, improves insulin resistance and
287 effectively alleviates both hepatic inflammatory infiltration and steatosis.

288 The substantial therapeutic benefits of PELNs in enhancing insulin sensitivity and improving
289 liver function are mediated by their regulation of multiple downstream targets of the PI3K
290 signaling pathway. Although this multifaceted therapeutic strategy of PELNs demonstrates great
291 potential, further clinical evidence is required to elucidate its therapeutic dosage, long-term effects,
292 drug-drug interactions, and applicability to different patient populations.

293 **3.2 Regulation of intestinal barrier function and intestinal flora**

294 The gut microbiome is a highly complex system that not only plays a crucial role in
295 fundamental physiological functions (including digestion, immunity, and metabolism) but also
296 indirectly regulates overall health by influencing various bodily systems, including the nervous
297 system. Imbalances in the gut microbiome have been strongly linked to various diseases, such as
298 obesity, T2DM, and depressive disorder. ^{73,74} Previous studies have coated GELNs onto the
299 surface of hollow mesoporous silica (HMS) loaded with ammonia borane to develop a biomimetic
300 oral nanoplatform known as HMS/A@GE. This platform significantly reduced fasting blood
301 glucose levels, improved glucose tolerance, and enhanced insulin sensitivity in T2DM mice after
302 oral administration. Additionally, it alleviated hepatic steatosis and reduced serum ALT/AST and
303 TG/TC levels. The mechanism underlying these effects involves the actions of GELNs in
304 reshaping the structure of the gut microbiome and significantly increasing the abundance of
305 beneficial bacteria such as *Lactobacillus*. The increased levels of tryptophan metabolites (e.g.,
306 indole and indoleacetic acid) produced by these beneficial bacteria subsequently increased the
307 expression of intestinal barrier proteins (e.g., Occludin) and systemically suppressed
308 inflammatory responses. When antibiotics were used to clear the gut microbiome in mice, many



309 beneficial effects of HMS/A@GE (such as improved insulin resistance) were diminished.⁷⁵

310 As a traditional Chinese medicine, dried tangerine peel has been observed to exhibit various
311 biological activities, including hypoglycemic, hypolipidemic, hepatoprotective, antioxidation,
312 and anti-inflammatory effects.^{76,77} Zou et al.⁷⁸ extracted tangerine nanovesicles (TNVs) from
313 fresh citrus peel juice and reported that the oral administration of TNVs could reshape the
314 disordered gut microbiome and increase the α -diversity of the gut microbiome in diabetic mice.
315 It also increased the abundance of beneficial bacteria such as Lactobacillaceae while reducing the
316 abundances of harmful bacteria such as Lachnospiraceae and Desulfovibrionaceae, thereby
317 improving insulin resistance and glucolipid metabolism disorders in diabetic model mice. TNVs
318 can also promote the repair of the intestinal mucosal barrier, as evidenced by increased colon
319 villus height and crypt depth, the recovery of goblet cell numbers, and the restoration of the
320 expression of tight junction proteins (including Claudin-1, ZO-1, and Occludin). Furthermore,
321 TNVs have been observed to regulate hepatic lipid metabolism and improve hepatic steatosis by
322 downregulating key genes for hepatic gluconeogenesis (such as PEPCK and G6Pase) and
323 lipogenesis (such as SREBP-1c, CD36, and PPAR- γ), along with upregulating genes related to
324 fatty acid β -oxidation (such as CPT1, PPAR- α , and UCP1). Furthermore, TNVs can regulate bile
325 acid metabolism, reduce the levels of various primary/secondary bile acids, and maintain bile acid
326 homeostasis by modulating the FXR/SHP/FGF19 signaling pathway and the expression of bile
327 acid transporters (such as NTCP and BSEP).⁷⁸

328 Although numerous *in vivo* and *in vitro* evidence has confirmed the potential of PELNs to
329 improve glucolipid metabolism by regulating intestinal flora, large amounts of data are still
330 needed to further determine their bioavailability, stability, and safety.

331 **3.3 Reduction of oxidative stress**

332 Oxidative stress is a key factor in the development of numerous diseases, including (but not
333 limited to) hypertension, atherosclerosis, chronic obstructive pulmonary disease, Alzheimer's
334 disease, and T2DM.⁷⁹ Free radical-mediated oxidative stress not only directly damages cells but
335 also interacts with inflammatory factors, thus jointly promoting the development of metabolic
336 diseases such as diabetes mellitus and NAFLD, along with their complications.⁸⁰ Nuclear factor
337 erythroid 2-related factor 2 (Nrf2) is a key transcription factor in the cellular antioxidant defense



338 system. The activation of Nrf2 can induce the expression of various antioxidant enzymes and
339 restore redox homeostasis.^{81,82} Studies have demonstrated that MELNs can promote the nuclear
340 translocation of the transcription factor Nrf2 by regulating the PI3K/Akt/GSK-3 β /Nrf2 signaling
341 pathway. They also upregulate the expression of antioxidant enzymes (such as heme oxygenase-
342 1 (HO-1), superoxide dismutase (SOD), and glutathione peroxidase (GSH-Px)), thereby
343 effectively clearing excess reactive oxygen species (ROS) under diabetic conditions and
344 alleviating oxidative stress-induced damage to the liver and islets.⁷²

345 Numerous studies have confirmed that nanoparticles isolated from ginger can mitigate tissue
346 damage caused by osteoarthritis⁸³ and periodontitis⁸⁴ by modulating oxidative stress and
347 inflammatory reactions. Zhang et al.⁸⁵ reported that GELNs can induce the nuclear translocation
348 of nuclear factor Nrf2 in hepatocytes through the pathway involving Toll-like receptor 4 (TLR4)
349 and its adaptor protein TRIF (but not MyD88). This process induces the upregulation of hepatic
350 antioxidant genes such as heme oxygenase-1 (HO-1) and NAD(P)H quinone dehydrogenase 1
351 (NQO1), along with reducing ROS levels. This effect is attributed to the high content of gingerol
352 in GELNs. GELNs can also reduce lipid peroxidation products such as malondialdehyde and
353 increase the levels of antioxidant substances such as glutathione (GSH), as well as enzymes such
354 as catalase. GELNs alleviate oxidative stress and inflammatory reactions, thereby reducing liver
355 injury and pancreatic β -cell destruction.⁶⁵

356 In addition to isolating exosomes from plant tissue homogenates or juices, Ambrosone et al.
357⁸⁶ developed an additional technique to isolate small extracellular vesicles from cardoon cell
358 suspension cultures. In an *in vitro* cellular model of NAFLD, vesicles derived from cardoon cells
359 significantly upregulated Sirt-1 protein expression and increased the phosphorylation of AMPK.
360 By activating the Sirt-1/AMPK signaling pathway, these vesicles significantly reduced ROS and
361 NO levels, enhanced cell viability, and decreased lipid accumulation in hepatocytes, with effects
362 being comparable to those of the lipid-lowering drug metformin.

363 Currently, numerous studies have documented the regulation of oxidative stress by specific
364 plant bioactives. However, research on PELNs in this field remains in the early stages. As a novel
365 delivery system and therapeutic carrier, PELNs encapsulate multiple components (phenolic acids,
366 flavonoids, miRNAs, etc.), thereby facilitating a multi-target therapeutic strategy for disorders of
367 glycolipid metabolism. However, the mechanisms underlying the synergistic interactions among



368 these components remain to be elucidated.

369 **3.4 Alleviation of the inflammatory response through regulation of M1/M2 phenotypic** 370 **transformation**

371 Research has demonstrated that macrophages (which are key and highly plastic members of
372 the immune system) have additional roles besides their traditional activities in pathogen defense;
373 specifically, they are deeply involved in tissue development, homeostasis maintenance, and
374 metabolic regulation.⁸⁷ Under metabolic stress conditions such as obesity, circulating monocytes
375 are systemically recruited to metabolic organs and differentiate into proinflammatory M1
376 macrophages, thereby acting as a central factor driving chronic inflammation and metabolic
377 dysfunction.⁸⁸ This recruitment process is particularly prominent in white adipose tissue (WAT),
378 where obesity induces a shift in macrophages from the anti-inflammatory M2 phenotype to the
379 proinflammatory M1 phenotype, thus representing a key driver of insulin resistance and tissue
380 fibrosis.⁸⁹

381 Studies have demonstrated that supplementation with garlic-derived exosomes (GDEs) can
382 modulate the levels of inflammatory cytokines in the blood and epididymal WAT of rats fed a
383 high-fat diet. This mechanism involves the activity of GDEs in downregulating the expression of
384 a glycolytic enzyme (PFKFB3) via the targeting of miRNA-396e and the promotion of
385 macrophage M2 polarization, thereby inhibiting the inflammatory response in adipocytes and
386 enhancing lipid metabolism.⁹⁰ Additionally, in pathological conditions including ulcerative
387 colitis, bone fractures, and liver fibrosis, ELNs derived from *Zanthoxylum bungeanum*, *Calendula*
388 *officinalis*, *Andrographis paniculata*, *Portulaca oleracea*, *Momordica charantia*, and *Camellia*
389 *sinensis* have been demonstrated to inhibit macrophage M1 polarization by regulating the
390 PI3K/AKT,⁹¹ NF- κ B,^{54,92} and HIF-1 α /p300-CBP⁹³ signaling pathways or by modulating
391 miRNAs.^{94,95}

392 However, not all plant-derived exosome-like vesicles promote macrophage conversion to
393 the M2 phenotype. For example, Cao et al.⁹⁶ used ginseng-derived nanoparticles (GDNPs) to
394 treat melanoma mice and reported that GDNPs significantly promoted polarization from the M2
395 phenotype to the M1 phenotype through the activation of TLR-4/MyD88 signaling, which
396 generates total ROS, thus leading to increased apoptosis in mouse melanoma cells. In a breast



397 cancer bone metastasis mouse model, fig-derived exosome-like nanoparticles also induced M1
398 polarization by activating the atypical NF- κ B pathway.⁹⁷ Pinellia-derived exosome-like vesicles
399 can also promote M1 polarization by activating the JAK/STAT pathway, thereby inhibiting lung
400 cancer cell proliferation.⁹⁸

401 The ability of PELNs to induce macrophage conversion to either the M1 or M2 phenotype
402 may depend on their microenvironment. For example, in diabetic wounds, GDNPs effectively
403 promote M2 macrophage polarization, accelerating the healing process.⁹⁹ However, in tumor
404 tissues, GDNPs reprogram tumor-associated macrophages from an M2 to M1 phenotype, thereby
405 inhibiting tumor growth. We speculate that the diversity of active components within PELNs may
406 lead to their varying reactivity across distinct microenvironments. Additionally, the types and
407 concentrations of bioactive substances vary significantly depending on the plant source and
408 processing methods. For instance, compared to white ginseng (WG), red ginseng (RG) exhibits a
409 total saponin content that is nearly 1.8-fold higher and contains several unique ginsenosides.^{100,101}
410 Variations in origin and composition may result in functional heterogeneity among PELNs of the
411 same type.

412 In conclusion, PELNs may precisely intervenes in the interconnected pathological networks
413 of insulin resistance, abnormal lipid metabolism, chronic low-grade inflammation, and gut
414 microbiota imbalance (Figure 3), thereby demonstrating comprehensive therapeutic potential that
415 surpasses single compounds or traditional extracts. The molecular mechanisms by which PELNs
416 regulate glucolipid metabolism are summarized in Table 2.

417

418


419

420

421

422





423 **Figure 3.** The molecular mechanisms of PELNs on disorders of glycolipid metabolism. PELNs
 424 ameliorate glucolipid metabolism through multi-targets and multi-pathways, including regulating
 425 IP3K/AKT signaling pathway, protecting gut microbiota and functions, reducing oxidative stress,
 426 and alleviating inflammatory response through driving M2 macrophage polarization.

427

428 4. Current bottlenecks and challenges

429 4.1 Technical Challenges: Precise Separation and Purification

430 Exosome-like vesicles produced by different plant tissues and cell types are highly
 431 heterogeneous in quantity, size, and composition, which complicates the effective isolation and
 432 concentration of all PELNs via the use of a single method. Currently, commonly used methods
 433 demonstrate their own advantages and disadvantages in terms of yield, purity, reproducibility,
 434 and other factors.^{27-30,32-36} This scenario is specifically due to the fact that these methods involve
 435 different principles and particular defects; for example, data from different laboratories are
 436 difficult to compare horizontally, and the difficulty of reproducing research results is increased.

437 In addition, unlike animal-derived exosomes, which possess a well-established and
 438 standardized set of surface protein markers (such as CD9, CD63, and CD81),¹⁰² PELNs typically
 439 lack homologs of these animal-characteristic proteins. The surface proteome of PELNs is enriched
 440 in proteins related to plant-specific physiological processes, such as chitinases involved in cell
 441 wall metabolism.¹⁰³ Animal exosome membranes are typically rich in cholesterol, sphingomyelin,



442 and phosphatidylserine.^{104,105} In contrast, PELNs membranes lack cholesterol and are primarily
443 composed of plant sterols (such as sitosterol and stigmasterol), which serve as the main sterol
444 components.¹⁰⁶ Membrane proteins of MDEs often undergo complex N-linked and O-linked
445 glycosylation, with glycan chains frequently terminating in acidic glycans, including sialic acid,
446^{107,108} whereas PELNs surface proteins are often enriched in high-mannose-type N-glycans.^{109,110}
447 The surface molecular composition of PELNs is highly heterogeneous, which complicates the
448 accurate determination of their origin and creates challenges for marker-based extraction and
449 identification methods. Therefore, the use of high-throughput proteomics, lipidomics, and
450 glycomics technologies to identify and validate specific surface markers of PELNs on a large
451 scale (particularly those that are conserved across species or exhibit tissue specificity) is a key
452 aspect of future research.

453 **4.2 Scientific Challenges: Precise Component Analysis**

454 In-depth analysis of the biomolecular composition of PELNs is crucial for understanding
455 their functional mechanisms. However, the endogenous protein abundance within vesicles is low,
456 and the isolation process is highly susceptible to contamination by high-abundance plant matrix
457 proteins. This scenario masks the characteristic signals of membrane proteins and luminal
458 proteins of the vesicles, thus leading to difficulties in identifying genuine functional proteins.¹¹¹
459 Therefore, the development of more sensitive and specific omics analysis technologies and the
460 establishment of standardized data analysis workflows are essential for comprehensively
461 decoding the molecular “cargo” and functional codes of PELNs.

462 Second, the specific process by which PELNs are recognized and internalized by mammalian
463 cells remains incompletely understood. Celery-derived exosome-like nanovesicles exhibit greater
464 cellular uptake efficiency compared to other common plant vesicles, which suggests that they
465 possess unique ligands, structures, or lipid fractions on their surface that interact more effectively
466 with animal cell membranes.¹¹² However, the specific interaction network between these surface
467 ligands and animal cell surface receptors has not been systematically mapped and validated,
468 which directly impacts our understanding of PELNs targeting.

469 Unlike those that are directly obtained from pure natural plants, traditional Chinese herbs
470 usually need to be processed to maximize their medicinal effects. For example, the medicinal
471 efficacies of the abovementioned fresh citrus peel and dried tangerine peel exhibit considerable



472 differences. The processing of adjuvants such as vinegar, wine, honey, and salt can significantly
473 alter the contents of active or toxic components in medicinal materials or change their
474 pharmacokinetic properties through chemical or physical transformations, thereby producing
475 synergistic effects or reducing toxicity.¹¹³ For example, vinegar processing can reduce toxicity
476 in *Euphorbia kansui* through transesterification reactions while enhancing the hepatoprotective
477 activity of saponins in *Bupleurum*. Wine processing, salt processing, and other methods can alter
478 the solubility, tissue targeting, or bioavailability of components. With the development of
479 advanced analytical tools such as mass spectrometry, nuclear magnetic resonance, high-
480 throughput screening, and omics technologies, the differences in active components before and
481 after the processing of traditional Chinese medicine are expected to be clearly identified. These
482 scientific advances will provide novel therapeutic strategies, such as the combination of
483 specifically engineered plant exosome membranes with more effective and less toxic active herbal
484 ingredients.

485 **4.3 Challenges in pharmacokinetics: Systematic and Reliable Data**

486 Currently, there is a lack of systematic and reliable data regarding the blood circulation half-
487 life, organ-specific distribution, potential immunogenicity, and final metabolic clearance
488 pathways of PELNs after they enter the mammalian body. The good biocompatibility and low
489 immunogenicity of PELNs enable them to evade rapid capture and clearance by the mononuclear
490 phagocyte system,¹¹⁴ thus potentially leading to longer circulation times and broader distribution
491 in vivo. However, their specific distribution, metabolism, and clearance pathways in different
492 organs and tissues remain to be explored.¹¹⁵ Additionally, the impacts of different administration
493 routes on the in vivo behavior of PELNs can significantly vary; however, systematic evaluation
494 is lacking. Oral administration is among the most attractive delivery methods for PELNs.
495 Although some studies have demonstrated that certain plant-derived vesicles (e.g., those obtained
496 from galangal and mulberry leaves) remain stable under acidic gastric conditions,^{116,117}
497 systematic assessment and thorough validation in more complex in vivo environments (e.g., pH
498 changes and digestive enzymes in the gastrointestinal tract) are still needed. Although intravenous
499 injection allows for direct entry into the systemic circulation, whether PELNs can effectively
500 penetrate important biological barriers (such as the blood-brain barrier) to treat diseases of the
501 central nervous system requires more evidence. The retention time, penetration efficiency, and



502 local versus systemic effects of PELNs via local administration (including effects on the skin and
503 joint cavity) also require further investigation. Systematic comparisons of the stability,
504 biodistribution, and final efficacy of PELNs across different administration routes are crucial for
505 determining their optimal clinical application strategies.

506 A major reason for this data scarcity is the lack of reliable, noninvasive in vivo real-time
507 imaging and tracking technologies. Existing research methods primarily utilize fluorescent dyes
508 (such as DiR and PKH67) or the radioactive isotope labeling of vesicles. However, these labeling
509 processes may alter the physicochemical properties of the vesicle surface, thus interfering with
510 their natural interactions with biomolecules and consequently affecting their authentic in vivo
511 distribution and behavior. The development of technologies that enable high-sensitivity and high-
512 specificity tracing of PELNs without altering their natural attributes is a prerequisite for accurately
513 assessing their in vivo biodistribution and targeting efficiency.

514 **4.4 Challenges in the application and transformation processes: standardization and safety** 515 **assessment**

516 Despite the encouraging potential of these substances, the clinical translation of PELNs and
517 their engineered products continues to demonstrate significant challenges related to
518 standardization and large-scale production. First, the research and production of PELNs are
519 currently largely dependent on extraction from the juice or tissue cultures of edible plants.¹¹ This
520 traditional acquisition method exhibits significant limitations. For example, the yield of this
521 method is not only low but also severely constrained by the plant's growing season, geographical
522 origin, and specific variety, thus leading to difficulties in maintaining consistency in vesicle yield,
523 size, and bioactive components across different batches, which poses a major challenge for
524 standardized production.¹¹⁸ Additionally, previous studies have utilized plant cell suspension
525 culture systems to produce vesicles, which is considered to be a potentially scalable method.⁸⁶
526 However, the application of this cultivation platform (analogous to animal cell bioreactors) in the
527 field of plant vesicle production is still in its infancy. The efficiency and cost-effectiveness of
528 vesicle production, as well as whether the biological functions of the produced vesicles are
529 consistent with those from natural sources, require systematic validation and optimization.

530 Although the natural characteristics of PELNs contribute to their good safety profile, this
531 safety is not absolute, and potential risks from plant allergen proteins still require vigilance and



532 monitoring. Extracellular nanovesicles isolated from germinating kiwifruit pollen have been
533 observed to carry allergens, thus indicating that when vesicles are developed from specific plant
534 sources, detailed proteomic analysis is necessary to assess sensitization risks.¹¹⁹ Strict testing for
535 potential pathogens in raw materials (such as plant viruses and bacterial endotoxins) is also
536 necessary. Furthermore, data on the stability and shelf life of PELNs under specific storage
537 conditions remain insufficient.

538 **5. Conclusions**

539 Research on natural plants and foods with medicinal properties is transitioning from
540 descriptive observations to mechanistic understanding. As an emerging and rapidly developing
541 field, the core value of PELNs does not involve the simple combination of plant extracts and
542 nanotechnology. Rather, it involves a novel application of a natural, biocompatible nanodelivery
543 system to encapsulate and deliver complex active components from plants (such as small-
544 molecule metabolites, nucleic acids, and proteins) in a highly organized form. This characteristic
545 enables it to mimic and enhance the plant's inherent 'multitarget, multipathway' synergistic
546 regulatory effects.

547 PELNs are expected to establish a new paradigm for the prevention and treatment of
548 metabolic diseases; however, several key issues need to be addressed in the future. In terms of
549 standardization, the systematic characterization of PELNs products is crucial, including
550 standardized assessments of their physicochemical properties (such as particle size and charge),
551 molecular composition (proteins, RNA, and lipids), and biological activity. The development and
552 application of technologies such as mass spectrometry, proteomics, spatial omics, and single-
553 particle analysis will provide strong support for solving this problem. In terms of targeting and
554 drug-loading efficiency, engineering modifications of PELNs represent a promising strategy.
555 Through physical, chemical, and peptide-based modification strategies, the heterogeneity and
556 lack of specificity of natural exosomes can be overcome; additionally, these exosomes can be
557 transformed into more advanced drug delivery carriers by increasing drug loading potential and
558 targeting capabilities. In terms of large-scale production, end-to-end quality control spanning
559 from raw materials to final products is a core requirement for drug production. Furthermore, the
560 assurance of compliance with good manufacturing practice (GMP) requirements and the
561 performance of rigorous preclinical safety validation and clinical trials are essential steps to



562 advance PELNs toward clinical application.

View Article Online
DOI: 10.1039/D6FO01315E

563 PELNs have demonstrated promising clinical efficacy in the treatment of tumors ^{120,121} and
564 inflammatory diseases, ¹²² but reports on their therapeutic effects in glycolipid metabolic diseases
565 such as T2DM and NAFLD are currently limited (Table 3). The abundance of natural medicinal
566 plants and dietary sources provide diverse options for treating metabolic diseases. For instance,
567 patients with low abundance of *Lactobacillus* in the gut microbiota may benefit from the use of
568 broccoli-derived ELNs; ¹²³ probiotic beverages rich in grape exosomes can be utilized for daily
569 health maintenance in individuals at high risk of cardiovascular disease. ¹²⁴ Through more
570 rigorous mechanistic studies and extensive systematic *in vivo* validation, the potential of PELNs
571 in treating glycolipid metabolic diseases can be fully realized and advanced toward clinical
572 translation.

573 **Author contributions**

574 **X.L. and R.A.:** Investigation, Data curation, Writing-original draft. **H.W. and S.Y. :**
575 Visualization, Formal analysis. **X.S:** Conceptualization, Writing- reviewing & editing.

576 **Conflicts of interest**

577 There are no conflicts to declare.

578 **Data Availability Statement**

579 No primary research results, software or code have been included and no new data were
580 generated or analysed as part of this review.

581 **Acknowledgments**

582 The authors would like to acknowledge the support of an online scientific drawing platform
583 (<https://jova.ai/>) for providing high-quality, customizable scientific illustrations used in this
584 review article.

585



586 **References**

- 587 1. Dong H, Sun Y, Nie L, et al. Metabolic memory: mechanisms and diseases. *Signal*
588 *transduction and targeted therapy*. 2024;9(1):38. doi:10.1038/s41392-024-01755-x
- 589 2. Tilg H, Petta S, Stefan N, Targher G. Metabolic Dysfunction-Associated Steatotic Liver
590 Disease in Adults: A Review. *Jama*. 2026;335(2):163-174. doi:10.1001/jama.2025.19615
- 591 3. Chen H, Rosen CE, González-Hernández JA, et al. Highly multiplexed bioactivity screening
592 reveals human and microbiota metabolome-GPCRome interactions. *Cell*.
593 2023;186(14):3095-3110.e19. doi:10.1016/j.cell.2023.05.024
- 594 4. Rehman FU, Liu Y, Zheng M, Shi B. Exosomes based strategies for brain drug delivery.
595 *Biomaterials*. 2023;293:121949. doi:10.1016/j.biomaterials.2022.121949
- 596 5. Aheget H, Tristán-Manzano M, Mazini L, et al. Exosome: A New Player in Translational
597 Nanomedicine. *Journal of clinical medicine*. 2020;9(8):2380. doi:10.3390/Jcm9082380
- 598 6. Zheng M, Chavda VP, Vaghela DA, et al. Plant-derived exosomes in therapeutic
599 nanomedicine, paving the path toward precision medicine. *Phytomedicine : international*
600 *journal of phytotherapy and phytopharmacology*. 2024;135:156087.
601 doi:10.1016/j.phymed.2024.156087
- 602 7. Chu K, Liu J, Zhang X, et al. Herbal Medicine-Derived Exosome-Like Nanovesicles: A
603 Rising Star in Cancer Therapy. *International journal of nanomedicine*. 2024;19:7585-7603.
604 doi:10.2147/ijn.s477270
- 605 8. Feng ZY, Huang JW, Fu JM, Li LX, Yu R, Li L. Medicinal Plant-Derived Exosome-Like
606 Nanovesicles as Regulatory Mediators in Microenvironment for Disease Treatment.
607 *International journal of nanomedicine*. 2025;20:8451-8479. doi:10.2147/Ijn.S526287
- 608 9. Liu RL, Zhang F, He XY, Huang KL. Plant Derived Exosome-Like Nanoparticles and Their
609 Therapeutic Applications in Glucolipid Metabolism Diseases. *Journal of agricultural and*
610 *food chemistry*. 2025;73(11):6385-6399. doi:10.1021/acs.jafc.4c12480
- 611 10. Xing HJ, Li GC, Qi CL, Zhang M, Ding N, Zhang XW. Emerging Role of Plant-Derived
612 Nanostructures in Nanomedicine. *International journal of nanomedicine*. 2025;20:12715-
613 12731. doi:10.2147/ijn.s547550
- 614 11. Rome S. Biological properties of plant-derived extracellular vesicles. *Food & function*.
615 2019;10(2):529-538. doi:10.1039/c8fo02295j



- 616 12. Li A, Li D, Gu Y, et al. Plant-derived nanovesicles: Further exploration of biomedical
617 function and application potential. *Acta pharmaceutica Sinica B*. Aug 2023;13(8):3300-3320.
618 doi:10.1016/j.apsb.2022.12.022
- 619 13. Xiao X, Guo Y, Msomi NZ, Islam MS, Chu M. Exosome-like Nanoparticles Extracted from
620 Plant Cells for Diabetes Therapy. *International journal of molecular sciences*.
621 2025;26(18):9155. doi:10.3390/Ijms26189155
- 622 14. Chuo ST, Chien JC, Lai CP. Imaging extracellular vesicles: current and emerging methods.
623 *Journal of biomedical science*. 2018;25(1):91. doi:10.1186/s12929-018-0494-5
- 624 15. Huang J, Chen L, Li W, Chang CJ. Anti-inflammatory and antioxidative effects of Perilla
625 frutescens-derived extracellular vesicles: Insights from Zebrafish models. *Mol Immunol*.
626 2025;182:126-138. doi:10.1016/j.molimm.2025.04.008
- 627 16. Zhu H, He W. Ginger: a representative material of herb-derived exosome-like nanoparticles.
628 *Frontiers in nutrition*. 2023;10:1223349. doi:10.3389/fnut.2023.1223349
- 629 17. Yang S, Lu S, Ren L, et al. Ginseng-derived nanoparticles induce skin cell proliferation and
630 promote wound healing. *Journal of ginseng research*. 2023;47(1):133-143.
631 doi:10.1016/j.jgr.2022.07.005
- 632 18. Tembo KM, Wang X, Bolideei M, et al. Exploring the bioactivity of MicroRNAs
633 Originated from Plant-derived Exosome-like Nanoparticles (PELNs): current perspectives.
634 *Journal of nanobiotechnology*. 2025;23(1):563. doi:10.1186/s12951-025-03602-9
- 635 19. Zhao Q, Wang T, Wang H, et al. Consensus statement on research and application of Chinese
636 herbal medicine derived extracellular vesicles-like particles (2023 edition). *Chinese herbal
637 medicines*. 2024;16(1):3-12. doi:10.1016/j.chmed.2023.11.002
- 638 20. Aqil F, Munagala R, Jeyabalan J, Agrawal AK, Gupta R. Exosomes for the Enhanced Tissue
639 Bioavailability and Efficacy of Curcumin. *The AAPS journal*. 2017;19(6):1691-1702.
640 doi:10.1208/s12248-017-0154-9
- 641 21. Abbasifarid E, Bolhassani A, Irani S, Sotoodehnejadnematalahi F. Synergistic effects of
642 exosomal crocin or curcumin compounds and HPV L1-E7 polypeptide vaccine construct on
643 tumor eradication in C57BL/6 mouse model. *PloS one*. 2021;16(10):e0258599.
644 doi:10.1371/journal.pone.0258599

View Article Online
DOI:10.1039/D6FO01315E



- 645 22. Ning H, Huang X, Deng N, et al. Plant-Derived Exosome-Like Nanovesicles: A Novel
646 Strategy for Targeted Oral Therapy in Ulcerative Colitis. *International journal of*
647 *nanomedicine*. 2025;20:10595-10611. doi:10.2147/ijn.S536056
- 648 23. U Stotz H, Brotherton D, Inal J. Communication is key: extracellular vesicles as mediators
649 of infection and defence during host-microbe interactions in animals and plants. *FEMS*
650 *Microbiol Rev*. 2022;46(1):fuab044. doi: 10.1093/femsre/fuab044.
- 651 24. Ambrosone A, Barbulova A, Cappetta E, et al. Plant Extracellular Vesicles: Current
652 Landscape and Future Directions. *Plants (Basel)*. 2023;12(24):4141. doi:
653 10.3390/plants12244141.
- 654 25. Zeng YB, Deng X, Shen LS, et al. Advances in plant-derived extracellular vesicles: isolation,
655 composition, and biological functions. *Food & function*. 2024;15(23):11319-11341.
656 doi:10.1039/d4fo04321a
- 657 26. Kim M, Jang H, Kim W, Kim D, Park JH. Therapeutic Applications of Plant-Derived
658 Extracellular Vesicles as Antioxidants for Oxidative Stress-Related Diseases. *Antioxidants*
659 *(Basel)*. 2023;12(6)doi:10.3390/antiox12061286
- 660 27. Tiszbein K, Koss-Mikołajczyk I, Martysiak-Żurowska D. Unlocking the Secrets of Human
661 Milk: Isolation and Characterization of Extracellular Vesicles. *Advances in nutrition*
662 *(Bethesda, Md)*. 2025;16(6):100430. doi:10.1016/j.advnut.2025.100430
- 663 28. Wei C, Zhang M, Cheng J, Tian J, Yang G, Jin Y. Plant-derived exosome-like nanoparticles
664 – from Laboratory to factory, a landscape of application, challenges and prospects. *Critical*
665 *Reviews in Food Science and Nutrition*. 2025;65(23):4510-4528.
666 doi:10.1080/10408398.2024.2388888
- 667 29. Kaddour H, Tranquille M, Okeoma CM. The Past, the Present, and the Future of the Size
668 Exclusion Chromatography in Extracellular Vesicles Separation. *Viruses-Basel*.
669 2021;13(11):2272. doi:10.3390/V13112272
- 670 30. Liu Y, Xiao SQ, Wang DB, Qin CY, Wei HL, Li DW. A review on separation and application
671 of plant-derived exosome-like nanoparticles. *Journal of separation science*.
672 2024;47(8):e2300669. doi:10.1002/Jssc.202300669



- 673 31. Sidhom K, Obi PO, Saleem A. A Review of Exosomal Isolation Methods: Is Size Exclusion
674 Chromatography the Best Option? *International journal of molecular sciences*.
675 2020;21(18)doi:10.3390/ijms21186466
- 676 32. Haraszti RA, Miller R, Stoppato M, et al. Exosomes Produced from 3D Cultures of MSCs
677 by Tangential Flow Filtration Show Higher Yield and Improved Activity. *Molecular therapy :
678 the journal of the American Society of Gene Therapy*. 2018;26(12):2838-2847.
679 doi:10.1016/j.ymthe.2018.09.015
- 680 33. Busatto S, Vilanilam G, Ticer T, et al. Tangential Flow Filtration for Highly Efficient
681 Concentration of Extracellular Vesicles from Large Volumes of Fluid. *Cells*. 2018;7(12):273.
682 doi:10.3390/Cells7120273
- 683 34. Liangsupree T, Multia E, Riekkola ML. Recent advances in modern extracellular vesicle
684 isolation and separation techniques. *Journal of chromatography A*. 2026;1767:466602.
685 doi:10.1016/j.chroma.2025.466602
- 686 35. Pratiwi FW, Shanthi KB, Makieieva O, et al. Biogenesis of Mesoporous Silica Nanoparticles
687 Enclosed in Extracellular Vesicles by Mouse Renal Adenocarcinoma Cells. *Methods in
688 molecular biology (Clifton, NJ)*. 2023;2668:241-256. doi:10.1007/978-1-0716-3203-1_17
- 689 36. Chee TM, O'Farrell HE, Lima LG, et al. Optimal isolation of extracellular vesicles from
690 pleural fluid and profiling of their microRNA cargo. *Journal of Extracellular Biology*.
691 2023;2(10):e119. doi: 10.1002/jex2.119
- 692 37. Rasmussen MK, Pedersen JN, Marie R. Size and surface charge characterization of
693 nanoparticles with a salt gradient. *Nature communications*. 2020;11(1):2337.
694 doi:10.1038/S41467-020-15889-3
- 695 38. Banerjee M, Rajeswari V D. Plant-Derived Exosome-Like Nanoparticles from Elettaria
696 cardamomum Inhibit Triple-Negative Breast Cancer Cell Lines. *ChemistrySelect*.
697 2025;10(23):e04734. doi:https://doi.org/10.1002/slct.202404734
- 698 39. Alshweiat A, Csóka I, Tömösi F, et al. Nasal delivery of nanosuspension-based
699 mucoadhesive formulation with improved bioavailability of loratadine: Preparation,
700 characterization, and in vivo evaluation. *International journal of pharmaceutics*.
701 2020;579:119166. doi:10.1016/j.ijpharm.2020.119166

View Article Online
DOI:10.1039/D6FO01315E



- 702 40. Salem D, Abdel-Ghany S, Mohamed E, et al. Natural Nanoparticles for Drug Delivery
703 Proteomic Insights and Anticancer Potential of Doxorubicin-Loaded Avocado Exosomes.
704 *Pharmaceuticals-Base*. 2025;18(6):844. doi:10.3390/Ph18060844
- 705 41. Taşkan E, Kırbaş OK, Sağraç D, et al. Celery root-plant derived vesicles: comprehensive
706 isolation, characterization and proteomic analysis. *Molecular biology reports*.
707 2025;52(1):974. doi:10.1007/s11033-025-11003-2
- 708 42. Ou X, Wang H, Tie H, et al. Novel plant-derived exosome-like nanovesicles from
709 *Catharanthus roseus*: preparation, characterization, and immunostimulatory effect via TNF-
710 α /NF- κ B/PU.1 axis. *Journal of nanobiotechnology*. 2023;21(1):160. doi:10.1186/s12951-
711 023-01919-x
- 712 43. Jain N, Pandey M, Sharma P, Gupta G, Gorain B, Dua K. Recent developments in plant-
713 derived edible nanoparticles as therapeutic nanomedicines. *J Food Biochem*.
714 2022;46(12):e14479. doi:10.1111/jfbc.14479
- 715 44. Yang LY, Li CQ, Zhang YL, Ma MW, Cheng W, Zhang GJ. Emerging Drug Delivery
716 Vectors: Engineering of Plant-Derived Nanovesicles and Their Applications in Biomedicine.
717 *International journal of nanomedicine*. 2024;19:2591-2610. doi:10.2147/ijn.S454794
- 718 45. Zhang R, Li H, Mu Y, et al. Turmeric-derived extracellular vesicles loaded microneedle
719 system attenuates rotator cuff degeneration by orchestrating energetic metabolism. *Mater*
720 *Today Bio*. 2025;35:102590. doi:10.1016/j.mtbio.2025.102590
- 721 46. Liu Z, Chen S, Dou B, et al. *Salvia miltiorrhiza*-Derived Vesicle-Like Nanoparticles
722 Functionalised Hydrogel With Excellent Ability of Oxidative Stress Modulation and Anti-
723 Cardiomyocyte Apoptosis for Sepsis-Induced Myocardial Injury. *Plant Biotechnol J*.
724 2026;doi:10.1111/pbi.70574
- 725 47. Zhang Y, Zhang R, Zhang T, et al. Restoration of tendon repair microenvironment by
726 grapefruit exosome-loaded microneedle system for tendinopathy therapy. *Front Bioeng*
727 *Biotechnol*. 2025;13:1615650. doi:10.3389/fbioe.2025.1615650
- 728 48. Zhou HH, Zhou X, Pei J, et al. A fibrin gel-loaded Gouqi-derived nanovesicle (GqDNV)
729 repairs the heart after myocardial infarction by inhibiting p38 MAPK/NF- κ B p65 pathway.
730 *Journal of nanobiotechnology*. 2025;23(1):535. doi:10.1186/s12951-025-03615-4
- 731



- 732 49. Han R, Zhou D, Ji N, et al. Folic acid-modified ginger-derived extracellular vesicles for
733 targeted treatment of rheumatoid arthritis by remodeling immune microenvironment via the
734 PI3K-AKT pathway. *Journal of nanobiotechnology*. 2025;23(1):41. doi:10.1186/s12951-
735 025-03096-5
- 736 50. Tardito S, Martinelli G, Soldano S, et al. Macrophage M1/M2 polarization and rheumatoid
737 arthritis: A systematic review. *Autoimmun Rev*. 2019;18(11):102397.
738 doi:10.1016/j.autrev.2019.102397
- 739 51. Yang Y, Guo L, Wang Z, et al. Targeted silver nanoparticles for rheumatoid arthritis therapy
740 via macrophage apoptosis and Re-polarization. *Biomaterials*. 2021;264:120390.
741 doi:10.1016/j.biomaterials.2020.120390
- 742 52. Warmink K, Siebelt M, Low PS, et al. Folate Receptor Expression by Human Monocyte-
743 Derived Macrophage Subtypes and Effects of Corticosteroids. *Cartilage*.
744 2022;13(1):19476035221081469. doi:10.1177/19476035221081469
- 745 53. Zhang G, Liang H, Zhang G, et al. Low-concentration cholesterol modification enhances
746 Clematis filamentosa Dunn-derived extracellular vesicle-mediated macrophage polarization
747 regulation for acute lung injury therapy. *RSC Adv*. 2026;16(7):5941-5955.
748 doi:10.1039/d5ra08928j
- 749 54. Wang X, Xie Z, Wu R, et al. Hydrogel-Wrapped Calendula officinalis L. extracellular
750 vesicles - A novel approach to enhance fracture healing by Macrophage reprogramming.
751 *Mater Today Bio*. 2025;35:102592. doi:10.1016/j.mtbio.2025.102592
- 752 55. Muhammad Z, Muhammad SA, Abbas AY, et al. Isolation and characterization of medicinal
753 plant-based extracellular vesicles as nano delivery systems for ascorbic acid. *J*
754 *Microencapsul*. 2025;42(2):120-131. doi:10.1080/02652048.2024.2443430
- 755 56. Guo Z, Zhang Y, Gong Y, et al. Antibody functionalized curcuma-derived extracellular
756 vesicles loaded with doxorubicin overcome therapy-induced senescence and enhance
757 chemotherapy. *J Control Release*. 2025;379:377-389. doi:10.1016/j.jconrel.2025.01.029
- 758 57. Steć A, Targońska M, Jaikishan S, et al. Incorporation of doxorubicin into plant-derived
759 nanovesicles: process monitoring and activity assessment. *Drug Delivery*.
760 2025;32(1):2439272. doi:10.1080/10717544.2024.2439272

View Article Online
DOI: 10.1039/D6FO01315E



- 761 58. Mammadova R, Maggio S, Fiume I, et al. Protein Biocargo and Anti-Inflammatory Effect of
762 Tomato Fruit-Derived Nanovesicles Separated by Density Gradient Ultracentrifugation and
763 Loaded with Curcumin. *Pharmaceutics*. 2023;15(2):333.
764 doi:10.3390/pharmaceutics15020333
- 765 59. Tinnirello V, Rabienezhad Ganji N, De Marcos Lousa C, Alessandro R, Raimondo S.
766 Exploiting the Opportunity to Use Plant-Derived Nanoparticles as Delivery Vehicles. *Plants*
767 (*Basel*). 2023;12(6)doi:10.3390/plants12061207
- 768 60. Wang J, Xie F, He Q, et al. Hybrid nanovesicles derived from grapes and tomatoes with
769 synergistic antioxidative activity. *Biomater Sci*. 2024;12(21):5631-5643.
770 doi:10.1039/d4bm00591k
- 771 61. Singh A, Shadangi S, Gupta PK, Rana S. Type 2 Diabetes Mellitus: A Comprehensive
772 Review of Pathophysiology, Comorbidities, and Emerging Therapies. *Compr Physiol*.
773 2025;15(1):e70003. doi:10.1002/cph4.70003
- 774 62. Barroso E, Jurado-Aguilar J, Wahli W, Palomer X, Vázquez-Carrera M. Increased hepatic
775 gluconeogenesis and type 2 diabetes mellitus. *Trends Endocrinol Metab*. 2024;35(12):1062-
776 1077. doi:10.1016/j.tem.2024.05.006
- 777 63. Su J, Luo Y, Hu S, Tang L, Ouyang S. Advances in Research on Type 2 Diabetes Mellitus
778 Targets and Therapeutic Agents. *International journal of molecular sciences*.
779 2023;24(17)doi:10.3390/ijms241713381
- 780 64. Samms RJ, Coghlan MP, Sloop KW. How May GIP Enhance the Therapeutic Efficacy of
781 GLP-1? *Trends Endocrinol Metab*. 2020;31(6):410-421. doi:10.1016/j.tem.2020.02.006
- 782 65. Bajaj G, Choudhary D, Singh V, et al. MicroRNAs Dependent G-ELNs Based Intervention
783 Improves Glucose and Fatty Acid Metabolism While Protecting Pancreatic β -Cells in Type
784 2 Diabetic Mice. *Small*. 2025;21(4):e2409501. doi:10.1002/smll.202409501
- 785 66. Chang YC, Chan MH, Yang YF, Li CH, Hsiao M. Glucose transporter 4: Insulin response
786 mastermind, glycolysis catalyst and treatment direction for cancer progression. *Cancer Lett*.
787 2023;563:216179. doi:10.1016/j.canlet.2023.216179
- 788 67. An J, He H, Yao W, Shang Y, Jiang Y, Yu Z. PI3K/Akt/FoxO pathway mediates glycolytic
789 metabolism in HepG2 cells exposed to triclosan (TCS). *Environ Int*. 2020;136:105428.
790 doi:10.1016/j.envint.2019.105428



- 791 68. Bai Q, Song D, Gu L, Verkhatsky A, Peng L. Bi-phasic regulation of glycogen content in
792 astrocytes via Cav-1/PTEN/PI3K/AKT/GSK-3 β pathway by fluoxetine.
793 *Psychopharmacology (Berl)*. 2017;234(7):1069-1077. doi:10.1007/s00213-017-4547-3
794 69. Wang D, Wei T, Cui X, et al. Fam3a-mediated prohormone convertase switch in α -cells
795 regulates pancreatic GLP-1 production in an Nr4a2-Foxa2-dependent manner. *Metabolism*.
796 2025;162:156042. doi:10.1016/j.metabol.2024.156042
797 70. Kumar A, Sundaram K, Teng Y, et al. Ginger nanoparticles mediated induction of Foxa2
798 prevents high-fat diet-induced insulin resistance. *Theranostics*. 2022;12(3):1388-1403.
799 doi:10.7150/thno.62514
800 71. Yao Y, Chen F, Wang M, Wang J, Ren G. Antidiabetic activity of Mung bean extracts in
801 diabetic KK-Ay mice. *Journal of agricultural and food chemistry*. 2008;56(19):8869-73.
802 doi:10.1021/jf8009238
803 72. He C, Wang K, Xia J, et al. Natural exosomes-like nanoparticles in mung bean sprouts
804 possesses anti-diabetic effects via activation of PI3K/Akt/GLUT4/GSK-3 β signaling
805 pathway. *Journal of nanobiotechnology*. 2023;21(1):349. doi:10.1186/s12951-023-02120-w
806 73. Adak A, Khan MR. An insight into gut microbiota and its functionalities. *Cellular and*
807 *Molecular Life Sciences*. 2019;76(3):473-493. doi:10.1007/s00018-018-2943-4
808 74. Ma Q, Li Y, Li P, et al. Research progress in the relationship between type 2 diabetes mellitus
809 and intestinal flora. *Biomedicine & Pharmacotherapy*. 2019;117:109138.
810 doi:https://doi.org/10.1016/j.biopha.2019.109138
811 75. Wang X, Tian R, Liang C, et al. Biomimetic nanoplatform with microbiome modulation and
812 antioxidant functions ameliorating insulin resistance and pancreatic β -cell dysfunction for
813 T2DM management. *Biomaterials*. 2025;313:122804.
814 doi:10.1016/j.biomaterials.2024.122804
815 76. Ling Y, Shi Z, Yang XL, et al. Hypolipidemic effect of pure total flavonoids from peel of
816 Citrus (PTFC) on hamsters of hyperlipidemia and its potential mechanism. *Exp Gerontol*.
817 2020;130:110786. doi: 10.1016/j.exger.2019.110786
818 77. Marwaha B, Gaur SS. Valorisation of citrus peels: implications for cardiovascular diseases
819 and diabetes management. *Inflammopharmacology*. 2026;34(1):205-226. doi:
820 10.1007/s10787-025-02089-y

View Article Online
DOI:10.1039/D6FO01315E



- 821 78. Zou J, Song Q, Shaw PC, et al. Tangerine Peel-Derived Exosome-Like Nanovesicles
822 Alleviate Hepatic Steatosis Induced by Type 2 Diabetes: Evidenced by Regulating Lipid
823 Metabolism and Intestinal Microflora. *Int J Nanomedicine*. 2024;19:10023-10043. doi:
824 10.2147/IJN.S478589
- 825 79. Singh H, Singh R, Singh A, et al. Role of oxidative stress in diabetes-induced complications
826 and their management with antioxidants. *Arch Physiol Biochem*. 2024;130(6):616-641. doi:
827 10.1080/13813455.2023.2243651
- 828 80. Ighodaro OM. Molecular pathways associated with oxidative stress in diabetes mellitus.
829 *Biomed Pharmacother*. 2018;108:656-662. doi: 10.1016/j.biopha.2018.09.058
- 830 81. Gu HY, Liu N, Lin FX, Yin J. Nrf2 signaling pathway: focus on oxidative stress in
831 osteoporosis. *Osteoporos Int*. 2025;36(10):1837-1854. doi: 10.1007/s00198-025-07592-0.
- 832 82. Xiao CL, Lai HT, Zhou JJ, Liu WY, Zhao M, Zhao K. Nrf2 Signaling Pathway: Focus on
833 Oxidative Stress in Spinal Cord Injury. *Mol Neurobiol*. 2025;62(2):2230-2249. doi:
834 10.1007/s12035-024-04394-z
- 835 83. Zeng Y, Yu S, Lu L, Zhang J, Xu C. Ginger-derived nanovesicles attenuate osteoarthritis
836 progression by inhibiting oxidative stress via the Nrf2 pathway. *Nanomedicine (Lond)*.
837 2024;19(28):2357-2373. doi: 10.1080/17435889.2024.2403324.
- 838 84. Xie Q, Gu J, Sun Y, et al. Therapeutic Potential of Ginger Exosome-Like Nanoparticles for
839 Alleviating Periodontitis-Induced Tissue Damage. *Int J Nanomedicine*. 2024;19:11941-
840 11956. doi: 10.2147/IJN.S483091.
- 841 85. Zhuang X, Deng ZB, Mu J, et al. Ginger-derived nanoparticles protect against alcohol-
842 induced liver damage. *J Extracell Vesicles*. 2015;4:28713. doi: 10.3402/jev.v4.28713.
- 843 86. Cappetta E, De Palma M, Vestuto V, et al. Cardoon cell cultures as a biofactory for
844 extracellular vesicles with antisteatotic activity in an in vitro model of non-alcoholic fatty
845 liver disease. *Food Res Int*. 2026;228:118395. doi: 10.1016/j.foodres.2026.118395.
- 846 87. Taranto D, Kloosterman DJ, Akkari L. Macrophages and T cells in metabolic disorder-
847 associated cancers. *Nat Rev Cancer*. 2024;24(11):744-767. doi:10.1038/s41568-024-00743-
848 1



- 849 88. Jaitin DA, Adlung L, Thaïss CA, et al. Lipid-Associated Macrophages Control Metabolic
850 Homeostasis in a Trem2-Dependent Manner. *Cell*. 2019;178(3):686-698.e14.
851 doi:10.1016/j.cell.2019.05.054
- 852 89. Lazarov T, Juarez-Carreño S, Cox N, Geissmann F. Physiology and diseases of tissue-
853 resident macrophages. *Nature*. 2023;618(7966):698-707. doi:10.1038/s41586-023-06002-x
- 854 90. Bian Y, Li W, Jiang X, et al. Garlic-derived exosomes carrying miR-396e shapes
855 macrophage metabolic reprogramming to mitigate the inflammatory response in obese adipose
856 tissue. *J Nutr Biochem*. 2023;113:109249. doi:10.1016/j.jnutbio.2022.109249
- 857 91. Zhu M, Wang S, Yue N, et al. "One stone three birds": Andrographis paniculata-derived
858 exosome-like nanoparticles mitigate dextran sulfate sodium-induced colitis. *Chin Med J
(Engl)*. 2025;doi:10.1097/cm9.00000000000003651
- 860 92. Long M, Li J, Zhu Y, et al. Microneedle-facilitated Portulaca oleracea L.-derived
861 nanovesicles ameliorate atopic dermatitis by modulating macrophage M1/M2 polarization
862 and inhibiting NF-κB and STING signaling pathways. *Acta pharmaceutica Sinica B*.
863 2025;15(11):5966-5987. doi:10.1016/j.apsb.2025.08.021
- 864 93. Liu H, He J, Xu L, et al. Echinocystic acid in Momordica charantia L. exosome-like
865 nanovesicles attenuates dengue virus-induced vascular leakage associated inflammatory
866 mediators through macrophage metabolic reprogramming and HIF-1α/p300-CBP interaction.
867 *Front Med (Lausanne)*. 2025;12:1653689. doi:10.3389/fmed.2025.1653689
- 868 94. Gong Q, Sun C, Jiang T, Guo Y. Zanthoxylum bungeanum-Derived Nanobiotics for
869 Effective Against Ulcerative Colitis in Mouse Model. *Int J Nanomedicine*. 2025;20:6317-
870 6331. doi: 10.2147/IJN.S515961
- 871 95. Gong Q, Zeng Z, Jiang T, et al. Anti-fibrotic effect of extracellular vesicles derived from tea
872 leaves in hepatic stellate cells and liver fibrosis mice. *Front Nutr*. 2022;9:1009139. doi:
873 10.3389/fnut.2022.1009139
- 874 96. Cao M, Yan H, Han X, et al. Ginseng-derived nanoparticles alter macrophage polarization
875 to inhibit melanoma growth. *J Immunother Cancer*. 2019;7(1):326. doi:10.1186/s40425-
876 019-0817-4



- 877 97. Wang DD, Qian J, Zou HZ, et al. Fig-derived exosome-like nanoparticles attenuating bone
878 metastasis of breast cancer through establishing an anti-tumor microenvironment.
879 *Pharmacol Res.* 2026;224:108088. doi:10.1016/j.phrs.2026.108088
- 880 98. Liu M, Wang T, Fu J, et al. Pinellia exosomal vesicles remodulate tumor-associated
881 macrophage polarization via the serine synthesis/JAK/STAT signaling pathway to inhibit
882 lung cancer growth. *Phytomedicine : international journal of phytotherapy and*
883 *phytopharmacology.* 2026;151:157759. doi:10.1016/j.phymed.2025.157759
- 884 99. Fan L, Jia X, Dong F, et al. Ginseng-derived nanoparticles accelerate diabetic wound healing
885 by modulating macrophage polarization and restoring endothelial cell function. *Mater Today*
886 *Bio.* 2025;34:102143. doi: 10.1016/j.mtbio.2025.102143
- 887 100. Zhou J, Zhang J, Jing P, et al. Ginseng in white and red processed forms: Ginsenosides and
888 cardiac side effects. *Food Sci Nutr.* 2023;12(3):1857-1868. doi: 10.1002/fsn3.3879
- 889 101. Zhou S, Auyeung KK, Yip K, et al. Stronger anti-obesity effect of white ginseng over red
890 ginseng and the potential mechanisms involving chemically structural/compositional
891 specificity to gut microbiota. *Phytomedicine.* 2020;74:152761. doi:
892 10.1016/j.phymed.2018.11.021
- 893 102. Krylova SV, Feng D. The Machinery of Exosomes: Biogenesis, Release, and Uptake.
894 *International journal of molecular sciences.* 2023;24(2)doi:10.3390/ijms24021337
- 895 103. Kim JS, Song BJ, Cho YE. Pomegranate-Derived Exosome-Like Nanovesicles Containing
896 Ellagic Acid Alleviate Gut Leakage and Liver Injury in MASLD. *Food Sci Nutr.*
897 2025;13(4):e70088. doi:10.1002/fsn3.70088
- 898 104. Skotland T, Llorente A, Sandvig K. Lipids in Extracellular Vesicles: What Can Be Learned
899 about Membrane Structure and Function? *Cold Spring Harb Perspect Biol.*
900 2023;15(8)doi:10.1101/cshperspect.a041415
- 901 105. Vanherle S, Guns J, Loix M, et al. Extracellular vesicle-associated cholesterol supports the
902 regenerative functions of macrophages in the brain. *J Extracell Vesicles.* 2023;12(12):e12394.
903 doi:10.1002/jev2.12394
- 904 106. Grosjean K, Mongrand S, Beney L, Simon-Plas F, Gerbeau-Pissot P. Differential effect of
905 plant lipids on membrane organization: specificities of phytosphingolipids and phytosterols.
906 *J Biol Chem.* 2015;290(9):5810-25. doi:10.1074/jbc.M114.598805



- 907 107. Macedo-da-Silva J, Santiago VF, Rosa-Fernandes L, Marinho CRF, Palmisano G. Protein
908 glycosylation in extracellular vesicles: Structural characterization and biological functions.
909 *Mol Immunol.* 2021;135:226-246. doi:10.1016/j.molimm.2021.04.017
- 910 108. Harada Y, Ohkawa Y, Maeda K, Kizuka Y, Taniguchi N. Extracellular Vesicles and
911 Glycosylation. *Adv Exp Med Biol.* 2021;1325:137-149. doi:10.1007/978-3-030-70115-4_6
- 912 109. Slivka EV, Shilova NV, Obraztsova EA, et al. Surface Glycans of Microvesicles Derived
913 from Endothelial Cells, as Probed Using Plant Lectins. *International journal of molecular*
914 *sciences.* 2024;25(11)doi:10.3390/ijms25115725
- 915 110. Cyboran-Mikołajczyk S, Jurkiewicz P, Hof M, Kleszczyńska H. The Impact of O-
916 Glycosylation on Cyanidin Interaction with POPC Membranes: Structure-Activity
917 Relationship. *Molecules.* 2018;23(11)doi:10.3390/molecules23112771
- 918 111. Sha A, Luo Y, Xiao W, et al. Plant-Derived Exosome-like Nanoparticles: A Comprehensive
919 Overview of Their Composition, Biogenesis, Isolation, and Biological Applications.
920 *International journal of molecular sciences.* 2024;25(22)doi:10.3390/ijms252212092
- 921 112. Lu X, Han Q, Chen J, et al. Celery (*Apium graveolens* L.) Exosome-like Nanovesicles as a
922 New-Generation Chemotherapy Drug Delivery Platform against Tumor Proliferation.
923 *Journal of agricultural and food chemistry.* 2023;71(22):8413-8424.
924 doi:10.1021/acs.jafc.2c07760
- 925 113. Chen LL, Verpoorte R, Yen HR, et al. Effects of processing adjuvants on traditional Chinese
926 herbs. *J Food Drug Anal.* 2018;26(2s):S96-s114. doi:10.1016/j.jfda.2018.02.004
- 927 114. Hume DA. The mononuclear phagocyte system. *Curr Opin Immunol.* 2006;18(1):49-53.
928 doi:10.1016/j.coi.2005.11.008
- 929 115. Zeng H, Wang Q, Wang F, et al. Extracellular vesicles derived from Citrus plants: innovative
930 applications in therapeutics and drug delivery. *Journal of nanobiotechnology.*
931 2026;doi:10.1186/s12951-026-04210-x
- 932 116. Nemidkanam V, Chaichanawongsaroj N. Characterizing *Kaempferia parviflora* extracellular
933 vesicles, a nanomedicine candidate. *PloS one.* 2022;17(1):e0262884.
934 doi:10.1371/journal.pone.0262884

View Article Online
DOI:10.1039/D6FO01315E



- 935 117. Chen L, Zu M, Cao Y, et al. Oral Plant-Derived Nanomedicines Mitigate Acetaminophen-
936 Induced Liver Injury by Modulating the Gut-Liver Axis and Intestinal Microbiota
937 Metabolism. *Small*. 2025;21(31):e2502001. doi:10.1002/smll.202502001
- 938 118. Bodala CK, Parveen A, Gogireddy BL, Priya S, Srinivas L. Plant-Based Exosomes: Insights
939 and Therapeutic Applications. *Crit Rev Ther Drug Carrier Syst*. 2026;43(1):1-55.
940 doi:10.1615/CritRevTherDrugCarrierSyst.2025057676
- 941 119. Suanno C, Tonoli E, Fornari E, et al. Small extracellular vesicles released from germinated
942 kiwi pollen (pollensomes) present characteristics similar to mammalian exosomes and carry
943 a plant homolog of ALIX. *Front Plant Sci*. 2023;14:1090026.
944 doi:10.3389/fpls.2023.1090026
- 945 120. Wang YY, Zhang X, Xu XY, et al. A Bioinspired Exosomal Nanoplatfor for Coordinated
946 Sorafenib and MicroRNA Delivery to Sensitize Ferroptosis and Induce Immunoactivation in
947 Triple-Negative Breast Cancer. *ACS Nano*. 2026. doi: 10.1021/acsnano.6c02310
- 948 121. An Y, Xu JZ, Ye GC, et al. Therapeutic translation of traditional Chinese medicine Huangqi
949 derived exosome like nanoparticles: targeting prostate cancer through ferroptosis activation,
950 immune reprogramming, and microbiome modulation. *J Nanobiotechnology*. 2026. doi:
951 10.1186/s12951-026-04160-4
- 952 122. Duan X, Li J, Gao R, et al. Antimicrobial hydrogel loaded with broccoli exosomes promotes
953 anti-scarring healing of MRSA-infected wounds. *Mater Today Bio*. 2025;35:102276. doi:
954 10.1016/j.mtbio.2025.102276
- 955 123. Filannino P, Bai Y, Di Cagno R, et al. Metabolism of phenolic compounds by *Lactobacillus*
956 spp. during fermentation of cherry juice and broccoli puree. *Food Microbiol*. 2015;46:272-
957 279. doi: 10.1016/j.fm.2014.08.018
- 958 124. Teng Y, He J, Zhong Q, et al. Grape exosome-like nanoparticles: A potential therapeutic
959 strategy for vascular calcification. *Front Pharmacol*. 2022;13:1025768. doi:
960 10.3389/fphar.2022.1025768



Table 1. Characteristics of the Main Separation and Purification Methods for PELNs.

Method of isolation		Basic principles	Purity	Quality	Advantages	Disadvantages	Ref
Ultracentrifugation (UC)	Differential ultracentrifugation (DUC)	Different settling velocities of particles with different sizes and densities	Limited	Vesicles are prone to aggregation or damage	No exogenous reagents	Complex operation; Time-consuming; Co-precipitation of impurities	27, 28
	Density gradient ultracentrifugation (DGUC)	Different densities of particles in a density gradient medium	Higher than that of the DUC method	High-concentration medium may affect vesicles' biological activity	High purity and recovery	Complex operation; Time-consuming	27, 28
Membrane-based filtration	Ultrafiltration (UF)	Micropores on the membrane surface screened particles with specific molecular weights	Lower than that of the DGUC method	No impact on vesicles' biological activity, but may cause structural deformation or damage	Simple operation; Suitable for large-volume samples	Co-precipitation of impurities with similar particle sizes; Prone to clogging membrane pores	31
	Tangential Flow Filtration (TFF)	Utilizing the fluid shear force parallel to the membrane surface	Lower than that of the DGUC method	Vesicles' biological activity and structure remain intact	Efficient and scalable production	Co-precipitation of impurities with similar particle sizes	31, 32
Size exclusion chromatography (SEC)		Particles of different sizes flow out of the chromatographic column packed with porous spheres at different rates	Higher than that of the DUC and DGUC method	Vesicles' biological activity and structure remain intact	Simple operation; High efficiency in removing impurities	Co-precipitation of impurities with similar particle sizes; Additional concentration steps; Prone to clogging chromatographic column	29
Polymer-based precipitation (PBP)		PEG competitively binds water molecules to separate PELNs from the solution	Low	Polymers may alter the physicochemical properties of the vesicle surface	Simple operation; Suitable for large-volume samples	Co-precipitation of other biological contaminants	30
Microfluidics		Combining physical filtration, electric fields or antibody capture	High	Vesicles' biological activity and structure remain intact	Highly automated processes; Fast separation speed	Expensive equipment; Limited in large-scale settings	34



Table 2. The mechanisms by which PELNs regulate glucolipid metabolism

PELNs	Disease model	Molecular mechanism	Regulatory factors	Therapeutic effects	Ref.
GELNs	T2DM mouse	Activating PI3K/Akt signaling via phosphorylating IRS-1; Regulating PI3K/Akt signaling via miRNAs;	PCK-1 ↓ G6PC ↓ GYS-2 ↑ SREBP-1c ↓ FAS ↓ PCK-1 ↓	Ameliorating insulin resistance; Reducing hepatic lipid deposition;	65
		Reshaping the structure of the gut microbiome; Increasing the abundance of beneficial bacteria;	<i>Lactobacillus</i> ↑ Indole ↑ indoleacetic acid ↑ Occludin ↑	Enhancing insulin sensitivity; Alleviating hepatic steatosis; Reducing serum ALT and AST;	75
GDNPs/ Phosphatidic acid in GDNPs	HFD-fed mouse	Inhibiting Akt-1-mediated Foxa2 phosphorylation and nuclear export; Increasing the proportion of phosphatidic acid and reducing the level of phosphatidylcholine in intestinal epithelial cell-derived exosomes;	Foxa2 ↑	Enhancing insulin sensitivity and glucolipid metabolism;	70
Gingerolin GELNs		Regulating TLR4/TRIF/Nrf2 signaling;	HO-1 ↑ NQO1 ↑ GSH ↑	Alleviating oxidative stress and inflammatory reactions	85



MELNs	HFD-induced diabetic mice	Activating PI3K/Akt signaling	GLUT4 ↑	Ameliorating insulin resistance;	72
		Regulating the PI3K/Akt/GSK-3β/Nrf2 signaling; Upregulating the expression of antioxidant enzymes;	HO-1 ↑ SOD ↑ GSH-Px ↑	Alleviating hepatic inflammatory infiltration and steatosis; Alleviating oxidative stress;	
TNVs	T2DM mouse	Reshaping the structure of the gut microbiome; Increasing the abundance of beneficial bacteria;	Claudin-1 ↑ ZO-1 ↑ Occludin ↑ <i>Lactobacillaceae</i> ↑ <i>Muribaculaceae</i> ↑ <i>Lachnospiraceae</i> ↓ <i>Desulfovibrionaceae</i> ↓	Enhancing insulin sensitivity; Repairing the intestinal mucosal barrier;	78
		Regulating genes relevant to glucolipid metabolism;	CD36 ↓ FASN ↓ LXR-α ↓ PPAR-γ ↓ SREBP-1c ↓ CPT1/2 ↑ FGFR4 ↑ PGC-1α ↑ PPAR-α ↑ UCP1 ↑	Alleviating hepatic steatosis;	



TNVs	T2DM mouse	Activating FXR/SHP/FGF19 signaling;	NTCP ↓ CYP7A1 ↓ HMG-COA ↓ MDR2 ↑ SBEP ↑ ABCG5/8 ↑	Maintain bile acid homeostasis;	78
Cardoon cell-derived vesicles	cellular model of NAFLD	Activating the Sirt-1/AMPK signaling;	Sirt-1 ↑ P-AMPK ↑ ROS ↓ NO ↓	Alleviating hepatic steatosis;	86
GDEs	HFD-fed mouse	Down-regulating PFKFB3 expression via miRNA-396e;	M2 macrophage ↑	Ameliorating insulin resistance; Inhibiting the inflammatory response; Enhancing lipid metabolism;	90

Abbreviations: GDEs, garlic-derived exosomes; GDNPs, ginger-derived nanoparticles; GELNs, ginger-derived exosome-like nanoparticles; HFD, high fat diet; IRS-1, insulin receptor substrate-1; MELNs, mung bean sprouts-derived exosome-like nanoparticles; TNVs, tangerine peel-derived exosome-like nanovesicles.

Table 3. The clinical application of PELNs in glycolipid metabolic disorders

PELNs	Disease model	Engineering strategies	Administration route	Dose	Control drug and dose	Therapeutic effects	Safety	Efficacy	Ref.
GELNs	T2DM mouse	NA	oral administration	1, 5, or 10 mg/kg	Met: 250 mg/kg;	Reduced fasting blood glucose levels of 6h fasted mice	No significant toxicity was observed based on histological and haematological assessments	Equally effective as Met	65
GDNPs	HFD-fed mouse	NA	dissolving administration	6 x 10 ⁸ /mL in the drinking water	NE	Ameliorating insulin resistance; Inhibiting inflammation	NE	NE	70
GELNs	T2DM mouse	coated GELNs onto the surface of HMS loaded with A (HMS/A@GE)	oral administration	5 mg/kg, once every two days;	Free A solution: 0.4 mg/kg, once every two days; HMS@GE: 5 mg/kg, once every two days;	Elevating insulin sensitivity; Alleviating liver steatosis; Ameliorating inflammatory response and oxidative stress	No significant toxicity was observed based on histological and haematological assessments	The therapeutic effect of HMS/A@GE was significantly better than that of HMS@GE or A alone	75
TNVs	T2DM mouse	NA	oral administration	200 mg/kg,	Met: 250 mg/kg;	Ameliorating insulin resistance; Restoring intestinal mucosal barrier;	NE	Equally effective as Met	78
GDEs	HFD-fed mouse	NA	intragastrically administrated	100, 200, or 400 mg/mL, once every two days;	NE	Ameliorating insulin resistance; Ameliorating inflammatory response;	NE	NE	90

Abbreviations: A, ammonia borane; GDEs, garlic-derived exosomes; GDNPs, ginger-derived nanoparticles; GELNs, ginger-derived exosome-like nanoparticles; HFD, high-fat diet; HMS, hollow mesoporous silica; Met, metformin; NA, not applicable; NE, not evaluated; TNVs, tangerine peel-derived exosome-like nanovesicles.



Data Availability Statement

View Article Online
DOI: 10.1039/D6FO01315E

No primary research results, software or code have been included and no new data were generated or analysed as part of this review.

