

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

DNA Based Multi-Copper Ions Assembly using Combined Pyrazole and Salen Ligandosides

+Cite this: DOI: 10.1039/x0xx00000x

Meng Su,^a María Tomás-Gamasa,^a and Thomas Carell^{*a},

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The DNA structure is an ideal building block for the construction of functional nano-objects. In this direction, metal coordinating base pairs (ligandosides) are an appealing tool for the future specific functionalization of such nano-objects. We present here a study, in which we combine the metal ion coordinating pyrazole ligandoside with the interstrand crosslinking salen ligandoside system. We show that both ligandosides, when combined, are able to create stable multi-copper ion complexing DNA double helix structures in a cooperative fashion.

Introduction

The Watson-Crick base pairing of DNA is a phenomenon that is more and more exploited for the construction of complex two and three dimensional self-assembled nano-objects.¹⁻³ Nowadays, sophisticated technologies provide scientists almost total control of the self-assembly process so that basically any desired nano-object can be created using DNA as the building block.⁴⁻⁶ The next phase in the field is the functionalization of these objects, thus facilitating access to materials of relevance, e.g. “intelligent” drug delivery devices⁷ or building blocks useful in nano-machines.⁸ One approach for the construction of alternative and functional DNA structures was based on the design of bases showing novel base-pairing schemes,⁹⁻¹¹ including those held together by hydrogen bonding patterns different from those of the natural base pairs¹² or by simply hydrophobic interactions.^{13, 14} The development of metal-mediated base pairs (ligandosides) represents a research direction of particular interest in respect to the desired functionalization of DNA.¹⁵⁻¹⁷ They can add properties to DNA, such as conductivity or magnetism,¹⁸⁻²⁰ or they can be used to construct DNA able to perform logic operations.^{21, 22}

Metal base pairs comprise natural or artificial nucleobases that are able to coordinate a central metal ion. The first group involves the conversion of a natural mismatched base pair into a metal

coordinated pair. Examples include the coordination of Hg²⁺ and Ag⁺ by T-T, U-U or C-A mismatches which goes in hand with the formation of stable metal base pairs.²³⁻²⁶

Regarding the artificial base pairs, a plethora of different metal modified double helices were generated depending on the choice of the synthesized ligandoside. Diverse systems, including monodentate,^{27, 28} bidentate²⁹⁻³⁴ as well as homo tridentate ligandosides,³⁵⁻³⁸ which can complex a variety of metal ions (e.g., Cu²⁺, Ni²⁺, Pd²⁺) have been reported to date. Fig. S1 summarizes current reported structures. Last examples comprise new bimetal base pairs,^{39,40} which in principle would allow to construct DNA structures with heavy metal ion loading.

In 2005, we introduced the salen concept for the construction of a new metal base pair.⁴¹⁻⁴⁵ This ligandoside involves formation of a covalent linkage between the two strands via a bridge established by ethylenediamine, as depicted in Fig. 1a. Due to the formed interstrand crosslink, the salen base pair was proven to be extremely stable after coordination of a copper ion.

In order to broaden this concept and to move away from the linking character of the salen system, we recently reported the pyrazole base pair (Pz) displayed in Fig. 1b.⁴⁶ This ligandoside construction lacks the bridge but still retains the same metal coordination geometry.





Fig. 1 Depiction of (a) the reversible salen (S) self-base pair; (b) the pyrazole (Pz) self base pair. R = deoxyribose.

Here we present stability and selectivity studies with the Pz ligand. We report that by combining the Pz and the salen ligand, the generation of stable multi-Cu²⁺ complexing DNA duplexes becomes possible.

Results and Discussion

We synthesized the pyrazole ligand Pz and Pm according to our published procedure.⁴⁶ Table 1 summarizes the modified DNA single strands prepared for this work.

Table 1 Depiction of the ligandosides Pz and Pm together with the synthesized oligonucleotides needed for this study.



No.	5' ----- 3' [a]	Mass (calc.)	Mass (exp.) [b]
0a	CAC ATT AGT GTT GTA	4579.8	4576.5
0b	CAC ATT ACT GTT GTA	4557.8	4553.7
1a	CAC ATT AXT GTT GTA	4588.8	4585.0
1b	TAC AAC AXT AAT GTG	4606.8	4602.9
2a	CAC ATT AMT GTT GTA	4602.8	4597.9
2b	TAC AAC AMT AAT GTG	4620.9	4616.3
3a	CAC ATT XXT GTT GTA	4613.8	4611.6
3b	TAC AAC AXX AAT GTG	4640.9	4636.1

4a	GCGCG XXXXX XXXXX GGCCG	6449.2	6452.4
4b	CGGCC XXXXX XXXXX CGCGC	6369.2	6370.8
5a	CAC STT AXT GTS GTA	4571.8	4570.0
5b	TAC SAC AXT AAS GTG	4589.8	4586.8
6a	CAC STT AMT GTS GTA	4585.8	4586.8
6b	TAC SAC AMT AAS GTG	4603.8	4604.2
7a	CATGST XGSA X CSTX CSTGCA	6451.1	6450.7
7b	TGCASG XASG X TSCX ASCATG	6500.1	6500.1
8a	GCGCG XXXXX GGCCG	4758.9	4757.6
8b	CGGCC XXXXX CGCGC	4678.9	4676.6

[a] X = Pz, M = Pm, S = salen base. [b] Data from MALDI-TOF mass spectrometry.

Complexation properties

To determine the thermal stability of DNA duplexes containing the ligands Pz and Pm, melting points were measured in the absence and in the presence of copper ions.

For duplexes **1a/1b** and **2a/2b** having one artificial base pair inserted, a decrease of the melting point value from 49 °C (T_M of the reference duplex **0a/0b** containing a GC base pair) to 41 °C and 42 °C was observed, respectively. Upon metal complexation, the duplex with the Pz-Cu-Pz complex furnished a melting temperature of 50 °C, slightly above the original T_M of the duplex **0a/0b**. In contrast, no increase of the T_M value was detected for duplex **2a/2b** (42 °C). This result shows that the presence of the phenolic hydroxyl groups and in particular, their ability to become deprotonated, are critical for metal complexation, which confirms the coordination geometry proposed in Fig 1b.

In order to investigate in more detail the need for deprotonation, a study of the relationship between duplex thermostability and the pH value was carried out, using duplexes **1a/1b** and **2a/2b**. The data are compiled in Fig. 2, Fig. S2, S3 and Table S1. At pH 6.0, no substantial differences between the presence and absence of copper ions were found. The T_M is approximately 8 °C lower than the T_M of the canonical duplex **0a/0b**. At this pH, the phenolic hydroxyl groups are obviously not deprotonated, which blocks Cu²⁺ complexation. Interestingly, at physiological pH 7.4, a slight difference (51 °C to 47 °C) regarding the melting point values between **1a/1b** without and with Cu²⁺ was observed. We speculate that, at this pH, half deprotonation takes place, which leads, in the absence of Cu²⁺, to the formation of a stabilizing H-bonding bridge between the Pz-Pz base pair. Upon addition of Cu²⁺, the copper base

pair is formed and is as stable as the H-bonded Pz-Pz self pair. At pH 9.0, the metal-free duplex is less stable, likely because of phenoxide formation followed by charge repulsion. Addition of Cu^{2+} stabilizes the system, in agreement with formation of the Pz-Cu-Pz base pair.




Fig. 2 Melting temperatures of duplexes **0a/0b** and **1a/1b** at different pH values. Conditions: 150 mM NaCl, 10 mM $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$ buffer pH 6.0 / 7.4 or CHES buffer pH 9.0, 1 μM oligonucleotide, with or without 1 μM Cu^{2+} , final volume of 200 μL .

The overall duplex conformation was further investigated by circular dichroism spectroscopy. All measured CD spectra of DNA containing the Pz ligand side in the absence and presence of Cu^{2+} support the presence of a B-form helix (Fig. S4, S5).

Selectivity of the Pz base

Next, the selectivity of the new Pz ligand side self pair was studied. Melting profiles of a Pz base containing strand **1a** with a complementary strand with a canonical base at the opposite position were measured at pH 9.0 (Fig. 3). The results showed that the Pz-C situation destabilizes the duplex dramatically while Pz-A/G/T arrangements give duplexes with lower but similar stability. In the presence of copper ions, the Pz-Cu-Pz metal base pair is by far the most stable structure which is the basis for selective $\text{Cu}(\text{II})$ -complexation.




Fig. 3 Melting temperatures of **1a** combined with different counter strands having A, G, C, T or Pz as the counter base. Conditions: 150 mM NaCl, 1 mM CHES buffer pH 9.0, 1 μM oligonucleotide, with or without 1 μM Cu^{2+} , final volume of 200 μL .

Multiple metal ion binding by (Pz-Pz)_n-DNA

To answer the question of whether the Pz ligand side allows construction of DNA structures containing more than one metal ion stacking on top of each other we prepared oligonucleotides with several consecutive Pz ligand sides (Table 1, duplexes **3a/3b** and **4a/4b**).

As a first approach, two complementary DNA single strands, each containing two neighbouring Pz pyrazole nucleosides, were hybridized to form the duplex **3a/3b**. Titration melting profiles are depicted in Fig. 4. In the absence of copper ions, the T_M of the duplex is decreased by about 11 $^\circ\text{C}$ ($T_M = 38$ $^\circ\text{C}$, Fig. 4, black line) with respect to the double strand **0a/0b** and by 3 $^\circ\text{C}$ with respect to the duplex **1a/1b** with only one artificial Pz base pair.

When one equivalent of Cu^{2+} was added, the system showed two sigmoidal transitions (Fig. 4, blue line). One transition was detectable at $T_M = 38$ $^\circ\text{C}$ (corresponding to the T_M for the duplex without Cu^{2+}) while the second transition appeared at a T_M of 70 $^\circ\text{C}$ (see also Fig. S6).





Fig. 4 Schematic depiction of the investigated duplex **3a/3b** and Cu^{2+} titration melting profiles of duplex **3a/3b** (Pz = Pyrazole). Conditions: 150 mM NaCl, 10 mM CHES buffer pH 9.0, 1 μM oligonucleotide, final volume of 200 μL .

These data indicate the presence of two well-defined species in solution. With one equivalent of Cu^{2+} ions in solution, half of the duplexes contains two metal ions ($T_M = 70\text{ }^\circ\text{C}$), while the other duplexes feature no metal ion ($T_M = 38\text{ }^\circ\text{C}$), arguing that Cu^{2+} complexation by the Pz ligand is a cooperative effect. This is in agreement with the observation of isosbestic points in the UV and CD titration experiments (Fig. S7, S8). When two or more equivalents of Cu^{2+} are present, the melting curves exhibit only the transition at $70\text{ }^\circ\text{C}$ (Fig. 4, green and red lines), showing full saturation of all Cu^{2+} binding sites.

In order to investigate if more than two Cu^{2+} ions can be arranged in a line, we next studied a duplex with ten consecutive Pz ligand sites, where GC rich DNA sequences were used as the terminus (Table 1, duplex **4a/4b**).

After hybridization in the presence of Cu^{2+} , the duplex **4a/4b** was again subjected to thermal UV analysis. However, due to its high stability, the melting point could not be accurately measured any more under the tested conditions. Therefore, CD titrations (Fig. 5, see also Fig. S9 for titrations at 240 nm) and UV titrations (Fig. S10) at the same concentration were employed to monitor the Cu^{2+} assembling process. The overlaid CD spectra revealed that the structure of the duplex changes significantly upon increasing Cu^{2+} complexation. A true isosbestic point is not present any more. Plotting of the ellipticity at 300 nm against the equivalents of Cu^{2+}

ions displayed however, a linearly decrease until a $[\text{Cu}^{2+}]/[\text{duplex}]$ ratio of about 10:1 was reached, in agreement with the expected complexation stoichiometry. From the experiments we conclude that the assembly of $10 \times (\text{Pz-Cu-Pz})$ is indeed possible. However, the duplex forms a distorted structure that appears to be tolerated by the flexible Pz-self base pair. The complexation process is certainly not a cooperative event, which explains the absence of clear isosbestic point. We also observe at 240 nm structural changes beyond the titration of 10 eq. of Cu^{2+} . We believe that these changes are due to additional association of Cu^{2+} with the multiple Pz-structure. These additional structural changes are not observable at longer wavelength. We therefore examined next how the presence of stiff, while crosslinking, salen ligand osides (S, Fig. 1a) would influence



the copper coordination process.

Fig. 5 (a) Schematic depiction of the assembly of 10 Cu^{2+} pyrazole ligand osides inside a DNA duplex **4a/4b**; (b) CD spectral changes of the duplex **4a/4b** at various concentrations of Cu^{2+} (from 0 to 15 eq., step of 1 eq.), spectra of 0 eq. and 10eq. are showed in red. Inset: plot of circular dichroic changes at 300 nm against the ratio of $[\text{Cu}^{2+}]/[\text{duplex}]$. Conditions: 150 mM NaCl, 10 mM CHES buffer pH 9.0, 1 μM oligonucleotide, final volume of 200 μL .

Multiple metal ion binding by hybrid Pz/S containing DNA

To this end mixed strands were prepared. For an initial study, oligonucleotides containing two salen nucleobases (S) and one additional Pz nucleobase in the middle were synthesized. In principle, this duplex can complex three metal ions (Table 1, duplex **5a/5b**).

The characteristic changes in the UV/Vis and CD spectrum of duplex **5a/5b** that occur upon titration with Cu^{2+} ions are presented in Fig. 6a, Fig. S11 and Fig. S12. The overlaid curves show now isosbestic points again at $\lambda = 340$ and 396 nm. A plot of the absorbance at 360 nm against the equivalents of Cu^{2+} ions is depicted in Fig. 6b. Initially, when one equivalent of Cu^{2+} was added, a value of 0.025 for the absorbance was measured. With increasing amounts of Cu^{2+} (up to three equivalents), the absorbance raised to 0.055 , which is a typical behaviour for copper complexation by a salen ligand.⁴² Further titration did not affect the absorbance any more. A similar trend was observed at 235 nm (Fig. S13). These results confirm the complexation of three metal ions and the data reflect that the first metal ion is complexed by the Pz-Pz pair, followed by complexation of the other two ions by the two salen ligandosides. It is clear that the complexation process by both ligandosides follows a different kinetic scheme.

More experimental support for this hypothesis was obtained when the Pz nucleoside was replaced by methylated Pz (Pm), which is unable to coordinate copper ions. UV titration of duplex **6a/6b** containing now the Pm instead of the Pz ligandosides showed that the absorbance at 360 nm increased up to a value of 0.050 after addition of 2 equivalents of Cu^{2+} , characteristic for salen copper complexes (Fig. 6c/d). Further addition of metal ions didn't change the absorbance. As Pm fails to coordinate copper, the ions go directly into the two salen base pairs. Because the increase at 360 nm for **6a/6b** is similar to those for **5a/5b**, we conclude that two equivalents of Cu^{2+} are complexed in this case. These observations support the hypothesis, that the Pz-Pz base pair is the first ligand side loaded with a metal ion.




Fig. 6 (a) Overlaid UV spectrum obtained at different concentrations of Cu^{2+} (from 0 to 4 eq.) of the duplex **5a/5b** and (b) Plot of UV absorbance at 360 nm against the ration of $[\text{Cu}^{2+}]/[\text{5a/5b}]$; (c) Overlaid UV spectrum at different concentrations of Cu^{2+} (from 0 to 4 eq.) of the duplex **6a/6b** and (d) Plot of UV absorbance at 360 nm against the ration of $[\text{Cu}^{2+}]/[\text{6a/6b}]$. Conditions: 150 mM NaCl, 10 mM CHES buffer pH 9.0 , 3 μM oligonucleotide, 30 eq. ethylenediamine, final volume of 200 μL .

For the characterization of these duplexes, ESI-Mass measurements were performed. Double strand **1a/1b**, hybridized with copper, provided always two main signals corresponding to the two single strands. We were unable to detect the desired mass peak of the duplex containing a copper ion inside, confirming that complexation and decomplexation of Pz-Pz \rightleftharpoons Pz-Cu-Pz is fast. In contrast, the same experiment performed with the duplex **5a/5b** with an excess of ethylenediamine and 3 equivalents of Cu^{2+} showed in the mass spectrum quantitative formation of the duplex containing two copper complexes (Fig. 7, Fig. S14). The third copper ion could not be detected in agreement with fast Pz-Cu-Pz decomplexation. Obviously, and in agreement with earlier data,⁴² in the MS experiment, only the Cu-salen complexes are stable enough to survive the ESI conditions.




Fig. 7 ESI-Mass spectrum and comparison of experimental data with calculated molecular weights of duplex **6a/6b** with 2 eq. Cu^{2+} , molecular formula

$C_{316}H_{384}O_{176}N_{96}P_{28}Cu_2$. Peaks contain 10 charges (top) compared with calculation (bottom). Conditions: 150 mM NH_4OAc , 10 mM CHES buffer pH 9.0, 30 μM oligonucleotide, 30 eq. ethylenediamine.

A more complicated scenario was found when four Pz pairs were mixed with three salen pairs in duplex **7a/7b**. The complexation of copper ions in this new situation was confirmed by CD titration. The CD data prove complexation of 7 equivalents of Cu^{2+} ions within the duplex (Fig. 8a/b; see also Fig. S15 for overlaid UV spectrum). The overlaid CD spectrum shows, despite the complex ligand composition of the duplex, two isosbestic points at 246 nm and 265 nm arguing for cooperative metal complexation. A plot of the UV absorbance at 405 nm during metal ion titration is displayed in Fig. 8d (see also Fig. S16 for absorbance at 360 nm). The UV data reveal two different trends which correspond to the two different coordination events established by the Pz and the salen base. Addition of three equivalents of Cu^{2+} led to a first rather sharp decrease in the absorbance at 405 nm, which corresponds to the first coordination process. With 3 Pz ligands we assume again initial complexation of Cu^{2+} by this ligand system. Thereafter, addition of another four equivalents of Cu^{2+} gave a further change of the UV data. Now, the reduction of the absorbance is slower. With all these evidences in hand, and accordingly to the previous hypothesis and results, we conclude that the Pz pairs complex the copper ions first while the 4 salen ligands are filled in the second process.




Fig. 8 Schematic depiction of the investigated duplex, S = salen, X = pyrazole. (a) Overlaid CD spectrum at different concentrations of Cu^{2+} (from 0 to 10 eq.) of the duplex **7a/7b**; (b) Plot of circular dichroic changes at 250 nm against the ration of $[Cu^{2+}]/[7a/7b]$; (c) Overlaid UV spectrum under the same conditions as in (a); (d) Plot of


UV absorbance at 405 nm against the ration of $[Cu^{2+}]/[7a/7b]$. Conditions: 150 mM NaCl, 10 mM CHES buffer pH 9.0, 3 μM oligonucleotide, 30 eq. ethylenediamine, final volume of 200 μL .

Catalysis of a Diels-Alder reaction by $(Pz-Cu^{2+}-Pz)_n$ -DNA

In order to investigate if the multiple Cu-containing DNA strands are able to exhibit some function e.g. as catalyst we started to investigate their ability to catalyze a Diels-Alder reaction. For other enantioselective Diels Alder reactions based on Cu^{2+} -ligand-DNA see ref. 47, 48 and for Cu^{2+} -G-quadruplex see ref. 49.

The D-A reaction between aza-chalcone (**1**) and cyclopentadiene (**2**), using duplex **8a/8b** (system with five Cu^{2+} ions complexed up in Pz containing DNA) as a catalyst was selected for this initial study and the results are summarized in Table 2. For us a central question was if the duplexes, which contain the metal ions in the middle are able to perform the catalysis and whether enantioselectivity can be obtained. We noted that both single and double strands with the Pz-ligands catalyze the reaction in the presence of Cu^{2+} and in both cases some chirality transfer is observed. At the same time, the complex accelerates the conversion. Copper counter ion and surrounding pH are also playing a role in the catalytic process. The result shows that the Pz-containing DNA strands can be turned into catalyst. Now, their efficacy and the ee-values obtained for the reaction need to be improved.

Table 2 Results of the catalytic Diels-Alder reaction with ds **8a/8b**.^[a]



Entry	Ligand	[Cu]	pH	ee (%)	endo/exo	conv. (%)
1	8a/8b	$Cu(NO_3)_2$	6.5	39	93	97
2	-	$Cu(NO_3)_2$	6.5	1	87	76
3	ss 8a	$Cu(NO_3)_2$	6.5	29	93	96
4	8a/8b	$Cu(NO_3)_2$	7.4	36	86	79
5	8a/8b	$Cu(NO_3)_2$	9.0	23	80	47
6	8a/8b	$CuSO_4$	6.5	32	87	52
7	8a/8b	$Cu(OTf)_2$	6.5	29	84	46
8	8a/8b	-	6.5	3	76	19

[a] See ESI for reaction details. All data are averaged over two experiments. OTf = trifluoromethanesulfonate.

Conclusion

The metal base pair, pyrazole ligand, was synthesized and incorporated into oligonucleotides. The ligand system shows a pH-dependent complexation behaviour due to the need for deprotonation of the phenolic groups. When the OH groups are replaced by a methoxy moiety as in the Pm ligand, no metal ion coordination is observed. Cu²⁺ ions stabilize a Pz-Pz containing duplex when compared to a canonical G-C pair, at pH 9.0.

The Pz-Pz ligand base pair allows complexation of up to 10 Cu²⁺ ions within a duplex. Better complexation of multiple Cu²⁺ ions is, however, observed when the unbridged Pz ligand is combined with the bridged salen (S) system which seems to add so much integrity and duplex stiffness that the metal ion complexation process is dominated by cooperative effects. The copper ions show a kinetic preference for complexing first into the Pz-Pz base pair while the salen complex is loaded in a second independent step. This stepwise complexation enables in principle the design of logic gates. The described complexation properties are indeed now the basis for the construction of defined metal ions clusters within oligonucleotides, with the goal to construct multi-metal ion based catalysts and logic gates.

Acknowledgements

We thank the Alexander von Humboldt Foundation for a post-doctoral fellowship to M. T.-G. Funding for this research was obtained from the Volkswagen Foundation and the DFG (SFB749, TP4 and SFB1032, TPA5).

Notes and references

^a M. Su, Dr. M. Tomás-Gamasa, and Dr. T. Carell, Department of Chemistry, Ludwig-Maximilians University, Butenandtstraße 5-13, 81377, Munich, Germany. E-mail: thomas.carell@cup.uni-muenchen.de;

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

1. N. C. Seeman, *Nature*, 2003, 421, 427-431.
2. S. M. Douglas, H. Dietz, T. Liedl, B. Hogberg, F. Graf and W. M. Shih, *Nature*, 2009, 459, 414-418.
3. M. Endo, K. Hidaka, T. Kato, K. Namba and H. Sugiyama, *J. Am. Chem. Soc.*, 2009, 131, 15570-15571.

4. S. H. Ko, M. Su, C. Zhang, A. E. Ribbe, W. Jiang and C. Mao, *Nat. Chem.*, 2010, 2, 1050-1055.
5. X. Bai, T. G. Martin, S. H. W. Scheres and H. Dietz, *Proc. Natl. Acad. Sci. USA*, 2012, 109, 20012-20017.
6. Z.-S. Wu, Z. Shen, K. Tram and Y. Li, *Nat. Commun.*, 2014, 5.
7. E. S. Andersen, M. Dong, M. M. Nielsen, K. Jahn, R. Subramani, W. Mamdouh, M. M. Golas, B. Sander, H. Stark, C. L. P. Oliveira, J. S. Pedersen, V. Birkedal, F. Besenbacher, K. V. Gothelf and J. Kjems, *Nature*, 2009, 459, 73-76.
8. H. Pei, L. Liang, G. Yao, J. Li, Q. Huang and C. Fan, *Angew. Chem., Int. Ed.*, 2012, 51, 9020-9024.
9. S. A. Benner, *Acc. Chem. Res.*, 2004, 37, 784-797.
10. F. Wojciechowski and C. J. Leumann, *Chem. Soc. Rev.*, 2011, 40, 5669-5679.
11. I. Hirao, M. Kimoto and R. Yamashige, *Acc. Chem. Res.*, 2012, 45, 2055-2065.
12. Z. Yang, F. Chen, J. B. Alvarado and S. A. Benner, *J. Am. Chem. Soc.*, 2011, 133, 15105-15112.
13. A. A. Henry and F. E. Romesberg, *Curr. Opin. Chem. Biol.*, 2003, 7, 727-733.
14. T. Lavergne, M. Degardin, D. A. Malyshev, H. T. Quach, K. Dhami, P. Ordoukhanian and F. E. Romesberg, *J. Am. Chem. Soc.*, 2013, 135, 5408-5419.
15. G. H. Clever and M. Shionoya, *Coord. Chem. Rev.*, 2010, 254, 2391-2402.
16. Y. Takezawa and M. Shionoya, *Acc. Chem. Res.*, 2012, 45, 2066-2076.
17. P. Scharf and J. Müller, *ChemPlusChem*, 2013, 78, 20-34.
18. K. Tanaka, A. Tengeji, T. Kato, N. Toyama and M. Shionoya, *Science*, 2003, 299, 1212-1213.
19. G. H. Clever, S. J. Reitmeier, T. Carell and O. Schiemann, *Angew. Chem., Int. Ed.*, 2010, 49, 4927-4929.
20. S. Liu, G. H. Clever, Y. Takezawa, M. Kaneko, K. Tanaka, X. Guo and M. Shionoya, *Angew. Chem., Int. Ed.*, 2011, 50, 8886-8890.
21. R. Freeman, T. Finder and I. Willner, *Angew. Chem., Int. Ed.*, 2009, 48, 7818-7821.
22. T. Carell, *Nature*, 2011, 469, 45-46.
23. Y. Miyake, H. Togashi, M. Tashiro, H. Yamaguchi, S. Oda, M. Kudo, Y. Tanaka, Y. Kondo, R. Sawa, T. Fujimoto, T. Machinami and A. Ono, *J. Am. Chem. Soc.*, 2006, 128, 2172-2173.
24. S. Johannsen, S. Paulus, N. Düpre, J. Müller and R. K. O. Sigel, *J. Inorg. Biochem.*, 2008, 102, 1141-1151.
25. A. Ono, S. Cao, H. Togashi, M. Tashiro, T. Fujimoto, T. Machinami, S. Oda, Y. Miyake, I. Okamoto and Y. Tanaka, *Chem. Commun.*, 2008, 44, 4825-4827.
26. T. Funai, Y. Miyazaki, M. Aotani, E. Yamaguchi, O. Nakagawa, S.-i. Wada, H. Torigoe, A. Ono and H. Urata, *Angew. Chem., Int. Ed.*, 2012, 51, 6464-6466.
27. K. Tanaka, Y. Yamada and M. Shionoya, *J. Am. Chem. Soc.*, 2002, 124, 8802-8803.
28. D. Böhme, N. Düpre, D. A. Megger and J. Müller, *Inorg. Chem.*, 2007, 46, 10114-10119.
29. K. Tanaka and M. Shionoya, *J. Org. Chem.*, 1999, 64, 5002-5003.
30. H. Weizman and Y. Tor, *J. Am. Chem. Soc.*, 2001, 123, 3375-3376.
31. K. Tanaka, A. Tengeji, T. Kato, N. Toyama, M. Shiro and M. Shionoya, *J. Am. Chem. Soc.*, 2002, 124, 12494-12498.
32. L. Zhang and E. Meggers, *J. Am. Chem. Soc.*, 2004, 127, 74-75.
33. C. Switzer and D. Shin, *Chem. Commun.*, 2005, 41, 1342-1344.
34. C. Switzer, S. Sinha, P. H. Kim and B. D. Heuberger, *Angew. Chem., Int. Ed.*, 2005, 44, 1529-1532.
35. E. Meggers, P. L. Holland, W. B. Tolman, F. E. Romesberg and P. G. Schultz, *J. Am. Chem. Soc.*, 2000, 122, 10714-10715.
36. N. Zimmermann, E. Meggers and P. G. Schultz, *J. Am. Chem. Soc.*, 2002, 124, 13684-13685.
37. D. Shin and C. Switzer, *Chem. Commun.*, 2007, 43, 4401-4403.
38. B. D. Heuberger, D. Shin and C. Switzer, *Org. Lett.*, 2008, 10, 1091-1094.
39. D. A. Megger, C. Fonseca Guerra, J. Hoffmann, B. Brutschy, F. M. Bickelhaupt and J. Müller, *Chem. Eur. J.*, 2011, 17, 6533-6544.
40. H. Mei, I. Röhl and F. Seela, *J. Org. Chem.*, 2013, 78, 9457-9463.
41. G. H. Clever, K. Polborn and T. Carell, *Angew. Chem., Int. Ed.*, 2005, 44, 7204-7208.
42. G. H. Clever, Y. Sörtl, H. Burks, W. Spahl and T. Carell, *Chem. Eur. J.*, 2006, 12, 8708-8718.
43. K. Tanaka, G. H. Clever, Y. Takezawa, Y. Yamada, C. Kaul, M. Shionoya and T. Carell, *Nat. Nano*, 2006, 1, 190-194.
44. G. H. Clever and T. Carell, *Angew. Chem., Int. Ed.*, 2007, 46, 250-253.
45. C. Kaul, M. Müller, M. Wagner, S. Schneider and T. Carell, *Nat. Chem.*, 2011, 3, 794-800.
46. M. Su, M. Tomás-Gamasa, S. Serdjukow, P. Mayer and T. Carell, *Chem. Commun.*, 2014, 50, 409-411.
47. G. Roelfes and B. L. Feringa, *Angew. Chem., Int. Ed.*, 2005, 44, 3230-3232.
48. J. Wang, E. Benedetti, L. Bethge, S. Vohhoff, S. Klussmann, J.-J. Vasseur, J. Cossy, M. Smetana and S. Arseniyadis, *Angew. Chem., Int. Ed.*, 2013, 52, 11546-11549.

49. C. Wang, G. Jia, J. Zhou, Y. Li, Y. Liu, S. Lu and C. Li, *Angew. Chem., Int. Ed.*, 2012, 51, 9352-9355.