

Cite this: *Chem. Sci.*, 2018, 9, 4488

Received 1st March 2018

Accepted 21st April 2018

DOI: 10.1039/c8sc00986d

rsc.li/chemical-science

# Gold-catalyzed (4 + 2)-annulations between $\alpha$ -alkyl alkenylgold carbenes and benzisoxazoles with reactive alkyl groups†

Bhanudas Dattatray Mokar,‡ Prakash D. Jadhav,‡ Y. B. Pandit and Rai-Shung Liu \*


This work reports new (4 + 2)-annulations of  $\alpha$ -alkyl vinylgold carbenes with benzisoxazoles to afford 3,4-dihydroquinoline derivatives with high *anti*-stereoselectivity. The annulations are operable with carbenes in both acyclic and cyclic forms. This reaction sequence involves an initial formation of imines from  $\alpha$ -alkylgold carbenes and benzisoxazoles, followed by a novel carbonyl-enamine reaction to yield 3,4-dihydroquinoline derivatives. This system presents the first alkyl C–H reactivity of  $\alpha$ -alkyl gold carbenes with an external substrate.

## Introduction



Metal carbenes are versatile intermediates to implement a vast number of useful reactions including cyclopropanation, X–H insertion (X = C, N and O), skeletal rearrangement and annulation reactions (eqn (1)).<sup>1</sup> Despite their widespread applications, applicable metal carbenes, derived from diazo precursors, are mainly restricted to donor/acceptor (D/A) types I (R = H, aryl and alkenyl; EWG = CN, ketones and esters) whereas highly desirable  $\alpha$ -alkyl metal carbenes II are less efficient because of a competitive 1,2-hydrogen shift to form olefins (eqn (2)).<sup>1</sup> This side reaction is particularly serious for gold carbenes because their LAu = C<sup>+</sup> carbons are highly cationic.<sup>2</sup> Few intermolecular reactions involving Ar–Pd(II) catalysts focused on  $\alpha$ -alkyl metal carbenes of D/A types.<sup>3</sup> The limited utility of  $\alpha$ -alkyl carbenoids features an unsolved and challenging task in metal carbene chemistry. We seek new  $\alpha$ -alkyl carbenoids beyond commonly used D/A carbenes II, aiming at two objectives: (i) suppression of a 1,2-H shift and (ii) an alkyl C–H reaction with an external substrate.

Interest in the reactions of benzisoxazoles is rapidly growing in gold catalysis because of their various annulation modes with gold  $\pi$ -alkynes.<sup>4–6</sup> To explore the reactivity of benzisoxazoles toward gold carbenes,<sup>7</sup> we first tested the reactions with D/A-type benzyl  $\alpha$ -oxogold carbene II' (R = Ph and EWG = CO<sub>2</sub>Et), yielding an olefin product III' efficiently (eqn (3)). We envisage that D/D type carbenes such as  $\alpha$ -alkyl alkenylgold carbenes IV

Currently used carbenes: D/A types



Our initial tests:

This work:  $\alpha$ -alkyl alkenylgold carbenes (D/D types)

might be viable species to achieve new annulations with benzisoxazoles because their gold-stabilized allyl cation character IV is unfavorable for a 1,2-H shift. According to this hypothesis, this work reports novel intermolecular (4 + 2)-annulations between  $\alpha$ -alkyl vinylgold carbenes and benzisoxazoles, thus manifesting an unprecedented C–H reactivity of  $\alpha$ -alkyl metal carbenes.

## Results and discussion


As shown in eqn (5), we further tested the reaction of acyclic  $\alpha$ -alkylgold carbenes A that were generated *in situ* from cyclopropene derivatives 1a–1b and gold catalysts.<sup>8</sup> With IPtAuCl/AgSbF<sub>6</sub>, quinoline derivatives 3a and 3b were isolated in satisfactory yields (72–75%), together with enones 1a-O and 1b-O in minor proportions (17–19%). A 1,2-hydrogen shift was

Frontier Research Centers for Materials Science and Technology, Department of Chemistry, National Tsing-Hua University, Hsinchu, Taiwan, Republic of China. E-mail: rslu@mx.nthu.edu.tw

† Electronic supplementary information (ESI) available. CCDC 1819135–1819138. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc00986d

‡ B. D. M. and P. D. J. contributed equally.

effectively suppressed with vinylcarbenes **A**, supporting our hypothetical role of gold-stabilized allyl cations **A**.



Our primary interest is to construct complicated frameworks *via* cascade reactions. Fig. 1 depicts several bioactive compounds (**VI-1**)–(**VI-6**) bearing a common tricyclic framework **VI**, which can be easily constructed from cyclopentenylgold carbene **A'** and benzisoxazole **2a** in dichloromethane (DCM) using various gold catalysts; species **4a** serves as a precursor for cyclopentenylgold carbene **A'** (Table 1).<sup>10</sup>

In this new task, we optimized the annulation cascades between vinylallene **4a** and benzisoxazole **2a** in dichloromethane (DCM) using various gold catalysts; species **4a** serves as a precursor for cyclopentenylgold carbene **A'** (Table 1).<sup>10</sup>

An initial test of IPrAuCl/AgSbF<sub>6</sub> at a 5 mol% loading afforded a new azacyclic product **5a** and cyclopentadiene **4a'** in 62% and 25% yields, respectively (entry 1); the latter was derived from a 1,2-H shift of gold carbenes **A'** that was generated from cyclizations of gold-stabilized pentadienyl cation **A-I**. Notably, an increased gold loading (10 mol%) enhanced the yield of desired **5a** up to 85%. Other gold catalysts L AuCl/AgSbF<sub>6</sub> (L = P(OPh)<sub>3</sub>, PPh<sub>3</sub> and P(*t*-Bu)<sub>2</sub>(*o*-biphenyl)) gave **5a** in 40–82% yields with L = P(OPh)<sub>3</sub> being the most effective (entries 3–5). For various silver salts as in IPrAuCl/AgX (X = OTf and NTf<sub>2</sub>), resulting **5a** was obtained in 65% and 71% yields, respectively (entries 6–7). AgNTf<sub>2</sub> was entirely inactive (entry 8). IPrAuCl/AgSbF<sub>6</sub> in various solvents gave **5a** in the following yields: DCE 70%, MeCN 20% and 1,4-dioxane 0 (entries 9–11). The molecular structure of compound **5a** was characterized with X-ray diffraction,<sup>11</sup> showing an *anti*-configuration between the alcohol and phenyl groups.





Fig. 1 Suitable alkylgold carbenes to access bioactive molecules.

Table 1 Catalytic reactions with various gold catalysts




Entry	Catalyst [mol%]	Solvent	<i>t</i> [h]	Yield <sup>b</sup> [%]		
				4a	5a	4a'
1	IPrAuCl/AgSbF <sub>6</sub> (5)	DCM	12	8	62	25
2	IPrAuCl/AgSbF <sub>6</sub> (10)	DCM	3	—	85	12
3	(PhO) <sub>3</sub> PAuCl/AgSbF <sub>6</sub> (10)	DCM	3	—	82	16
4	Ph <sub>3</sub> PAuCl/AgSbF <sub>6</sub> (10)	DCM	4	—	55	36
5	LAuCl/AgSbF <sub>6</sub> (10) <sup>c</sup>	DCM	3	—	40	52
6	IPrAuCl/AgOTf (10)	DCM	4	—	65	26
7	IPrAuCl/AgNTf <sub>2</sub> (10)	DCM	4	—	71	20
8	AgSbF <sub>6</sub> (10)	DCM	24	95	—	—
9	IPrAuCl/AgSbF <sub>6</sub> (10)	DCE	5	—	70	24
10	IPrAuCl/AgSbF <sub>6</sub> (10)	MeCN	12	—	20	65
11	IPrAuCl/AgSbF <sub>6</sub> (10)	Dioxane	10	—	—	90

<sup>a</sup> [4a] = 0.05 M. <sup>b</sup> Product yields are reported after purification from a silica column. <sup>c</sup> L = P(*t*-Bu)<sub>2</sub>(*o*-biphenyl). IPr = 1,3-bis(diisopropylphenyl)imidazole-2-ylidene, DCE = 1,2-dichloroethane.

Table 2 assesses the generality of these gold-catalyzed reactions using various vinylallenes **4b–4t** catalyzed with IPrAuCl/AgSbF<sub>6</sub> (10 mol%) in DCM. All resulting products **5b–5t** assumed *anti*-configurations with the alcohol and R<sup>1</sup> groups being mutually *trans*. We tested the reaction of trisubstituted

Table 2 Catalytic annulations with various alkenylallenes



<sup>a</sup> [4] = 0.05 M. <sup>b</sup> Product yields are reported after purification from a silica column.



vinylallenes **4b–4f** bearing  $R^1 = 4\text{-MePh}$ ,  $4\text{-OMePh}$ ,  $4\text{-ClPh}$ ,  $4\text{-CF}_3\text{Ph}$  and  $n\text{-Bu}$ , yielding desired **5b–5f** in 78–88% yields (entries 1–5). For species **4g** and **4h** bearing 3-phenyl substituents ( $X = \text{OMe}$  and  $\text{Cl}$ ), their corresponding products **5g** and **5h** were obtained in 84% and 87% yields, respectively (entries 6 and 7). The reactions were extensible to other vinylallenes **4i–4k** bearing 2-naphthyl, 2-furan and 2-thiophene, further delivering desired products **5i–5k** in 82–84% yields (entries 8–10). We tested the reaction on vinylallene **4l** bearing distinct  $R^1 = \text{Me}$  and  $R^2 = \text{Ph}$ , which yielded compound **5l** with an anti-configuration in which the hydroxy and methyl groups are mutually *trans* (entry 11); this configuration was established by the  $^1\text{H}$  NOE effect. Additional alkyl-substituted vinylallenes **4m–4p** yielded desired **5m–5p** in satisfactory yields (80–85%, entries 12–15). Variations of the  $R^2$  group with an *n*-butyl group as in species **4q** gave expected product **5q** in 86% yield (entry 16). We prepared species **4r** bearing varied  $R^2 = \text{Ph}$  and  $R^3 = n\text{-butyl}$ , producing compound **5r** in 80% yield (entry 17). For 1,3-disubstituted vinylallenes **4s** and **4t** ( $R^3 = \text{H}$ ), their resulting compounds **5s** and **5t** were obtained in 82–83% yields (entries 18 and 19).

We tested these new annulations on distinct substrates such as enynyl acetates **6a–6g**, bearing varied phenyl ( $R = 4\text{-XC}_6\text{H}_4$ ,  $X = \text{H}$ ,  $\text{Cl}$ ,  $\text{Br}$ ,  $\text{Me}$ , and  $\text{OMe}$ ), 2-thienyl and isopropyl substituents; these enyne acetates can be catalyzed with the same gold catalyst to yield distinct  $\alpha$ -alkylgold carbenes **A'** (see Table 3).<sup>12</sup> To our pleasure, new alkylgold carbenes **A'**, generated from these enynyl acetates, were trapped efficiently with benzisoxazole **2a** to afford the desired (4 + 2)-annulation products **7a–7g** in satisfactory yields (61–74%), further manifesting the reaction generality (entries 1–7). For unsubstituted propargyl acetate **6h** ( $R = \text{H}$ ), its reaction led to a 68% recovery of initial **6h** (entry 8). Even if the reaction is successful, a dehydration of compound **7h** would occur to give quinoline products. The molecular structure of compound **7a** ( $R = \text{Ph}$ ) was confirmed with

Table 4 Catalytic annulations with various benzisoxazoles

(1) <b>8b</b> ( $R^1 = \text{Me}$ , 1.5 h, 84%) <sup>a,b</sup> (2) <b>8c</b> ( $R^1 = \text{OMe}$ , 1.5 h, 84%) (3) <b>8d</b> ( $R^1 = \text{Br}$ , 2.5 h, 90%) (4) <b>8e</b> ( $R^1 = \text{Cl}$ , 2.5 h, 85%) (5) <b>8f</b> ( $R^1 = \text{OCO}_2\text{Et}$ , 3 h, 80%)	(6) <b>8g</b> ( $R^2 = \text{Me}$ , 4 h, 88%) (7) <b>8h</b> ( $R^2 = \text{Br}$ , 3 h, 86%) (8) <b>8i</b> ( $R^2 = \text{Cl}$ , 3 h, 92%)	(9) <b>8j</b> (2.5 h, 82%)

<sup>a</sup> **4a** = 0.05. <sup>b</sup> Product yields are reported after purification from a silica column.

X-ray diffraction analysis that revealed an *anti*-configuration (Table 3).<sup>11</sup>


The scope of these catalytic reactions is further expanded with various applicable benzisoxazoles **2b–2j** substituted with the C(3), C(5) and C(6) carbons. Other C(5)-substituted benzisoxazoles **2b–2f** ( $R^1 = \text{Me}$ ,  $\text{OMe}$ ,  $\text{Br}$ ,  $\text{Cl}$ , and  $-\text{OCO}_2\text{Et}$ ) maintained high efficiencies to deliver *anti*-configured products **8b–8f** in 80–90% yields (entries 1–5). High reaction efficiencies were maintained also for C(6)-substituted benzisoxazoles **2g–2i** that furnished products **8g–8i** in 86–92% yields (entries 6–8). A final applicable reaction with a C(3)-substituted benzisoxazole **2j** enabled the production of a tertiary alcohol **8j**, reflecting the reaction feasibility (entry 9).  $^1\text{H}$  NOE spectra were recorded to verify the stereochemistry of compound **8j** (Table 4).

Gold-catalyzed reactions of 3,5-dimethylisoxazole **2a'** with vinylallenes **4a** and **4u** delivered 2-aminocyclopentadienes **9a** and **9b** in 72% and 64% yields, respectively (eqn (6)).<sup>5a,13,14</sup> The molecular structure of compound **9b** was characterized with X-ray diffraction.<sup>11</sup> Cyclizations of compounds **9a** and **9b** with a gold catalyst were unsuccessful because of the two different forms of the enol imines (eqn (6)). To rationalize the origin of the stereoselectivity, compound **5a** was treated with  $\text{Zn}(\text{OTf})_2$  (20 mol%) in refluxing DCE to examine the hydroxyl epimerization that turned out to be slow. An equilibrium, *anti*/*syn* = 4 : 1, was attained for species **5a** after reflux in DCE for 48 h (eqn (7)).

Table 3 Annulation reactions with enynyl acetates


(1) <b>7a</b> (48 h, 74%, X-ray) <sup>a,b</sup>	(2) <b>7b</b> ( $X = \text{Cl}$ , 48 h, 67%) (3) <b>7c</b> ( $X = \text{Br}$ , 48 h, 66%)	(4) <b>7d</b> ( $X = \text{Me}$ , 46 h, 71%) (5) <b>7e</b> ( $X = \text{OMe}$ , 45 h, 73%)
(6) <b>7f</b> (50 h, 74%)	(7) <b>7g</b> (47 h, 61%)	(8) <b>7h</b> (48 h, 0%) <sup>c</sup>

<sup>a</sup> **6** = 0.05 M. <sup>b</sup> Product yields are reported after purification from a silica column. <sup>c</sup> A 68% recovery of initial **6h** is found in entry 8.




Scheme 1 shows the stereoselective functionalizations of *anti*-**5a** via  $\text{NaBH}_4$  reductions and *m*-CPBA oxidations,





Scheme 1 Chemical functionalizations.



Scheme 2 A plausible reaction mechanism.

respectively yielding compounds **5a-H** and **5a-O** as single diastereomeric products. The stereochemistries of compounds **5a-H** and **5a-O** were established with  $^1\text{H}$  NOE spectra. Likewise, the acetate species **7a** was readily removed under basic conditions, yielding the enol form **7a'** as shown by its NMR in  $\text{CD}_3\text{COCD}_3$  and  $\text{CDCl}_3$ . We also studied an  $\text{O}_3$ -induced oxidative cleavage of the acetate derivative **5a-OAc** to cleave the olefin group, yielding the peroxide **5a-O<sub>3</sub>** in 85% yield. The molecular structure of species **5a-O<sub>3</sub>** has been characterized by X-ray diffraction.<sup>11</sup>

As depicted in Scheme 2, we postulate an initial formation of imines between alkylgold carbene **A** and benzisoxazole, yielding 2-iminoyl benzaldehyde **C**. This hypothesis is supported by our observation of 3,5-dimethylisoxazole, depicted in eqn (6). A tautomerization of imine species **C** is expected to form enamines **D** bearing an  $\text{NH}\cdots\text{O}=\text{C}$  hydrogen bond. We believe that this enamine form, unlike other enamine-carbonyl couplings,<sup>15</sup> is stabilized with the  $\text{NH}\cdots\text{O}=\text{C}$  bond to enable a concerted process, analogous to the well-known carbonyl-ene reactions. A boat-like conformation **D** is favorable to yield *anti*-5 stereoselectively.

## Conclusions

This work reports novel gold-catalyzed (4 + 2)-annulations between alkylgold carbenes and benzisoxazoles **2** to form 3,4-dihydroquinoline derivatives. Gold carbenes in cyclic and acyclic forms are both applicable. In this reaction sequence, the gold complex catalyzes an initial formation of imines between alkylgold carbenes<sup>13,14</sup> and benzisoxazoles; the resulting intermediates bear an enamine moiety that is bound to an aldehyde *via* a hydrogen bond to induce a carbonyl-enamine reaction. Control experiments with 3,5-dimethylisoxazoles supported this postulated mechanism. This new synthetic design involving  $\alpha$ -

alkyl metal carbenes of **D/D** types will attract growing interest because of its distinct utility.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements


We thank the Ministry of Education (MOE 106N506CE1) and Ministry of Science and Technology (MOST 107-3017-F-007-002), Taiwan, for financial support of this work.

## Notes and references

- (a) H. M. L. Davies and E. G. Antoulinakis, *Intermolecular Metal-Catalyzed Carbenoid Cyclopropanations in Organic Reactions*, ed. L. E. Overman, John Wiley & Sons, Inc., New York, NY, 2001, vol. 57, pp. 1–326; (b) H. M. L. Davies and R. E. J. Beckwith, *Chem. Rev.*, 2003, **103**, 2861; (c) M. P. Doyle, R. Duffy, M. Ratnikov and L. Zhou, *Chem. Rev.*, 2010, **110**, 704; (d) Q.-Q. Cheng, Y. Yu, J. Yedoyan and M. P. Doyle, *ChemCatChem*, 2018, **10**, 488.
- For selected reviews for gold carbenes, see: (a) D. Qian and J. Zhang, *Chem. Soc. Rev.*, 2015, **44**, 677; (b) L. Liu and J. Zhang, *Chem. Soc. Rev.*, 2016, **45**, 506; (c) E. López, S. Gonzalez-Pelayo and L. A. López, *Chem. Rec.*, 2017, **17**, 312; (d) C. Obradors and A. M. Echavarren, *Chem. Commun.*, 2014, **50**, 16; (e) L. Zhang, *Acc. Chem. Res.*, 2014, **47**, 877; (f) D. P. Day and P. W. H. Chan, *Adv. Synth. Catal.*, 2016, **358**, 1368.
- The reactions were only reported for  $\text{Ar-Pd(II)}$  species; see: (a) C. Peng, Y. Wang and J. Wang, *J. Am. Chem. Soc.*, 2008, **130**, 1566; (b) Z. Zhang, Y. Liu, M. Gong, X. Zhao, Y. Zhang and J. Wang, *Angew. Chem., Int. Ed.*, 2010, **49**, 1139.
- (a) L. Li, T.-D. Tan, Y.-D. Tan, Y.-Q. Zhang, X. Liu and L.-W. Ye, *Org. Biomol. Chem.*, 2017, **15**, 8483; (b) D. B. Huple, S. Ghorpade and R.-S. Liu, *Adv. Synth. Catal.*, 2016, **358**, 1348; (c) S. S. Giri and R.-S. Liu, *Chem. Sci.*, 2018, **9**, 2991.
- (a) A.-H. Zhou, Q. He, C. Shu, Y.-F. Yu, S. Liu, T. Zhao, W. Zhang, X. Lu and L.-W. Ye, *Chem. Sci.*, 2015, **6**, 1265; (b) X.-Y. Xiao, A.-H. Zhou, C. Shu, F. Pan, T. Li and L.-W. Ye, *Chem.-Asian J.*, 2015, **10**, 1854; (c) H. Jin, L. Huang, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 794; (d) H. Jin, B. Tian, X. Song, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2016, **55**, 12688.
- (a) Z. Zeng, H. Jin, J. Xie, B. Tian, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Org. Lett.*, 2017, **19**, 1020; (b) M. Chen, N. Sun, H. Chen and Y. Liu, *Chem. Commun.*, 2016, **52**, 6324; (c) W. Xu, G. Wang, N. Sun and Y. Liu, *Org. Lett.*, 2017, **19**, 3307.
- The reactions of isoxazoles with rhodium carbenes were noted by Davies' group; distinct [3 + 3]-annulations were reported for  $\alpha$ -alkenylrhodium esters as shown below. (a) J. R. Manning and H. M. L. Davies, *Tetrahedron*, 2008, **64**,



6901; (b) J. R. Manning and H. M. L. Davies, *J. Am. Chem. Soc.*, 2008, **130**, 8602.



- 8 (a) A. Archambeau, F. Miege, J. Crossy and C. Meyer, in *Patai's Chemistry of Functional Groups*, ed. Z. Rappoport, J. F. Liebman and I. Marek, John Wiley & Sons Ltd., Hoboken, NJ, 2014, pp. 631–700; (b) Y. Deng and M. P. Doyle, *Isr. J. Chem.*, 2016, **56**, 399–408; (c) F. Miege, C. Meyer and J. Crossy, *Org. Lett.*, 2010, **12**, 4144; (d) F. Miege, C. Meyer and J. Crossy, *Chem.-Eur. J.*, 2012, **18**, 7810; (e) C. Li, Y. Zeng and J. Wang, *Tetrahedron Lett.*, 2009, **50**, 2956; (f) Z.-B. Zhu and M. Shi, *Chem.-Eur. J.*, 2008, **14**, 10219; (g) S. B. Wagh, Y.-C. Hsu and R.-S. Liu, *ACS Catal.*, 2016, **6**, 7160; (h) Z. Liu, Q. Li, P. Liao and X. Bi, *Chem.-Eur. J.*, 2017, **23**, 4756; (i) W. Rao, M. J. Koh, D. Li, H. Hirao and P. W. H. Chan, *J. Am. Chem. Soc.*, 2013, **135**, 7926; (j) W. Rao and P. W. H. Chan, *Chem.-Eur. J.*, 2014, **20**, 713; (k) J. Yan, G. L. Tay, C. Neo, B. R. Lee and P. W. H. Chan, *Org. Lett.*, 2015, **17**, 4176.
- 9 (a) S. Chakrabarty, M. S. Croft, M. G. Marko and G. Moyna, *Bioorg. Med. Chem.*, 2013, **21**, 1143; (b) Merck Sharp and Dohme Corp, WO 2012/174176 A1, 2012; (c) P. S. Ramamoorthy and R. E. McDevitt, *US Pat.*, US 2004/0019040, 2004; (d) J. Wöfling, É. Frank, G. Schneider, M. T. Bes and L. F. Tietze, *Synlett*, 1998, 1205; (e) L. F. Tietze and A. Modi, *Med. Res. Rev.*, 2000, **20**, 304; (f) R. M. Kariba, P. J. Houghton and A. Yenesew, *J. Nat. Prod.*, 2002, **65**, 566; (g) J. L. Hubbs and C. H. Heathcock, *Org. Lett.*, 1999, **1**, 1315.
- 10 (a) S. Bhunia and R.-S. Liu, *J. Am. Chem. Soc.*, 2008, **130**, 16488; (b) J. H. Lee and F. D. Toste, *Angew. Chem., Int. Ed.*, 2007, **46**, 912; (c) H. Funami, H. Kasama and N. Iwasawa, *Angew. Chem., Int. Ed.*, 2007, **46**, 909; (d) G. Lemiere, V. Gandon, K. Cariou, T. Fukuyama, A. L. Dhiman, L. Fensterbank and M. Malacria, *Org. Lett.*, 2007, **9**, 2207; (e) V. Gandon, G. Lemiere, A. Hours, L. Fensterbank and M. Malacria, *Angew. Chem., Int. Ed.*, 2008, **47**, 7534; (f) M. R. Fructos, M. Besora, A. A. C. Braga, M. M. Díaz-Requejo, F. Maseras and P. J. Pérez, *Organometallics*, 2017, **36**, 172; (g) F.-Q. Shi, X. Li, Y. Xia, L. Zhang and Z.-X. Yu, *J. Am. Chem. Soc.*, 2007, **129**, 15503; (h) W. Rao, D. Susanti, B. J. Ayers and P. W. H. Chan, *J. Am. Chem. Soc.*, 2015, **137**, 6350; (i) W. Rao, J. W. Boyle and P. W. H. Chan, *Chem.-Eur. J.*, 2016, **22**, 6532.
- 11 Crystallographic data of compounds **5a**, **5a-O<sub>3</sub>**, **7a**, and **9b** were deposited at the Cambridge Crystallographic Data Center: **5a** (CCDC 1819135); **5a-O<sub>3</sub>** (CCDC 1819137); **7a** (CCDC 1819138); and **9b** (CCDC 1819136).†
- 12 L. Zhang and S. Wang, *J. Am. Chem. Soc.*, 2006, **128**, 1442.
- 13 For gold-catalyzed nitrene reactions of alkynes; see selected examples: (a) D. J. Gorin, N. R. Davies and F. D. Toste, *J. Am. Chem. Soc.*, 2005, **127**, 11260; (b) A. Wetzel and F. Gagosz, *Angew. Chem., Int. Ed.*, 2011, **50**, 7354; (c) B. Lu, Y. Luo, L. Liu, L. Ye, Y. Wang and L. Zhang, *Angew. Chem., Int. Ed.*, 2011, **50**, 8358; (d) C. Shu, Y.-H. Wang, B. Zhou, X.-L. Li, Y.-L. Li, Y.-F. Ping, X. Lu and L.-W. Ye, *J. Am. Chem. Soc.*, 2015, **137**, 9567; (e) H.-H. Hung, Y.-C. Liao and R.-S. Liu, *J. Org. Chem.*, 2013, **78**, 7970.
- 14 Benzisoxazoles serve as nitrene sources in rhodium-catalyzed C–H functionalizations, see selected examples: (a) S. Yu, G. Tang, Y. Li, X. Zhou, Y. Lan and X. Li, *Angew. Chem., Int. Ed.*, 2016, **55**, 8696; (b) M. Zou, J. Liu, C. Tang and N. Jiao, *Org. Lett.*, 2016, **18**, 3030; (c) S. Yu, Y. Li, X. Zhou, H. Wang, L. Kong and X. Li, *Org. Lett.*, 2016, **18**, 2812.
- 15 The reactions of enamines and aldehydes are generally implemented with Lewis acids or a base, see: (a) R. Matsubara, N. Kawai and S. Kobayashi, *Angew. Chem., Int. Ed.*, 2006, **45**, 3814; (b) T. Kochi, T. P. Tang and J. A. Ellman, *J. Am. Chem. Soc.*, 2003, **125**, 11276; (c) R. Matsubara, P. Vital, Y. Nakamura, H. Kiyohara and S. Kobayashi, *Tetrahedron*, 2004, **60**, 9769.

