

Cite this: *Chem. Sci.*, 2023, 14, 7559

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Synthesis of oligosaccharides to identify an immunologically active epitope against *Candida auris* infection†

Rajat Kumar Singh,<sup>ab</sup> Emelie E. Reuber,<sup>ab</sup> Mariolina Bruno,<sup>c</sup> Mihai G. Netea<sup>cd</sup> and Peter H. Seeberger<sup>ib</sup>\*<sup>ab</sup>

*Candida auris* (*C. auris*) is an emerging multidrug-resistant fungal pathogen that represents a significant public health challenge as it can spread rapidly and result in high mortality rates. The mannans on the *C. auris* cell surface are potent immunogens and attractive targets for developing a glycoconjugate vaccine. We synthesized the oligosaccharides resembling cell surface mannans of *C. auris* and printed them onto microarray slides that were used to screen plasma from mice infected with *C. auris*. IgM antibodies in mouse plasma recognize the  $\beta$ -1,2 linkage present in *C. auris* surface mannans. Disaccharide **19** emerged from glycan array screening as a lead for developing a vaccine against *C. auris*, as the majority of patient plasma samples showed antibodies against this glycan. The synthetic oligosaccharides can be used for the early detection of *C. auris* infections.

Received 7th March 2023  
Accepted 16th June 2023

DOI: 10.1039/d3sc01242e

rsc.li/chemical-science

## Introduction

*Candida auris* (*C. auris*) is a species of yeast that was first reported<sup>1</sup> in 2009 and is now on the World Health Organization (WHO) list as a “critical priority group” pathogen.<sup>2</sup> The fungus causes multidrug-resistant and potentially life-threatening infections that can result in sepsis, meningitis, and other serious conditions if left untreated. *C. auris* infections spread most commonly in healthcare settings from person to person through contact with contaminated surfaces or medical equipment.<sup>3</sup> *C. auris* infections are typically diagnosed by a combination of laboratory testing and medical history.<sup>4</sup> Treatment of *C. auris* infections is often difficult as resistance to anti-fungal drugs such as fluconazole, amphotericin, voriconazole and caspofungin is emerging.<sup>5</sup> Anti-fungal vaccines are urgently needed.

The outer membrane of the fungus contains a variety of proteins, including outer membrane proteins, lipoproteins, and glycoproteins. The polysaccharide-rich cell membrane is composed of chitin and  $\beta$ -glucans in the inner layer, and mannoprotein in the outer layer of cell wall.<sup>6</sup> *C. auris*



Fig. 1 (A) Structure of *C. auris* mannans; (B) structure of synthetic mannans resembling the *C. auris* cell surface.

<sup>a</sup>Department of Biomolecular System, Max Planck Institute of Colloids and Interfaces, 14476, Potsdam, Germany. E-mail: peter.seeberger@mpikg.mpg.de

<sup>b</sup>Institute of Chemistry and Biochemistry, Freie Universität Berlin, 14195, Berlin, Germany

<sup>c</sup>Department of Internal Medicine, Radboud Institute for Molecular Life Sciences, Radboud University Medical Center, Nijmegen, The Netherlands

<sup>d</sup>Department of Immunology and Metabolism, Life & Medical Sciences Institute (LIMES), University of Bonn, Bonn, Germany

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3sc01242e>






Fig. 2 Glycan microarray analysis of *Candida auris* mannans. (A) Printing pattern of glycan microarray. (B) Exemplary binding pattern of mice plasma to immobilized synthetic glycans. (C) Mean fluorescence intensity of IgM antibody binding to synthetic glycans. A plasma dilution of 1:50 was used. Values represent mean  $\pm$  SEM. Differences were tested for significance using one-way ANOVA followed by Tukey's post hoc test with (\*\*\*\*)  $p < 0.0001$ .

mannoproteins are composed of *O*- and *N*-linked mannans with  $\alpha$ -1,2-,  $\alpha$ -1,3-,  $\alpha$ -1,6- and  $\beta$ -1,2-linked mannosyl units as well as phosphodiester-linked mannosyl residues (Fig. 1).<sup>6–8</sup> The cell membrane is important for protecting the fungus from environmental stress, providing a barrier to the entry of antimicrobial agents, and aiding in the adherence to surfaces. The cell membrane polysaccharides are recognized by pattern recognition receptors (PRRs) on the surface of the immune cell and in some cases may contribute to the pathogenicity of *C. auris* by allowing it to evade the host immune system.<sup>9</sup>

Isolated mixtures of *Candida* mannans and mannoproteins were explored as potential vaccine antigens in animal models of candidiasis two decades ago without follow-up.<sup>10,11</sup> Rather than using isolated mixtures of glycans, defined, synthetic antigens resembling cell surface mannans may constitute attractive vaccine candidates. As a first step in the vaccine design and development process, we report the synthesis of a series of conjugate-ready glycans related to the *C. auris* mannans (Fig. 1) that were used to screen plasma of infected mice for antibodies on glycan microarrays (Fig. 2) to identify a minimal epitope.<sup>12</sup>

## Results and discussion

Glycan assembly is based on orthogonally protected building blocks. *H*-phosphonate building block 13 (Scheme 1), was



Scheme 1 Synthesis of *H*-phosphonate building block 13.

prepared by the formation of a Nap ether at the C-2 hydroxyl of commercially available thioglycoside 5 to form 7 in 96% yield.<sup>13</sup> The benzylidene acetal in 7 was hydrolysed using ethanethiol and camphorsulfonic acid (CSA) to give dihydroxyl mannose thioglycoside 8. Dibenylation of 8 afforded fully protected thioglycoside 9 in 91% yield. Oxidative removal of naphthyl ether using DDQ afforded thioglycoside 10 in 76% yield. Levulinoylation of the C2-OH in 10 using 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC·HCl) and DMAP produced 11 in 86% yield.<sup>15,16</sup> Hydrolysis of thioglycoside 11 using *N*-bromosuccinimide (NBS) and subsequent treatment of hemiacetal<sup>16</sup> 11 with diphenyl phosphite provided *H*-phosphonate 13.<sup>17</sup>

Union of *D*-thiomannoside<sup>18</sup> 15 and aminopentyl linker<sup>19</sup> 14 promoted by NIS/TfOH afforded desired  $\alpha$ -linked glycoside 16 in 85% yield (Scheme 2). The linker is placed in anticipation of the conjugation to a carrier protein or a microarray surface. Benzoyl ester cleavage using sodium methoxide gave differentially protected monosaccharide 17.<sup>20</sup> The synthesis of  $\beta$ -(1 $\rightarrow$ 2)-linked disaccharide mannan<sup>21</sup> 19 began with the activation of thioglycoside 6 by addition of 1-benzene-sulfonyl piperidine (BSP), 2,4,6-tri-*tert*-butylpyrimidine (TTBP) and triflic anhydride (Tf<sub>2</sub>O) at  $-60$  °C followed by addition of acceptor 17 at  $-70$  °C to get exclusively fully protected  $\beta$ -(1 $\rightarrow$ 2)-linked mannan 18. The newly formed  $\beta$ -glycosidic bond was confirmed by coupled HSQC NMR. The global deprotection<sup>23</sup> of protected mannan 18 using Pd/C provided  $\beta$ -(1 $\rightarrow$ 2)-linked disaccharide 19 with 70% yield. Glycosylation of monosaccharide 17 with building block 15 using NIS/TfOH followed by cleavage of the benzoyl ester using sodium methoxide provided disaccharide 21. Hydrogenation of disaccharide 21 using Pd/C yielded 64% of deprotected disaccharide 24. For the synthesis of trisaccharide 23, disaccharide acceptor 21 was glycosylated with building block 6 using BSP, TTBP, and triflic anhydride to afford  $\beta$ -(1 $\rightarrow$ 2)-linked trisaccharide 22 in 84% yield. Hydrogenation of trisaccharide 22 furnished deprotected trisaccharide 23 in 70% yield. Synthesis of tetrasaccharide 28 began with the union of disaccharide 21 with donor 15 using







Scheme 3 Synthesis of oligosaccharides having phosphodiester linkage.

## Conclusions

We synthesized a library of oligosaccharides resembling the cell surface mannans of *C. auris*. The challenging phosphodiester linkage was synthesized using *H*-phosphonate building block 13. Glycan array analysis of plasma from mice infected with *C. auris* for three days were screened for antibodies to the synthetic glycans. Disaccharide 19 is recognized by IgM and appears to be a promising vaccine lead for further development of a glyco-conjugate vaccine candidate.

## Data availability

Data supporting this article have been uploaded as ESI.†

## Author contributions

P. H. S. conceived and supervised this project. R. K. S. synthesized all the target compounds. E. E. R. performed glycan microarray experiments. M. B. and M. G. N. provided plasma for

glycan microarray experiments. R. K. S. and E. E. R. wrote the original draft of the manuscript which was edited by all authors.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the Max-Planck Society for generous financial support. Open Access funding provided by the Max Planck Society.

## Notes and references

- 1 K. Satoh, K. Makimura, Y. Hasumi, Y. Nishiyama, K. Uchida and H. Yamaguchi, *Microbiol. Immunol.*, 2009, **53**, 41–44.
- 2 P. Beyer, V. Gigante and H. Getahun, *WHO fungal priority pathogens list to guide research, development and public health action*, World Health Organization, Geneva, p. 2022.



- 3 G. Tronchin, M. Pihet, L. M. Lopes-Bezerra and J.-P. Bouchara, *Med. Mycol.*, 2008, **46**, 749–772.
- 4 C. Keighley, K. Garnham, S. A. J. Harch, M. Robertson, K. Chaw, J. C. Teng and S. C. A. Chen, *Curr. Fungal Infect. Rep.*, 2021, **15**, 116–126.
- 5 M. G. Frías-De-León, R. Hernández-Castro, T. Vite-Garín, R. Arenas, A. Bonifaz, L. Castañón-Olivares, G. Acosta-Altamirano and E. Martínez-Herrera, *Antibiotics*, 2020, **9**, 568.
- 6 L. Yan, K. Xia, Y. Yu, A. Miliakos, S. Chaturvedi, F. Zhang, S. Chen, V. Chaturvedi and R. J. Linhardt, *ACS Infect. Dis.*, 2020, **6**, 1018–1031.
- 7 M. Bruno, S. Kersten, J. M. Bain, M. Jaeger, D. Rosati, M. D. Kruppa, D. W. Lowman, P. J. Rice, B. Graves, Z. Ma, Y. N. Jiao, A. Chowdhary, G. Renieris, F. L. van de Veerdonk, B.-J. Kullberg, E. J. Giamarellos-Bourboulis, A. Hoischen, N. A. R. Gow, A. J. P. Brown, J. F. Meis, D. L. Williams and M. G. Netea, *Nat. Microbiol.*, 2020, **5**, 1516–1531.
- 8 N. Shibata, H. Kobayashi and S. Suzuki, *Proc. Jpn. Acad., Ser. B*, 2012, **88**, 250–265.
- 9 Y. Wang, Y. Zou, X. Chen, H. Li, Z. Yin, B. Zhang, Y. Xu, Y. Zhang, R. Zhang, X. Huang, W. Yang, C. Xu, T. Jiang, Q. Tang, Z. Zhou, Y. Ji, Y. Liu, L. Hu, J. Zhou, Y. Zhou, J. Zhao, N. Liu, G. Huang, H. Chang, W. Fang, C. Chen and D. Zhou, *Nat. Commun.*, 2022, **13**, 3553.
- 10 B. Sendid, J. L. Poirot, M. Tabouret, A. Bonnin, D. Caillot, D. Camus and D. Poulain, *J. Med. Microbiol.*, 2002, **51**, 433–442.
- 11 M. H. Weiner and W. J. Yount, *J. Clin. Invest.*, 1976, **58**, 1045–1053.
- 12 P. H. Seeberger, *Chem. Rev.*, 2021, **121**, 3598–3626.
- 13 M. P. Lisboa, N. Khan, C. Martin, F.-F. Xu, K. Reppe, A. Geissner, S. Govindan, M. Witzentrath, C. L. Pereira and P. H. Seeberger, *Proc. Natl. Acad. Sci.*, 2017, **114**, 11063–11068.
- 14 M. Rajesh, E. Mathias, B. Afshan, F. Timmy, K. Peter, N. Sofie and H. Per, *Biochim. Biophys. Acta, Proteins Proteomics*, 2019, **1867**, 909–921.
- 15 R. K. Singh, J. Sianturi and P. H. Seeberger, *Org. Lett.*, 2022, **24**, 2371–2375.
- 16 J. Wu and Z. Guo, *J. Org. Chem.*, 2006, **71**, 7067–7070.
- 17 S. R. Sanapala, B. M. S. Seco, J. Y. Baek, S. I. Awan, C. L. Pereira and P. H. Seeberger, *Chem. Sci.*, 2020, **11**, 7401–7407.
- 18 P. Karen, *Carbohydr. Res.*, 2003, **338**, 1441–1454.
- 19 R. Castelli, H. S. Overkleeft, G. A. van der Marel and J. D. C. Codée, *Org. Lett.*, 2013, **15**, 2270–2273.
- 20 C. E. Martin, M. W. Weishaupt and P. H. Seeberger, *Chem. Commun.*, 2011, **47**, 10260–10262.
- 21 D. Crich and S. Sun, *J. Org. Chem.*, 1997, **62**, 1198–1199.
- 22 C. J. Capicciotti, J. D. R. Kurach, T. R. Turner, R. S. Mancini, J. P. Acker and R. N. Ben, *Sci. Rep.*, 2015, **5**, 9692.
- 23 C. J. Crawford, Y. Qiao, Y. Liu, D. Huang, W. Yan, P. H. Seeberger, S. Oscarson and S. Chen, *Org. Process Res. Dev.*, 2021, **25**, 1573–1578.
- 24 F. Broecker and P. H. Seeberger, *Small molecule microarrays: methods and protocols*, 2017, pp. 227–240.

