

Showcasing research from Professor Tim Jackson's laboratory, Department of Chemistry and Center for Environmentally Beneficial Catalysis, University of Kansas, Lawrence, KS, United States.

Characterization and chemical reactivity of room-temperature-stable Mn^{III}-alkylperoxo complexes

Manganese is an earth-abundant metal used both in ancient cave art and in modern oxidation catalysis. This work describes two new Mn^{III}-alkylperoxo complexes that are stable at room temperature. Crystallographic, spectroscopic, and kinetic methods are used to define the properties and reactivity of these complexes. Image created by Abhishek Singh of SlydS.





Chemical Science



EDGE ARTICLE

View Article Online



Cite this: Chem. Sci., 2021, 12, 12564

d All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 8th April 2021 Accepted 20th August 2021

DOI: 10.1039/d1sc01976q

rsc.li/chemical-science

Characterization and chemical reactivity of roomtemperature-stable Mn^{III}-alkylperoxo complexes†

Adedamola A. Opalade, Joshua D. Parham, Victor W. Day and Timothy A. Jackson 🕞 *

While alkylperoxomanganese(III) (Mn^{III}-OOR) intermediates are proposed in the catalytic cycles of several manganese-dependent enzymes, their characterization has proven to be a challenge due to their inherent thermal instability. Fundamental understanding of the structural and electronic properties of these important intermediates is limited to a series of complexes with thiolate-containing N₄S⁻ ligands. These well-characterized complexes are metastable yet unreactive in the direct oxidation of organic substrates. Because the stability and reactivity of Mn^{III}-OOR complexes are likely to be highly dependent on their local coordination environment, we have generated two new Mn^{III}-OOR complexes using a new amidecontaining N₅⁻ ligand. Using the 2-(bis((6-methylpyridin-2-yl)methyl)amino)-N-(quinolin-8-yl)acetamide $(H^{6Me}dpaq)$ ligand, we generated the $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]OTf$ and $[Mn^{III}(OOCm)(^{6Me}dpaq)]OTf$ complexes through reaction of their Mn^{II} or Mn^{III} precursors with ^tBuOOH and CmOOH, respectively. Both of the new Mn^{III}-OOR complexes are stable at room-temperature ($t_{1/2} = 5$ and 8 days, respectively, at 298 K in CH₃CN) and capable of reacting directly with phosphine substrates. The stability of these Mn^{III}-OOR adducts render them amenable for detailed characterization, including by X-ray crystallography for [Mn^{III}(OOCm)(^{6Me}dpaq)]OTf. Thermal decomposition studies support a decay pathway of the Mn^{III}-OOR complexes by O-O bond homolysis. In contrast, direct reaction of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ with PPh₃ provided evidence of heterolytic cleavage of the O-O bond. These studies reveal that both the stability and chemical reactivity of Mn^{III}-OOR complexes can be tuned by the local coordination sphere.


Introduction

Metal-alkylperoxo adducts are essential species in industrial and biological oxidation reactions. 1-3 For example, Co^{III}-alkylperoxo adducts are proposed as intermediates in the industrial oxidation of cyclohexane to adipic acid.1,4-7 In the oxidation mechanism, homolytic cleavage of the O-O bond of a Co^{III}cyclohexylperoxo species leads to the production of cyclohexanol and cyclohexanone.4,8 Further radical-induced oxidation of cyclohexanone by C-C bond cleavage yields adipic acid.4 In biological systems, metal-alkylperoxo adducts are common intermediates in a variety of oxygenase enzymes, where they can be directly involved in substrate oxidation or precede the formation of high-valent metal-oxo species.9,10 Given the importance of metal-alkylperoxo species in such reactions, there are now many examples of synthetic Fe-,11,12 Co-,1,4,5 and Cu-alkylperoxo^{3,13,14} adducts, and these complexes are capable of oxidizing substrates such as 1,4-cyclohexadiene, 2-phenylpropionaldehyde, and triphenylphosphine.

The University of Kansas, Department of Chemistry, Center for Environmentally Beneficial Catalysis, 1567 Irving Hill Road, Lawrence, KS 66045, USA. E-mail: taj@ ku.edu; Tel: +1-785-864-3968

† Electronic supplementary information (ESI) available. CCDC 2048663 2049911 2048664. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1sc01976g

While such studies of synthetic model complexes have probed the properties and reactivity of many types of metalalkylperoxo complexes, examples of Mn-alkylperoxo adducts are more limited, and there remain many open questions concerning the factors governing the decay and reactivity of these complexes. Kovacs and co-workers have performed pioneering investigations of Mn^{III}-alkylperoxo adducts, including structural characterization of a family of complexes by X-ray crystallography. 15,16 These studies employed pentadentate, thiolate-containing N₄S⁻ ligands, which in the corresponding [Mn^{III}(OOR)(N₄S)]⁺ complexes placed the thiolate



[MnIII(OOfBu)(SN4(6-MeDPEN))]*

 $R = H: [Mn^{III}(OO^tBu)(dpag)]^+$ $R = Me: [Mn^{III}(OO^tBu)(dpaq^{2Me})]^+$

Fig. 1 Schematic structures of [Mn^{III}(OO^tBu)(SN₄(6-MeDPEN))]⁺ (left) and [Mn^{III}(OO^tBu)(dpaq^R)]⁺ (right). The [Mn^{III}(OO^tBu)(SN₄(6-MeDPEN))]⁺ complex is a representative example of a Mn^{III}-alkylperoxo adduct supported by an N₄S⁻ ligand.

Edge Article Chemical Science

donor cis to the alkylperoxo ligand, with bulky quinolinyl or 6methylpyridyl substituents trans to each other and cis to the alkylperoxo moiety (Fig. 1, left).^{15,16} The crystallographically observed Mn-N distances for the quinolinyl and 6-methylpyridyl donors range from 2.35 to 2.52 Å, which are quite long for Mn^{III}-N interactions. Interestingly, these long Mn-N distances are correlated with the alkylperoxo O-O bond lengths, which vary from 1.43 to 1.47 Å. As shorter Mn-N distances gave longer O-O bonds, it was proposed that less Lewis acidic Mn^{III} centers yielded more activated Mn^{III}alkylperoxo adducts. By using variable-temperature kinetic studies, the O-O bond lengths for these Mn^{III}-alkylperoxo complexes were in turn related to their thermal decay rates. Mn^{III}-alkylperoxo adducts with longer O-O bonds decayed more rapidly, with lower ΔH^{\ddagger} values and ΔS^{\ddagger} values that were more negative. 16 Because of the correlation between the Mn-N distances and the O-O bond lengths, these results suggest that activation of Mn^{III}-alkylperoxo complexes can be controlled by the donor strength of groups cis to the alkylperoxo unit. Thermal decomposition studies and analysis of the decay products of the Mn^{III}-cumylperoxo adduct supported a decay by homolytic cleavage of the alkylperoxo O-O bond.16 Very recently, Kovacs et al. reported a roomtemperature stable Mn^{III}-alkylperoxo complex supported by an alkoxide analogue of the N₄S⁻ ligands.¹⁷ DFT computations for the $[Mn^{III}(OO^tBu)(N_4O)]^+$ and $[Mn^{III}(OO^tBu)(N_4S)]^+$ pair support the notion that the enhanced stability of the alkoxideligated complex arises from greater Lewis acidity of the Mn^{III}

While these studies provide structure-reactivity correlations with regards to the thermal decay pathway, the Mn^{III}-alkylperoxo adducts of the N₄S⁻ ligands failed to show any direct reaction towards a range of substrates. 16 Product analysis of the reaction solutions provided evidence of the oxidation of a variety of substrates (i.e., PEt₃, TEMPOH, and cyclohexane carboxaldehyde) following the decay of the MnIII-alkylperoxo adducts, implying that a product of the decay pathway is a capable oxidant. In contrast, Mn^{III}-hydroperoxo complexes supported by neutral, macrocyclic N₄ ligands are known to react directly with aldehydes, sulfides, and hydrocarbons possessing weak C-H bonds. 18,19 Given that metal-alkylperoxo adducts are often taken as analogues of metal-hydroxoperoxo species, 18-25 the stark difference in reactivity between Mn^{III}-alkylperoxo and Mn^{III}-hydroperoxo adducts in substrate oxidation reactions is striking. The disparate reactivities of these complexes might reflect the differences in the properties of the supporting ligands employed (N_4S^- for Mn^{III} -alkylperoxo versus neutral N_4 for MnIII-hydroperoxo). There is a clear need to understand better the role of non-thiolate-containing supporting ligands in influencing the properties and reactivity of Mn^{III}-alkylperoxo complexes.

We previously generated a pair of Mn^{III}-alkylperoxo complexes with ligands lacking thiolate ligation.²⁶ These Mn^{III}-alkylperoxo complexes were supported by the pentadentate dpaq and dpaq^{2Me} ligands, both of which feature strongly donating amide groups trans to the alkylperoxo moiety (Fig. 1, right; dpaq = 2-[bis(pyridin-2-ylmethyl)]amino-*N*-quinolin-8-yl-

acetamidate, dpaq^{2Me} = 2-[bis(pyridin-2-ylmethyl)]amino-N-2-methyl-quinolin-8-yl-acetamidate). Both [Mn^{III}(OO t Bu)(dpaq)]⁺ and [Mn^{III}(OO t Bu)(dpaq^{2Me})]⁺ were unstable ($t_{1/2} = 3200$ and 3600 s, respectively, for 2 mM solution in CH₃CN at -15 °C) but were characterized by electronic absorption, Mn K-edge X-ray absorption, and FT-IR spectroscopies. The observation of tBuOO in EPR spectra of the complexes following their thermal decay provided support for a decay pathway involving Mn–O bond homolysis. While these data suggest differences in decay pathways for the thiolate- versus non-thiolate-ligated complexes, the large excess of t BuOOH (ca. 100 equivalents) required to form the [Mn^{III}(OO t Bu)(dpaq)]⁺ and [Mn^{III}(OO t Bu)(dpaq^{2Me})]⁺ complexes made a complete analysis of their reactivity and decay pathways unfeasible. The large excess of t BuOOH also complicated any investigations of substrate oxidation.

Given the limitations of the Mn^{III}-alkylperoxo complexes of the dpaq and dpaq^{2Me} ligands, we sought to develop a derivative of the dpaq ligand that would better stabilize the MnIII-alkylperoxo adduct. Herein, we report MnIII-alkylperoxo adducts supported by ^{6Me}dpag (Fig. 2). This new ligand incorporates steric bulk at the 6 position of the pyridyl substituents in the equatorial plane. This choice was inspired by the higher stability of the Mn^{III}-alkylperoxo complexes supported by N₄S⁻ ligands with two bulky N-donor ligands cis to the alkylperoxo ligand (Fig. 1, left). Xray crystallographic characterization of the Mn^{III}-hydroxo adduct [Mn^{III}(OH)(^{6Me}dpaq)](OTf) reveals that the 6-methylpyridyl groups cause elongations in the Mn-N distance of 0.11 Å relative to [Mn^{III}(OH)(dpaq)](OTf).²⁷ The [Mn^{III}(OH)(^{6Me}dpaq)](OTf) complex reacts with stoichiometric amounts of ^tBuOOH and CmOOH in CH₃CN to generate the [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complexes. These Mn^{III}-alkylperoxo complexes are stable in solution at room temperature with half-lives of ca. 5 and 8 days, respectively. Structural characterization for [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ was obtained by X-ray crystallography. Despite their relatively high thermal stabilities, kinetic studies of [MnIII(OOtBu)(6Medpaq)]+ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ provide evidence for the direct reaction of these Mn^{III}-alkylperoxo adducts with phosphines. These results show that the ligand sphere of Mn^{III}-alkylperoxo adducts is critically important in governing their reactivity in substrate oxidation reactions.




Fig. 2 H^{6Me} dpaq ligand (left), and $[Mn^{III}(OH)(^{6Me}$ dpaq)]⁺, $[Mn^{III}(OO^{t-}Bu)(^{6Me}$ dpaq)]⁺, and $[Mn^{III}(OOCm)(^{6Me}$ dpaq)]⁺ complexes (right). The 6-Me-pyridyl substituents have been highlighted in blue in the structure of the metal complexes.

Results and analysis

Formation and characterization of [Mn $^{\rm II}({\rm OH_2})(^{\rm 6Me}{\rm dpaq})]({\rm OTf})$ and [Mn $^{\rm III}({\rm OH})(^{\rm 6Me}{\rm dpaq})]({\rm OTf})$

[Mn^{II}(H₂O)(^{6Me}dpaq)]OTf was prepared by the metalation of H^{6Me}dpag with Mn^{II}(OTf)₂. The X-ray crystal structure of [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf reveals a monomeric, six-coordinate Mn^{II} center coordinated by the pentadentate ^{6Me}dpag ligand and a water molecule (Fig. 3, left). Previous X-ray crystal structures for [Mn^{II}(dpaq)](OTf)²⁷ and [Mn^{II}(dpaq^{2Me})](OTf)²⁸ showed polymeric species, where the coordination site trans to the amide function was occupied by a carbonyl oxygen from a second [Mn^{II}(dpaq)]⁺ (or [Mn^{II}(dpaq^{2Me})]⁺) cation (Fig. S4†). The 6-Me-pyridyl functions in [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf give rise to elongated bonds (Mn-N4 and Mn-N5) when compared to those of [Mn^{II}(dpaq)](OTf) and [Mn^{II}(dpaq)^{2Me}](OTf) (Tables 1 and S1†). The bonds involving the amide and amine functions (Mn-N2 and Mn-N3, respectively) of [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf are slightly contracted compared to the corresponding distances in [Mn^{II}(dpaq)](OTf) and [Mn^{II}(dpaq)^{2Me}](OTf) (Table 1), which could reflect some compensation for the longer bond lengths with the 6-Me-pyridyl donors. In the solid-state structure, the aqua ligand of [Mn^{II}(H₂O)(6Medpaq)]OTf hydrogen bonds with both an amide oxygen of a neighboring [Mn^{II}(H₂-O)(6Me dpaq)]OTf molecule in the unit cell and with an oxygen of a triflate counter anion (H···O separations of 1.79(7) and 1.86(6)




Fig. 3 X-ray crystal structures of $[Mn^{II}(H_2O)]^{6Me}$ dpaq)]OTf and $[Mn^{III}(OH)]^{6Me}$ dpaq)](OTf) showing 50% probability thermal ellipsoid. Triflate counter ion and non-aqua and non-hydroxo hydrogen atoms were removed for clarity.

Å, respectively). Solution-phase characterization of $[Mn^{II}(H_2-O)]^{6Me}$ dpaq)]OTf in CH₃CN by EPR (Fig. S5†), ESI-MS (Fig. S6†), and Evans NMR (Fig. S7†) provide evidence that the mononuclear structure observed in the solid state is retained in solution.

[Mn^{II}(H₂O)(6Medpaq)]OTf reacts very slowly with dioxygen. Electronic absorption data collected during the oxygenation of a 2.5 mM solution of [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf in CH₃CN shows the appearance of a single feature at 510 nm, but this new chromophore is still forming even after 48 hours (Fig. S8,† left). This reactivity contrasts with that of [Mn^{II}(dpaq)](OTf) and the majority of its derivatives, as these Mn^{II} complexes reacted with dioxygen with full conversion to Mn^{III} products over the course of several hours. 27-29 Presumably, the elongation of the Mn-N4 and Mn-N5 bonds in [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf leads to a more electron-deficient MnII center with muted reactivity with dioxygen. In contrast, the reaction between [Mn^{II}(H₂-O)(6Medpaq)]OTf and 0.5 equiv. PhIO is rapid, resulting in the formation of a bronze colored solution with a single electronic absorption feature at ca. 510 nm (Fig. S8,† right), which can be attributed to [Mn^{III}(OH)(^{6Me}dpaq)](OTf) (vide infra).

X-ray diffraction studies of crystals obtained from this dark orange solution establish the oxidation product as [Mn^{III}(OH)(^{6Me}dpaq)](OTf) (Fig. 3, right). In this complex, a sixcoordinate Mn^{III} center is in a distorted octahedral geometry with the hydroxo ligand trans to the amide function. This coordination mode is identical to that observed in Mn^{III}hydroxo complexes of dpag and its derivatives. 27-29 The Mn-OH distance of 1.806(6) Å is within error of that observed for [Mn^{III}(OH)(dpaq)](OTf) (1.806(13); see Table 1)²⁷ and is on the low end of the range of Mn-OH bond lengths reported for other Mn^{III}-hydroxo complexes (1.81-1.86 Å).^{28,30-38} Further comparisons of the X-ray structure of [Mn^{III}(OH)(^{6Me}dpaq)](OTf) with those of related Mn^{III}-hydroxo species reveals that the 6-Mepyridyl groups give rise to substantial bond elongations. Specifically, the Mn-N4 and Mn-N5 distances of [MnIII(OH)(6-Medpaq)](OTf) are 2.322(6) and 2.381(7) Å, while the corresponding distances in [Mn^{III}(OH)(dpaq)](OTf) are 2.260(14) and 2.216(15) Å (Table 1).27 Mn^{III} complexes of the N₄S⁻ class of ligands (Fig. 1) with 6-Me-pyridyl or quinolinyl substituents in the equatorial field had Mn-N distances ranging from 2.352-

 $\begin{tabular}{ll} \textbf{Table 1} & Manganese-ligand bond lengths (Å) and angles from X-ray crystal structures of $[Mn^{II}(OH_2)(^{6Me}dpaq)](OTf)$, $[Mn^{II}(OH_2)(^{6Me}dpaq)]$, $[Mn^{II}(OH_2)(^{6Me}$

	$\boxed{[Mn^{II}(OH_2)(L)](OTf)}$	$\underline{[Mn^{II}(L)](OTf)}$	[Mn ^{III} (OH)(L)](OTi	f)	$\underline{[Mn^{III}(OOCm)(L)](OTf)}$
Bond	$L = {}^{6Me}dpaq$	L = dpaq	$L = {}^{6Me}dpaq$	L = dpaq	$L = {}^{6Me}dpaq$
Mn-O1	2.108(3)	$2.079(2)^a$	1.806(6)	1.806(13)	1.849(3)
Mn-N1	2.233(3)	2.214(3)	2.041(7)	2.072(14)	2.044(4)
Mn-N2	2.152(4)	2.191(3)	1.962(6)	1.975(14)	1.955(4)
Mn-N3	2.280(3)	2.314(3)	2.130(6)	2.173(14)	2.100(4)
Mn-N4	2.354(4)	2.244(3)	2.322(6)	2.260(14)	2.284(4)
Mn-N5	2.417(3)	2.286(3)	2.381(7)	2.216(15)	2.394(4)
O1-O2					1.466(4)
Mn-O1-O2					110.4(2)

^a For [Mn^{II}(dpaq)](OTf), the oxygen atom derives from a carbonyl unit of a separate [Mn^{II}(dpaq)]⁺ cation.

Edge Article Chemical Science




Fig. 4 1 H NMR spectra of solutions of 20 mM [Mn^{III}(OH)($^{6\text{Me}}$ dpaq)]⁺ (red), 15 mM [Mn^{III}(OO 6 Bu)($^{6\text{Me}}$ dpaq)]⁺ (green), 15 mM [Mn^{III}(OOCm)($^{6\text{Me}}$ dpaq)]⁺ (purple), and 15 mM [Mn^{III}(OH)(dpaq)]⁺ (blue) with 45 μ L added D₂O. All samples were prepared in CD₃CN at 298 K. The Mn^{III}-alkylperoxo complexes were prepared by treating [Mn^{III}(OH)($^{6\text{Me}}$ dpaq)]⁺ with 1 equiv. $^{6\text{Me}}$ BuOOH (green) or CmOOH (purple). Inset: expanded view of the 0 to 13 ppm region.

2.581 Å.¹6,³9 The 6-Me-pyridyl Mn–N distances of [Mn^{III}(OH)(^{6-Me}dpaq)](OTf) are thus on the short end of this range. For [Mn^{III}(OH)(^{6Me}dpaq)](OTf), the longer Mn–N4 and Mn–N5 distances are accompanied by modest contractions of Mn–N1, Mn–N2, and Mn–N3 by 0.031, 0.013, and 0.043 Å, respectively, relative to [Mn^{III}(OH)(dpaq)](OTf) (Table 1). The extended X-ray structure of [Mn^{III}(OH)(^{6Me}dpaq)](OTf) reveals a hydrogen bond between the hydroxo ligand and the amide oxygen of a neighboring [Mn^{III}(OH)(^{6Me}dpaq)]⁺ cation (H···O distance of 2.006 Å,

see Fig. S11†). This kind of interaction was also observed in the crystal structure of [Mn^{III}(OH)($^{2\text{Me}}$ dpaq)](OTf) (H···O distance of 1.982 Å, see Fig. S11†). 28 A free triflate ion is also present in the asymmetric unit of [Mn^{III}(OH)($^{6\text{Me}}$ dpaq)](OTf), though there is no interaction with the Mn center (closest Mn–O distance of *ca.* 7.5 Å).

Previous investigations of [MnIII(OH)(dpaq)](OTf) and a subset of its derivatives have shown that dissolution of the salts of these Mn^{III}-hydroxo complexes in dried CH₃CN leads to the formation of an equilibrium mixture of MnIII-hydroxo and (μ-oxo)dimanganese(III,III) complexes that can both be detected by ¹H NMR spectroscopy.^{29,40} The ¹H NMR spectrum of [Mn^{III}(OH)(6Medpaq)](OTf) in CD₃CN at 298 K exhibits seven hyperfine-shifted peaks that lie well outside 0-20 ppm (the diamagnetic region), as well as two well-resolved peaks in the 0-10 ppm region (Fig. 4, red trace). The lack of a large number of peaks in the diamagnetic region suggests that dissolution of [Mn^{III}(OH)(^{6Me}dpaq)](OTf) in CD₃CN does not result in the formation of (μ-oxo)dimanganese(III,III) species. The chemical shifts for the ¹H NMR signals of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ are quite similar to those of [Mn^{III}(OH)(dpaq)]⁺ (Fig. 4 and Table 2).^{29,40} The four most upfield-shifted peaks in the ¹H NMR spectrum of [Mn^{III}(OH)(dpaq)]⁺ were assigned to protons from the quinolinyl moiety.40 The upfield region of the 1H NMR spectrum of $[Mn^{III}(OH)(^{6Me}dpaq)]^{+}$ shows three sharp peaks at -19.3, -45.0,and −61.6 ppm that resemble the peaks of [Mn^{III}(OH)(dpaq)]⁺ at -15.5, -33.7, -53.8 ppm (Fig. 4). The ¹H NMR spectrum of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ lacks a broad, highly upfield-shifted peak analogous to the weak, broad -63.4 ppm signal of [Mn^{III}(OH)(dpaq)]⁺, but the breadth of this signal renders it difficult to resolve. The overall similarities between the upfield regions of the ¹H NMR spectra of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ and [Mn^{III}(OH)(dpaq)]⁺ are expected given the lack of changes to the quinolinyl group in the former complex. The downfield ¹H NMR signals of $[Mn^{III}(OH)(dpaq)]^+$ and the resonance at -4.6 ppm were attributed to pyridyl protons.40 Consequently, the larger relative perturbations in the downfield regions of the ¹H NMR spectra of [Mn^{III}(OH)(dpaq)]⁺ and [Mn^{III}(OH)(^{6Me}dpaq)]⁺ can be rationalized by changes in chemical shifts of pyridyl protons. The peak at 130.5 ppm for [Mn^{III}(OH)(dpaq)]⁺ was assigned to

Table 2 1 H NMR chemical shifts (ppm) for [Mn^{III}(OH)(6 Medpaq)] $^{+}$, [Mn^{III}(OO^tBu)(6 Medpaq)] $^{+}$, [Mn^{III}(OOCm)(6 Medpaq)] $^{+}$, and [Mn^{III}(OH)(dpaq)] $^{+}$ in CD₃CN at 298 K $^{\alpha}$

[Mn ^{III} (OH)(^{6Me} dpaq)] ⁺	[Mn ^{III} (OO ^t Bu)(^{6Me} dpaq)] ⁺	[Mn ^{III} (OOCm)(^{6Me} dpaq)] ⁺	[Mn ^{III} (OH)(dpaq)] ⁺
			130.5 (H-py)
66.0	66.8	67.8	62.7 (H-qn)
51.4	46.1	46.7	60.9 (H-py)
44.8	42.2	41.8	54.3 (H-py)
			40.5
8.9	9.2	8.9, 7.38, 7.26	
5.5		6.05, 5.3	
-9.6			-4.6 (H-py)
-19.3	-22	-22.3	-15.5 (H-qn)
			-33.7 (H-qn)
-45	-46.9	-47.6	-53.8 (H-qn)
-61.6	-60	-60.3	-63.4 (H-qn)

^a Data and assignments (in parentheses) for [Mn^{III}(OH)(dpaq)]⁺ in CD₃CN with 880 equivalents of D₂O from ref. 40 (py = pyridine, qn = quinoline).

Chemical Science

the α -H of the pyridine substituent.⁴⁰ The lack of a corresponding peak in the ¹H NMR spectrum of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ is consistent with the functionalization of the pyridyl functions in the 6-position. The ¹H NMR spectrum of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ contains a broad, upfield peak at -9.5 ppm, lacking in the ¹H NMR spectrum of [Mn^{III}(OH)(dpaq)]⁺, that we attribute to protons of the 6-methyl-substituents. Overall, the ¹H NMR spectra of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ and [Mn^{III}(OH)(dpaq)]⁺ are quite similar, and the observed differences can be rationalized in terms of the presence of the 6-Me-pyridyl groups in the former complex.

Formation of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺

The addition of a slight excess of *BuOOH to [MnII(H2-O)(6Medpaq)]OTf in CH₃CN at 298 K results in the formation of a green chromophore with a prominent electronic absorption band at 650 nm ($\varepsilon = 240 \text{ M}^{-1} \text{ cm}^{-1}$) and a shoulder near 500 nm ($\varepsilon = 263 \text{ M}^{-1} \text{ cm}^{-1}$; see Fig. 5, left). (Extinction coefficients were determined assuming full conversion under these conditions.) Maximal formation of the green chromophore from [Mn^{II}(H₂-O)(6Medpaq)]OTf requires 1.5 equiv. BuOOH (Fig. S12†). A similar reaction is observed upon the addition of CmOOH to [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf (Fig. S13†). Spectroscopic data provided below and in the ESI (Fig. S14-S16†) support the formulation of these green chromophores as $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ and $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$. The half-lives of the 2 mM solutions of the [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ are ca. 5 and 8 days, respectively, in CH₃CN at 298 K.

The time course for the formation of $[Mn^{III}(OO^tBu)]^{6Me}$ dpaq $]^{-1}$ from [Mn^{II}(H₂O)(6Medpaq)]+ shows an initial rise in absorbance intensity at 510 nm that maximizes near ca. 60 minutes and then drops and levels by 120 minutes. In contrast, the absorbance intensity at 650 nm shows a steep rise from 0 to 40 minutes, grows more slowly from 40-100 minutes, and then rises quickly and levels by 120 minutes (Fig. 5, left inset). On the basis of this





Fig. 5 Left: electronic absorption spectra showing the formation of the green 1.5 mM [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ species (blue trace) from the oxidation of [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf (red trace) with 1.5 equiv. ^tBuOOH. Right: electronic absorption spectra showing the formation of 2.0 mM $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ from the reaction of $[\mathrm{Mn^{III}}(\mathrm{OH})(^{\mathrm{6Me}}\mathrm{dpaq})]^+$ (red trace) with 1.0 equiv. $^t\mathrm{BuOOH}$ (blue trace is the final spectrum). Time courses for each reaction are shown in the insets.



reaction profile, we propose the formation of the Mn^{III}-hydroxo adduct [Mn^{III}(OH)(^{6Me}dpaq)]⁺ as an intermediate in this reaction, as this species shows an absorption maximum at 510 nm. This proposal is consistent with the 1.5:1 *BuOOH: Mn^{II} stoichiometry, where initial oxidation of [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf to [Mn^{III}(OH)(^{6Me}dpaq)]⁺ consumes 0.5 equiv. ^tBuOOH, and the remaining 1.0 equiv. 'BuOOH converts [Mn^{III}(OH)(^{6Me}dpaq)]⁺ to [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ by a ligand substitution reaction that yields water as a co-product (Scheme 1). To test this mechanism, we added 1.0 equiv. 'BuOOH to [Mn^{III}(OH)(^{6Me}dpaq)]⁺ in CH₃CN at 298 K and observed the formation of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ in maximal yield (Fig. 5, right). In this case, the conversion is accompanied with isosbestic points at 505 and 555 nm, indicating the lack of an accumulating intermediate. The formation of a Mn^{III}-hydroxo intermediate during the reaction of ^tBuOOH with [Mn^{II}(dpaq)](OTf) and [Mn^{II}(dpaq^{2Me})](OTf) was observed previously.26

Structural properties of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺

While all attempts at obtaining crystalline material for $[Mn^{III}(OO^tBu)(^{6Me}dpaq)](OTf)$ were unsuccessful, we were able to obtain diffraction-quality crystals for [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf), which confirmed the formulation of this complex (Fig. 6). The cumylperoxo ligand of [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf) is bound trans to the amide nitrogen (N2-Mn-O1 angle of 176.6°), occupying the position of the hydroxo ligand in [Mn^{III}(OH)(^{6Me}dpaq)](OTf). The Mn-O1 bond length for [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf) is longer than the Mn-O1 bond in the Mn^{III}-hydroxo analogue (1.849(3) and 1.806(6) Å, respectively), but similar to crystallographic Mn-OOCm distances for the [Mn^{III}(OOCm)(N₄S)]⁺ complexes (1.848(4) and 1.84(1) Å).15,16 The O1-O2 bond is oriented such that the projection of this bond onto the equatorial plane bisects the N1-Mn-N4 bond angle. The aryl ring of the cumyl moiety is parallel to the plane of the pyridines coordinated to the Mn^{III} center, as opposed to the out-of-plane orientation observed in the [Mn^{III}(S^{Me2}N₄(6-Me-DPEN))(OOCm)](BPh₄) complex.¹⁶ This orientation of the cumyl moiety is stabilized by π -CH interactions between the aryl ring and a methylene group of the ^{6Me}dpaq ligand, as evidenced by short H···C contacts of *ca.* 2.8

Edge Article Chemical Science




Fig. 6 ORTEP diagram of cationic portion of [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf) showing 50% probability thermal ellipsoids. Hydrogen atoms of the 6Medpaq ligand, solvent of crystallization, and the triflate counterion were omitted for clarity.

Å. The O1-O2 bond length of 1.466(4) Å for [Mn^{III}(OOCm)(⁶⁻ Medpaq)](OTf) is consistent with that expected for an alkylperoxo moiety4,5,41,42 and is within the range of values observed for the $[Mn^{III}(OOCm)(N_4S)]^+$ complexes (1.457(5) and 1.51(2) Å). 15,16 The nitrogen atoms of the 6-Me-pyridyl groups (N4 and N5 of [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf)) are 2.284(4) and 2.394(4) Å, respectively, from the Mn center.

Spectroscopic properties of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺

Although we were unable to obtain crystallographic information concerning [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺, the ¹H NMR spectrum of this species in CD₃CN is essentially identical to that of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ (Fig. 4 and Table 2). Each spectrum shows six peaks outside the diamagnetic region, three downfield and three upfield. The only notable difference between the spectra of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ and $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ are the number of peaks from 10-5 ppm (Table 2). While [Mn^{III}(OO^{t-} Bu)(6Medpaq)]+ shows a single prominent peak at 9.2 ppm, [Mn^{III}(OOCm)(6Medpaq)]+ shows five peaks in this region. We tentatively attributed these peaks to protons from the t-butyl and cumyl moieties, respectively. The ¹H NMR spectra of $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ and $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ are also very similar to that of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ (Fig. 4 and Table 2), which is consistent with the same binding mode of the ^{6Me}dpag ligand in each of these complexes. In particular, the three upfield peaks of the Mn^{III}-alkylperoxo adducts at ca. -22, -47, and -60 ppm have chemical shifts very similar to the upfield resonances observed for [Mn^{III}(OH)(6Medpaq)]+ (-19.3, -45, and -61.6 ppm; see Table 2). Similarly, the downfield peaks at *ca.* 67, 46, and 42 ppm in the ¹H NMR spectrum of the Mn^{III}-alkylperoxo adducts show only slight deviations from the corresponding resonances of [Mn^{III}(OH)(6Medpaq)]⁺ (66.0, 51.4, and 44.8 ppm; see Table 2). The solution FT-IR spectra of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ show features at 875 cm⁻¹ and 861 cm⁻¹, respectively, that are absent in the FT-IR spectrum of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ (Fig. 7). These features have energies similar to those of O-O vibrations reported for Mn^{III}-alkylperoxo complexes (872-895 cm⁻¹; Table S2†). 15,16,26 To evaluate this assignment, we prepared [Mn^{III}(¹⁸O¹⁸O^tBu)(^{6Me}dpaq)]⁺, which was




Fig. 7 Solution FT-IR spectra obtained following the reactions of 2 mM solutions of [Mn^{III}(OH)(^{6Me}dpaq)]⁺ in CH₃CN and with 1.0 equiv. ^tBuOOH (red trace) and 1.0 equiv. CmOOH (black trace). The FT-IR spectrum of [Mn^{III}(OH)(^{6Me}dpag)]⁺ (blue trace) is shown for comparison.

confirmed by ESI-MS experiments (Fig. S17†). In the IR spectrum of the ¹⁸O isotopologue, the 875 cm⁻¹ feature found in the spectrum of [Mn^{III}(¹⁶O¹⁶O^tBu)(^{6Me}dpaq)]⁺ has disappeared, and we observe two new features at 854 cm⁻¹ and 818 cm⁻¹ (Fig. S18†). The lack of the 875 cm⁻¹ band supports its assignment as the O-O vibration. The appearance of two new peaks suggests that the O-O vibration is coupled with other vibrations. The IR spectrum of ¹⁸O-labeled [Mn^{III}(OO^tBu)(dpaq)]⁺ also showed two new features relative to the ¹⁶O sample. ²⁶ A normal coordinate analysis of high-spin Fe^{III}alkylperoxo adducts revealed coupling of O-O and t-butyl C-C vibrations that gave rise to unusual isotope shift patterns,43 and vibrational coupling has also been observed for Fe^{III}-hydroperoxo adducts.24,44 Vibrational coupling could explain the multiple peaks observed in the IR spectrum of [Mn^{III}(¹⁸O¹⁸O^tBu)(^{6Me}dpaq)]⁺ and account for deviations in isotope shifts from Hooke's law, as coupled modes would not follow a simple harmonic oscillator model. Under this proposal, one might expect to observe multiple isotope-sensitive peaks in the IR spectrum of [Mn^{III}(¹⁶O¹⁶O^tBu)(⁶ Medpaq)]+. However, such peaks could easily be obscured by the solvent-derived vibrations at ca. 895 cm⁻¹ and from ca. 845-820 cm⁻¹ (Fig. 7).

EPR analysis of frozen 5 mM CH₃CN solutions of [Mn^{III}(OO^{t-} Bu)(6Medpaq)]+ and [MnIII(OOCm)(6Medpaq)]+ collected at 10 K showed no signals in either perpendicular- or parallel-mode (Fig. S19†). This observation is consistent with the lack of X-band EPR signals for many Mn^{III} complexes due to moderate to large zero-field splitting relative to the microwave energy.⁴⁵⁻⁴⁹ The magnetic moments of these complexes $(4.8\mu_B)$ are consistent with $S = 2 \text{ Mn}^{\text{III}}$ centers (Fig. S15 and S16†). Additional ESI-MS data and solution-phase magnetic moments further support the formulations for these Mn^{III}-alkylperoxo complexes (Fig. S14†).

Thermal decay pathways of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺

The thermal decay of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ in CH₃CN at 323 K under anaerobic This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 20 2021. Downloaded on 16.10.25 11:51:50.

Chemical Science

conditions was monitored by electronic absorption spectroscopy. Each decay reaction progressed with the disappearance of the 650 nm feature associated with the MnIII-alkylperoxo adduct and the appearance of a feature at 510 nm (Fig. 8). The feature at 510 nm, along with ESI-MS analyses of the product solutions (Fig. S20†), marks [Mn^{III}(OH)(^{6Me}dpaq)]⁺ as the major product of these decay reactions (99 and 92% formation from [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ and $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ respectively, on the basis of the extinction coefficient of the Mn^{III}-hydroxo complex). For [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺, the decay proceeds with isosbestic behavior, and the respective decay and formation rates of the 650 and 510 nm electronic absorption signals are identical (Fig. 8, right; $k_{\rm obs} = 0.016 \, \rm min^{-1}$ for both processes). In contrast, the thermal decay for $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ is not isosbestic, and the rate of formation of the 510 nm chromophore lags behind the decay of the 650 nm band (Fig. 8, left). We tentatively attribute this difference to the higher purity of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ used in these experiments, as this complex was obtained as a recrystallized solid. In support, thermal decay studies of crude [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ also failed to show isosbestic behavior (Fig. S22†). Additional product analysis following the thermal decay of 18 mM [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ in CH₃CN at 323 K revealed 61.3 \pm 0.1% 2-phenyl-2-propanol and 25.7 \pm 0.1% acetophenone formed relative to the [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ concentration. (The organic products from the thermal decay of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ were not quantified, because the volatility of acetone, a potential product, renders quantification unreliable.)

When the decay of 6 mM $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ was performed in CD₃CN, we observed 50 \pm 0.3% 2-phenyl-2propanol and 40 \pm 0.3% acetophenone relative to the initial [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺, a marked change in the product distribution. In addition, when [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ was allowed to decay in benzonitrile (PhCN; see Fig. S23†), we observed 30 \pm 1.1% 2-phenyl-2-propanol and 70 \pm 1.1% acetophenone. The implications of these results with respect to the decay mechanism are explored in the Discussion section (vide infra).




Fig. 8 Electronic absorption spectra showing the decay of anaerobic CH₃CN solutions of 2 mM [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and 6 mM [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ (red traces) at 323 K to give [Mn^{III}(OH)(⁶ Medpaq)]⁺ (blue traces). The inset shows the change in absorbance at 650 and 510 nm over the course of the decay reaction.

Fig. 9 Eyring plot from variable-temperature thermal decay kinetics of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ (left) and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ (right) in CH₃CN.

To further probe the decay reactions, we monitored decay kinetics for 1.25 mM CH₃CN solutions of each complex from 303-348 K in CH₃CN under anaerobic conditions. At each temperature, the decay could be fit to a pseudo-first-order process, and the $k_{\rm obs}$ values at different temperatures were fit to the Eyring equation to obtain activation parameters (Fig. 9). This analysis yielded $\Delta H^{\ddagger} = 21.4 \pm 1.5 \text{ kcal mol}^{-1}$, $\Delta S^{\ddagger} = -9.5 \pm$ 4.9 cal mol⁻¹ K⁻¹, and $\Delta G^{\ddagger} = 24.2 \pm 3.0 \text{ kcal mol}^{-1}$ at 298 K for $[\mathrm{Mn^{III}}(\mathrm{OO}^t\mathrm{Bu})(^{6\mathrm{Me}}\mathrm{dpaq})]^+$; and $\Delta H^{\ddagger}=23.5\pm1.2~\mathrm{kcal~mol}^{-1},\Delta S^{\ddagger}$ $=-1.5\pm3.6~{
m cal~mol}^{-1}~{
m K}^{-1}$, and $\Delta G^{\ddagger}=23.9\pm2.2~{
m kcal~mol}^{-1}$ at 298 K for [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺. Both complexes display ΔH^{\ddagger} values significantly higher than those of the $[Mn^{III}(OOR)(N_4S)]^+$ complexes $(\Delta H^{\ddagger} = 15.9 - 10.5 \text{ kcal mol}^{-1}),^{16}$ which is in line with the greater thermal stabilities of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ and $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$. The entropies of activation of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ are slightly negative and smaller in magnitude than those of the $[Mn^{III}(OOR)(N_4S)]^+$ complexes (ΔS^{\ddagger}) $= -15 \text{ to } -34 \text{ cal mol}^{-1} \text{ K}^{-1}$).

Substrate oxidation by Mn^{III}-alkylperoxo adducts

1 Direct oxidation of triphenylphosphine. The addition of 100 equiv. PPh₃ to an anaerobic solution of [Mn^{III}(OO^tBu)(⁶⁻ Medpaq)]+ (2.0 mM in CH₃CN) at 298 K resulted in the loss of intensity at 650 nm over the course of two hours, resulting in an electronic absorption spectrum consistent with the generation of Mn^{II} products (Fig. 10, left). The decay of the 650 nm absorption signal could be well-fit to a first-order model, yielding a pseudo-first- order rate constant $k_{\rm obs}$ (Fig. 10, left inset). A ³¹P NMR analysis of the organic products revealed the formation of Ph₃PO in 70% yield relative to the Mn^{III}-alkylperoxo concentration (Fig. S24†). EPR analysis of the reaction mixture shows a signal centered at g = 2.03 that is similar in appearance to that of the [Mn^{II}(H₂O)(^{6Me}dpaq)]OTf starting material, albeit with the lack of apparent hyperfine splitting (Fig. S25†). 28,39,50 This evidence is in accordance with the featureless UV-vis spectrum of the final reaction mixture, which is characteristic of a $\mathbf{Mn}^{\mathrm{II}}$ product (Fig. 10, left). The rate of decay of [MnIII(OOtBu)(6Medpaq)]+ increased linearly with increasing concentrations of PPh3 (Fig. 10, right). A linear fit of kobs versus PPh3 concentration yields a second-order rate constant for PPh₃ oxidation by [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ of **Edge Article**

1.4 Data
1.2 Data
1.0 Data
1.0

Fig. 10 Left: electronic absorption spectra monitoring the reaction of a 2.0 mM anaerobic solution of $[Mn^{III}(OO^tBu)]^{6Me}$ dpaq)]⁺ (red trace) in CH₃CN with 100 equivalents of PPh₃ at 298 K. Left inset: time course for the spectral changes. Right: pseudo-first-order rate constants, k_{obs} (s⁻¹), versus PPh₃ concentration (M) for a 1.25 mM solution of $[Mn^{III}(OO^tBu)]^{6Me}$ dpaq)]⁺ at 298 K.

0.0035 M $^{-1}$ s $^{-1}$ at 298 K in CH $_3$ CN. The [Mn $^{\rm III}$ (OOCm)($^{6\rm Me}$ d-paq)] $^+$ complex displayed similar reactivity with PPh $_3$ ($k_2=0.0033~{\rm M}^{-1}{\rm s}^{-1}$ at 298 K; see Fig. S26†). These rates show that the nature of the alkylperoxo group does not have a significant influence on the reactivity with PPh $_3$.

Further insight into the reaction of PPh₃ with [Mn^{III}(OO^{t-}Bu)(^{6Me}dpaq)]⁺ was obtained through an Eyring analysis of variable-temperature kinetic experiments (Fig. 11). These experiments yielded $\Delta H^{\ddagger}=17.6\pm1.4$ kcal mol⁻¹, $\Delta S^{\ddagger}=-12.6\pm4.6$ cal mol⁻¹ K⁻¹, and $\Delta G^{\ddagger}=21.3\pm2.8$ kcal mol⁻¹ at 298 K. The high activation enthalpy and Gibbs free energy of activation account for the sluggishness of this reaction, and the negative entropy of activation is consistent with a bimolecular reaction.

Mechanistic insight into the reaction between PPh₃ and the Mn^{III}-alkylperoxo was obtained by the reaction of 22 mM [Mn^{III}(OOCm)($^{6\text{Me}}$ dpaq)]⁺ with 5 equiv. of PPh₃ in CH₃CN at 298 K. Quantification of the organic products of this reaction by GC-MS revealed 88.5 \pm 0.3% 2-phenyl-2-propanol and 1.5 \pm 0.3% acetophenone based on the initial concentration of [Mn^{III}(OOCm)($^{6\text{Me}}$ dpaq)]⁺. ESI-MS analysis of the final reaction mixtures for the reaction of [Mn^{III}(OOCm)($^{6\text{Me}}$ dpaq)]⁺ with PPh₃ showed the presence of peaks at m/z = 465.14, 482.14 and 743.23, which are consistent with expected m/z values for




Fig. 11 Eyring plot from variable-temperature reaction kinetics of $[Mn]^{II}(OO^tBu)(^{6Me}dpaq)]^+$ with PPh₃ in CH₃CN.

 $[Mn^{II}(^{6Me}dpaq)]^+$, $[Mn^{III}(OH)(^{6Me}dpaq)]^+$ and $[Mn(OPPh_3)(^{6-Me}dpaq)]^+$, respectively (Fig. S27†).

2 Indirect oxidation of 9,10-dihydroanthracene (DHA). The reaction of the Mn^{III}-alkylperoxo complexes with the hydrogenatom donor DHA was also explored. In this case, the addition of a large excess of DHA (100 equiv. relative to MnIII) to $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ or $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ CH₃CN at 323 K did not result in any change in the rate of decay of the Mn^{III}-alkylperoxo complexes (Fig. S28†). We also observed no change in the thermal decay rate with d_4 -DHA (Fig. S29†). Nonetheless, an analysis of the reaction solution following the full thermal decay of [MnIII(OOtBu)(6Medpaq)]+ or [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ (after ca. 7 hours) revealed the formation of ca. 1.4 equiv. anthracene relative to the starting Mn^{III}-alkylperoxo concentration (Fig. S30†). Thus, while neither $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ nor $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ is capable of direct oxidation of DHA, a product of the thermal decay of each complex is an effective oxidant of DHA.

Discussion

Ligand-sphere influence on the structure-property correlations of Mn^{III}-alkylperoxo complexes

The generation of the room-temperature stable [Mn^{III}(OO^tBu)(⁶⁻ Medpaq)]+ and [MnIII(OOCm)(6Medpaq)]+ complexes relied upon previous observations that Mn^{III} centers with higher Lewis acidity give rise to corresponding Mn^{III}-alkylperoxo adducts with shorter and more stable O-O bonds. 16 The basis of this correlation rests on the π -donating properties of the alkylperoxo ligand. According to this model, which was originally proposed by Kovacs, DeBeer, and co-workers, 16 a more Lewis acidic Mn^{III} center fosters greater πdonation from the alkylperoxo π^* MO, which strengths the O-O bond. The 6-Me-pyridyl groups of the structurally characterized [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex give two elongated Mn-N distances of 2,284(4) and 2,394(4) Å. These weak metal-ligand interactions increase the Lewis acidity of the MnIII center, stabilizing the Mn^{III}-alkylperoxo unit. DFT computations lend credence to this model, showing a reduction in the Mulliken charge of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ relative to $[Mn^{III}(OO^tBu)(dpaq)]^+$ (0.48) and 0.52, respectively). The computations also predict a greater admixture of alkylperoxo character in the Mn–OO^tBu π -antibonding MO for [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ (Fig. S33†), which supports stronger π -donation in this complex.

The crystal structure of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ allows us to determine how closely this complex follows previously observed correlations based on the [Mn^{III}(OOR)(N₄S)]²⁺ complexes. The [Mn^{III}(OOR)(N₄S)]⁺ complexes showed a linear correlation between the elongated Mn–N distances and the O–O bond lengths, with the shortest Mn–N distances of *ca.* 2.40 Å giving rise to the longest O–O bonds of *ca.* 1.47 Å (Fig. 12). The metric parameters for [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ follow the spirit of this correlation; that is, this complex has a short average Mn–N distance of 2.34 Å and an O–O bond of 1.466(4) Å, on the long end of that observed for Mn^{III}–alkylperoxo adducts (1.43–1.47 Å). However, if we use the previous correlation as a guide, the Mn–N distances observed for [Mn^{III}(OOCm)(^{6Me}d-paq)]⁺ would predict a O–O bond length far longer than that




Fig. 12 Correlation between average Mn–N distance and alkylperoxo O–O bond length for [Mn III (OOR)(N₄S)] $^+$ complexes (see ref. 16) and corresponding point for [Mn III (OOCm)(6Me dpaq)](OTf) described in this work

observed, making this complex a clear outlier (Fig. 12). It is not completely surprising that the markedly different coordination spheres of $[Mn^{\rm III}(OOCm)(^{6Me}dpaq)]^+$ and the $[Mn^{\rm III}(OOR)(N_4S)]^+$ series would cause such a deviation, as the O–O distance should be a reporter of the entire coordination sphere. It is additionally possible that there is a limit to the extent to which the O–O bond can be elongated in $Mn^{\rm III}$ –OOR complexes, and that the limit is near 1.47 Å.

The [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex also breaks the previously observed correlation that Mn^{III}-alkylperoxo adducts with longer O-O bonds are less stable than those with shorter O-O bonds. 15 [Mn^{III}(OOCm)(6Medpaq)] has an O-O distance at the long end of the [Mn^{III}(OOR)(N₄S)]⁺ series but has a roomtemperature half-life of ca. 5 days. In contrast, the most stable [Mn^{III}(OOR)(N₄S)]⁺ complex has a half-life of ca. 5 minutes at 293 K.16 One caveat that must be noted in comparing the thermal stability of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ with the [Mn^{III}(OOR)(N₄S)]⁺ series is the difference in solvents (CH₃CN and CH2Cl2, respectively).16 To address this complication, we determined the half-life of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ in CH₂Cl₂ at 298 K. The [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex did decay more rapidly in CH₂Cl₂ than in CH₃CN (half-life of 3 vs. 8 days, respectively). While solvent does have some effect on the stability of the Mn^{III}-alkylperoxo complex, the solvent change alone cannot account for the dramatic increase in stability of [Mn^{III}(OOR)(^{6Me}dpaq)]⁺ complexes relative to the [Mn^{III}(OOR)(N₄S)]⁺ series. On this basis, while also noting the limited sample size, it is tempting to speculate that the presence of thiolate ligands in the [Mn^{III}(OOR)(N₄S)]²⁺ series severely reduces the stability of the Mn^{III}-alkylperoxo adducts. This conclusion makes it all the more remarkable that the first isolable Mn^{III}-alkylperoxo adducts contained thiolate ligands.


Thermal decay mechanism of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺

The organic products observed upon the decay of cumylperoxometal complexes are often used to infer the nature of the decay

pathway.⁵¹ Homolytic cleavage of the O–O bond produces cumyloxyl radical that can rearrange by β-scission to produce acetophenone and 'CH₃ (Scheme 2, path b). Alternatively, heterolytic cleavage of the O–O bond produces cumyl oxyanion that can deprotonate solvent to produce 2-phenyl-2-propanol (Scheme 2, path a).^{51–53} Previous studies of Mn^{III}–alkylperoxo¹⁶ and some Fe^{III}–alkylperoxo¹² complexes showed organic products exclusively attributable to O–O homolysis. In contrast, there are examples of Cu^{II}–alkylperoxo⁵⁴ adducts that decay exclusively by heterolytic cleavage of the O–O bond, and there are a handful of examples where the decay of a metal–alkylperoxo adduct yields a mixture of products characteristic of both pathways.^{2,5,11,13,55}

Analysis of the organic decay products of [Mn^{III}(OOCm)(^{6-Me}dpaq)]⁺ reveal both 2-phenyl-2-propanol and acetophenone (61 and 26% yield relative to the [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ adduct). Electronic absorption, ¹H NMR, and ESI-MS data identify [Mn^{III}(OH)(^{6Me}dpaq)]⁺ as a dominant decay product (Fig. S20 and S21†). The distribution of 2-phenyl-2-propanol and acetophenone could suggest that both homolysis and heterolysis of the O-O bond occur in the thermal decay of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ in CH₃CN, with a preference for the heterolytic pathway. Either O-O cleavage pathway would yield a high-valent Mn-oxo intermediate that could be reduced to the observed Mn^{III}-hydroxo product. There are recent reports of Mn^{IV}-oxo adducts of the closely related dpaq ligand that react with C-H bonds to yield a Mn^{III}-hydroxo product. ^{56,57}

To clarify the decay pathway of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺, we examined the products formed when the complex decayed in CD₃CN. In this case, we observed increased formation of acetophenone and decreased formation of 2-phenyl-2-propanol (40:50%) compared to the decay in CH₃CN (26:61%). A change in product distribution in deuterated solvent was also observed by Cho *et al.* in their investigations of a Cu^{II}–alkylperoxo complex.¹³ Itoh^{2,3} and others^{51,52} have rationalized a change in the acetophenone: 2-phenyl-2-propanol distribution in terms of solvent involvement in the decay pathway. In CH₃CN, the cumyloxyl radical decays by competing reactions: (1) β-scission to yield acetophenone and methyl radical, and (2) hydrogen-atom abstraction from solvent to yield 2-phenyl-2-



Scheme 2 Possible decay pathways for [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf).

Edge Article Chemical Science

propanol (Scheme 2, paths c and d).2,3,51,54,55,58 In CD₃CN, the rate of the hydrogen-atom abstraction reaction from the solvent is decreased, yielding a marked increase in acetophenone formation by β-scission. When [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ decays in PhCN, we observe an even greater increase in acetophenone formation (70%), with only 30% formation of 2-phenyl-2propanol. We attribute this change to the strong C-H bonds in PhCN that suppress the reaction of solvent with the cumyloxyl radical (Scheme 2, path d). Taken together, the change in the distribution of acetophenone: 2-phenyl-2-propanol in CD₃CN and PhCN is strong evidence that [Mn^{III}(OOCm)(^{6Me}dpaq)] decays, at least to some extent, by O-O homolysis. However, the persistence of a significant fraction of 2-phenyl-2propanol product, even in CD₃CN and PhCN, suggests that the thermal decay of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ also involves O-O heterolysis. Thus, our data provide evidence for both decay pathways in Scheme 2, each of which yields [Mn^{III}(OH)(^{6Me}dpaq]]⁺ as the Mn-containing product. This situation is distinct from that observed for [Mn^{III}(OOR)(N₄S)]⁺ complexes, ¹⁶ where the decay proceeded exclusively by O-O homolysis and yielded a mixture of Mn-containing products.

A second line of evidence for the production of cumyloxyl radicals by O-O homolysis of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ comes from changes to the decay kinetics of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ in CH₃CN and CD₃CN. The [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex decays slowly in CH₃CN, and this rate of decay matches the rate of formation of the [Mn^{III}(OH)(^{6Me}dpaq)]⁺ product (Fig. 8). In contrast, the decay rate of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ increases in CD₃CN by about eight-fold relative to that in CH₃CN (Fig. S34†). In addition, the decay rate of [Mn^{III}(OOCm)(^{6Me}dpaq)] in CD₃CN is five-fold faster than the rate of formation of [Mn^{III}(OH)(^{6Me}dpaq)]⁺. These observations are consistent with our proposal that a fraction of the [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex decays by homolytic O-O cleavage to give a Mn^{IV}-oxo adduct and cumyloxyl radical. In CH₃CN, the cumyloxyl radical and Mn^{IV}-oxo intermediates react rapidly and preferentially with solvent to give the observed [Mn^{III}(OH)(^{6Me}dpaq)]⁺ and 2phenyl-2-propanol products. Under these conditions, a relatively small amount of cumyloxyl radical undergoes β-scission to yield acetophenone. In CD₃CN, the Mn^{IV}-oxo adduct and cumyloxyl radical decay products have slower rates of reaction with solvent, allowing for reaction with [MnIII(OOCm)(6Medpaq)]⁺, which hastens its decay.

Reaction mechanism of [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ with PPh₃

To the best of our knowledge, the reactions of [Mn^{III}(OO'Bu)($^{6-Me}$ dpaq)]⁺ and [Mn^{III}(OOCm)(6Me dpaq)]⁺ with PPh₃ at 298 K are the first observations of direct substrate oxidation by Mn^{III}-alkylperoxo complexes. The reaction of [Mn^{III}(OOCm)(6Me d-paq)]⁺ with PPh₃ showed the near exclusive formation of 2-phenyl-2-propanol, with only a trace amount of acetophenone (Fig. S35†). This distribution suggests a change to O–O heterolysis under these conditions. An Eyring analysis for the reaction of [Mn^{III}(OO'Bu)(6Me dpaq)]⁺ with PPh₃ gives $\Delta S^{\ddagger} = -12.6 \pm 4.6$ cal mol $^{-1}$ K $^{-1}$, which is consistent with a bimolecular reaction involving the association of [Mn^{III}(OO'Bu)(6Me dpaq)]⁺ with PPh₃

to form the activated complex. The change in reaction rate as a function of PPh₃ concentration is further evidence of a direct reaction between the [Mn^{III}(OOR)(^{6Me}dpaq)]⁺ complexes and PPh₃. In addition, an ESI-MS analysis of the products of the reaction of [Mn^{III}(OO^tBu)(^{6Me}dpaq)]⁺ with PPh₃ revealed a peak for [Mn(OPPh₃)(^{6Me}dpaq)]⁺ that shifts by +2 mass units when the Mn^{III}-alkylperoxo adduct is prepared using ^tBu¹⁸O¹⁸OH (Fig. S27†). Thus, the oxygen in the OPPh₃ product derives from the Mn^{III}-alkylperoxo unit.

We propose a reaction mechanism where [Mn^{III}(OOR)(^{6-Me}dpaq)]⁺ and PPh₃ form an activated complex, with PPh₃ interacting with the proximal oxygen of the alkylperoxo ligand (Scheme 3). Recent reports show that Brønsted and Lewis acids, or the introduction of secondary coordination interaction through pendant amines which act as hydrogen-bond acceptor in an Fe^{III}–OOR (R = H, acyl) adduct could direct heterolytic cleavage. ^{11,59-61} This interaction between [Mn^{III}(OOR)(^{6Me}dpaq)]⁺ and PPh₃ may also be able to instigate heterolytic cleavage of the Mn^{III}–alkylperoxo O–O bond. For the [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ complex, this decay will lead to the formation of cumyloxy anion, which gives 2-phenyl-2-propanol after protonation^{2,54,55} and a Mn^{III}-species that is reduced to the Mn^{II} product observed by UV-vis and EPR spectroscopy (Scheme 3). The identity of the reductant for the Mn^{III} center is unclear.

Reaction mechanism of [Mn^{III}(OOCm)(^{6Me}dpaq)](OTf) with DHA

In contrast to the direct oxidation of PPh₃, the reaction of $[Mn^{III}(OO^tBu)(^{6Me}dpaq)]^+$ and $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$ with DHA is an indirect oxidation caused by products of the Mn^{III} -alkylperoxo decay process (Scheme 4). Neither $[Mn^{III}(OOR)(^{6-Me}dpaq)]^+$ complex shows any change in decay rate in the presence of an excess amount of DHA (Fig. S29†), although the decay solutions reveal the formation of 1.4 equiv. anthracene relative to the initial Mn^{III} -OOR concentration. The notion that the reaction is indirect is further supported by the lack of any change in decay rate when d_4 -DHA is used as substrate.

The formation of 1.4 equiv. anthracene is consistent with the thermal decay of the $[Mn^{III}(OOR)(^{6Me}dpaq)]^+$ complexes by O–O homolysis (Scheme 4). The Mn^{IV} -oxo decay product should be

Scheme 3 Proposed mechanisms for PPh_3 oxidation by $[Mn^{III}(OOCm)(^{6Me}dpaq)]^+$.

Scheme 4 Mechanistic proposal for the reaction of DHA with $[Mn]^{II}(OO^tBu)(^{6Me}dpaq)]^+$.

capable of DHA oxidation, as observed for several oxo–manganese complexes. ^{56,57} This reaction will result in a Mn^{II}–aqua complex, consistent with the observation of a Mn^{II} signal in the EPR spectrum of the final reaction mixture (Fig. S30 and S31†). The cumyloxyl radical also generated by O–O homolysis could be responsible for the remaining 0.4 equiv. anthracene.

Conclusions

Inspired by previously developed structure-reactivity correlations, we developed a new ligand derivative (6Me dpaq) that provides remarkable stability to Mn^{III}-alkylperoxo complexes. A simple change of two pyridyl groups to 6-Me-pyridyl groups results in new Mn^{III}-alkylperoxo complexes that (i) can be generated using stoichiometric amounts of oxidant rather than large excesses, and (ii) are stable at room temperature. This enhanced stability allowed us to structurally characterize a Mn^{III}-cumylperoxo adduct by X-ray diffraction. In spite of the unusual stability of these Mn^{III}-alkylperoxo adducts, these complexes are the first members of their class to show direct reactivity with a substrate (triphenylphosphine). This result demonstrates that the ligand-sphere of Mn^{III}-alkylperoxo adducts has great control over reactivity. Examination of the thermal decay of these N₅--ligated Mn^{III}-alkylperoxo adducts provides evidence from both homolytic and heterolytic O-O bond cleavage, which is distinct from that observed for Mn^{III}alkylperoxo adducts bound by thiolate-containing N₄S⁻ ligands.

While the basis of the enhanced stability of [Mn^{III}(OO^{f-}Bu)(^{6Me}dpaq)]⁺ and [Mn^{III}(OOCm)(^{6Me}dpaq)]⁺ will be the subject of future investigations, it is tempting to speculate that the thiolate ligands in the [Mn^{III}(OOR)(N₄S)]⁺ complexes serve to lower the activation energy for decay. In addition, while the new Mn^{III}-alkylperoxo adducts generally follow a previously identified structural correlation between Mn–N and O–O distances, the observed O–O distance for the Mn^{III}-cumylperoxo adduct is far shorter than expected on the basis of the Mn–N distances. Thus, perturbations to the primary coordination sphere not only affect reactivity but also render this complex an outlier compared to previous compounds. Future work will be aimed at understanding the basis for this outlier status in terms of both structural correlations and chemical reactivity.

Data availability

Crystallographic data are available through the Cambridge Crystallographic Data Centre (CCDC) at https://www.ccdc.cam.ac.uk/ with structure codes 2048663, 2049911, and 2048664.

Author contributions

A. A. O., J. D. P. and T. A. J. conceived and planned the experiments. A. A. O. and J. D. P. performed all experiments and computations, except those involving X-ray crystallography, which were performed by V. W. D. All authors contributed to data analysis and provided contributions to writing of the final manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the U.S. National Science Foundation (CHE-1900384 to T. A. J.). Support for the NMR instrumentation was provided by NIH Shared Instrumentation Grant # S100D016360. The U.S. NSF is also acknowledged for funds used for the purchase of X-ray instruments (CHE-0079282) and the EPR spectrometer (CHE-0946883). We will also acknowledge Samuel R. Crowell for the provision of some [Mn^{III}(OH)(^{6Me}dpaq)]OTf complex, and we thank Prof. Mikhail Barybin (University of Kansas) for helpful discussions regarding the synthesis of ^tBu¹⁸O¹⁸OH.

Notes and references

- 1 F. A. Chavez and P. K. Mascharak, Acc. Chem. Res., 2000, 33, 539-545.
- 2 T. Tano, M. Z. Ertem, S. Yamaguchi, A. Kunishita, H. Sugimoto, N. Fujieda, T. Ogura, C. J. Cramer and S. Itoh, *Dalton Trans.*, 2011, **40**, 10326–10336.
- 3 S. Itoh, Acc. Chem. Res., 2015, 48, 2066-2074.
- 4 F. A. Chavez, C. V. Nguyen, M. M. Olmstead and P. K. Mascharak, *Inorg. Chem.*, 1996, 35, 6282–6291.
- 5 F. A. Chavez, J. M. Rowland, M. M. Olmstead and P. K. Mascharak, J. Am. Chem. Soc., 1998, 120, 9015–9027.
- 6 R. Chakrabarty, P. Sarmah, B. Saha, S. Chakravorty and B. K. Das, *Inorg. Chem.*, 2009, **48**, 6371–6379.
- 7 G. W. Parshall and S. D. Ittel, *Homogeneous Catalysis. The Applications and Chemistry of Catalysis by Soluble Transition Metal Complexes*, Wiley-Interscience, New York, 1992.
- 8 J. Ebner and D. Riley, in *Active Oxygen in Chemistry*, ed. C. S. Foot, J. S. Valentine, A. Greenberg, and J. F. Liebman, Chapman & Hall, London, 1995, pp. 205–248.
- 9 P. F. Fitzpatrick, Biochemistry, 2003, 42, 14083-14091.
- 10 S. C. Peck and W. A. van der Donk, *JBIC*, *J. Biol. Inorg. Chem.*, 2017, 22, 381–394.
- 11 I. Ghosh, S. Banerjee, S. Paul, T. Corona and T. K. Paine, *Angew. Chem., Int. Ed.*, 2019, **58**, 12534–12539.
- 12 S. Hong, Y.-M. Lee, K.-B. Cho, M. S. Seo, D. Song, J. Yoon, R. Garcia-Serres, M. Clémancey, T. Ogura, W. Shin, J.-M. Latour and W. Nam, *Chem. Sci.*, 2014, 5, 156–162.
- 13 B. Kim, D. Jeong and J. Cho, Chem. Commun., 2017, 53, 9328–9331.
- 14 H. Oh, W.-M. Ching, J. Kim, W.-Z. Lee and S. Hong, *Inorg. Chem.*, 2019, **58**, 12964–12974.
- 15 M. K. Coggins and J. A. Kovacs, *J. Am. Chem. Soc.*, 2011, **133**, 12470–12473.
- M. K. Coggins, V. Martin-Diaconescu, S. DeBeer and J. A. Kovacs, *J. Am. Chem. Soc.*, 2013, 135, 4260–4272.
- 17 A. N. Downing, M. K. Coggins, P. C. Y. Poon and J. A. Kovacs, *J. Am. Chem. Soc.*, 2021, **143**, 6104–6113.
- 18 M. Sankaralingam, Y.-M. Lee, S. H. Jeon, M. S. Seo, K.-B. Cho and W. Nam, *Chem. Commun.*, 2018, **54**, 1209–1212.

10 H So V I Park K-R Cho V-M Lee M S Seo I Cho 42 N Lehnert P V N Ho I Ou

19 H. So, Y. J. Park, K.-B. Cho, Y.-M. Lee, M. S. Seo, J. Cho, R. Sarangi and W. Nam, J. Am. Chem. Soc., 2014, 136, 12229–12232.

Edge Article

- 20 N. Lehnert, F. Neese, R. Y. N. Ho, L. Que and E. I. Solomon, *J. Am. Chem. Soc.*, 2002, **124**, 10810–10822.
- 21 P. Chen, K. Fujisawa and E. I. Solomon, *J. Am. Chem. Soc.*, 2000, **122**, 10177–10193.
- 22 B. Kim, D. Jeong, T. Ohta and J. Cho, Commun. Chem., 2019, 2, 81.
- 23 A. S. Faponle, M. G. Quesne, C. V. Sastri, F. Banse and S. P. de Visser, *Chemistry*, 2015, **21**, 1221–1236.
- 24 L. R. Widger, Y. Jiang, A. C. McQuilken, T. Yang, M. A. Siegler, H. Matsumura, P. Moënne-Loccoz, D. Kumar, S. P. de Visser and D. P. Goldberg, *Dalton Trans.*, 2014, 43, 7522–7532.
- 25 A. W. Stubbs and M. Dincă, *Inorg. Chem.*, 2019, 58, 13221–13228.
- 26 J. D. Parham, G. B. Wijeratne, D. B. Rice and T. A. Jackson, *Inorg. Chem.*, 2018, 57, 2489–2502.
- 27 G. B. Wijeratne, B. Corzine, V. W. Day and T. A. Jackson, *Inorg. Chem.*, 2014, 53, 7622–7634.
- 28 D. B. Rice, G. B. Wijeratne, A. D. Burr, J. D. Parham, V. W. Day and T. A. Jackson, *Inorg. Chem.*, 2016, 55, 8110–8120.
- 29 D. B. Rice, A. Munasinghe, E. N. Grotemeyer, A. D. Burr, V. W. Day and T. A. Jackson, *Inorg. Chem.*, 2019, 58, 622–636.
- 30 C. R. Goldsmith, A. P. Cole and T. D. P. Stack, J. Am. Chem. Soc., 2005, 127, 9904–9912.
- 31 S. El Ghachtouli, B. Lassalle-Kaiser, P. Dorlet, R. Guillot, E. Anxolabéhère-Mallart, C. Costentin and A. Aukauloo, Energy Environ. Sci., 2011, 4, 2041–2044.
- 32 Z. Shirin, B. S. Hammes, V. G. Young and A. S. Borovik, *J. Am. Chem. Soc.*, 2000, **122**, 1836–1837.
- 33 B. Cheng, P. H. Fries, J.-C. Marchon and W. R. Scheidt, *Inorg. Chem.*, 1996, 35, 1024–1032.
- 34 A. A. Eroy-Reveles, Y. Leung, C. M. Beavers, M. M. Olmstead and P. K. Mascharak, *J. Am. Chem. Soc.*, 2008, **130**, 4447– 4458.
- 35 D. M. Eichhorn and W. H. Armstrong, *J. Chem. Soc., Chem. Commun.*, 1992, 85–87, DOI: 10.1039/C39920000085.
- 36 T. J. Hubin, J. M. McCormick, N. W. Alcock and D. H. Busch, *Inorg. Chem.*, 2001, **40**, 435–444.
- 37 M. K. Coggins, L. M. Brines and J. A. Kovacs, *Inorg. Chem.*, 2013, **52**, 12383–12393.
- 38 Z. Shirin, A. S. Borovik and V. G. Young Jr, *Chem. Commun.*, 1997, 1967–1968, DOI: 10.1039/A703395H.
- 39 M. K. Coggins, S. Toledo, E. Shaffer, W. Kaminsky, J. Shearer and J. A. Kovacs, *Inorg. Chem.*, 2012, **51**, 6633–6644.
- 40 D. B. Rice, S. D. Jones, J. T. Douglas and T. A. Jackson, *Inorg. Chem.*, 2018, **57**, 7825–7837.
- 41 C. Fontaine, K. N. V. Duong, C. Merienne, A. Gaudemer and C. Giannotti, *J. Organomet. Chem.*, 1972, 38, 167–178.
- 42 L. Saussine, E. Brazi, A. Robine, H. Mimoun, J. Fischer and R. Weiss, *J. Am. Chem. Soc.*, 1985, **107**, 3534–3540.

- 43 N. Lehnert, R. Y. N. Ho, L. Que and E. I. Solomon, *J. Am. Chem. Soc.*, 2001, **123**, 12802–12816.
- 44 G. Roelfes, V. Vrajmasu, K. Chen, R. Y. N. Ho, J.-U. Rohde, C. Zondervan, R. M. la Crois, E. P. Schudde, M. Lutz, A. L. Spek, R. Hage, B. L. Feringa, E. Münck and L. Que, *Inorg. Chem.*, 2003, 42, 2639–2653.
- 45 A. Abragam and B. Bleaney, *Electron Paramagnetic Resonance of Transition Ions*, Dover Publications Inc., New York, 1986.
- 46 A.-L. Barra, D. Gatteschi, R. Sessoli, G. L. Abbati, A. Cornia, A. C. Fabretti and M. G. Uytterhoeven, *Angew. Chem., Int. Ed. Engl.*, 1997, 36, 2329–2331.
- 47 D. P. Goldberg, J. Telser, J. Krzystek, A. G. Montalban, L.-C. Brunel, A. G. M. Barrett and B. M. Hoffman, *J. Am. Chem. Soc.*, 1997, 119, 8722–8723.
- 48 J. Krzystek, J. Telser, B. M. Hoffman, L.-C. Brunel and S. Licoccia, *J. Am. Chem. Soc.*, 2001, **123**, 7890–7897.
- 49 J. Limburg, J. S. Vrettos, R. H. Crabtree, G. W. Brudvig, J. C. de Paula, A. Hassan, A.-L. Barra, C. Duboc-Toia and M.-N. Collomb, *Inorg. Chem.*, 2001, 40, 1698–1703.
- 50 The breadth of the EPR signal supports its assignment to a Mn^{II} species rather than an organic radical. We have observed Mn^{II} centers lacking resolved hyperfine structure for similar complexes, as described in ref. 26 and 27.
- 51 D. V. Avila, C. E. Brown, K. U. Ingold and J. Lusztyk, *J. Am. Chem. Soc.*, 1993, **115**, 466–470.
- 52 I. Chodák, D. Bakoš and V. Mihálov, *J. Chem. Soc., Perkin Trans.* 2, 1980, 1457–1459, DOI: 10.1039/P29800001457.
- 53 R. Oyama and M. Abe, J. Org. Chem., 2020, 85, 8627–8638.
- 54 T. Tano, K. Mieda, H. Sugimoto, T. Ogura and S. Itoh, *Dalton Trans.*, 2014, **43**, 4871–4877.
- 55 T. Tano, H. Sugimoto, N. Fujieda and S. Itoh, Eur. J. Inorg. Chem., 2012, 2012, 4099–4103.
- 56 S. Biswas, A. Mitra, S. Banerjee, R. Singh, A. Das, T. K. Paine, P. Bandyopadhyay, S. Paul and A. N. Biswas, *Inorg. Chem.*, 2019, 58, 9713–9722.
- 57 M. Sankaralingam, Y.-M. Lee, Y. Pineda-Galvan, D. G. Karmalkar, M. S. Seo, S. H. Jeon, Y. Pushkar, S. Fukuzumi and W. Nam, J. Am. Chem. Soc., 2019, 141, 1324–1336.
- 58 E. Baciocchi, M. Bietti, M. Salamone and S. Steenken, *J. Org. Chem.*, 2002, **67**, 2266–2270.
- 59 S. Kal, A. Draksharapu and L. Que, J. Am. Chem. Soc., 2018, 140, 5798–5804.
- 60 J. Serrano-Plana, F. Acuña-Parés, V. Dantignana, W. N. Oloo, E. Castillo, A. Draksharapu, C. J. Whiteoak, V. Martin-Diaconescu, M. G. Basallote, J. M. Luis, L. Que Jr, M. Costas and A. Company, *Chem.-Eur. J.*, 2018, 24, 5331– 5340.
- 61 K. Cheaib, M. Q. E. Mubarak, K. Sénéchal-David, C. Herrero, R. Guillot, M. Clémancey, J.-M. Latour, S. P. de Visser, J.-P. Mahy, F. Banse and F. Avenier, *Angew. Chem., Int. Ed.*, 2019, 58, 854–858.