

PAPER

View Article Online
View Journal | View Issue

Cite this: *Anal. Methods*, 2024, 16, 5082

Comprehensive assessment of clean-up strategies for optimizing an analytical multi-method to determine pesticides and mycotoxins in Brazilian medicinal herbs using QuEChERS-LC-TQ-MS/MS

Marlos Eduardo Zorzella Fontana,^a Rosselei Caiel da Silva,^a Ingrid Duarte dos Santos,^{ab} Júlia Paula Neu,^{id a} Robson Dias Wouters,^{id a} Paola Jennifer Babinski,^{id a} Jessica Fernanda Hoffmann,^c Rochele Cassanta Rossi,^c Liliana Essi^d and Ionara Regina Pizzutti^{id *a}

The use of medicinal herbs has increased significantly. However, the presence of pesticide residues and mycotoxins in medicinal herbs has generated constant discussion and concern among regulatory agencies. Developing and validating an analytical method for determining pesticides and mycotoxins in medicinal plants is challenging due to the naturally occurring substances in these plants. The purpose of this work was to develop and to optimize a sensitive, accurate, precise, effective QuEChERS method for simultaneous determination of over 160 pesticide and mycotoxin residues in complex medicinal plant matrices using LC-TQ-MS/MS. A comprehensive comparison of clean-up procedures and other parameters was conducted to achieve this goal. The validation procedure was performed according to SANTE 11312/2021. More polar analytes, such as acephate, methamidophos and omethoate, presented a higher negative matrix effect in both *Melissa officinalis* L. and *Malva sylvestris* L. However, other molecules, such as spirodiclofen, showed a 24% signal enhancement in *M. officinalis* and a 46% signal suppression in *M. sylvestris*, indicating that a representative matrix-matched calibration would lead to inaccurate quantification of the analyte. Accuracy and precision were satisfactory according to SANTE 11312/2021 for 157 pesticide residues and mycotoxins in *M. officinalis* and for 152 molecules in *M. sylvestris*. LOQs at 10 $\mu\text{g kg}^{-1}$ were achieved for 117 pesticides in *M. officinalis* and 99 pesticides in *M. sylvestris*. Among the mycotoxins, all four aflatoxins (B1, B2, G1 and G2) presented LOQs of 5 $\mu\text{g kg}^{-1}$, and ochratoxin A had an LOQ of 10 $\mu\text{g kg}^{-1}$ in *M. officinalis*. The same LOQ values were shown for these mycotoxins in *M. sylvestris*, except for aflatoxin B2 and ochratoxin A, which had LOQs of 20 $\mu\text{g kg}^{-1}$. Moreover, in Southern Brazil, there has been no previous study on mycotoxin and pesticide contamination in medicinal herbs. Therefore, the application of this method was assessed through the analysis of forty-two real samples. Imidacloprid was found in *M. officinalis*, and methyl pirimiphos was found in *M. sylvestris*. The proposed method not only serves as a helpful tool for routine monitoring but also offers a basis for further research on risk assessment and control in food safety.

Received 2nd April 2024
Accepted 26th June 2024

DOI: 10.1039/d4ay00599f

rsc.li/methods

Introduction

Brazil is one of the most biodiverse countries in the world, with plant species which have been widely used by the population for

medicinal purposes as well as providing material for research into the search for new drugs against different diseases.¹ During recent years, the consumption of natural resources has gained notoriety for its increase along with national policies related to traditional and complementary medicine. Medicinal plants have been an essential part of ancient healthcare practices and have become a valuable resource in the treatment of illnesses and pathologies.²

Melissa officinalis L., popularly known as lemon balm, is an edible and medicinal plant belonging to the Lamiaceae. It has been traditionally used as a sedative, analgesic, and hypnotic,³ and with its antioxidant effects being beneficial to the brain, as a treatment for memory disorders and Alzheimers.^{4,5}

^aUFSM – Federal University of Santa Maria, Chemistry Department, Center of Research and Analysis of Residues and Contaminants (CEPARC), 97105-900, Santa Maria/RS, Brazil. E-mail: ionara.pizzutti@ceparc.com.br; Tel: +55 55 3220 9458

^bUFSM – Federal University of Santa Maria, Food Science and Technology Department, 97105-900, Santa Maria/RS, Brazil

^cUNISINOS – University of Vale do Rio dos Sinos, Health School – Professional Master's in Food, Nutrition and Health, 93022-000, São Leopoldo/RS, Brazil

^dUFSM – Federal University of Santa Maria, Biology Department, 97105-900, Santa Maria/RS, Brazil

Another plant that presents therapeutic properties is *Malva sylvestris* L., known as high mallow, which is another important medicinal plant and has been considered a good candidate for drug discovery.⁶ Currently distributed worldwide, *M. sylvestris* presents anti-inflammatory properties mainly due to the presence of some flavonoids and mucilage. *M. sylvestris* has been used to treat many diseases, such as gingivitis, toothache, abdominal pain, gastrointestinal disorders, and diarrhea. In addition, its flowers are recommended for acne, the treatment of eczema, and inflammatory diseases.⁷

The growing demand for medicinal plants requires an increase in production and thus, it is necessary to protect them from pests, increase their production and shelf life whilst reducing post-harvest and storage losses. Therefore, like other plants, medicinal herbs can not only be exposed to pesticides during agricultural practices but also contaminated by mycotoxins during processing and storage.^{8,9}

According to Sedova,¹⁰ mycotoxins, pesticide residues, and toxic heavy metals are the most common chemical pollutants found in tea and medicinal herbs during production, storage, and consumption. Through eating polluted foods, chemical pollutants may cause significant health issues, such as carcinogenesis, immunosuppression, teratogenicity, as well as hepatotoxic, genotoxic, and nephrotoxic effects^{11,12} and result in huge commercial losses. For these reasons, the quality and safety of medicinal plants are of big concern¹³ and specific legislation for these matrices need to be created in order to control contamination by pesticide residues and mycotoxins. Since 2018, Brazilian legislation recommends the determination of pesticides according to RDC n° 105/2016 (ref. 14) and mycotoxins on herbal products, in all registration requests and post-registration petitions.

Several analytical methods can be used to identify and quantify this large variety of chemical compounds.¹⁵ In an effort to reduce the number of methods needed to perform a complete chemical analysis, recent trends have focused on the development of multi-residue^{16,17} and multi-class methods.^{18–20} Development of improved methods for multi-mycotoxin and multi-pesticide analysis, including sample preparation and extraction and detection parameters, has become an increasingly large research field due to co-occurrence processes while still responding to the wide range of physicochemical properties and low residue levels found in different matrices.²¹ These analyses are difficult since the analytes have varied properties and polarity. As a result, selecting the best extraction process can be difficult.^{13,22}

Different methods for multi-compound analysis have been proposed for the analysis of mycotoxins and pesticides, in which ultra-performance liquid chromatography (UHPLC) coupled with tandem mass spectrometry (MS/MS) has become the technique of choice for the analysis of a wide range of contaminants in food. It allows the simultaneous determination and accurate quantification of several analytes at very low concentrations in complex matrices in a short chromatographic run time.^{23,24} It is important to have effective and reliable analytical methods for the determination of mycotoxins and pesticides at the legislated levels in representative samples, not

only to perform accurate risk assessments, but also to enforce the regulatory limits established worldwide.²¹

A QuEChERS (quick, easy, cheap, effective, rugged and safe) method originally used just for pesticide residue analysis in vegetables and fruits^{16,17,25} has been further modified for pesticide determination in several matrices. Currently, this method is quickly becoming one of the most popular dispersive solid-phase extraction (d-SPE) methods in food safety.²⁶ Parameters such as time, solvent consumption, simplicity, selectivity, and sensitivity are crucial when considering an appropriate extraction/clean-up strategy.²¹ According to recent investigations, different types of adsorbents, such as primary secondary amines (PSA), octadecyl (C18), and graphitized carbon black (GCB) have been used based on their physical and chemical properties.^{27,28}

However, while many analytical methods have been reported for the determination of pesticides and mycotoxins in different foodstuffs,^{29,30} there is a lack of a simple and generic method for the simultaneous determination of such residues in medicinal plants due to matrices complexity as well as the diversity of species. Due to low water content, natural pigments, essential oils, and a high number of undesired components such as sugars, phenolics, and flavonoids, medicinal plants present more complicated interference when compared with other matrices, like fruits and vegetables.³¹ In addition, different species and parts of plants can affect analyte responses, making the development of analytical procedures a challenging task. Thus, it is necessary to develop a general multiclass-residue method to monitor different kinds of residues in medicinal plants, such as *M. officinalis* and *M. sylvestris*.

So far, there are no representative matrices for different medicinal parts and families, indicating that it is necessary to validate each medicinal plant separately. Additionally, even employing LC-MS/MS techniques for quantification, the present work is very significant considering that it is necessary to apply sample preparation for two distinct complex matrices whilst being able to minimize interference effects in addition to extracting with acceptable accuracy and precision the distinct classes of compounds (pesticides and mycotoxins).

The purpose of this work was to develop and optimize a sensitive, precise, effective QuEChERS method for the analysis of over 160 compounds in medicinal plant matrices by LC-MS/MS. As far as we know, the present study is the first method for simultaneous analysis of pesticides and mycotoxins in complex matrices such as *M. sylvestris* (flowers) and *M. officinalis* (leaves). In this matter, a comprehensive comparison of clean-up procedure efficiencies and other parameters were evaluated to achieve this goal. To ensure the adequate analysis of the selected mycotoxins and pesticides in medicinal plant samples, a validation process was ultimately performed for the most efficient extraction procedure. Moreover, in South Brazil, there has been no study on mycotoxin and pesticide contamination in medicinal herbs and an application of the method was assessed through the analysis of forty-two real samples. The proposed method not only works as a helpful tool for routine and surveillance monitoring but also offers a basis for further research on risk assessment and control in food safety.

Experimental

Chemicals and reagents

All reagents used were of at least analytical grade purity. Acetonitrile and acetone were obtained from Merck (Darmstadt, Germany), while methanol and toluene were purchased from Honeywell Chromasolv (Seelze, Germany). Anhydrous magnesium sulfate and sodium chloride were obtained from Êxodo Científica (São Paulo, Brazil), and formic acid from JT Baker (Deventer, Netherlands). Ultrapure water (resistivity of 18.2 MΩ cm) was obtained using a Milli-Q purification system (Millipore, Bedford, MA, USA).

Two dispersive SPE (d-SPE) kits (Agilent Technologies, Santa Clara, CA, USA) were used for clean-up purposes. These kits contained 25 mg of primary-secondary amine (PSA), 2.5 mg of graphitized carbon (GCB) and 150 mg of MgSO₄ (pigmented fruits and vegetables) (tests B and C – Table 1), or 25 mg of PSA, 7.5 mg of GCB and 150 mg of MgSO₄ (highly pigmented fruits and vegetables) (tests D and E – Table 1).

Reference standards

Reference standards of pesticides (purity > 97%) were obtained from Dr Ehrenstorfer (Augsburg, Germany), while the mycotoxin standards (purity > 98%) were obtained from Fermentek Biotechnology (Jerusalem, Israel) and Sigma-Aldrich (St. Louis, USA).

Individual stock solutions of pesticides (1000 mg L⁻¹) were prepared by dissolving the reference standards in toluene, methanol, or acetone, depending on their solubility. Similarly, individual stock solutions of mycotoxins (500 or 1000 mg L⁻¹) were prepared in acetonitrile or methanol. A standard mixture solution of 150 pesticides (1 mg L⁻¹) was prepared by diluting 100 µL of each stock solution in 100 mL of 0.1% formic acid in methanol (v/v). The 11 mycotoxins were divided into two groups based on their sensitivity in the liquid chromatographer-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS) system. Group M1 included aflatoxins B1, B2, G1, G2, and ochratoxin A, while Group M2 included diacetoxyscirpenol (DAS), deoxynivalenol (DON), fumonisins B1 and B2, T2-toxin, zearalenone (ZEN). A solution containing 0.5 mg L⁻¹ of standard mixture M1 and 25 mg L⁻¹ of M2 was prepared by appropriately diluting the stock solutions with acetonitrile.

Analytical work solutions of pesticides and mycotoxins were prepared by suitably diluting the mixture solutions with acetonitrile. All solutions were stored at -18 °C in amber glass.

LC-MS/MS

Chromatographic analysis was performed using an Agilent 1260 prime II Liquid chromatography system coupled to a triple quadrupole mass spectrometer (LC-TQ-MS/MS) (ULTIVO, Agilent technologies, USA) with an Agilent Jet Stream Technology ion source (AJS), operating in dynamic multiple reaction monitoring (dMRM) mode. Chromatographic separations were carried out on an Infinity Lab Poroshell 120 EC1-C18 (2.1 mm i.d. O 100 mm O2.7 µm) reverse phase analytical column coupled to a pre-column (UHPLC GUARD Infinity Lab Poroshell) of the same stationary phase. Water (A) and acetonitrile (B), both acidified with 0.1% (v/v) formic acid, were used as the mobile phase at a constant flow rate of 0.3 mL min⁻¹. The gradient elution program ranged from 20 to 90% B from 0 to 5 min. This condition was maintained for 4 min, then changed to 95% B from 9 to 9.25 min and maintained for 2 min. Finally, the mobile phase was changed to the initial composition from 11.25 to 14 min. The chromatographic column was maintained at 45 °C (±0.5 °C) and the injection volume was 2 µL.

All mass spectrometer parameters were optimized using the Optimizer software version 1.1 (Agilent, USA).

Sample preparation

According to the Brazilian pharmacopeia,³² the pharmacologically active parts of *M. officinalis* are the dried leaves, while for *M. sylvestris*, they are the entire or fragmented dried flowers. Therefore, all samples used as blank samples (free of pesticides and mycotoxins) were in accordance with Brazilian pharmacopeia criteria.

Commercially available organic samples identified by sellers as *M. officinalis* and *M. sylvestris* were purchased from local pharmacies in Santa Maria city, Brazil. These samples were checked for the absence of pesticides and mycotoxins before being used as blank samples for method optimization and validation.

Samples were obtained individually or in groups of packages with the same lot number, containing a minimum amount of 200 g, as recommended by sampling methods.³³ The dried

Table 1 Extraction protocols tested prior to validation

Extraction protocol	A	B	C	D	E
Slurry portion	10 g of <i>M. officinalis</i> (1 : 4 ratio) or 14 g of <i>M. sylvestris</i> (1 : 6 ratio)				
Extraction solvent	10 mL of acetonitrile		10 mL of acetonitrile + 1% formic acid		
Partitioning salts	4 g MgSO ₄ + 1 g NaCl				
Clean-up	—	1 mL upper layer to 25 mg of PSA, 2.5 mg of GCB and 150 mg of MgSO ₄	1 mL upper layer to 50 mg of PSA, 5 mg of GCB and 300 mg of MgSO ₄	1 mL upper layer to 25 mg of PSA, 7.5 mg of GCB and 150 mg of MgSO ₄	
Dilution	1 : 1				
Analysis	LC-MS/MS				

leaves and flowers were ground separately in a multiprocessor and sieved (granulometry 1 μm). Before the extraction procedure, the samples were hydrated for 30 min with ultrapure water 1:4 and 1:6 (w/w) for *M. officinalis* and *M. sylvestris*, respectively, at 8 °C, forming a slurry.

Extraction procedure

The extraction procedure employed was a modification of the QuEChERS method using the highly pigmented fruits and vegetables clean-up kit from Agilent Technologies. A slurry of 10 g (1:4 ratio) of *M. officinalis* leaves or 14 g (1:6 ratio) of *M. sylvestris* flowers was weighed in a 50 mL PTFE centrifuge tube. Subsequently, 10 mL of acetonitrile acidified with 1% formic acid was added, along with 40 μL of propoxur, which served as the internal standard solution. It should be noted that the concentration of propoxur in this study was 20 ng mL^{-1} . The tubes were shaken using an automatic mechanical shaker (Orbital Shaker 3016, Gesellschaft für Labortechnik mbH, Germany) for 1 minute. Following this, 4 g of magnesium sulfate and 1 g of sodium chloride were added, and the samples were vortexed for an additional 1 minute. The extracts were then centrifuged at 4000 rpm for 4 minutes, and 1 mL of the supernatant was transferred to a dispersive clean-up kit. After homogenizing the tubes in a vortex for 1 minute, they were centrifuged again (4000 rpm, 4 minutes), and 0.5 mL of the extract was transferred to a vial and diluted with 0.5 mL of acetonitrile/water (1:1, v/v) containing the injection internal standard solution of PCB-153 at a concentration of 100 ng mL^{-1} .

To develop a fast extraction protocol that causes less damage to the chromatographic system, which is robust and reliable, and still presents acceptable recovery rates, five preliminary studies were conducted to evaluate the accuracy, precision and matrix effects. For all tests, the slurries of *M. officinalis* and *M. sylvestris* samples were spiked ($n = 3$) at two different levels with pesticides (10 and 70 $\mu\text{g kg}^{-1}$) and mycotoxins (group 1: 2 and 20 $\mu\text{g kg}^{-1}$; group 2: 100 and 1000 $\mu\text{g kg}^{-1}$), simultaneously.

Solvent extraction evaluation

Since the proposed method aims to extract a variety of target analytes with different polarities, pK_{a} , and other chemical properties, two approaches were tested to evaluate the recovery rate of analytes. The first approach used pure acetonitrile according to the original QuEChERS method. The second approach employed acetonitrile acidified with 1% (v/v) formic acid to improve recovery, especially for mycotoxins.

Sorbent evaluation for clean-up

The absence of and different proportions of dispersive solid-phase extraction (d-SPE) sorbents (Table 1) were tested for selectivity, sensitivity, reliability, acceptable accuracy and precision, and to achieve less damage to the chromatographic system. Mixtures of primary secondary amine (PSA) and graphitized carbon black (GCB) were tested to remove pigments (mostly chlorophyll), sugars, lipids, flavonoids, acids, and carotenoids.³⁴

Method performance

Analyte identification and confirmation were conducted according to SANTE document 11312/2021,³⁵ including retention time standard (± 0.1 min), and at least two product ions with fully overlapping peaks and ion ratio within $\pm 30\%$.

Method validation

A validation protocol in accordance with SANTE document 11312/2021 (ref. 35) was conducted for the simultaneous determination of pesticides and mycotoxins in *M. officinalis* and *M. sylvestris*. The analytical method validation assessed the following parameters: sensitivity, selectivity, linearity of the analytical curves, matrix effects, trueness (expressed as recovery percentage), precision as repeatability RSD_{r} and reproducibility (RSD_{WR}), limit of detection (LOD), and limit of quantification (LOQ).

For linearity, sensitivity, and matrix effect evaluation, seven different solutions for each concentration were prepared. For pesticides and mycotoxins of group 1, the concentrations of the solutions were 0.1, 0.5, 1, 5, 10, 25, 50 and 100 ng mL^{-1} . For mycotoxins of group 2 the concentrations of analytical solutions prepared in neat organic solvent (acetonitrile) and in blank *M. officinalis* and *M. sylvestris* extracts were 5, 25, 50, 250, 500, 1250, 2500 and 5000 ng mL^{-1} . Each solution was injected seven times.

The LOD was considered the lowest concentration level, injected repeatedly, obtained from 7 injections of an analytical solution prepared in blank matrix extract with a signal-to-noise ratio (S/N) ≥ 3 . The LOQ was considered the lowest concentration level spiked with acceptable accuracy (70–120%) and precision ($\text{RSD} \leq 20\%$) obtained by the proposed analytical method.

Spiking/recovery experiments were performed by two different analysts on two different days to evaluate method reproducibility (RSD_{WR}). Matrix effects were calculated as described by Dias *et al.*³⁶ For accuracy (trueness and precision), recovery experiments were conducted by spiking blank *M. officinalis* and *M. sylvestris* at concentration levels of 10, 20, 50, and 70 $\mu\text{g kg}^{-1}$ for pesticides; 2, 5, 10, and 20 $\mu\text{g kg}^{-1}$ for mycotoxins of group 1; and 100, 250, 500, and 1000 $\mu\text{g kg}^{-1}$ for mycotoxins of group 2. Seven replicates for each spiked level ($n = 7$) were performed by each analyst on two different days, totaling fourteen replicates ($n = 14$). All samples were extracted as mentioned in the section 'Extraction Procedure'.

Repeatability (RSD_{r}) was calculated for each analyst from recovery experiments performed using the same extraction protocol, quantification method, system, and blank sample on the same day. Reproducibility (RSD_{WR}) was obtained *via* intermediate precision assessment by executing the same recovery experiments with different analysts, with a one-week interval between recovery experiments.

Sampling

The medicinal herb samples were obtained from the Public Market in Porto Alegre city, Rio Grande do Sul State, Brazil, due to the commercialization, consumer turnover, and location.

The samples were collected from 10 commercial stores between May 2021 and July 2022. Each sample consisted of at least 200 g of medicinal herbs, comprising 23 samples of *M. officinalis* leaves and 19 samples of *M. sylvestris* flowers, totaling 42 samples over the course of the study.

Results and discussion

Over the years, with the rise in food inspection and the escalating demand for quality control analyses, coupled with the need for promptly delivering results, multianalyte methods have garnered attention for their ability to analyze a diverse range of substances in a single operation. Methods enabling the simultaneous detection of pesticides and mycotoxins are available for various matrices, including fruits,³⁷ cereals,^{38–40} wine,⁴¹ eggs,⁴² feed,^{15,43,44} raw coffee,⁴⁵ and even some teas,^{18,46} spices, medicinal herbs,⁴⁷ and infant milk formulae.⁴⁸

While there are methods available for analyzing teas and spices, these primarily focus on green and black teas. Additionally, there is currently no validated method for the simultaneous analysis of mycotoxins and pesticides in medicinal herbs, specifically comparing the dried flowers of *M. sylvestris* and the dried leaves of *M. officinalis*.

Clean-up optimization

Each matrix submitted to an extraction protocol for the analysis of residues and contaminants must undergo an optimization process. This optimization improves the selectivity of the analytes, reduces the matrix effect, and achieves quantification limits at low concentration levels while maintaining the accuracy and precision required in an analytical method.

Medicinal herbs are particularly challenging matrices containing various extractable compounds such as pigments, essential oils, and flavonoids that may cause notable matrix effects in chromatographic analysis. The concern is not only about signal suppression caused by the co-extracts but also the potential damage caused to the systems, reducing the overall lifespan of the consumables. In addition, in long injection

sequences, dirt can accumulate in the ionization source, decreasing the detectability along the sequence. Thus, a sample injected at the beginning and at the end of the sequence can present significant deviations in results, decreasing the accuracy and precision of the method.³⁴

To improve method performance, different sorbent quantities were compared *via* recovery experiments, applying the following spike levels for mycotoxins groups: group 1: 2 and 20 $\mu\text{g kg}^{-1}$; group 2: 100 and 1000 $\mu\text{g kg}^{-1}$; and for pesticides: 10 and 70 $\mu\text{g kg}^{-1}$, $n = 3$.

No clean-up step and two d-SPE kits (25 mg PSA + 2.5 mg GCB + 150 mg MgSO_4 (tests B and C), and 25 mg PSA, 7.5 mg GCB + 150 mg of MgSO_4 (tests D and E)) were tested (Table 1). The results are shown in Fig. 1 and 2, respectively, for the mycotoxins and pesticides. The concentration levels 1 and 2 were, respectively, 10 and 50 $\mu\text{g kg}^{-1}$ for the pesticides; 2 and 10 $\mu\text{g kg}^{-1}$ for the mycotoxins of group 1; and 100 and 500 $\mu\text{g kg}^{-1}$ for the mycotoxins of group 2.

When no clean-up step was conducted, a highly pigmented extract was obtained for both matrices, causing the extensive deposition of co-extracts in the ion source, decreasing precision and causing a significant loss in detectability within the same injection sequence.

To efficiently remove pigment interferences from the extracts, graphitized carbon black (GCB) is a worthy option. However, it might also retain specific analytes, such as aromatic compounds and/or planar pesticides, due to π - π interactions.⁴⁹ To mitigate this problem, small quantities of GCB were tested (2.5, 5 and 7.5 mg), with the latter being able to remove enough pigment while maintaining acceptable method accuracy and precision.

In this study, the final combination of PSA (25 mg) and GCB (7.5 mg) plus 150 mg of MgSO_4 was the most effective for removing matrix co-extracts while maintaining acceptable recoveries and avoiding significant damage to the LC-TQ-MS/MS system. For instance, cyprodinil presented recoveries ranging from 71% to 83% and proper precision ($\text{RSD} < 18\%$) despite the use of GCB. These results were also verified by Ly *et al.*⁵⁰ who used GCB in green tea extraction and obtained




Fig. 1 Number of mycotoxins presenting recoveries within the range of 70–120% in assays A, B, C, D and E, for *M. officinalis* and *M. sylvestris*.




Fig. 2 Number of pesticides presenting recoveries within the range of 70–120% in assays A, B, C, D and E, for *M. officinalis* and *M. sylvestris*.




Fig. 3 Mycotoxin Aflatoxin B1 and pesticide Fenamiphos chromatograms obtained by analysis of: (a)(f) *M. sylvestris* blank extract, (b)(g) melissa blank extract, (c)(h) 1 ng mL^{-1} analytical solution in organic solvent, (d)(i) 1 ng mL^{-1} analytical solution in malva blank extract, (e)(j) 1 ng mL^{-1} analytical solution in *M. officinalis* blank extract.

satisfactory results for this pesticide. Fig. 3 represents a total ion chromatogram of two injections of the same vial, at the beginning and the end of a work list of over 100 injections and 16 h difference between those two injections, of fenamiphos and aflatoxin B1. No significant loss in precision was verified when comparing those two injections of both analytes.

Sample preparation optimization

Furthermore, the wide range of polarities, acidities and solubilities of pesticides and mycotoxins makes it challenging to develop and validate an appropriate analytical method for simultaneous determination. Additionally, representative food matrices belonging to the same food group (SANTE 11312/2021)

Table 2 Linear range, matrix effect (ME), LOD and LOQ for all analytes in *M. officinalis* and *M. sylvestris*

Pesticide/Mycotoxin	<i>Melissa officinalis</i>				<i>Malva sylvestris</i>			
	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)
Acephate	5–1000	−74	5	10	5–1000	−80	1	20
Acetamiprid	5–1000	−10	5	10	1–1000	50	1	10
Acetochlor	10–1000	−5	5	20	10–1000	−16	5	10
Aflatoxin B1^a	5–500	−70	1	5	5–500	−20	1	5
Aflatoxin B2^a	5–500	−71	3	5	10–500	−17	1	20
Aflatoxin G1^a	5–500	−53	3	5	5–1000	31	4	5
Aflatoxin G2^a	5–500	−12	3	5	5–1000	9	1	5
Aldicarb sulfone	1–500	−24	5	10	5–1000	−29	1	10
Aldicarb sulfoxide	5–500	−77	1	10	5–500	−76	5	10
Atrazine	5–1000	−17	1	10	5–500	−10	1	10
Azamethiphos	5–500	0	1	10	5–1000	−4	1	10
Azinphos-methyl	10–500	23	5	10	50–1000	−15	1	50
Azoxystrobin	5–500	55	1	10	10–500	29	1	10
Bifenazate	10–500	11	1	10	10–500	11	1	n.f.r. ^b
Bitertanol	5–1000	20	1	10	5–1000	−12	1	20
Boscalid	10–500	−8	1	10	5–500	−10	1	10
Bupirimate	5–500	−26	1	10	5–500	−21	1	10
Buprofezin	5–500	−32	1	10	5–500	−37	1	10
Cadusafos	10–500	8	1	10	5–1000	47	1	10
Carbaryl	5–500	−22	1	10	5–1000	−18	1	10
Carbendazin	5–500	−67	1	10	10–1000	−75	1	70
Carbofuran	5–500	−11	1	10	10–500	7	1	10
Carpropamid	5–500	−13	5	10	5–500	−28	1	20
Chlorantraniliprol	5–500	−5	5	20	5–1000	6	5	20
Chlorfenvinphos	10–500	13	1	10	5–500	3	5	10
Chlorpyrifos	5–500	−3	5	10	5–1000	−19	5	10
Clofentezine	10–1000	−25	8	10	10–1000	−7	8	10
Clomazone	10–500	−14	1	10	5–1000	−13	1	10
Clothianidin	10–1000	−14	8	10	5–1000	−2	5	10
Cyazofamid	50–500	−29	5	10	10–500	−36	5	20
Cyproconazol	10–500	7	5	10	5–1000	−14	5	10
Cyprodinil	5–500	−48	5	10	10–500	−48	5	10
Demeton-S-methyl sulfone	5–1000	9	5	10	5–1000	21	1	10
Demeton-S-methyl sulfoxide	10–1000	−72	5	10	5–1000	−71	1	20
Deoxynivalenol^a	250–5000	−67	50	250	250–5000	−65	250	500
Diacetoxyscirpenol^a	250–5000	−15	25	250	500–2500	21	250	500
Diazinon	10–500	−12	1	10	5–1000	2	1	10
Diethofencarb	10–500	17	1	10	50–1000	n.f.r. ^b	1	n.f.r. ^b
Difenoconazole	5–1000	−15	1	10	10–1000	−7	1	10
Diffubenzuron	50–1000	−18	10	n.f.r. ^b	50–1000	15	10	70
Dimethoate	1–1000	−23	1	10	1–1000	−3	1	10
Dimethomorph	10–500	42	1	10	50–500	50	1	10
Diniconazol	5–500	−8	5	10	5–1000	−10	1	10
Diphenylamine	5–1000	−17	1	10	5–500	−34	5	10
Diuron	10–500	−42	5	10	5–1000	−8	1	10
DMST	10–1000	−7	10	50	10–1000	5	10	50
Epoxiconazol	10–1000	−4	1	10	10–1000	15	1	10
Ethion	10–500	58	1	10	10–500	−5	1	10
Ethiprole	10–500	−15	1	10	10–500	−7	5	10
Ethoprophos	10–500	10	1	10	5–500	−7	5	10
Etofenprox	10–500	−10	1	10	5–1000	−6	1	10
Etoxazol	10–1000	−19	1	10	5–1000	−44	1	10
Fenamidone	5–500	3	1	10	5–500	5	1	10
Fenamiphos	10–500	78	1	10	5–1000	29	1	20
Fenarimol	5–1000	−17	5	20	5–500	15	1	50
Fenazaquin	5–500	30	1	10	5–1000	−29	1	10
Fenbuconazol	5–500	−2	1	10	5–500	7	5	20
Fenhexamid	10–500	−37	1	10	10–500	9	1	20
Fenobucarb	10–500	−19	1	10	10–500	−32	1	10
Fenoxycarb	10–500	−31	5	20	10–1000	−40	1	20

Table 2 (Contd.)

Pesticide/Mycotoxin	Melissa officinalis				Malva sylvestris			
	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)
Fenpropimorph	5–1000	–13	1	10	5–1000	–14	1	10
Fenpyroximate	5–500	–4	1	10	5–1000	–16	1	10
Fensulfothion	50–1000	39	5	50	50–1000	23	10	50
Fluazifop-butyl	10–500	–5	1	10	5–1000	7	1	10
Fludioxonil	5–1000	–17	5	10	10–500	–20	1	10
Flufenoxuron	10–500	–42	1	10	50–1000	–24	20	50
Fluquinconazol	10–1000	–25	10	20	5–500	–25	1	20
Flusilazol	5–500	–8	1	10	5–500	–26	1	10
Flutolanil	50–1000	–2	1	50	5–1000	–10	1	20
Flutriafol	10–1000	–9	8	10	5–500	2	1	10
Fosthiazate	10–1000	7	1	10	5–1000	–6	1	10
Fumonisin B1^a	250–5000	51	250	500	250–5000	–1	250	n.f.r.^b
Fumonisin B2^a	250–5000	–16	250	500	250–5000	–31	500	n.f.r.^b
Furalaxyl	10–500	18	1	10	5–500	20	1	20
Furathiocarb	5–1000	8	1	10	5–1000	4	1	10
Halofenozide	50–1000	–23	10	50	10–1000	–33	10	n.f.r. ^b
Haloxypop-2-ethoxyethyl	5–1000	–8	1	10	5–1000	–41	1	n.f.r. ^b
Hexaconazol	5–1000	–16	5	10	5–1000	–31	5	10
Hexytiazox	5–1000	–6	1	10	1–1000	–42	1	10
Imazalil	10–1000	–26	10	20	10–1000	–22	5	10
Imazapic	5–500	–32	5	10	10–500	5	5	10
Imazetapyr	5–1000	3	1	10	5–1000	35	5	10
Imidacloprid	5–1000	31	1	10	10–1000	98	8	10
Indoxacarb	5–1000	34	1	20	5–1000	97	5	10
Iprovalicarb	5–1000	20	1	10	5–1000	–6	1	10
Isoxaflutole	5–1000	–30	5	50	50–1000	–20	10	50
Kresoxim-methyl	5–1000	–16	1	10	5–1000	–16	5	20
Linuron	50–1000	–40	10	50	50–1000	–32	20	50
Lufenuron	10–500	–58	8	10	5–500	–84	5	10
Malathion	5–1000	14	1	10	5–1000	11	5	20
Mecarbam	5–1000	5	5	10	5–1000	4	1	10
Mepanipyrim	5–500	–44	1	20	10–1000	–17	10	50
Metalaxyl	5–1000	23	1	20	10–1000	33	1	10
Metconazole	5–1000	–75	5	20	5–1000	–29	5	20
Methamidophos	50–1000	–77	10	50	10–1000	–76	5	70
Methidathion	10–1000	–5	10	20	50–1000	–31	5	n.f.r. ^b
Methiocarb	5–1000	–33	5	20	5–1000	–30	5	20
Methomyl	5–1000	–10	1	10	5–500	–21	1	10
Methoxyfenozide	5–1000	31	1	10	5–500	15	1	20
Monocrotophos	1–1000	–62	1	10	10–1000	–57	1	20
Myclobutanil	5–1000	–45	1	10	5–1000	–1	1	10
Nitenpyram	10–1000	–76	5	50	50–1000	–82	1	70
Ochratoxin A^a	10–1000	20	8	10	10–1000	6	10	20
Ofurace	10–500	46	1	10	5–1000	59	1	10
Omethoate	5–1000	–76	1	10	n.f.r. ^b	–76	n.f.r. ^b	n.f.r. ^b
Oxadixyl	10–500	5	5	10	5–1000	–10	1	10
Oxamyl	5–1000	–20	1	10	5–1000	–35	1	10
Paclobutrazol	10–1000	2	10	20	5–500	–4	1	10
Penconazole	5–500	–22	1	10	10–1000	–29	1	10
Pencycuron	5–1000	6	1	10	5–1000	64	1	10
Pendimethalin	5–500	–4	5	10	5–1000	–21	1	10
Phenothrin	50–1000	–19	20	50	50–1000	2	20	50
Phenthoate	5–500	–28	5	10	5–500	–21	1	10
Phosalone	50–1000	–4	10	50	50–1000	2	20	50
Phosmet	50–1000	–8	5	50	5–1000	–21	5	20
Picoxystrobin	5–500	–18	1	10	10–1000	–22	1	10
Piperonyl butoxide	5–1000	–25	1	10	5–1000	–27	1	10
Pirimicarb	5–1000	–21	1	10	5–1000	–17	1	10
Pirimiphos-ethyl	1–1000	–15	1	10	5–1000	–25	1	10
Pirimiphos-methyl	5–1000	–20	1	10	5–1000	–15	1	10

Table 2 (Contd.)

Pesticide/Mycotoxin	Melissa officinalis				Malva sylvestris			
	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)	Linear range ($\mu\text{g kg}^{-1}$)	ME (%)	LOD ($\mu\text{g kg}^{-1}$)	LOQ ($\mu\text{g kg}^{-1}$)
Prochloraz	1–1000	11	1	10	5–1000	6	1	10
Profenofos	5–500	–29	1	10	5–1000	–45	1	20
Prometryn	5–1000	–22	1	10	5–1000	–20	1	10
Propamocarb	10–500	–36	1	n.f.r. ^b	10–500	–61	1	70
Propanil	1–1000	–26	10	50	5–1000	–21	5	50
Prophan	10–1000	–33	8	10	50–1000	–8	20	50
Propiconazol	5–500	–13	1	10	5–500	5	1	10
Propyzamide	10–1000	–19	8	10	5–1000	22	5	10
Pyraclostrobin	5–1000	–27	1	10	5–1000	–25	1	10
Pyrazophos	10–500	35	5	10	5–1000	50	1	10
Pyridaben	5–1000	–9	1	10	5–1000	–58	1	10
Pyrimethanil	5–500	–38	1	10	5–1000	–31	1	10
Pyriproxyfen	5–500	–24	1	10	5–1000	–30	1	10
Quinalphos	1–1000	–42	1	10	5–500	–18	5	10
Quinoxifen	1–1000	–16	1	10	5–1000	–15	1	10
Simazine	5–500	–35	1	10	10–1000	–23	5	10
Spinosyn A	5–1000	–39	1	10	5–1000	–25	1	20
Spinosyn D	5–1000	–35	5	10	5–1000	–35	5	10
Spirodiclofen	5–500	24	5	50	5–500	–46	5	10
Spiromesifen	5–500	19	5	20	10–1000	–58	10	20
Spiroxamine	1–1000	–11	1	10	1–1000	–11	1	10
Tau-fluvalinate	50–1000	19	50	70	50–1000	–55	10	50
Tebuconazol	5–500	8	5	10	5–500	14	1	10
Tebufenozide	5–1000	–12	1	20	5–500	–14	1	10
Tebufenpyrad	5–500	–15	5	10	5–1000	–35	1	10
Terbutryn	5–1000	–37	1	10	5–1000	–20	1	10
Tetrachlorvinphos	10–1000	–13	10	50	50–1000	–58	10	50
Tetraconazole	5–500	0	5	10	10–1000	–4	5	10
Tetramethrin	10–1000	–4	8	10	10–500	–13	5	10
Thiacloprid	1–500	–8	1	10	5–500	2	1	10
Thiamethoxam	5–1000	–7	5	70	5–500	17	1	10
Thiodicarb	5–500	–2	5	10	5–1000	55	1	n.f.r. ^b
Toxin T2^a	50–2500	–12	25	500	250–50 000	3	5	1000
Triadimefon	10–1000	9	8	10	10–1000	49	1	10
Triadimenol	10–1000	21	8	n.f.r. ^b	1–500	23	1	70
Triazophos	5–500	20	1	10	10–1000	–22	1	20
Trifloxystrobin	5–500	–8	1	10	5–1000	–45	1	10
Triflumizol	5–1000	–40	1	10	10–1000	–40	1	10
Triticonazol	1–1000	4	1	10	5–500	–6	1	10
Zearalenone^a	25–2500	–31	25	250	250–25 000	–35	25	500
Zoxamide	5–500	–19	5	10	5–500	–21	5	10

^a Mycotoxins. ^b n.f.r.: not fulfill requirements of SANTE document.

are often used for optimization of time, reagents, and other parameters. However, in the case of dry medicinal plants, which contain a larger number of secondary metabolites (such as flavonoids, saponins and alkaloids), using a single representative matrix may present weaknesses in quantification due to differences in analytical signal suppression and enhancement in the LC-TQ-MS/MS system.

For matrices with low water content, it is recommended to add water to increase the extraction efficiency. Therefore, a slurry was prepared with cold water (8 °C) for matrix rehydration (≈ 30 minutes) to facilitate the extraction of the analytes

and prevent matrix components from being extracted and interfering with the instrumental analysis.

Analytical method validation

The validation data summarized in Table 2 show the linear range, matrix effect, LOQ, and LOD. Tables 3 and 4 demonstrate recoveries, precision (RSD_r) and intermediate precision (RSD_{WR}) for *M. officinalis* and *M. sylvestris* obtained from the method validation procedure for all spike levels studied.

For all pesticides and mycotoxins, the criterion for linearity was $r^2 \geq 0.99$ and the deviation of back-calculated

Table 3 Average recoveries, precision (RSD_{WR}) and intermediate precision (RSD_{WR}) obtained for *M. officinalis* from the method validation procedure

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%) (n = 7))		Average recovery (%) (RSD _{WR} (%) (n = 14))		Average recovery (%) (RSDr (%) (n = 7))		Average recovery (%) (RSD _{WR} (%) (n = 14))		Average recovery (%) (RSDr (%) (n = 7))		Average recovery (%) (RSD _{WR} (%) (n = 14))		Average recovery (%) (RSDr (%) (n = 7))		Average recovery (%) (RSD _{WR} (%) (n = 14))	
	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2
	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value	P value
Acephate	82 (16)	76 (15)	79 (15)	74 (11)	82 (8)	74 (11)	78 (11)	81 (14)	81 (5)	80 (20)	81 (14)	89 (16)	91 (8)	85 (10)	88 (9)	0.109
Acetamidiprid	99 (15)	85 (7)	92 (14)	76 (6)	87 (14)	76 (6)	81 (13)	89 (16)	0.075	95 (16)	83 (11)	89 (16)	0.102	83 (16)	79 (5)	0.496
Acetochlor	90 (27)	81 (12)	86 (22)	91 (14)	78 (12)	84 (15)	88 (15)	88 (15)	0.139	88 (14)	87 (16)	88 (15)	0.926	87 (9)	82 (15)	0.526
Aflatoxin B1 ^a	60 (60)	114 (26)	87 (48)	95 (19)	95 (19)	97 (17)	97 (17)	96 (17)	0.785	88 (16)	104 (13)	96 (17)	0.112	84 (17)	97 (8)	0.138
Aflatoxin B2 ^a	77 (19)	31 (0)	54 (47)	95 (15)	83 (12)	95 (15)	89 (15)	90 (13)	0.196	87 (18)	92 (8)	90 (13)	0.406	83 (7)	89 (8)	0.056
Aflatoxin G1 ^a	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	83 (18)	86 (19)	83 (18)	84 (18)	87 (8)	0.543	85 (9)	88 (6)	87 (8)	0.418	91 (14)	99 (5)	0.083
Aflatoxin G2 ^a	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	79 (6)	102 (19)	79 (6)	90 (20)	89 (16)	0.035	97 (15)	81 (11)	89 (16)	0.093	80 (12)	73 (6)	0.060
Aldicarb sulfone	98 (15)	91 (14)	95 (15)	91 (11)	98 (8)	91 (11)	95 (10)	113 (16)	0.064	106 (5)	113 (16)	110 (12)	0.361	103 (14)	106 (11)	0.547
Aldicarb sulfoxide	94 (19)	81 (15)	87 (18)	75 (16)	83 (13)	75 (16)	79 (14)	77 (11)	0.334	80 (4)	75 (5)	77 (11)	0.224	81 (4)	82 (8)	0.762
Arazine	100 (10)	89 (12)	95 (12)	83 (11)	91 (8)	83 (11)	87 (10)	83 (10)	0.083	88 (4)	78 (12)	83 (10)	0.019	83 (9)	76 (9)	0.052
Azamephos	85 (15)	75 (3)	80 (13)	75 (7)	87 (11)	75 (7)	81 (12)	82 (17)	0.060	86 (12)	78 (22)	82 (17)	0.217	96 (7)	84 (15)	0.155
Azinphos-methyl	88 (16)	81 (16)	85 (16)	102 (6)	92 (18)	102 (6)	97 (13)	77 (7)	0.196	80 (8)	74 (4)	77 (7)	0.141	96 (14)	105 (4)	0.087
Azoxystrobin	95 (13)	99 (6)	97 (10)	81 (8)	91 (9)	81 (8)	86 (10)	81 (14)	0.060	85 (10)	78 (18)	81 (14)	0.236	83 (18)	71 (5)	0.091
Bifenazate	78 (9)	86 (9)	82 (10)	91 (1)	84 (16)	91 (1)	88 (11)	71 (6)	0.283	72 (3)	69 (8)	71 (6)	0.260	86 (15)	99 (8)	0.072
Bitertanol	85 (13)	81 (16)	83 (14)	77 (15)	85 (13)	77 (15)	81 (14)	89 (14)	0.076	91 (7)	87 (20)	89 (14)	0.490	90 (8)	96 (20)	0.480
Boscalid	96 (12)	107 (12)	101 (13)	99 (5)	94 (7)	99 (5)	96 (6)	84 (9)	0.256	87 (3)	80 (11)	84 (9)	0.108	83 (8)	80 (20)	0.764
Bupirimate	98 (9)	91 (11)	94 (10)	84 (15)	89 (7)	84 (15)	86 (12)	82 (13)	0.303	85 (3)	80 (18)	82 (13)	0.409	80 (8)	74 (10)	0.117
Buprofezin	90 (11)	79 (19)	85 (16)	76 (19)	86 (5)	76 (19)	81 (14)	79 (13)	0.153	83 (2)	75 (18)	79 (13)	0.177	74 (15)	86 (16)	0.196
Cadusafos	96 (10)	98 (8)	97 (9)	101 (4)	91 (6)	101 (4)	96 (12)	82 (13)	0.053	87 (5)	77 (16)	82 (13)	0.141	87 (9)	83 (16)	0.496
Carbaryl	93 (11)	78 (18)	85 (16)	83 (7)	89 (11)	83 (7)	86 (10)	81 (12)	0.197	85 (10)	77 (11)	81 (12)	0.061	78 (18)	87 (2)	0.159
Carbendazim	79 (9)	72 (14)	75 (12)	77 (19)	74 (11)	77 (19)	76 (15)	76 (14)	0.517	72 (15)	80 (11)	76 (14)	0.137	71 (6)	74 (6)	0.35
Carbofuran	111 (9)	100 (13)	106 (12)	99 (8)	111 (11)	99 (8)	105 (11)	113 (13)	0.073	118 (8)	108 (16)	113 (13)	0.080	113 (11)	103 (5)	0.083
Carpropamid	84 (18)	94 (17)	89 (18)	82 (14)	82 (14)	82 (14)	88 (18)	89 (14)	0.161	83 (12)	96 (12)	89 (14)	0.115	97 (13)	103 (12)	0.387
Chlorantraniliprole	56 (41)	39 (37)	48 (43)	86 (19)	82 (18)	86 (19)	84 (18)	90 (17)	0.579	96 (15)	85 (19)	90 (17)	0.240	92 (9)	77 (17)	0.074
Chlorfenvinphos	92 (18)	108 (10)	100 (16)	101 (4)	94 (13)	101 (4)	97 (10)	80 (7)	0.156	80 (9)	79 (5)	80 (7)	0.749	85 (16)	85 (5)	0.931
Chlorpyrifos	101 (15)	92 (19)	96 (17)	97 (20)	97 (20)	106 (19)	102 (19)	74 (9)	0.335	76 (10)	72 (8)	74 (9)	0.336	84 (12)	79 (8)	0.272
Clofentezine	83 (17)	99 (7)	91 (15)	100 (9)	100 (9)	110 (14)	105 (13)	75 (13)	0.170	78 (14)	73 (10)	75 (13)	0.370	91 (15)	79 (10)	0.054
Clomazone	88 (15)	80 (6)	84 (12)	90 (8)	83 (7)	83 (7)	87 (9)	81 (12)	0.060	85 (5)	77 (16)	81 (12)	0.216	83 (16)	75 (15)	0.388
Clothianidin	89 (18)	77 (8)	83 (16)	85 (22)	98 (7)	85 (22)	91 (16)	93 (14)	0.145	99 (11)	87 (14)	93 (14)	0.07	90 (11)	79 (9)	0.134
Cyazofamid	103 (16)	100 (14)	101 (15)	76 (10)	83 (13)	76 (10)	79 (12)	82 (15)	0.132	88 (9)	76 (19)	82 (15)	0.172	95 (8)	82 (19)	0.095
Cyproconazole	102 (10)	105 (10)	103 (10)	81 (10)	82 (12)	81 (10)	81 (10)	84 (10)	0.808	87 (4)	80 (13)	84 (10)	0.159	83 (13)	72 (5)	0.057
Cyprodinil	83 (10)	79 (4)	81 (8)	72 (13)	77 (6)	72 (13)	75 (10)	73 (9)	0.407	71 (6)	75 (10)	73 (9)	0.476	73 (4)	79 (8)	0.063
Demeton-S-methyl sulfone	103 (17)	95 (3)	99 (13)	89 (12)	96 (8)	89 (12)	93 (11)	89 (13)	0.176	94 (4)	83 (18)	89 (13)	0.095	96 (20)	80 (9)	0.075
Demeton-S-methyl sulfoxide	86 (9)	97 (17)	91 (15)	78 (4)	90 (18)	78 (4)	84 (16)	77 (16)	0.083	79 (12)	75 (20)	77 (16)	0.455	79 (12)	73 (11)	0.359
Deoxynivalenol ^a	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	101 (16)	89 (9)	101 (16)	95 (15)	91 (12)	0.092	86 (14)	97 (5)	91 (12)	0.077	84 (9)	78 (2)	0.054
Diactoxyscirpenol ^a	67 (17)	128 (161)	98 (148)	90 (1)	85 (15)	90 (1)	88 (11)	79 (8)	0.353	81 (11)	76 (1)	79 (8)	0.144	97 (10)	99 (9)	0.697
Diazinon	95 (10)	101 (8)	98 (10)	99 (10)	94 (5)	99 (10)	97 (8)	88 (7)	0.322	92 (4)	84 (6)	88 (7)	0.052	87 (8)	84 (6)	0.467

Table 3 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Average recovery (%) (RSD _{wr} (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSD _{wr} (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSD _{wr} (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSD _{wr} (%))	
	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2
	91 (14)	79 (9)	85 (14)	85 (10)	80 (7)	85 (10)	0.124	0.127	87 (7)	88 (7)	88 (7)	88 (7)	81 (15)	88 (7)	0.648	81 (15)
Diethofencarb	94 (16)	81 (13)	87 (16)	87 (9)	83 (3)	87 (9)	0.073	0.051	81 (6)	72 (15)	76 (12)	76 (12)	80 (12)	73 (1)	0.094	80 (12)
Difenoconazole	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r.	n.f.r.
Diflubenzuron	103 (16)	87 (10)	95 (16)	100 (11)	91 (9)	95 (11)	0.073	0.180	98 (9)	85 (13)	91 (13)	91 (13)	89 (17)	88 (3)	0.091	89 (17)
Dimethoate	96 (20)	81 (15)	89 (19)	101 (9)	94 (12)	98 (11)	0.059	0.185	87 (15)	77 (12)	82 (15)	82 (15)	91 (17)	81 (17)	0.110	91 (17)
Dimethomorph	89 (19)	78 (9)	83 (16)	159 (8)	83 (13)	86 (11)	0.159	0.141	86 (8)	76 (11)	81 (11)	81 (11)	83 (14)	72 (8)	0.071	83 (14)
Diniconazole	86 (19)	83 (1)	84 (13)	590 (8)	78 (16)	83 (17)	0.590	0.196	90 (16)	78 (15)	84 (17)	84 (17)	82 (11)	76 (14)	0.112	82 (11)
Diphenylamine	83 (19)	81 (10)	82 (15)	687 (1)	72 (12)	76 (13)	0.687	0.220	78 (12)	74 (15)	76 (13)	76 (13)	82 (12)	70 (11)	0.567	82 (12)
Diuron	64 (57)	90 (21)	77 (40)	188 (3)	62 (27)	75 (36)	0.188	0.072	97 (16)	83 (14)	90 (17)	90 (17)	107 (13)	93 (15)	0.149	107 (13)
DMST	113 (19)	93 (7)	103 (17)	058 (2)	82 (4)	88 (16)	0.058	0.148	86 (9)	74 (19)	80 (16)	80 (16)	89 (8)	81 (13)	0.116	89 (8)
Epoxiconazole	84 (16)	79 (15)	81 (15)	502 (3)	95 (8)	89 (15)	0.502	0.130	77 (11)	78 (0)	77 (8)	77 (8)	84 (16)	78 (0)	0.831	84 (16)
Ethion	95 (14)	97 (6)	96 (11)	689 (0)	103 (17)	97 (18)	0.689	0.238	90 (17)	93 (7)	92 (13)	92 (13)	91 (10)	82 (19)	0.632	91 (10)
Ethoprophos	101 (13)	86 (7)	94 (13)	060 (9)	84 (12)	90 (12)	0.060	0.080	89 (6)	81 (14)	85 (11)	85 (11)	88 (12)	80 (14)	0.106	88 (12)
Etofenprox	93 (8)	86 (11)	89 (10)	124 (8)	86 (5)	87 (6)	0.124	0.215	75 (8)	83 (6)	79 (9)	79 (9)	82 (11)	82 (6)	0.072	82 (11)
Etiozazole	89 (10)	75 (16)	82 (15)	101 (8)	79 (11)	81 (8)	0.101	0.112	81 (4)	75 (15)	78 (11)	78 (11)	80 (11)	75 (15)	0.302	80 (11)
Fenamidone	94 (14)	91 (19)	97 (17)	373 (8)	77 (11)	92 (10)	0.373	0.069	91 (15)	93 (7)	92 (11)	92 (11)	86 (12)	95 (7)	0.638	86 (12)
Fenamidophos	91 (14)	94 (1)	93 (10)	560 (8)	84 (11)	86 (9)	0.560	0.381	83 (8)	81 (10)	82 (9)	82 (9)	82 (15)	73 (17)	0.581	82 (15)
Fenarimol	98 (47)	89 (1)	94 (34)	610 (8)	101 (16)	94 (16)	0.610	0.077	92 (13)	86 (19)	89 (16)	89 (16)	85 (8)	81 (4)	0.188	85 (8)
Fenazaquin	89 (11)	94 (6)	91 (9)	427 (8)	100 (16)	93 (15)	0.427	0.056	81 (9)	77 (14)	79 (12)	79 (12)	85 (11)	76 (14)	0.443	85 (11)
Fenbuconazole	115 (15)	104 (5)	110 (13)	199 (5)	109 (19)	102 (18)	0.199	0.280	89 (7)	86 (10)	88 (9)	88 (9)	91 (20)	86 (10)	0.419	91 (20)
Fenhexamid	107 (19)	108 (15)	107 (16)	892 (9)	85 (20)	92 (16)	0.892	0.071	87 (9)	74 (15)	80 (14)	80 (14)	87 (10)	78 (9)	0.072	87 (10)
Fenobucarb	86 (17)	75 (18)	81 (18)	070 (4)	86 (12)	89 (9)	0.070	0.136	92 (5)	89 (17)	90 (12)	90 (12)	80 (18)	71 (11)	0.696	81 (17)
Fenoxycarb	74 (43)	11 (444)	42 (121)	002 (4)	94 (15)	91 (16)	0.002	0.498	83 (18)	73 (11)	78 (16)	78 (16)	80 (18)	71 (11)	0.140	80 (18)
Fenprothimorph	84 (15)	76 (6)	80 (13)	112 (7)	74 (5)	76 (5)	0.112	0.213	77 (3)	73 (8)	75 (7)	75 (7)	77 (5)	75 (9)	0.093	77 (5)
Fenpyroximate	88 (12)	92 (4)	90 (9)	406 (8)	103 (12)	95 (16)	0.406	0.079	78 (15)	74 (11)	76 (13)	76 (13)	83 (20)	73 (11)	0.543	83 (20)
Fensulfotion	41 (0)	941 (3)	450 (113)	000 (20)	493 (2)	236 (113)	0.000	0.000	97 (15)	105 (8)	101 (12)	101 (12)	92 (16)	108 (8)	0.234	92 (16)
Fluazifop-butyl	92 (13)	95 (8)	94 (10)	652 (10)	109 (13)	105 (12)	0.652	0.318	87 (8)	74 (14)	80 (13)	80 (13)	89 (15)	74 (14)	0.064	89 (15)
Fludioxonil	85 (16)	92 (17)	88 (17)	349 (10)	100 (13)	100 (15)	0.349	0.983	84 (19)	89 (13)	86 (16)	86 (16)	79 (18)	89 (13)	0.372	79 (18)
Flufenoxuron	105 (9)	89 (18)	97 (16)	072 (6)	103 (7)	91 (12)	0.072	0.151	95 (12)	85 (12)	90 (13)	90 (13)	81 (14)	72 (8)	0.058	81 (14)
Fluquinconazole	82 (13)	78 (17)	80 (15)	592 (10)	107 (6)	102 (11)	0.592	0.757	90 (14)	79 (12)	84 (14)	84 (14)	77 (11)	68 (12)	0.103	77 (11)
Flusilazole	96 (12)	112 (12)	104 (14)	107 (9)	85 (15)	88 (12)	0.107	0.320	87 (3)	84 (4)	86 (4)	86 (4)	83 (8)	72 (19)	0.258	83 (8)
Flutolanil	98 (15)	108 (7)	103 (12)	124 (9)	89 (15)	90 (11)	0.124	0.857	88 (8)	80 (13)	84 (11)	84 (11)	86 (14)	80 (13)	0.135	86 (14)
Flutriafol	98 (17)	107 (10)	103 (14)	303 (8)	100 (16)	94 (17)	0.303	0.226	87 (9)	88 (10)	87 (9)	87 (9)	82 (9)	90 (10)	0.889	82 (9)
Fosthiazate	92 (12)	81 (8)	87 (12)	067 (9)	112 (17)	86 (15)	0.067	0.146	87 (11)	76 (8)	81 (12)	81 (12)	92 (17)	77 (10)	0.014	92 (17)
Funonisin B1 ^a	378 (2)	378 (2)	378 (1)	n.f.r. ^b	152 (3)	152 (3)	n.f.r. ^b	n.f.r. ^b	77 (4)	78 (3)	78 (3)	78 (3)	94 (5)	93 (5)	0.083	94 (5)
Funonisin B2 ^a	351 (1)	351 (1)	351 (1)	819 (3)	149 (3)	148 (3)	0.819	0.085	77 (5)	78 (3)	78 (4)	78 (4)	75 (3)	83 (11)	0.356	75 (3)
Furalaxyl	93 (9)	79 (19)	86 (16)	055 (5)	85 (13)	88 (10)	0.055	0.266	88 (6)	77 (13)	82 (12)	82 (12)	88 (11)	75 (10)	0.081	88 (11)
Furathiocarb	93 (12)	98 (7)	96 (10)	437 (8)	85 (17)	87 (14)	0.437	0.668	84 (8)	75 (16)	80 (13)	80 (13)	88 (17)	75 (16)	0.100	88 (17)
Halofenozide	143 (23)	441 (60)	292 (81)	000 (15)	161 (81)	156 (59)	0.000	1.000	91 (10)	93 (18)	92 (14)	92 (14)	106 (9)	115 (15)	0.697	106 (9)

Table 3 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Analyst 2	Analyst 1	Average recovery (%) (RSDr (%))		Analyst 2	Analyst 1	Average recovery (%) (RSDr (%))		Analyst 2	Analyst 1	Average recovery (%) (RSDr (%))		Analyst 2	Analyst 1
	Average recovery (%) (RSD _{WR} (%))				Average recovery (%) (RSD _{WR} (%))				Average recovery (%) (RSD _{WR} (%))				Average recovery (%) (RSD _{WR} (%))			
	Analyst 1	Analyst 2		P value 1	Analyst 1	Analyst 2		P value 1	Analyst 1	Analyst 2		P value 1	Analyst 1	Analyst 2		P value 1
Haloxypop-2-ethoxyethyl	93 (19)	102 (11)	98 (15)	0.188	83 (11)	93 (12)	88 (13)	0.051	80 (10)	79 (15)	79 (12)	0.913	79 (18)	79 (15)	79 (16)	0.939
Hexaconazole	92 (16)	77 (14)	84 (17)	0.12	84 (11)	73 (13)	78 (13)	0.069	83 (6)	77 (10)	80 (9)	0.140	78 (8)	74 (17)	76 (13)	0.565
Hexythiazox	87 (15)	95 (11)	91 (13)	0.368	81 (13)	87 (19)	84 (17)	0.516	81 (10)	73 (15)	77 (13)	0.064	76 (15)	73 (15)	75 (15)	0.451
Imazalil	90 (42)	−268 (−3)	−89 (−211)	0.000	84 (16)	95 (12)	90 (15)	0.086	70 (9)	69 (9)	70 (9)	0.809	80 (12)	90 (7)	85 (11)	0.127
Imazapic	88 (13)	97 (8)	93 (11)	0.141	83 (6)	97 (19)	90 (17)	0.154	81 (6)	75 (11)	78 (9)	0.197	81 (3)	75 (11)	78 (9)	0.127
Imazetapyr	83 (13)	78 (19)	81 (16)	0.318	81 (4)	78 (8)	79 (6)	0.103	77 (4)	72 (17)	75 (12)	0.288	74 (14)	79 (2)	77 (10)	0.203
Imidacloprid	95 (12)	83 (12)	89 (14)	0.099	114 (16)	119 (12)	117 (14)	0.441	100 (12)	105 (8)	102 (10)	0.518	97 (8)	107 (10)	102 (10)	0.069
Indoxacarb	106 (9)	92 (20)	99 (16)	0.078	88 (19)	98 (18)	93 (19)	0.067	81 (14)	75 (16)	78 (15)	0.291	84 (18)	76 (16)	80 (17)	0.183
Iprovalicarb	95 (13)	101 (18)	98 (16)	0.519	92 (15)	109 (17)	100 (18)	0.060	87 (10)	95 (8)	91 (10)	0.078	92 (16)	95 (8)	94 (12)	0.666
Isoxaflutole	−30 (0)	903 (59)	437 (139)	0.000	95 (41)	105 (143)	100 (106)	1.000	84 (16)	85 (21)	85 (18)	0.896	98 (16)	93 (9)	96 (13)	0.548
Kresoxim-methyl	74 (18)	72 (20)	73 (18)	0.716	90 (13)	97 (10)	94 (12)	0.121	78 (14)	77 (12)	77 (13)	0.811	82 (21)	77 (12)	79 (17)	0.577
Linuron	111 (76)	20 (216)	66 (121)	0.000	80 (26)	78 (40)	79 (33)	0.884	73 (19)	75 (19)	74 (18)	0.816	84 (13)	85 (18)	84 (15)	0.862
Lufenuron	94 (15)	96 (9)	95 (12)	0.730	88 (16)	94 (3)	91 (11)	0.381	82 (6)	79 (12)	81 (9)	0.369	83 (20)	79 (12)	81 (16)	0.608
Malathion	91 (15)	97 (9)	94 (13)	0.380	85 (9)	95 (8)	90 (10)	0.051	85 (11)	77 (7)	81 (10)	0.111	84 (15)	79 (7)	81 (12)	0.44
Mecarbam	79 (12)	92 (13)	85 (14)	0.098	84 (4)	88 (4)	86 (5)	0.078	81 (11)	88 (8)	85 (10)	0.289	82 (19)	89 (8)	86 (14)	0.424
Mepanipyrim	86 (18)	92 (11)	89 (14)	0.189	85 (18)	100 (13)	92 (17)	0.062	77 (15)	78 (8)	78 (12)	0.622	86 (18)	80 (8)	83 (14)	0.374
Metalaxyl	116 (8)	117 (18)	117 (14)	0.905	114 (16)	110 (15)	112 (15)	0.252	96 (8)	92 (14)	94 (11)	0.249	94 (11)	84 (17)	89 (15)	0.188
Metconazole	320 (136)	123 (17)	222 (141)	0.000	97 (13)	86 (14)	91 (14)	0.078	85 (9)	77 (13)	81 (12)	0.264	80 (6)	72 (15)	76 (12)	0.186
Methamidophos	84 (17)	73 (15)	79 (17)	0.191	83 (12)	90 (17)	86 (15)	0.294	72 (8)	79 (12)	76 (11)	0.064	81 (12)	93 (10)	87 (13)	0.102
Methidathion	17 (422)	17 (422)	17 (406)	n.f.r. ^b	88 (16)	76 (5)	82 (14)	0.057	84 (4)	84 (6)	84 (5)	0.811	75 (15)	76 (15)	76 (14)	0.853
Methiocarb	87 (20)	85 (3)	86 (14)	0.751	88 (14)	99 (19)	93 (17)	0.152	81 (8)	88 (9)	84 (9)	0.098	80 (12)	88 (9)	84 (11)	0.113
Methomyl	90 (11)	81 (7)	85 (11)	0.136	89 (9)	81 (8)	85 (10)	0.066	88 (4)	80 (12)	84 (10)	0.071	85 (13)	74 (5)	80 (13)	0.066
Methoxyfenozide	99 (15)	103 (7)	101 (11)	0.601	91 (9)	109 (18)	100 (17)	0.060	89 (8)	79 (12)	84 (12)	0.140	90 (14)	78 (12)	84 (15)	0.109
Monocrotophos	85 (12)	73 (18)	79 (16)	0.159	90 (9)	77 (15)	84 (14)	0.090	87 (4)	81 (15)	84 (11)	0.263	91 (19)	81 (8)	86 (16)	0.181
Myclobutanil	96 (18)	81 (11)	88 (17)	0.129	92 (7)	81 (12)	86 (12)	0.077	88 (2)	82 (18)	85 (12)	0.350	84 (9)	74 (14)	79 (13)	0.111
Nitenpyram	−267 (0)	−78 (−35)	−173 (−58)	0.000	−134 (0)	−20 (−145)	−77 (−81)	0.000	100 (19)	99 (7)	100 (14)	0.933	92 (19)	98 (9)	95 (15)	0.200
Ochratoxin A ^a	53 (0)	53 (0)	53 (0)	n.f.r. ^b	39 (15)	40 (4)	39 (10)	0.772	106 (19)	109 (8)	107 (14)	0.686	81 (18)	89 (5)	85 (13)	0.220
Ofurace	99 (13)	93 (17)	96 (15)	0.244	90 (12)	76 (11)	83 (14)	0.067	89 (8)	82 (17)	85 (13)	0.300	99 (9)	88 (18)	93 (15)	0.171
Omethoate	102 (7)	98 (19)	100 (14)	0.629	84 (4)	80 (20)	82 (14)	0.466	79 (8)	76 (12)	78 (10)	0.194	84 (13)	73 (8)	78 (13)	0.057
Oxadixyl	93 (16)	81 (9)	87 (15)	0.109	91 (13)	80 (13)	86 (14)	0.115	87 (7)	80 (10)	83 (9)	0.055	88 (8)	84 (11)	86 (10)	0.266
Oxamyl	96 (12)	82 (12)	89 (14)	0.098	94 (8)	89 (14)	92 (11)	0.328	89 (3)	83 (11)	86 (8)	0.120	93 (6)	89 (11)	91 (9)	0.389
Paclobutrazol	109 (15)	100 (19)	105 (17)	0.354	99 (12)	92 (14)	96 (13)	0.429	89 (5)	83 (13)	86 (10)	0.102	83 (5)	75 (13)	79 (11)	0.053
Penconazole	94 (14)	77 (19)	86 (18)	0.071	85 (9)	79 (11)	82 (10)	0.127	85 (8)	74 (17)	80 (14)	0.080	79 (12)	70 (6)	75 (11)	0.086
Pencycuron	83 (14)	87 (8)	85 (11)	0.511	89 (15)	102 (13)	95 (15)	0.075	87 (14)	94 (4)	90 (10)	0.192	88 (13)	93 (4)	91 (9)	0.352
Pendimethalin	83 (19)	79 (5)	81 (14)	0.588	72 (11)	71 (3)	72 (8)	0.630	82 (17)	78 (15)	80 (15)	0.544	88 (6)	79 (15)	83 (12)	0.165
Phenothrin	n.f.r. ^b	49 (72)	n.f.r. ^b	n.f.r. ^b	87 (12)	88 (6)	87 (9)	0.774	81 (18)	97 (16)	89 (18)	0.133	91 (6)	98 (16)	95 (12)	0.315
Phenthoate	91 (15)	95 (15)	93 (15)	0.520	85 (9)	91 (16)	88 (13)	0.195	85 (11)	81 (11)	83 (11)	0.412	84 (15)	83 (11)	83 (13)	0.862
Phosalone	43 (113)	15 (215)	29 (146)	0.000	115 (12)	105 (18)	110 (15)	0.390	93 (8)	85 (14)	89 (12)	0.219	81 (20)	88 (13)	85 (17)	0.433

Table 3 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))	
	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2
	Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))		Average recovery (%) (RSD _{WR} (%))	
	(n = 7)		(n = 14)		(n = 7)		(n = 14)		(n = 7)		(n = 14)		(n = 7)		(n = 14)	
	P-value		P-value		P-value		P-value		P-value		P-value		P-value		P-value	
Phosmet	84 (17)	102 (14)	93 (18)	92 (10)	99 (13)	96 (12)	0.080	0.293	80 (19)	83 (12)	81 (15)	0.655	84 (15)	84 (12)	84 (13)	0.968
Picoxystrobin	103 (11)	110 (13)	106 (12)	87 (8)	89 (7)	88 (7)	0.201	0.617	84 (10)	82 (4)	83 (7)	0.311	86 (17)	81 (4)	84 (13)	0.403
Piperonyl butoxide	94 (11)	103 (11)	98 (12)	88 (8)	96 (9)	92 (9)	0.129	0.067	84 (7)	76 (9)	80 (9)	0.063	83 (14)	76 (9)	79 (12)	0.202
Pirimicarb	94 (11)	87 (9)	90 (11)	93 (7)	82 (9)	88 (10)	0.067	0.06	89 (3)	82 (11)	86 (8)	0.090	85 (3)	82 (14)	83 (10)	0.458
Pirimiphos-ethyl	94 (12)	84 (10)	89 (12)	89 (4)	81 (9)	85 (8)	0.128	0.077	87 (1)	77 (15)	82 (12)	0.057	83 (9)	77 (16)	80 (13)	0.269
Pirimiphos-methyl	95 (13)	96 (19)	96 (16)	92 (4)	80 (19)	86 (14)	0.926	0.091	87 (3)	77 (13)	82 (11)	0.051	84 (10)	73 (11)	79 (12)	0.070
Prochloraz	78 (11)	93 (14)	86 (15)	79 (9)	90 (13)	85 (13)	0.053	0.103	76 (9)	81 (4)	78 (7)	0.211	72 (13)	80 (4)	76 (11)	0.103
Profenofos	93 (11)	83 (8)	88 (11)	83 (14)	105 (17)	94 (19)	0.096	0.058	77 (19)	71 (7)	74 (15)	0.273	77 (16)	72 (7)	75 (12)	0.299
Prometryn	89 (13)	80 (4)	85 (11)	88 (5)	83 (11)	86 (9)	0.082	0.079	85 (2)	80 (12)	82 (9)	0.284	82 (6)	78 (13)	80 (10)	0.274
Propamocarb	262 (7)	218 (4)	240 (11)	212 (4)	171 (5)	191 (12)	0.000	0.000	181 (2)	157 (16)	169 (13)	0.059	176 (9)	147 (6)	162 (12)	0.002
Propanil	50 (193)	28 (252)	39 (210)	100 (41)	75 (81)	89 (59)	1.000	0.368	102 (13)	97 (13)	99 (13)	0.597	85 (17)	102 (13)	93 (17)	0.059
Propam	96 (11)	102 (11)	99 (11)	102 (12)	105 (5)	104 (9)	0.340	0.567	93 (17)	94 (8)	94 (13)	0.963	94 (12)	95 (8)	95 (10)	0.840
Propiconazole	89 (10)	76 (19)	82 (16)	84 (14)	65 (14)	74 (19)	0.152	0.025	79 (8)	70 (17)	74 (13)	0.150	83 (10)	75 (17)	79 (14)	0.136
Propyzamide	88 (17)	82 (19)	85 (18)	90 (10)	82 (16)	86 (14)	0.597	0.342	88 (18)	76 (16)	82 (18)	0.247	82 (15)	75 (15)	78 (15)	0.104
Pyraclostrobin	98 (13)	109 (9)	104 (12)	95 (13)	102 (2)	98 (9)	0.062	0.201	76 (11)	82 (7)	79 (9)	0.069	84 (6)	83 (7)	84 (6)	0.643
Pyrazophos	103 (12)	112 (8)	107 (11)	92 (15)	101 (2)	96 (11)	0.097	0.151	82 (14)	78 (5)	80 (11)	0.384	83 (18)	78 (5)	81 (14)	0.410
Pyridaben	91 (16)	79 (19)	85 (18)	78 (12)	86 (11)	82 (12)	0.223	0.101	73 (15)	71 (5)	72 (11)	0.711	78 (17)	71 (5)	74 (13)	0.190
Pyrimethanil	84 (13)	77 (10)	81 (12)	80 (6)	73 (11)	77 (9)	0.293	0.099	77 (3)	71 (13)	74 (10)	0.071	73 (11)	70 (18)	71 (14)	0.622
Pyriproxyfen	87 (12)	96 (7)	91 (10)	82 (8)	83 (11)	82 (9)	0.137	0.862	79 (8)	78 (13)	79 (10)	0.869	82 (4)	78 (13)	80 (9)	0.371
Quinalphos	86 (20)	76 (15)	81 (18)	83 (20)	75 (9)	79 (16)	0.241	0.341	81 (9)	87 (9)	84 (9)	0.223	90 (7)	87 (9)	89 (8)	0.452
Quinoxifen	97 (15)	108 (4)	102 (12)	87 (11)	97 (11)	92 (12)	0.100	0.146	83 (15)	70 (4)	77 (14)	0.055	73 (6)	69 (4)	71 (6)	0.198
Simazine	88 (18)	77 (12)	82 (16)	89 (10)	75 (16)	82 (15)	0.289	0.076	90 (7)	81 (15)	86 (12)	0.157	87 (2)	81 (8)	84 (7)	0.082
Spinosyn A	79 (16)	74 (13)	77 (14)	72 (8)	72 (4)	72 (6)	0.408	0.748	71 (6)	73 (2)	72 (5)	0.221	74 (6)	72 (14)	73 (10)	0.646
Spinosyn D	70 (16)	74 (19)	72 (17)	78 (18)	84 (6)	81 (13)	0.502	0.348	70 (9)	74 (10)	72 (10)	0.394	72 (6)	74 (10)	73 (8)	0.442
Spirodiclofen	77 (53)	66 (23)	71 (42)	72 (36)	81 (17)	77 (27)	0.517	0.395	79 (18)	78 (16)	78 (17)	0.896	84 (7)	79 (16)	81 (12)	0.270
Spiromesifen	98 (7)	97 (19)	97 (14)	101 (17)	99 (16)	100 (16)	0.877	0.798	85 (11)	96 (19)	90 (16)	0.213	107 (17)	96 (19)	102 (18)	0.248
Spiroxamine	89 (11)	81 (10)	85 (11)	83 (4)	78 (8)	81 (7)	0.212	0.087	82 (2)	77 (11)	80 (8)	0.123	81 (2)	77 (19)	79 (13)	0.588
Tau-fluvalinate	64 (114)	334 (52)	199 (95)	112 (63)	247 (33)	180 (56)	0.000	0.000	76 (30)	123 (17)	100 (33)	0.010	85 (15)	85 (12)	85 (13)	0.999
Tebuconazole	97 (12)	85 (13)	91 (14)	79 (13)	89 (4)	84 (11)	0.068	0.089	85 (3)	81 (1)	83 (3)	0.017	83 (7)	76 (19)	79 (14)	0.324
Tebuflufenozide	115 (29)	143 (3)	129 (21)	105 (6)	111 (5)	109 (4)	0.056	0.276	101 (5)	93 (16)	97 (12)	0.251	95 (9)	83 (11)	89 (12)	0.051
Tebuflufenpyrad	94 (18)	106 (7)	100 (14)	87 (12)	92 (1)	89 (9)	0.133	0.320	83 (13)	71 (7)	77 (13)	0.074	80 (16)	71 (7)	75 (14)	0.081
Terbutryn	91 (11)	81 (13)	86 (13)	88 (4)	81 (10)	85 (8)	0.142	0.069	85 (2)	80 (12)	83 (8)	0.167	82 (6)	77 (9)	80 (8)	0.157
Tetrachlorvinphos	94 (9)	90 (18)	92 (14)	91 (32)	57 (31)	74 (39)	0.473	0.039	86 (12)	89 (13)	88 (12)	0.472	98 (16)	89 (13)	94 (15)	0.302
Tetraconazole	97 (14)	81 (14)	89 (17)	85 (10)	73 (14)	79 (14)	0.067	0.078	93 (11)	91 (13)	92 (12)	0.61	84 (10)	81 (14)	82 (12)	0.354
Tetramethrin	98 (12)	90 (20)	94 (16)	80 (13)	95 (8)	90 (12)	0.309	0.113	86 (4)	81 (13)	83 (10)	0.312	84 (15)	80 (13)	82 (14)	0.548
Thiabendazole	45 (16)	39 (8)	42 (15)	38 (4)	36 (8)	37 (7)	0.070	0.144	37 (4)	39 (17)	38 (12)	0.348	38 (8)	38 (12)	38 (10)	0.952
Thiacloprid	88 (16)	76 (8)	82 (15)	81 (14)	70 (12)	76 (15)	0.085	0.059	78 (14)	74 (26)	76 (20)	0.526	82 (15)	71 (13)	76 (15)	0.222
Thiamethoxam	139 (24)	42 (35)	90 (63)	118 (42)	71 (24)	94 (46)	0.000	0.091	104 (28)	96 (33)	100 (30)	0.155	100 (7)	89 (17)	95 (13)	0.177
Thiodicarb	97 (13)	81 (13)	89 (16)	89 (11)	82 (19)	85 (15)	0.103	0.381	84 (8)	74 (16)	79 (13)	0.101	86 (7)	77 (19)	81 (14)	0.176

Table 4 Average recoveries, precision (RSDr) and intermediate precision (RSD_{WR}) obtained for *M. sylvestris* from the method validation procedure

	Concentration 1					Concentration 2					Concentration 3					Concentration 4				
	Average recovery (%)					Average recovery (%)					Average recovery (%)					Average recovery (%)				
	(RSDr (%))					(RSDr (%))					(RSDr (%))					(RSDr (%))				
	Analyst	Analyst	2		Average recovery (%)	Analyst	Analyst	2		Average recovery (%)	Analyst	Analyst	2		Average recovery (%)	Analyst	Analyst	2		Average recovery (%)
	1	2			(n = 14)	1	2			(n = 14)	1	2			(n = 14)	1	2			(n = 14)
Acephate	76 (20)	84 (9)			80 (15)	0.164	84 (13)	101 (12)		93 (15)	0.061	86 (16)	90 (13)		88 (14)	0.619	94 (17)	108 (10)		101 (15)
Acetamidiprid	107 (9)	102 (5)			105 (7)	0.273	94 (6)	89 (11)		92 (9)	0.396	86 (17)	78 (5)		82 (14)	0.295	95 (11)	98 (2)		97 (8)
Acetochlor	106 (17)	95 (11)			100 (15)	0.127	99 (8)	98 (16)		99 (12)	0.811	87 (6)	83 (9)		85 (8)	0.371	99 (5)	97 (10)		98 (8)
Aflatoxin B1 ^a	51 (21)	84 (15)			67 (31)	0.001	76 (8)	71 (19)		73 (14)	0.353	88 (18)	76 (14)		82 (18)	0.237	95 (10)	91 (12)		93 (11)
Aflatoxin B2 ^a	–111 (–11)	–120 (–20)			–115 (–17)	0.438	9 (220)	8 (195)		8 (203)	0.883	57 (18)	62 (10)		59 (15)	0.167	75 (18)	75 (6)		75 (13)
Aflatoxin G1 ^a	66 (58)	83 (38)			75 (47)	0.191	78 (13)	84 (16)		81 (15)	0.223	96 (14)	85 (14)		90 (15)	0.143	84 (15)	85 (11)		85 (12)
Aflatoxin G2 ^a	n.f.r.	39 (228)			n.f.r. ^b	n.f.r. ^b	90 (20)	85 (11)		88 (16)	0.434	89 (19)	88 (7)		89 (14)	0.801	85 (11)	96 (11)		90 (12)
Aldicarb sulfone	90 (16)	93 (18)			91 (16)	0.726	90 (13)	84 (13)		87 (13)	0.416	95 (4)	98 (9)		98 (7)	0.728	103 (6)	111 (5)		107 (7)
Aldicarb sulfoxide	101 (17)	86 (20)			94 (20)	0.180	90 (13)	106 (13)		98 (15)	0.087	86 (16)	91 (19)		89 (17)	0.616	83 (5)	94 (13)		89 (12)
Arazine	104 (13)	114 (8)			109 (11)	0.184	107 (12)	116 (3)		111 (9)	0.155	90 (10)	83 (2)		87 (8)	0.118	87 (10)	82 (2)		84 (7)
Azinphos	80 (16)	89 (14)			85 (16)	0.298	89 (20)	102 (6)		95 (15)	0.135	87 (19)	96 (3)		91 (14)	0.214	87 (12)	98 (3)		93 (10)
Azinphos-methyl	75 (49)	123 (57)			99 (60)	0.059	81 (51)	168 (19)		124 (46)	0.008	82 (17)	93 (18)		88 (18)	0.294	92 (14)	79 (19)		85 (18)
Azoxystrobin	95 (19)	110 (4)			103 (14)	0.089	93 (11)	102 (6)		98 (9)	0.063	90 (9)	98 (4)		94 (8)	0.054	93 (12)	96 (1)		94 (8)
Bifenazate	23 (48)	–10 (–52)			6 (301)	0.000	42 (12)	11 (97)		26 (68)	0.000	56 (15)	16 (56)		36 (63)	0.000	56 (16)	19 (27)		38 (53)
Bitertanol	83 (18)	79 (11)			81 (15)	0.518	89 (19)	81 (12)		85 (17)	0.281	95 (16)	85 (9)		90 (14)	0.195	94 (16)	92 (5)		93 (11)
Boscalid	87 (10)	110 (24)			98 (23)	0.052	83 (11)	75 (14)		79 (13)	0.219	88 (8)	82 (13)		85 (11)	0.282	91 (15)	80 (6)		85 (13)
Bupirimate	102 (10)	96 (7)			99 (9)	0.224	98 (12)	89 (5)		93 (10)	0.119	93 (9)	93 (5)		93 (7)	0.907	93 (8)	88 (3)		91 (7)
Buprofezin	101 (15)	114 (5)			108 (12)	0.091	111 (15)	111 (5)		111 (11)	0.968	86 (9)	103 (24)		94 (20)	0.172	106 (16)	110 (3)		108 (11)
Cadusafos	100 (17)	112 (7)			106 (13)	0.133	96 (13)	107 (12)		102 (13)	0.254	103 (13)	113 (8)		108 (11)	0.081	103 (19)	116 (2)		109 (14)
Carbaryl	99 (18)	104 (6)			102 (13)	0.592	96 (11)	108 (7)		102 (10)	0.087	106 (15)	113 (3)		109 (10)	0.318	99 (11)	110 (5)		105 (10)
Carbendazim	65 (54)	48 (11)			57 (45)	0.263	53 (40)	75 (11)		64 (30)	0.067	62 (11)	90 (4)		76 (20)	0.000	82 (11)	91 (3)		86 (9)
Carbofuran	118 (11)	115 (5)			116 (8)	0.632	115 (5)	107 (10)		111 (8)	0.139	113 (10)	104 (5)		108 (9)	0.093	116 (13)	104 (2)		110 (11)
Carpropamid	–40 (0)	34 (24)			–3 (–1288)	0.000	108 (10)	92 (19)		100 (16)	0.050	88 (9)	75 (17)		81 (15)	0.133	100 (8)	97 (5)		98 (7)
Chlorantraniliprole	78 (52)	56 (26)			67 (47)	0.199	78 (20)	87 (19)		82 (19)	0.488	84 (10)	91 (13)		87 (12)	0.172	97 (12)	93 (9)		95 (10)
Chlorfenvinphos	97 (19)	102 (9)			100 (14)	0.599	92 (8)	100 (9)		96 (9)	0.066	89 (11)	80 (19)		84 (16)	0.340	93 (16)	108 (4)		101 (13)
Chlorpyrifos	112 (5)	101 (18)			107 (13)	0.174	102 (14)	103 (16)		102 (15)	0.915	83 (7)	90 (13)		87 (11)	0.116	81 (16)	81 (6)		81 (11)
Clofentezine	90 (20)	79 (16)			84 (19)	0.096	76 (12)	76 (18)		76 (15)	0.998	75 (11)	93 (18)		84 (19)	0.066	89 (19)	79 (19)		84 (19)
Clomazone	83 (20)	73 (13)			78 (18)	0.271	92 (6)	98 (8)		95 (7)	0.192	93 (10)	105 (14)		99 (13)	0.209	105 (10)	111 (12)		108 (11)
Clothianidin	109 (8)	95 (12)			102 (12)	0.051	101 (11)	88 (13)		94 (14)	0.055	92 (19)	78 (6)		85 (17)	0.049	89 (17)	84 (5)		87 (13)
Cyazofamid	76 (47)	41 (54)			59 (58)	0.136	91 (17)	93 (19)		92 (17)	0.746	96 (10)	105 (7)		101 (10)	0.091	99 (11)	107 (10)		103 (11)
Cyproconazole	92 (19)	94 (7)			93 (14)	0.866	94 (9)	103 (9)		99 (10)	0.101	90 (9)	90 (6)		90 (8)	0.946	94 (10)	88 (12)		91 (11)
Cyprodinil	75 (17)	71 (10)			73 (14)	0.469	71 (5)	73 (7)		72 (6)	0.401	70 (9)	72 (8)		71 (8)	0.678	71 (18)	78 (5)		75 (13)
Demeton-S-methyl sulfone	94 (17)	93 (6)			94 (13)	0.836	93 (9)	99 (4)		96 (7)	0.146	101 (10)	107 (5)		104 (8)	0.147	104 (8)	103 (4)		104 (6)
Demeton-S-methyl sulfoxide	77 (60)	79 (16)			78 (42)	0.913	87 (10)	102 (15)		95 (15)	0.128	89 (16)	105 (20)		97 (20)	0.118	96 (11)	107 (3)		102 (9)
Deoxynivalenol ^a	202 (358)	1010 (67)			606 (131)	0.147	657 (68)	109 (283)		383 (122)	0.015	25 (11)	32 (14)		28 (18)	0.038	26 (8)	25 (9)		26 (9)

Table 4 (Contd.)

	Concentration 1					Concentration 2					Concentration 3					Concentration 4								
	Average recovery (%) (RSDr (%))		Analyst		P value	Average recovery (%) (RSDr (%))		Analyst		P value	Average recovery (%) (RSDr (%))		Analyst		P value	Average recovery (%) (RSDr (%))		Analyst		P value				
	Average recovery (%) (RSDr (%))		Analyst			Average recovery (%) (RSDr (%))		Analyst			Average recovery (%) (RSDr (%))		Analyst			Average recovery (%) (RSDr (%))		Analyst						
	1	2	1	2		1	2	1	2		1	2	1	2		1	2	1	2					
Diacetoxyscirpenol^a	2 (291)	–4 (–167)	–1 (–728)	0.036	0.036	16 (60)	17 (20)	16 (41)	0.699	0.699	23 (7)	23 (8)	23 (7)	23 (8)	0.547	22 (12)	25 (4)	24 (11)	0.038					
Diazinon	98 (12)	108 (15)	103 (14)	0.330	0.330	99 (6)	107 (15)	103 (12)	0.226	0.226	108 (19)	113 (9)	111 (14)	111 (14)	0.509	93 (9)	108 (15)	100 (15)	0.106					
Diethofencarb	82 (37)	39 (27)	61 (51)	0.013	0.013	109 (20)	98 (18)	104 (19)	0.140	0.140	89 (14)	95 (6)	92 (11)	92 (11)	0.357	100 (13)	94 (5)	97 (10)	0.155					
Difenoconazole	91 (17)	81 (6)	86 (14)	0.109	0.109	94 (10)	96 (7)	95 (8)	0.594	0.594	89 (10)	100 (5)	94 (10)	94 (10)	0.063	105 (17)	101 (4)	103 (12)	0.641					
Diflubenzuron	49 (168)	14 (560)	32 (253)	0.347	0.347	91 (44)	54 (92)	73 (66)	0.147	0.147	89 (11)	101 (13)	95 (13)	95 (13)	0.170	95 (14)	115 (19)	105 (19)	0.092					
Dimethoate	101 (11)	100 (9)	101 (9)	0.831	0.831	93 (7)	98 (6)	95 (7)	0.161	0.161	88 (18)	92 (3)	90 (12)	90 (12)	0.464	89 (18)	91 (2)	90 (12)	0.816					
Dimethomorph	93 (12)	106 (8)	100 (12)	0.125	0.125	102 (12)	110 (6)	106 (10)	0.197	0.197	91 (12)	87 (4)	89 (9)	89 (9)	0.398	94 (13)	86 (3)	90 (10)	0.189					
Diniconazole	91 (13)	83 (14)	87 (14)	0.129	0.129	91 (16)	80 (7)	85 (14)	0.061	0.061	85 (8)	79 (9)	82 (9)	82 (11)	0.189	89 (10)	82 (11)	85 (11)	0.275					
Diphenylamine	115 (13)	96 (15)	106 (16)	0.114	0.114	98 (11)	87 (10)	92 (12)	0.071	0.071	89 (11)	84 (8)	86 (10)	86 (10)	0.388	86 (10)	86 (8)	86 (9)	0.944					
Diuron	86 (17)	74 (7)	80 (16)	0.096	0.096	90 (8)	93 (7)	91 (7)	0.248	0.248	92 (9)	102 (5)	97 (8)	97 (8)	0.055	90 (9)	89 (20)	90 (15)	0.850					
DMST	86 (16)	100 (18)	93 (19)	0.110	0.110	105 (20)	97 (13)	101 (17)	0.532	0.532	97 (9)	98 (16)	97 (13)	97 (13)	0.963	94 (13)	82 (10)	88 (13)	0.088					
Epoxiconazole	89 (5)	102 (17)	96 (14)	0.088	0.088	96 (10)	111 (14)	104 (14)	0.061	0.061	97 (6)	100 (15)	98 (12)	98 (12)	0.587	89 (15)	103 (9)	96 (14)	0.059					
Ethion	87 (14)	108 (17)	98 (19)	0.051	0.051	92 (8)	104 (14)	98 (13)	0.063	0.063	89 (9)	96 (5)	93 (8)	93 (8)	0.155	88 (13)	94 (6)	91 (10)	0.170					
Ethiprole	86 (14)	79 (19)	82 (16)	0.393	0.393	102 (12)	86 (17)	94 (16)	0.093	0.093	95 (6)	97 (17)	96 (13)	96 (13)	0.735	102 (14)	103 (5)	103 (10)	0.827					
Ethoprophos	98 (16)	112 (9)	105 (14)	0.213	0.213	87 (13)	86 (8)	87 (10)	0.952	0.952	88 (9)	81 (20)	84 (15)	84 (15)	0.427	98 (10)	106 (10)	102 (11)	0.200					
Etofenprox	95 (19)	87 (20)	91 (19)	0.478	0.478	107 (18)	98 (20)	103 (19)	0.452	0.452	86 (9)	80 (15)	83 (13)	83 (13)	0.366	85 (15)	87 (6)	86 (11)	0.821					
Etoxazole	92 (16)	88 (3)	90 (11)	0.464	0.464	89 (9)	86 (5)	87 (7)	0.177	0.177	86 (11)	88 (2)	87 (8)	87 (8)	0.572	87 (6)	87 (3)	87 (4)	0.817					
Fenamidone	97 (12)	108 (8)	102 (11)	0.111	0.111	100 (7)	97 (20)	98 (14)	0.761	0.761	93 (11)	92 (13)	93 (12)	93 (12)	0.689	91 (11)	86 (4)	88 (9)	0.164					
Fenamiphos	105 (12)	98 (11)	102 (11)	0.352	0.352	91 (13)	91 (5)	91 (9)	0.948	0.948	87 (10)	91 (3)	89 (8)	89 (8)	0.475	90 (14)	92 (2)	91 (9)	0.654					
Fenarimol	76 (35)	48 (24)	62 (39)	0.097	0.097	89 (11)	102 (15)	96 (15)	0.118	0.118	76 (19)	74 (10)	75 (15)	75 (15)	0.739	89 (16)	80 (10)	85 (15)	0.117					
Fenazaquin	89 (18)	84 (7)	86 (14)	0.499	0.499	89 (16)	94 (7)	91 (12)	0.384	0.384	84 (14)	93 (3)	88 (11)	88 (11)	0.062	83 (15)	91 (2)	87 (11)	0.161					
Fenbuconazole	65 (27)	40 (47)	53 (42)	0.025	0.025	90 (19)	110 (16)	100 (20)	0.158	0.158	98 (19)	106 (11)	102 (15)	102 (15)	0.403	98 (6)	99 (21)	98 (15)	0.908					
Fenhexamid	81 (34)	78 (20)	79 (27)	0.854	0.854	95 (15)	94 (19)	94 (16)	0.910	0.910	93 (10)	103 (13)	98 (13)	98 (13)	0.099	99 (14)	104 (15)	102 (15)	0.684					
Fenobucarb	89 (20)	103 (7)	96 (16)	0.116	0.116	99 (13)	111 (9)	105 (12)	0.107	0.107	91 (10)	106 (16)	98 (15)	98 (15)	0.084	101 (7)	113 (13)	107 (12)	0.074					
Fenoxycarb	72 (55)	52 (0)	62 (47)	0.231	0.231	81 (19)	73 (14)	77 (17)	0.191	0.191	86 (15)	79 (18)	82 (16)	82 (16)	0.457	94 (13)	84 (19)	89 (17)	0.259					
Fenpropimorph	100 (13)	107 (7)	103 (11)	0.341	0.341	96 (6)	105 (12)	100 (11)	0.139	0.139	93 (5)	96 (4)	94 (5)	94 (5)	0.401	86 (11)	95 (2)	90 (9)	0.068					
Fenpyroximate	85 (17)	93 (4)	89 (12)	0.252	0.252	83 (11)	94 (8)	88 (11)	0.058	0.058	85 (15)	95 (5)	90 (12)	90 (12)	0.151	85 (7)	91 (9)	88 (9)	0.174					
Fensulfthion	104 (13)	101 (19)	103 (16)	0.752	0.752	100 (14)	99 (9)	100 (12)	0.871	0.871	113 (14)	101 (11)	107 (14)	107 (14)	0.079	119 (5)	111 (6)	115 (7)	0.068					
Fluazifop-butyl	89 (13)	99 (10)	94 (12)	0.117	0.117	87 (9)	99 (9)	93 (11)	0.066	0.066	88 (10)	93 (18)	90 (14)	90 (14)	0.361	87 (12)	98 (14)	93 (14)	0.195					
Fludioxonil	108 (14)	92 (15)	100 (16)	0.085	0.085	90 (18)	89 (15)	89 (16)	0.817	0.817	97 (11)	98 (12)	97 (11)	97 (11)	0.853	95 (9)	95 (8)	95 (8)	0.992					
Flufenoxuron	–22 (0)	57 (34)	18 (245)	0.000	0.000	105 (17)	101 (12)	103 (15)	0.522	0.522	78 (11)	79 (12)	78 (11)	78 (11)	0.911	96 (14)	98 (19)	97 (16)	0.833					
Fluquinconazole	89 (53)	65 (40)	77 (51)	0.308	0.308	94 (15)	94 (20)	94 (17)	0.968	0.968	93 (17)	89 (17)	91 (16)	91 (16)	0.715	94 (14)	86 (12)	90 (14)	0.125					
Flusilazol	84 (19)	104 (15)	94 (20)	0.055	0.055	94 (18)	104 (10)	99 (15)	0.172	0.172	88 (8)	78 (10)	83 (11)	83 (11)	0.057	96 (10)	85 (14)	91 (13)	0.157					
Flutolanil	56 (39)	25 (22)	41 (54)	0.016	0.016	81 (19)	99 (18)	90 (20)	0.106	0.106	85 (13)	80 (17)	82 (15)	82 (15)	0.502	97 (18)	84 (16)	91 (19)	0.066					
Flutriafol	99 (16)	87 (9)	93 (15)	0.118	0.118	93 (9)	84 (12)	89 (11)	0.122	0.122	87 (11)	90 (5)	88 (8)	88 (8)	0.509	91 (15)	82 (7)	87 (13)	0.223					
Fosthiazate	105 (15)	95 (7)	100 (12)	0.172	0.172	98 (6)	94 (10)	96 (8)	0.269	0.269	89 (13)	90 (4)	89 (9)	89 (9)	0.788	91 (13)	89 (4)	90 (9)	0.722					
Fumonisin B1^a	156 (2)	163 (14)	159 (10)	0.490	0.490	72 (17)	61 (0)	67 (15)	0.047	0.047	49 (39)	30 (1)	40 (41)	40 (41)	0.044	23 (31)	15 (1)	19 (32)	0.030					
Fumonisin B2^a	186 (4)	186 (1)	186 (3)	0.996	0.996	86 (8)	72 (1)	79 (11)	0.002	0.002	51 (8)	36 (0)	44 (19)	44 (19)	0.000	26 (28)	18 (1)	22 (29)	0.032					
Furalaxyl	81 (33)	124 (5)	102 (28)	0.003	0.003	79 (14)	131 (4)	105 (27)	0.000	0.000	83 (10)	89 (2)	86 (8)	86 (8)	0.146	95 (11)	89 (2)	92 (9)	0.154					
Furathiocarb	92 (19)	108 (6)	100 (15)	0.118	0.118	92 (9)	103 (10)	97 (11)	0.151	0.151	88 (9)	100 (13)	94 (13)	94 (13)	0.072	87 (12)	98 (11)	93 (12)	0.086					

Table 4 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))	
	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2	Analyst 1	Analyst 2
Halofenozide	–177 (0)	–177 (0)	–177 (0)	–177 (0)	n.f.r. ^b	–88 (0)	127 (34)	19 (603)	0.000	91 (20)	103 (20)	97 (20)	0.161	103 (15)	97 (20)	100 (17)
Haloxyp-2-ethoxyethyl	87 (18)	92 (12)	89 (15)	89 (15)	0.464	87 (10)	91 (9)	89 (9)	0.312	86 (7)	98 (11)	92 (11)	0.072	93 (8)	102 (8)	97 (9)
Hexaconazol	97 (15)	95 (12)	96 (13)	96 (13)	0.677	97 (6)	93 (13)	95 (10)	0.464	90 (12)	85 (6)	88 (9)	0.393	86 (10)	84 (8)	85 (9)
Hexythiazox	100 (18)	81 (17)	91 (20)	91 (20)	0.059	90 (15)	87 (19)	88 (16)	0.712	86 (8)	83 (9)	85 (8)	0.507	86 (14)	92 (9)	89 (12)
Imazalil	96 (18)	83 (5)	89 (16)	89 (16)	0.101	83 (9)	88 (8)	85 (9)	0.337	85 (9)	93 (5)	89 (9)	0.087	86 (9)	89 (4)	87 (7)
Imazapic	111 (7)	75 (10)	93 (22)	93 (22)	0.000	91 (8)	82 (16)	87 (13)	0.083	86 (20)	71 (10)	78 (19)	0.114	81 (18)	74 (6)	77 (14)
Imazetapyr	94 (14)	86 (12)	90 (13)	90 (13)	0.118	84 (5)	89 (8)	86 (7)	0.119	72 (17)	79 (4)	76 (12)	0.220	78 (16)	79 (3)	78 (11)
Imidacloprid	100 (13)	92 (15)	96 (14)	96 (14)	0.428	101 (18)	88 (12)	95 (17)	0.058	99 (8)	89 (14)	94 (12)	0.062	89 (19)	92 (21)	91 (19)
Indoxacarb	82 (17)	102 (18)	92 (21)	92 (21)	0.091	87 (12)	90 (14)	89 (12)	0.543	93 (14)	107 (18)	100 (17)	0.054	101 (8)	109 (8)	105 (9)
Iprovalicarb	72 (13)	76 (14)	74 (13)	74 (13)	0.381	79 (15)	92 (10)	86 (14)	0.050	81 (11)	97 (10)	89 (13)	0.055	94 (15)	101 (1)	98 (11)
Isoxaflutole	281 (110)	203 (94)	242 (103)	242 (103)	0.324	168 (35)	225 (85)	196 (71)	0.474	96 (18)	106 (21)	101 (19)	0.076	100 (15)	92 (10)	96 (13)
Kresoxim-methyl	86 (17)	93 (19)	89 (18)	89 (18)	0.468	92 (13)	95 (17)	94 (15)	0.646	100 (19)	117 (6)	108 (15)	0.067	96 (11)	109 (7)	103 (11)
Linuron	–14 (0)	55 (29)	21 (183)	21 (183)	0.000	93 (13)	101 (16)	97 (15)	0.187	118 (12)	119 (7)	119 (9)	0.803	113 (10)	116 (10)	114 (10)
Lufenuron	86 (20)	97 (11)	91 (17)	91 (17)	0.169	93 (15)	84 (17)	89 (16)	0.130	81 (14)	81 (9)	81 (11)	0.931	92 (11)	84 (6)	88 (10)
Malathion	103 (15)	104 (12)	104 (13)	104 (13)	0.962	81 (13)	91 (14)	86 (14)	0.200	84 (6)	79 (17)	82 (12)	0.378	106 (8)	114 (5)	110 (8)
Mecarban	91 (19)	89 (14)	90 (16)	90 (16)	0.732	98 (11)	87 (14)	92 (14)	0.103	93 (12)	101 (11)	97 (12)	0.096	99 (7)	100 (9)	100 (7)
Mepanipyridin	56 (44)	186 (31)	121 (66)	121 (66)	0.002	59 (31)	78 (40)	69 (39)	0.147	77 (13)	77 (14)	77 (13)	0.919	91 (7)	101 (10)	96 (10)
Metalaxyl	109 (11)	100 (4)	105 (9)	105 (9)	0.111	96 (5)	94 (6)	95 (5)	0.438	89 (15)	88 (2)	88 (11)	0.850	88 (13)	87 (2)	88 (9)
Metconazol	92 (20)	110 (8)	101 (16)	101 (16)	0.102	90 (15)	103 (5)	96 (13)	0.060	86 (18)	96 (6)	91 (14)	0.134	88 (14)	97 (5)	92 (11)
Methamidophos	51 (107)	166 (79)	109 (104)	109 (104)	0.063	104 (15)	101 (20)	102 (17)	0.711	44 (18)	64 (11)	54 (24)	0.003	78 (17)	93 (14)	86 (17)
Methidathion	–73 (–89)	65 (65)	–4 (–2073)	–4 (–2073)	0.005	92 (18)	97 (17)	94 (17)	0.616	89 (18)	84 (15)	86 (16)	0.309	104 (13)	113 (10)	109 (12)
Methiocarb	56 (43)	17 (62)	37 (74)	37 (74)	0.005	112 (17)	113 (13)	112 (15)	0.957	86 (9)	78 (8)	82 (10)	0.073	98 (13)	103 (16)	100 (14)
Methomyl	103 (13)	113 (7)	108 (11)	108 (11)	0.146	100 (14)	104 (13)	102 (13)	0.679	97 (19)	107 (18)	102 (19)	0.119	102 (10)	105 (19)	103 (15)
Methoxyfenozide	89 (31)	164 (8)	127 (35)	127 (35)	0.001	89 (17)	105 (18)	97 (19)	0.089	85 (13)	86 (19)	85 (16)	0.748	98 (9)	107 (14)	103 (13)
Monocrotophos	94 (13)	104 (11)	99 (13)	99 (13)	0.241	75 (14)	87 (17)	81 (17)	0.077	91 (13)	99 (8)	95 (11)	0.133	89 (10)	96 (7)	93 (9)
Myclobutanil	99 (17)	82 (10)	91 (17)	91 (17)	0.060	103 (6)	92 (15)	98 (12)	0.130	96 (15)	112 (8)	104 (14)	0.075	91 (12)	91 (4)	91 (9)
Nitenpyran	290 (41)	871 (28)	580 (61)	580 (61)	0.002	141 (92)	666 (37)	404 (82)	0.002	93 (47)	244 (46)	169 (67)	0.011	109 (14)	187 (70)	148 (66)
Ochloroxin A ^a	101 (8)	109 (4)	105 (7)	105 (7)	0.106	70 (11)	69 (15)	70 (13)	0.760	81 (14)	73 (8)	77 (12)	0.137	70 (5)	71 (10)	70 (7)
Oflurace	97 (19)	95 (13)	96 (16)	96 (16)	0.783	92 (9)	85 (3)	88 (8)	0.059	91 (17)	77 (5)	84 (15)	0.079	90 (12)	81 (5)	85 (11)
Omethoate	980 (72)	697 (58)	839 (68)	839 (68)	0.377	210 (83)	739 (16)	475 (65)	0.001	96 (20)	107 (19)	102 (20)	0.228	89 (4)	84 (18)	86 (13)
Oxadixyl	90 (17)	83 (5)	86 (13)	86 (13)	0.318	88 (7)	88 (8)	88 (7)	0.945	90 (17)	93 (1)	92 (12)	0.639	94 (11)	92 (3)	93 (8)
Oxamyl	91 (16)	80 (6)	85 (14)	85 (14)	0.086	82 (14)	88 (9)	85 (12)	0.337	84 (5)	90 (12)	87 (9)	0.331	85 (10)	89 (4)	87 (7)
Paclobutrazol	105 (15)	119 (12)	112 (15)	112 (15)	0.163	98 (11)	111 (9)	105 (12)	0.054	96 (6)	107 (9)	102 (9)	0.063	93 (10)	108 (14)	100 (14)
Penconazol	92 (17)	102 (11)	97 (14)	97 (14)	0.086	90 (9)	92 (6)	91 (7)	0.715	92 (9)	96 (6)	94 (8)	0.303	88 (9)	98 (3)	93 (8)
Pencycuron	94 (17)	106 (11)	100 (15)	100 (15)	0.103	91 (10)	107 (13)	99 (14)	0.063	95 (11)	108 (12)	101 (13)	0.113	99 (7)	106 (4)	103 (6)
Pendimethalin	85 (18)	80 (18)	83 (17)	83 (17)	0.689	88 (11)	92 (11)	90 (11)	0.517	86 (9)	94 (6)	90 (9)	0.081	81 (7)	88 (13)	84 (11)
Phenothrin	113 (18)	107 (14)	110 (16)	110 (16)	0.548	97 (13)	113 (13)	105 (15)	0.082	83 (12)	90 (9)	86 (11)	0.154	88 (15)	82 (7)	85 (13)

Table 4 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))		Average recovery (%) (RSDr (%))	
	Analyst 1	Analyst 2	P value	Average recovery (%) (RSD _{wr} (%))	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	Analyst 1	Analyst 2	P value	Average recovery (%) (RSD _{wr} (%))
Phenthoate	416 (266)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	141 (216)	n.f.r. ^b	0.027	n.f.r. ^b
Phosalone	86 (46)	−67 (−205)	0.029	9 (1328)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	516 (207)	86 (10)	0.510	92 (14)
Phosmet	−79 (0)	31 (71)	0.000	−24 (−247)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	94 (15)	87 (4)	0.241	91 (12)
Picoxystrobin	101 (17)	115 (2)	0.068	108 (13)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	95 (15)	104 (18)	0.176	99 (17)
Piperonyl butoxide	92 (16)	103 (6)	0.173	97 (13)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	86 (8)	95 (11)	0.140	91 (10)
Prirnicarb	101 (12)	97 (5)	0.420	99 (9)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	85 (15)	96 (4)	0.356	91 (12)
Pririmiphos-ethyl	98 (13)	104 (6)	0.405	101 (10)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	92 (7)	99 (12)	0.245	95 (10)
Pririmiphos-methyl	102 (13)	104 (9)	0.761	103 (11)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	99 (13)	110 (6)	0.011	104 (11)
Prochloraz	86 (19)	105 (9)	0.074	95 (17)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	85 (10)	92 (4)	0.728	89 (8)
Profenofos	100 (19)	88 (12)	0.193	94 (17)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	92 (6)	102 (9)	0.956	97 (9)
Prometryn	97 (12)	101 (4)	0.445	99 (9)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	96 (3)	95 (5)	0.477	91 (7)
Propamocarb	44 (73)	−4 (−707)	0.030	20 (197)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	59 (16)	18 (8)	0.002	38 (58)
Propanil	71 (59)	−80 (0)	0.000	−5 (−1818)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	73 (45)	70 (56)	0.811	100 (18)
Propham	77 (18)	88 (20)	0.269	82 (20)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	84 (16)	90 (16)	0.475	96 (18)
Propiconazole	82 (15)	88 (13)	0.258	85 (14)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	84 (11)	96 (17)	0.229	87 (16)
Propyzamide	93 (20)	102 (12)	0.361	98 (16)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	84 (20)	102 (20)	0.196	90 (16)
Pyraclostrobin	88 (16)	71 (13)	0.056	79 (18)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	88 (10)	93 (10)	0.375	85 (15)
Pyrazophos	275 (11)	296 (7)	0.097	286 (10)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	236 (14)	257 (2)	0.151	200 (10)
Pyridaben	88 (18)	84 (10)	0.649	86 (14)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	87 (5)	89 (5)	0.146	85 (9)
Pyrimethanil	85 (16)	73 (4)	0.064	79 (14)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	78 (5)	77 (5)	0.367	75 (7)
Pyriproxyfen	96 (16)	88 (6)	0.331	92 (13)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	90 (8)	91 (7)	0.569	87 (10)
Quinalphos	95 (19)	101 (16)	0.457	98 (17)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	90 (12)	97 (17)	0.171	105 (8)
Quinoxifen	95 (16)	89 (11)	0.483	92 (14)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	81 (13)	93 (11)	0.087	88 (5)
Simazine	98 (15)	96 (10)	0.714	97 (12)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	92 (9)	89 (9)	0.480	88 (10)
Spinosyn A	92 (16)	79 (14)	0.142	86 (16)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	89 (8)	84 (7)	0.097	85 (8)
Spinosyn D	100 (17)	84 (16)	0.139	92 (19)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	94 (10)	93 (13)	0.789	87 (6)
Spirodiclofen	78 (17)	86 (17)	0.324	82 (17)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	80 (15)	99 (12)	0.055	95 (10)
Spiromesifen	74 (33)	83 (23)	0.359	79 (28)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	78 (6)	91 (17)	0.127	78 (10)
Spiroxamine	104 (12)	97 (5)	0.208	101 (10)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	97 (4)	100 (3)	0.067	93 (6)
Tau-fluvalinate	161 (41)	80 (119)	0.091	121 (74)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	102 (10)	103 (5)	0.720	92 (16)
Tebuconazole	97 (17)	110 (6)	0.110	104 (13)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	92 (23)	112 (7)	0.069	86 (15)
Tebufenozide	98 (12)	95 (9)	0.444	97 (10)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	104 (5)	107 (17)	0.667	97 (8)
Tebufenpyrad	95 (18)	95 (12)	0.974	95 (15)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	90 (15)	101 (11)	0.236	107 (6)
Terbutryn	100 (12)	101 (3)	0.746	100 (9)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	94 (4)	95 (4)	0.465	91 (7)
Tetrachlorvinphos	77 (99)	72 (110)	0.858	74 (100)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	68 (44)	64 (25)	0.759	83 (16)
Tetraconazole	100 (18)	115 (9)	0.106	108 (15)	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	n.f.r. ^b	100 (10)	112 (11)	0.158	103 (12)

Table 4 (Contd.)

	Concentration 1				Concentration 2				Concentration 3				Concentration 4			
	Average recovery (%) (RSDr (%))		Analyst 2	P value	Average recovery (%) (RSDr (%))		Analyst 2	P value	Average recovery (%) (RSDr (%))		Analyst 2	P value	Average recovery (%) (RSDr (%))		Analyst 2	P value
	Average recovery (%) (RSD _{wrk} (%))				Average recovery (%) (RSD _{wrk} (%))				Average recovery (%) (RSD _{wrk} (%))				Average recovery (%) (RSD _{wrk} (%))			
	Analyst 1	Analyst 2	Average recovery (%) (RSD _{wrk} (%))	Average recovery (%) (RSD _{wrk} (%))	Analyst 1	Analyst 2	Average recovery (%) (RSD _{wrk} (%))	Average recovery (%) (RSD _{wrk} (%))	Analyst 1	Analyst 2	Average recovery (%) (RSD _{wrk} (%))	Average recovery (%) (RSD _{wrk} (%))	Analyst 1	Analyst 2	Average recovery (%) (RSD _{wrk} (%))	Average recovery (%) (RSD _{wrk} (%))
Tetramethrin	97 (14)	97 (16)	97 (14)	97 (15)	90 (14)	104 (12)	0.969	97 (15)	83 (9)	92 (9)	88 (10)	100 (10)	88 (12)	100 (10)	94 (12)	0.068
Thiabendazol	26 (82)	37 (7)	31 (50)	45 (39)	33 (49)	58 (12)	0.226	45 (39)	0.009	65 (9)	54 (24)	68 (6)	57 (14)	68 (6)	62 (14)	0.012
Thiacloprid	96 (13)	88 (11)	92 (12)	90 (8)	89 (9)	90 (8)	0.157	90 (8)	0.762	92 (1)	88 (8)	90 (4)	96 (11)	90 (4)	93 (9)	0.135
Thiamethoxan	96 (21)	113 (4)	104 (16)	90 (15)	86 (13)	93 (17)	0.076	90 (15)	0.437	97 (10)	94 (9)	99 (10)	93 (9)	99 (10)	96 (10)	0.093
Thiodicarb	86 (14)	77 (6)	82 (12)	76 (8)	78 (9)	74 (5)	0.141	76 (8)	0.183	70 (5)	70 (7)	74 (20)	65 (13)	74 (20)	70 (18)	0.202
Thiophanate-methyl	40 (28)	23 (55)	32 (45)	46 (23)	54 (12)	37 (15)	0.044	46 (23)	0.002	85 (18)	78 (19)	72 (18)	73 (6)	72 (18)	72 (13)	0.935
Toxin T2 ^a	15 (40)	17 (47)	16 (43)	23 (22)	24 (25)	22 (19)	0.623	23 (22)	0.585	109 (4)	109 (6)	84 (3)	87 (3)	84 (3)	86 (3)	0.157
Triadimefon	97 (17)	83 (10)	90 (16)	100 (11)	97 (5)	104 (14)	0.129	97 (5)	0.325	90 (9)	91 (14)	105 (4)	92 (11)	105 (4)	99 (10)	0.052
Triadimenol	193 (92)	-32 (-66)	80 (210)	84 (23)	96 (19)	73 (19)	0.018	84 (23)	0.071	98 (13)	95 (16)	86 (20)	96 (20)	77 (11)	86 (20)	0.075
Triazophos	90 (16)	103 (7)	97 (13)	80 (13)	82 (17)	77 (7)	0.094	80 (13)	0.449	90 (14)	88 (18)	97 (5)	108 (13)	97 (5)	102 (11)	0.065
Trifloxystrobin	89 (19)	84 (10)	86 (15)	94 (14)	98 (10)	89 (16)	0.497	94 (14)	0.104	88 (7)	98 (15)	93 (13)	94 (7)	95 (12)	94 (10)	0.818
Triflumizole	97 (18)	92 (4)	95 (13)	93 (8)	94 (9)	92 (7)	0.554	93 (8)	0.527	86 (7)	93 (5)	96 (4)	89 (7)	96 (4)	92 (7)	0.097
Triticonazole	90 (20)	104 (16)	97 (19)	85 (10)	89 (11)	82 (6)	0.113	85 (10)	0.121	83 (9)	83 (7)	84 (5)	91 (12)	84 (5)	87 (10)	0.138
Zearelenone ^a	45 (109)	-180 (-45)	-67 (-198)	3 (975)	23 (123)	-17 (-50)	0.001	3 (975)	0.011	29 (8)	15 (101)	8 (22)	22 (12)	8 (22)	15 (49)	0.000
Zoxamide	86 (17)	96 (11)	91 (15)	93 (8)	92 (9)	95 (8)	0.092	93 (8)	0.584	89 (13)	95 (12)	96 (5)	94 (7)	96 (5)	95 (6)	0.475

^a Mycotoxins. ^b n.f.r.: not fulfill requirements of SANTE document.




Fig. 4 Percentage of analytes presenting a matrix effect in the ranges of $\pm 20\%$, between $\pm 20\%$ and $\pm 50\%$ and higher than $\pm 50\%$.

that a representative matrix-matched calibration would lead to inaccurate quantification of the analyte. No analyte had a matrix effect within the $\pm 20\%$ range nor was the same matrix effect observed for an analyte in both matrices. Therefore, individual analytical curves for each matrix were used. Table 2 presents a summary of the analytes' matrix effect for *M. officinalis* and *M. sylvestris*.

Method accuracy was determined by assessing trueness (as recovery) and precision (as repeatability and as reproducibility – RSD_r and RSD_{WR} , respectively). *M. officinalis* and *M. sylvestris* were spiked at 12, 20, 50, and $75 \mu\text{g kg}^{-1}$ for pesticides, 2, 5, 10 and $20 \mu\text{g kg}^{-1}$ for mycotoxins of group 1, and 100, 250, 500 and $1000 \mu\text{g kg}^{-1}$ for mycotoxins of group 2, with seven replicates at each level. As shown in Tables 3 and 4, the recovery percentages obtained (70–120%) and the standard deviations associated with the replicates showed $RSD < 20\%$, which are acceptable according to the SANTE document 11312/2021 (ref. 35) for the 157 pesticide residues and mycotoxins in *M. officinalis* and the 152 in *M. sylvestris*.

The LOD and LOQ were established as the lowest tested solution with a $S/N > 3$ and the lowest spiked concentration with acceptable accuracy and precision (RSD_r and RSD_{WR}), respectively, fulfilling the requirements of SANTE document 11312/2021 (ref. 35) for a quantitative method. When the data were analyzed, 117 pesticides presented an LOQ at $10 \mu\text{g kg}^{-1}$, and 15, 14 and 2 pesticides presented an LOQ at 20, 50, and $70 \mu\text{g kg}^{-1}$, respectively, for *M. officinalis*. For *M. sylvestris*, 99 pesticides presented an LOQ at $10 \mu\text{g kg}^{-1}$, and 20, 14 and 6 pesticides presented an LOQ at 20, 50, and $70 \mu\text{g kg}^{-1}$, respectively,

showing that most pesticides met the accuracy and precision requirements at the lowest spiked level.

In some cases, such as diflubenzuron, propamocarb, and triadimenol, an LOQ ($70 \mu\text{g kg}^{-1}$) was achieved in *M. sylvestris* that did not fulfill validation requirements (n.f.r.) for *M. officinalis*. Conversely, analytes validated in *M. officinalis* but not in *M. sylvestris* included bifentazate, diethofencarb, fumonisin B1 and B2, halofenozide, haloxyfop-2-ethoxyethyl, methidathion, omethoate and thiodicarb. Most of these analytes had recovery fluctuations between all 14 replicates, leading to a low precision, indicating the method was not repeatable nor reproducible for these analytes in this specific matrix. For mycotoxins, all four aflatoxins presented an LOQ at $5 \mu\text{g kg}^{-1}$ and ochratoxin A at $10 \mu\text{g kg}^{-1}$.

In this study, two different medicinal herbs, from distinct families and genera, with different pharmacological parts were used for method validation. When comparing the two matrices for all compounds, it is evident that significant deviation in results can occur due to the unique matrix effect caused by each matrix on each analyte. The matrix-matched calibration for both matrices presented similar matrix effects for 111 analytes. Most mycotoxins presented a difference higher than 20% in matrix effect between the two matrices. Fenazaquin, fenhexamid, imazapic, and propyzamid showed signal suppression in *M. officinalis* while in *M. sylvestris*, an enhancement in the analytical signal was observed. More polar compounds, such as acephate, methamidophos, and omethoate, presented the same matrix effect in both matrices, indicating that

a representative matrix could be used without compromising the results.

Commercial sample

Imidacloprid residues ($13 \mu\text{g kg}^{-1}$) were found in a *M. officinalis* sample. However, there is no MRL for this pesticide, meaning there should be no residues in medicinal herbs sold in the country. Only one sample of *M. sylvestris* showed residues of methyl pirimiphos, at a concentration of $11.6 \mu\text{g kg}^{-1}$, which is within the MRL ($4000 \mu\text{g kg}^{-1}$) set by Brazilian legislation.³²

Sample comparisons were carried out with herbarium reference material (SMDB) and *via* anatomical analysis of samples that showed pesticide residues. These evaluations were carried out in the herbarium of the Botanical Garden (SMDB) and in the Laboratory of Plant Taxonomy (Biology Department/UFSM). The sample sold as *M. officinalis* was not confirmed to be this species but was compatible with species of Lamiaceae and Verbenaceae. Thus, the consumer used a species other than *M. officinalis*, and in addition to not having its pharmacological properties, they were also exposed to pesticide residue. The *Malva sylvestris* sample was identified as partially compatible with *Malva* sp., mostly mixed with other Malvaceae species.

Despite the limited sampling, the results obtained suggest the non-application of pesticides or the conscious use of pesticides on the medicinal herbs analyzed. In China, in green tea samples analyzed by Y. Huang *et al.*,⁵⁴ 67% of the samples contained some pesticide residue, and the majority contained more than five pesticides.

Regarding the presence of mycotoxins, none of those studied were detected in the analyzed samples, indicating correct drying and storage. In the study by N. Pallarés *et al.*,⁵⁵ 224 samples of herbal medicines and their infusions were analyzed. The results revealed that aflatoxins B₂, G₁, and G₂ as well as zearalenone, were detected in infusions with incidences $\leq 6\%$ and at concentrations below the limit of quantification up to $82.2 \mu\text{g L}^{-1}$. Even though in this study the majority of samples were not positive for the target compounds, investigations need to continue so that more data can be collected to guide national public policies.

Conclusion

This study presents the first reported method for the determination of over 160 mycotoxins/pesticides in medicinal herbs. The developed approach involves a rapid, simple, and effective extraction applying QuEChERS coupled with dSPE clean-up and LC-TQ-MS/MS quantification, which proved to be sufficiently sensitive to meet the diverse analytical requirements for multi-mycotoxin and multi-pesticide analysis. Through a comprehensive clean-up study, it was determined that a combination of GCB, PSA, and MgSO_4 provided the optimal conditions for the simultaneous determination of mycotoxins and pesticides. Validation of the method was conducted using two complex matrices, *M. officinalis* and *M. sylvestris*, demonstrating that the majority of analytes met the criteria outlined in the EU SANTE/11312/2021 method validation guidelines. The method

demonstrates reliable recoveries, as well as excellent accuracy and precision. Additionally, quality controls were implemented for both the extraction process and equipment injection to identify any potential method deviations during the analysis of commercial samples. Analysis of forty-two commercial samples from Southern Brazil revealed the presence of imidacloprid in *M. officinalis* and methyl pirimiphos in *M. sylvestris* underscoring the efficacy of the method for routine analysis of medicinal plants.

Importantly, this method addresses a significant gap in the literature, as specific analytical methods for mycotoxins and pesticides in *M. officinalis* and *M. sylvestris* are currently limited. Consequently, this method represents a valuable tool for monitoring programs aimed at generating data on residue and contaminants in medicinal plants, thereby aiding in the establishment of maximum residue levels (MRLs) and facilitating risk assessment procedures.

Data availability

At this moment, the raw data generated from this study are only available from computers located at the Center of Research and Analysis of Residues and Contaminants (CEPARC) – Chemistry Department – Federal University of Santa Maria, Santa Maria, Brazil. However, the great majority of secondary data obtained are already present in the tables, figures and text submitted here.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to acknowledge Agilent Technologies; The Brazilian Ministry of Science, Technology, and Innovation (MCTI); The Ministry of Agriculture, Livestock and Food Supply (MAPA); The Studies and Projects Finance Organization (FINEP); The National Council for Scientific and Technological Development (CNPq); The Coordination for the Improvement of Higher-Level Personnel (CAPES); Rio Grande do Sul State Research Support Foundation (FAPERGS) – PPSUS 2020 call and Federal University of Santa Maria (UFSM).

References

- 1 C. S. P. Della Pasqua, R. D. Iwamoto, E. Antunes, A. A. Borghi, A. C. H. F. Sawaya and E. C. T. Landucci, *J. Ethnopharmacol.*, 2019, **231**, 50–56, DOI: [10.1016/j.jep.2018.11.012](https://doi.org/10.1016/j.jep.2018.11.012).
- 2 M. K. Purkait, D. Haldar and P. Duarah, in *Advances in Extraction and Applications of Bioactive Phytochemicals*, Elsevier, 2023, pp. 197–217, DOI: [10.1016/B978-0-443-18535-9.00002-8](https://doi.org/10.1016/B978-0-443-18535-9.00002-8).

- 3 B. Motahareh, H. Shahin, M. Masoud and S. Tabandeh, *J. Herb. Med.*, 2022, **31**, 100532, DOI: [10.1016/j.hermed.2021.100532](https://doi.org/10.1016/j.hermed.2021.100532).
- 4 W. A. Abdel-Naime, J. R. Fahim, M. A. Fouad and M. S. Kamel, *S. Afr. J. Bot.*, 2019, **124**, 228–234, DOI: [10.1016/j.sajb.2019.05.011](https://doi.org/10.1016/j.sajb.2019.05.011).
- 5 M. Meneses, A. L. Antonio and S. Cabo Verde, *Appl. Radiat. Isot.*, 2021, **168**, 109391, DOI: [10.1016/j.apradiso.2020.109391](https://doi.org/10.1016/j.apradiso.2020.109391).
- 6 A. Irfan, M. Imran, M. Khalid, M. Sami Ullah, N. Khalid, M. A. Assiri, R. Thomas, S. Muthu, M. A. Raza Basra, M. Hussein, A. G. Al-Sehemi and M. Shahzad, *J. Saudi Chem. Soc.*, 2021, **25**, 101277, DOI: [10.1016/j.jscs.2021.101277](https://doi.org/10.1016/j.jscs.2021.101277).
- 7 M. Seddighfar, S. M. Mirghazanfari and M. Dadpay, *J. Integr. Med.*, 2020, **18**, 181–188, DOI: [10.1016/j.joim.2020.02.003](https://doi.org/10.1016/j.joim.2020.02.003).
- 8 L. Luo, L. Dong, Q. Huang, S. Ma, P. Fantke, J. Li, J. Jiang, M. Fitzgerald, J. Yang, Z. Jia, J. Zhang, H. Wang, Y. Dai, G. Zhu, Z. Xing, Y. Liang, M. Li, G. Wei, J. Song, J. Wei, C. Peng, H. Zhang, W. Zhang, S. Wang, K. Mizuno, A. A. G. Marco, L. Wu, J. Xu, C. Xiong and S. Chen, *Chemosphere*, 2021, **262**, 127477, DOI: [10.1016/j.chemosphere.2020.127477](https://doi.org/10.1016/j.chemosphere.2020.127477).
- 9 N. S. Shaban, K. A. Abdou and N. E.-H. Y. Hassan, *Beni-Suef Univ. J. Basic Appl. Sci.*, 2016, **5**, 102–106, DOI: [10.1016/j.bjbas.2015.10.001](https://doi.org/10.1016/j.bjbas.2015.10.001).
- 10 I. Sedova, M. Kiseleva and V. Tutelyan, *Toxins*, 2018, **10**, 444, DOI: [10.3390/toxins10110444](https://doi.org/10.3390/toxins10110444).
- 11 S. Agriopoulou, E. Stamatelopoulou and T. Varzakas, *Foods*, 2020, **9**, 137, DOI: [10.3390/toxins13060399](https://doi.org/10.3390/toxins13060399).
- 12 D. Pickova, V. Ostry, J. Toman and F. Malir, *Toxins*, 2021, **13**, 399, DOI: [10.3390/toxins13060399](https://doi.org/10.3390/toxins13060399).
- 13 O. M. Areo, O. A. Abafe, S. Gbashi and P. B. Njobeh, *Food Control*, 2023, **143**, 109255, DOI: [10.1016/j.foodcont.2022.109255](https://doi.org/10.1016/j.foodcont.2022.109255).
- 14 Brasil. Ministério da Saúde, Agência Nacional de Vigilância Sanitária, RDC no 105 de 31 de agosto de 2016. Altera a Resolução da Diretoria Colegiada - RDC no 26, de 13 de maio de 2014, que dispõe sobre o registro de medicamentos fitoterápicos e o registro e a notificação de produtos tradicionais fitoterápicos, 2016, available at, <https://www.gov.br/saude/pt-br/composicao/sectics/pnpmf/orientacao-ao-prescritor/Publicacoes/resolucao-rdc-no-26-de-13-de-maio-de-2014.pdf/view>.
- 15 P. Eyring, M. Tienstra, H. Mol, S. S. Herrmann, P. H. Rasmussen, H. L. Frandsen and M. E. Poulsen, *Food Chem.*, 2021, **356**, 129653, DOI: [10.1016/j.foodchem.2021.129653](https://doi.org/10.1016/j.foodchem.2021.129653).
- 16 M. Anastassiades, S. J. Lehotay, D. Stajnbaher and F. J. Schenck, *J. AOAC Int.*, 2003, **86**, 412–431, DOI: [10.1093/jaoac/86.2.412](https://doi.org/10.1093/jaoac/86.2.412).
- 17 S. J. Lehotay, K. Mastovská and A. R. Lightfield, *J. AOAC Int.*, 2005, **88**, 615–629.
- 18 G. Martínez-Domínguez, R. Romero-González and A. Garrido Frenich, *Food Chem.*, 2016, **197**, 907–915, DOI: [10.1016/j.foodchem.2015.11.070](https://doi.org/10.1016/j.foodchem.2015.11.070).
- 19 H. G. J. Mol, P. Plaza-Bolaños, P. Zomer, T. C. de Rijk, A. A. M. Stolker and P. P. J. Mulder, *Anal. Chem.*, 2008, **80**, 9450–9459, DOI: [10.1016/j.foodcont.2022.109255](https://doi.org/10.1016/j.foodcont.2022.109255).
- 20 Y. Sapozhnikova and S. J. Lehotay, *Anal. Chim. Acta*, 2013, **758**, 80–92, DOI: [10.1016/j.aca.2012.10.034](https://doi.org/10.1016/j.aca.2012.10.034).
- 21 M. Leite, A. Freitas, J. Barbosa and F. Ramos, *Food Chem. Adv.*, 2023, **2**, 100145, DOI: [10.1016/j.focha.2022.100145](https://doi.org/10.1016/j.focha.2022.100145).
- 22 J. Rubert, Z. Dzuman, M. Vaclavikova, M. Zachariasova, C. Soler and J. Hajslova, *Talanta*, 2012, **99**, 712–719, DOI: [10.1016/j.talanta.2012.07.010](https://doi.org/10.1016/j.talanta.2012.07.010).
- 23 I. Rodríguez-Cañás, J. M. González-Jartín, R. Alvarino, A. Alfonso, M. R. Vieytes and L. M. Botana, *Food Chem.*, 2023, **408**, 135182, DOI: [10.1016/B978-0-12-805392-8.00016-5](https://doi.org/10.1016/B978-0-12-805392-8.00016-5).
- 24 I. Rodríguez, J. M. González, A. M. Botana, M. J. Sainz, M. R. Vieytes, A. Alfonso and L. M. Botana, in *Liquid Chromatography*, Elsevier, 2017, pp. 479–514, DOI: [10.1016/B978-0-12-805392-8.00016-5](https://doi.org/10.1016/B978-0-12-805392-8.00016-5).
- 25 M. Anastassiades, E. Scherbaum, B. Taşdelen and D. Štajnbaher, in *Pesticide Chemistry*, Wiley, 2007, pp. 439–458, DOI: [10.1002/9783527611249.ch46](https://doi.org/10.1002/9783527611249.ch46).
- 26 L. Zhang, X.-W. Dou, C. Zhang, A. Logrieco and M.-H. Yang, *Toxins*, 2018, **10**, 65, DOI: [10.3390/toxins10020065](https://doi.org/10.3390/toxins10020065).
- 27 Y.-B. Luo, H.-B. Zheng, X.-Y. Jiang, X. Li, H.-F. Zhang, F.-P. Zhu, Y.-Q. Pang and Y.-Q. Feng, *Chin. J. Anal. Chem.*, 2015, **43**, 1538–1544, DOI: [10.1016/S1872-2040\(15\)60870-2](https://doi.org/10.1016/S1872-2040(15)60870-2).
- 28 X. Xu, X. Xu, M. Han, S. Qiu and X. Hou, *Food Chem.*, 2019, **276**, 419–426, DOI: [10.1016/j.foodchem.2018.10.051](https://doi.org/10.1016/j.foodchem.2018.10.051).
- 29 Y.-B. Luo, Z.-G. Shi, Q. Gao and Y.-Q. Feng, *J. Chromatogr. A*, 2011, **1218**, 1353–1358, DOI: [10.1016/j.foodchem.2018.10.051](https://doi.org/10.1016/j.foodchem.2018.10.051).
- 30 X.-T. Peng, L. Jiang, Y. Gong, X.-Z. Hu, L.-J. Peng and Y.-Q. Feng, *Talanta*, 2015, **132**, 118–125, DOI: [10.1016/j.talanta.2014.08.069](https://doi.org/10.1016/j.talanta.2014.08.069).
- 31 E. Rutkowska, B. Łozowicka and P. Kaczynski, *Food Chem.*, 2019, **279**, 20–29, DOI: [10.1016/j.foodchem.2018.11.130](https://doi.org/10.1016/j.foodchem.2018.11.130).
- 32 Brasil. Agência Nacional de Vigilância Sanitária, Farmacopéia Brasileira, plantas medicinais, Anvisa 6a edição, 2019, pp. 1–744, available at <https://www.gov.br/anvisa/pt-br/assuntos/farmacopeia/farmacopeia-brasileira>.
- 33 Codex Alimentarius, *Recommended Methods of Sampling for the Determination of Pesticide Residues for Compliance with Maximum Residue Levels (MRLs)*, 1999, available in: https://apeda.gov.in/apedawebsite/Announcements/CODEX_method_for_sampling_for_determination_of_pesticides.pdf & gt, accessed August 30th, 2021.
- 34 R. C. da Silva, I. D. dos Santos, J. P. Neu, R. D. Wouters, M. E. Z. Fontana, P. D. R. Balbinot, R. Wagner and I. R. Pizzutti, *Food Chem.*, 2022, **394**, 133513, DOI: [10.1016/j.foodchem.2022.133513](https://doi.org/10.1016/j.foodchem.2022.133513).
- 35 European Commission, *Analytical Quality Control and Method Validation Procedures for Pesticide Residues Analysis in Food and Feed Document n° SANTE/11312/2021 v.2*, European Commission Directorate-General for Health and Food Safety, 2024, available at <https://www.eurl->

- pesticides.eu/docs/public/tmpl_article.asp?CntID=727&LabID=100&Lang=EN.
- 36 J. V. Dias, M. da G. P. Nunes, I. R. Pizzutti, B. Reichert, A. A. Jung and C. D. Cardoso, *Food Chem.*, 2019, **293**, 83–91, DOI: [10.1016/j.foodchem.2019.04.088](https://doi.org/10.1016/j.foodchem.2019.04.088).
 - 37 O. Lacina, M. Zachariasova, J. Urbanova, M. Vaclavikova, T. Cajka and J. Hajslova, *J. Chromatogr. A*, 2012, **1262**, 8–18, DOI: [10.1016/j.chroma.2012.08.097](https://doi.org/10.1016/j.chroma.2012.08.097).
 - 38 L. P. da Silva, F. Madureira, E. de Azevedo Vargas, A. F. Faria and R. Augusti, *Food Chem.*, 2019, **270**, 420–427, DOI: [10.1016/j.foodchem.2018.07.126](https://doi.org/10.1016/j.foodchem.2018.07.126).
 - 39 M. Kresse, H. Drinda, A. Romanotto and K. Speer, *J. Chromatogr. B*, 2019, **1117**, 86–102, DOI: [10.1016/j.jchromb.2019.04.013](https://doi.org/10.1016/j.jchromb.2019.04.013).
 - 40 A.-K. Rausch, R. Brockmeyer and T. Schwerdtle, *Anal. Bioanal. Chem.*, 2021, **413**, 3041–3054, DOI: [10.1007/s00216-021-03239-1](https://doi.org/10.1007/s00216-021-03239-1).
 - 41 J. V. Dias, R. C. da Silva, I. R. Pizzutti, I. D. dos Santos, M. Dassi and C. D. Cardoso, *J. Food Compos. Anal.*, 2019, **82**, 103242, DOI: [10.1016/j.jfca.2019.103242](https://doi.org/10.1016/j.jfca.2019.103242).
 - 42 X. Xu, X. Xu, M. Han, S. Qiu and X. Hou, *Food Chem.*, 2019, **276**, 419–426, DOI: [10.1016/j.foodchem.2018.10.051](https://doi.org/10.1016/j.foodchem.2018.10.051).
 - 43 T. W. Na, H.-J. Seo, S.-N. Jang, H. Kim, H. Yun, H. Kim, J. Ahn, H. Cho, S.-H. Hong, H. J. Kim and S. H. Lee, *J. Chromatogr. A*, 2022, **1676**, 463257, DOI: [10.1016/j.chroma.2022.463257](https://doi.org/10.1016/j.chroma.2022.463257).
 - 44 Y. Sapozhnikova, P. Zomer, A. Gerssen, A. Nuñez and H. G. J. Mol, *Food Control*, 2020, **116**, 107323, DOI: [10.1016/j.foodcont.2020.107323](https://doi.org/10.1016/j.foodcont.2020.107323).
 - 45 B. Reichert, A. de Kok, I. R. Pizzutti, J. Scholten, C. D. Cardoso and M. Spanjer, *Anal. Chim. Acta*, 2018, **1004**, 40–50, DOI: [10.1016/j.aca.2017.11.077](https://doi.org/10.1016/j.aca.2017.11.077).
 - 46 M. Cladière, G. Delaporte, E. Le Roux and V. Camel, *Food Chem.*, 2018, **242**, 113–121, DOI: [10.1016/j.foodchem.2017.08.108](https://doi.org/10.1016/j.foodchem.2017.08.108).
 - 47 K. Russo, D. Lucchetti, D. Triolone, P. di Giustino, M. Mancuso, D. Delfino and B. Neri, *Phytochem. Lett.*, 2021, **46**, 153–161, DOI: [10.1016/j.phytol.2021.10.002](https://doi.org/10.1016/j.phytol.2021.10.002).
 - 48 L. Izzo, A. Narváez, L. Castaldo, A. Gaspari, Y. Rodríguez-Carrasco, M. Grosso and A. Ritieni, *J. Dairy Sci.*, 2022, **105**, 2948–2962, DOI: [10.3168/jds.2021-21123](https://doi.org/10.3168/jds.2021-21123).
 - 49 S. Bhattacharyya, R. Poi, M. Baskey Sen, D. Kumar Hazra, R. Ghosh, S. Mandal and R. Karmakar, *Microchem. J.*, 2022, **179**, 107444, DOI: [10.1016/j.microc.2022.107444](https://doi.org/10.1016/j.microc.2022.107444).
 - 50 T.-K. Ly, T.-D. Ho, P. Behra and T.-T. Nhu-Trang, *Food Chem.*, 2020, **326**, 126928, DOI: [10.1016/j.foodchem.2020.126928](https://doi.org/10.1016/j.foodchem.2020.126928).
 - 51 C. S. Vareli, I. R. Pizzutti, L. Gebler, C. D. Cardoso, D. S. H. Gai and M. E. Z. Fontana, *Talanta*, 2018, **184**, 202–209, DOI: [10.1016/j.talanta.2018.03.009](https://doi.org/10.1016/j.talanta.2018.03.009).
 - 52 X. Wu and Z. Ding, *Food Chem.*, 2023, **405**, 134755, DOI: [10.1016/j.foodchem.2022.134755](https://doi.org/10.1016/j.foodchem.2022.134755).
 - 53 W. Li, Y. Liu, J. Duan, C. P. Saint and D. Mulcahy, *J. Chromatogr. A*, 2015, **1389**, 76–84, DOI: [10.1016/j.chroma.2015.02.044](https://doi.org/10.1016/j.chroma.2015.02.044).
 - 54 Y. Huang, T. Shi, X. Luo, H. Xiong, F. Min, Y. Chen, S. Nie and M. Xie, *Food Chem.*, 2019, **275**, 255–264, DOI: [10.1016/j.foodchem.2018.09.094](https://doi.org/10.1016/j.foodchem.2018.09.094).
 - 55 N. Pallarés, H. Berrada, M. Fernández-Franzón and E. Ferrer, *Plant Foods Hum. Nutr.*, 2020, **75**, 362–368, DOI: [10.1007/s11130-020-00820-4](https://doi.org/10.1007/s11130-020-00820-4).