

PAPER

View Article Online
View Journal | View Issue



Cite this: *Environ. Sci.: Nano*, 2025, 12, 3139

Polystyrene nanoplastics trigger changes in cell surface properties of freshwater and marine cyanobacteria†

Nigarsan Kokilathanan, ^a Basirath Raoof^a and Maria Dittrich ^{★ab}

Anthropogenic pressures on aquatic ecosystems have led to threats, including nanoplastics, defined as plastic particles less than 1 μm in diameter. This is concerning as cyanobacterial abundance is expected to increase under rising temperatures and carbon dioxide levels. Cyanobacterial cell surfaces are significant in nutrient and metal uptake and cycling; negatively charged functional groups and extracellular polymeric substances (EPS) help promote calcium carbonate formation along cell surfaces. Yet, the impacts of nanoplastics on cyanobacterial physiology and surface properties remain unclear. Here, we investigated the effects of polystyrene nanoplastics on cell growth, morphology, and surface properties of marine *Synechococcus* sp. PCC8806 and freshwater *Spirulina platensis*, crucial primary producers and players in many biogeochemical processes, under environmentally relevant nanoplastic concentrations. Nanoplastic aggregation and elevated secretion of EPS minimized negative impacts on growth and morphology by reducing potential nanoplastic–cell interactions. Polystyrene nanoplastics modified cell surface compositions by increasing polysaccharide and lipid contents by up to 35% and 37% in *Synechococcus* and *Spirulina*, respectively. Additionally, cell wall thickness increased by 15–20 nm for both cyanobacterial species. Together, these findings demonstrated the impacts of nanoplastics on cell surfaces and the coping mechanisms cyanobacteria employ against polystyrene nanoplastics.

Received 29th December 2024,
Accepted 16th April 2025

DOI: 10.1039/d4en01223b

rscl.es-nano

Environmental significance

With the rapid rise of plastic production globally, there are ecological and human health concerns about the increased presence of nanoplastics within aquatic environments. Cyanobacteria and their cell surfaces provide key ecosystem services as primary producers. In this study, we examine the impacts of polystyrene nanoplastics at environmentally relevant concentrations on the physiology and cell surface properties of freshwater and marine cyanobacteria. Our findings showcased the role of extracellular polymeric substances (EPS) secretion and cell surface modifications as coping mechanisms cyanobacteria employ to mitigate any negative impacts of polystyrene nanoplastics on cyanobacterial physiology. Studying the impacts of polystyrene nanoplastics on cell surface properties can provide insights into the physiological and biogeochemical functions that cyanobacteria contribute to aquatic environments.

Introduction

Plastic pollution, a pressing global threat, has far-reaching adverse consequences. The plastic debris, often discarded in landfills, can end up in aquatic environments through processes like surface runoff and wastewater treatment plant (WWTP) effluent.¹ Within aquatic systems, plastic debris can undergo weathering processes, forming nanoplastics (NPLs), plastic particles less than 1 μm in diameter.^{2,3} When

transported into aquatic systems, NPLs can also be generated from the progressive degradation of bulk plastic particles and microplastics (MPs).⁴ The formation processes of NPLs can lead to physical and chemical heterogeneity of NPLs and their environmental fate.⁵ The size, shape, surface charge, and chemical reactivity of NPLs are influenced by their formation processes and environmental factors, *i.e.*, pH, ionic strength, and natural organic matter. For example, weathering and fragmentation of bulk plastics increase the abundance of oxygen and hydroxyl radicals (OH $\cdot$) on the plastic surface, making bulk plastics more susceptible to photodegradation and further degradation to NPLs.⁶ Weathering of NPLs by ultraviolet radiation can lead to cracks due to the oxidation of active functional groups.⁷ Conversely, the charge of surface functional groups on NPLs and interacting environmental factors can promote or hinder

^a Biogeochemistry Group, Department of Physical and Environmental Sciences, University of Toronto Scarborough, 1065 Military Trail, Toronto, ON, M1C 1A4, Canada. E-mail: m.dittrich@utoronto.ca

^b Department of Earth Sciences, University of Toronto St. George, 22 Ursula Franklin Street, Toronto, ON, M5S 3B1, Canada

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4en01223b>



cyanobacterial cell surfaces. Any slight reduction in cyanobacterial populations, primary production, or nutrient cycling by NPLs will severely affect ecosystem services and functioning, zooplankton, and young fish populations.

The effects of NPLs on cell surfaces remain understudied despite the importance of cyanobacterial cell surfaces. To the best of our knowledge, no studies have been documented on the detailed impact of NPLs on the cell surface properties of *Synechococcus* and *Spirulina*. Furthermore, an essential aspect of interpreting laboratory studies and upscaling the findings for environmental systems is the choice of NPL concentrations reported in nature. So far, most laboratory studies have used high NPL concentrations, including studies exposing *Synechococcus* to NPLs.² Such concentrations would only be applicable based on future trends of plastic waste in aquatic environments. These laboratory studies may not accurately represent the environmental impact of NPLs since field studies on NPLs documented a range of concentrations obtained with different protocols. The lack of a defined set of protocols limits the determination of environmentally relevant NPL concentrations. However, multiple studies suggested that environmentally relevant concentrations are 2–20 mg L⁻¹ NPLs.^{28–31} Accordingly, we tested a concentration range of 2–20 mg L⁻¹ at the laboratory scale to reflect concentrations that could be found in natural systems.

Picocyanobacteria (0.2–2.0 μm) genera *Synechococcus* are major primary producers in oceans and large freshwater lakes.²¹ Cyanobacteria genera *Spirulina* are widely distributed in freshwater and marine environments and are an essential source of algal biomass, pigments, and antioxidants.²² *Synechococcus* and *Spirulina* are important, particularly as abundances of both cyanobacterial species are expected to increase under rising temperatures and CO_2 levels.²³ In addition, *Synechococcus* and *Spirulina* can combat CO_2 increases in freshwater and marine environments through photosynthetic activity and CaCO_3 formation along their cell surfaces, which serves as a buffer against the acidification of water bodies.^{24,25} With the combination and increasing nano-fractions of plastics based on future trends of bulk plastic production and degradation^{26,27} and increased cyanobacterial populations due to climate change, there will potentially be more interactions between NPLs and

that comprise more of the total plastic produced globally than PS.²⁸ Nevertheless, any observed impacts on cell physiology and surface properties can be generalized to other NPL polymers as the primary mode of interaction is *via* sorption of NPLs onto cell surfaces.

Methods

Characterization of PS NPLs

Synthetic 600 nm diameter PS NPLs were purchased from Sigma-Aldrich as 10% solids aqueous solutions with a density of 1.05 g mL⁻¹. The stock PS NPLs underwent dialysis under ambient conditions against 400 mL deionized (DI) water for five days using benzoylated dialysis tubing with a molecular weight cut-off of 2000 in 500 mL beakers. This was done to circumvent any potential influences from the presence of additives and/or surfactants that could amplify the impacts of PS NPLs. The DI water was replaced daily throughout the dialysis process.

The morphology of PS NPLs was characterized using TEM (Hitachi H-7500, Tokyo, Japan) at an accelerating voltage of 80 kV. Samples were sonicated, vortexed, and suspended in DI water and then dried at room temperature on carbon-coated 200 mesh copper grids. Particle size analysis of the PS NPLs was performed using ImageJ software on the TEM images.

Zeta potential (ZP) and hydrodynamic diameter (HD) values of the three PS NPL concentrations (2, 10, and 20 mg L⁻¹) in the two growth media (ASN-III and Zarrouk) and DI water were measured using a NanoBrook Omni (Brookhaven Instruments, Holtsville, NY, USA). Furthermore, ZP measurements of 2, 10, and 20 mg L⁻¹ PS NPLs in 0.1 M NaNO₃ at pH values ranging from 4 to 10 were performed. The pH values were chosen to reflect natural and predicted conditions.³³ The pH was adjusted using 0.1 M HCl or 0.1 M NaOH. Samples were vortexed before each measurement and were measured at 25 °C. Measurements were performed as triplicates with three measurements per replicate ($n = 9$). Particle number concentrations of 2, 10, and 20 mg L⁻¹ PS NPLs were calculated (see ESI†).

Cyanobacterial cell cultures

Synechococcus sp. PCC8806 (NCBI: txid195249; accession number: AF448077), isolated from a lagoon in Port-Gentil, Gabon, was received from the Pasteur Culture Collection. *Spirulina platensis* LB 2340 (NCBI: txid1521533; accession number: DQ39280), isolated from Natron Lake in Chad, was obtained from the University of Texas (UTEX) Austin Culture Collection of Algae. PCC8806 was grown in ASN-III medium under a constant light intensity of $38 \pm 3 \mu\text{E m}^{-2} \text{s}^{-1}$ (HOBO Micro Station H21-002 light sensor), room temperature ($24 \pm 1 \text{ }^{\circ}\text{C}$), and shaken at 130 rpm continuously. *Spirulina* was grown in Zarrouk medium under continuous light intensity ($29 \pm 7 \mu\text{E m}^{-2} \text{s}^{-1}$), temperature ($33 \pm 1 \text{ }^{\circ}\text{C}$), and shaken at the same speed as PCC8806. See ESI† for culture media composition. All

culture media were prepared using autoclaved DI water; glassware was washed with 5% (v/v) nitric acid (HNO₃), followed by rinses with Milli-Q grade water (18.2 MΩ cm). Rinsed glassware was autoclaved before exposure experiments.

Cell count calibration curves were constructed for PCC8806 and *Spirulina* based on optical density (OD) readings (Fig. S1†). OD readings were measured at 680 nm for *Spirulina* and 650 nm for PCC8806 using a GENESYS 10S ultraviolet-visible (UV-Vis) spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). OD 650 and OD 680 were chosen as they represent the number of living cyanobacterial cells with chlorophyll pigments on their intact cell surfaces.¹² The cell count calibration curves were based on previous studies of the strains from our research group.^{24,34,35} The validation arises from plotting the cell numbers against the OD readings.

Exposure of cyanobacteria to PS NPLs

PCC8806 and *Spirulina* were grown in 125 mL flasks containing 75 mL ASN-III and Zarrouk media with an initial cell count of 2.8×10^6 cells per L and 1.1×10^5 cells per L, respectively. Cultures were inoculated from parent cultures at the early exponential phase. The cyanobacteria were then exposed to environmentally relevant PS NPL concentrations (2, 10, and 20 mg L⁻¹), with a negative control of 0 mg L⁻¹. We chose these environmentally relevant NPL concentrations based on predicted and previous MP and NPL studies ranging from 0.01 to 100 mg L⁻¹.²⁸ Each experimental condition had three independent replicates with separate cultures and was exposed to the same growing conditions described above. Each replicate was spaced sufficiently to avoid overshadowing or interferences amongst the flasks. Experiments were conducted at the Biogeochemistry Lab at the Department of Physical and Environmental Sciences at the University of Toronto, Scarborough (43°47'14" N, 79°11'26" W).

OD readings were conducted daily until the cells reached the stationary phase, at 60 days for PCC8806 and 27 days for *Spirulina*. The OD readings of the PS NPLs in the growth media were subtracted from the OD readings of the cells exposed to PS NPLs. Average specific growth rates were obtained from Gompertz model curve fitting using cell counts. Visualization of morphology and ultrastructure and measurement of cell surface properties were performed at the exponential growth phase, at 48 and 12 days of exposure for PCC8806 and *Spirulina*, respectively.

To avoid potential contamination from environmental NPLs or MPs, cotton lab coats and powder-free nitrile gloves were worn throughout the experiments. In a biosafety cabinet, measurements and sample preparation involving cyanobacteria were performed under sterile conditions (using sterile glassware) within a biosafety cabinet. Benches and surfaces were cleaned and disinfected with 70% (v/v) ethanol.



Scanning electron microscopy and transmission electron microscopy

To visualize cell morphology under the presence of PS NPLs, cell suspensions were centrifuged at $10\,610 \times g$ for 10 min at room temperature; the growth medium was discarded. The pellets were washed four times with Milli-Q water before being fixed with 2% (wt/vol) glutaraldehyde in 0.1 M sodium cacodylate (pH = 7.4) for two hours at 4 °C in the dark. After the incubation, the pellets were washed twice with Milli-Q water and filtered through 0.2 μm black polycarbonate (PC) filters.^{13,36} The filters were air-dried overnight, mounted on metal stubs, and sputter-coated with gold. SEM images were obtained on a field-emission SEM (FESEM; Hitachi SU-7000, Tokyo, Japan) using an acceleration voltage of 1.5 kV.

To visualize the ultrastructure, the cell suspensions were centrifuged and washed four times in Milli-Q water before being fixed with 2% (wt/vol) glutaraldehyde in 0.1 M sodium cacodylate (pH = 7.4) overnight at 4 °C in the dark.³⁷ The cell pellets were then embedded in 12% (wt/vol) gelatin and rinsed three times in $1\times$ phosphate buffer saline (PBS) washing buffer (pH = 7.4). After the rinses, the cell pellets were fixed with 1% (wt/vol) osmium tetroxide in the washing buffer for two hours at room temperature. Following the fixation, the cell pellets underwent a graded ethanol dehydration series (50%, 70%, 95%, and 100% ethanol). The pellets were infiltrated under various ratios of 100% ethanol–Spurr resin (5:1, 3:1, 1:1, 1:3, 1:3, pure resin, and pure resin) before being embedded in fresh resin. After embedding, thin sections were cut by a DiATOME Ultra 45° diamond knife under a Leica Ultracut UCT ultramicrotome microscope and mounted on copper grids. The grids were stained with uranyl acetate and lead citrate and dried for electron contrast in TEM images. TEM images were taken on a Hitachi H-7500 TEM, using an accelerating voltage of 80 kV. As a control, 2 mL of stock PS NPLs underwent the same sample preparation. Cell wall thickness was measured from the TEM images using ImageJ.

Filters and grids were stored within plates to mitigate any dust and/or NPLs/MP deposition from the surroundings.

Analysis of cyanobacterial cell surface properties

Once the OD readings were taken, cell pellets at the exponential phase were harvested and centrifuged at $10\,610 \times g$ for 10 min at room temperature; the growth medium was discarded. Afterward, the pellets were washed once in Milli-Q water and resuspended in 0.001 M ethylenediaminetetraacetic acid (EDTA). The washing procedure was repeated four times under the same conditions. The washed pellets were transferred to sterile ceramic crucibles and freeze-dried overnight at $-50\text{ }^{\circ}\text{C}$ using a FreeZone 2 freeze dryer (Labconco, Kansas City, MO, USA). The freeze-dried biomass was stored at $-20\text{ }^{\circ}\text{C}$ before ZP measurements were performed.

ZP measurements were conducted by resuspending 0.1 mg dried bacteria per mL in 0.1 M sodium nitrate (NaNO_3); 0.1

M NaNO_3 was selected as the background electrolyte. After the OD readings were measured, the pH was adjusted (pH 4–10) using 0.1 M HCl or 0.1 M NaOH, and the bacterial solution was equilibrated for 1–1.5 hours.³⁸ Measurements were performed in triplicate with five measurements per replicate (100 runs at a temperature of 25 °C) at pH 4–10. Samples were shaken and vortexed before each measurement. Based on the observation under epifluorescence microscopy, most freeze-dried cells maintain their morphology. Therefore, the ZP measured originated from freeze-dried cells.

ATR-FTIR spectra serve as a valuable tool in our research, providing crucial information on the presence of functional groups associated with macromolecules present on cell surfaces. These macromolecules, mainly comprised of carbohydrates/polysaccharides, nucleic acids, proteins, and lipids, are pivotal in determining cell surface functionality and stability. ATR-FTIR (Bruker Ltd., ON, CA) equipped with a single reflection diamond crystal and operating with the OPUS software (v. 7.5 and 8.5) was used to analyze the impacts of NPLs on cell surfaces. The diamond crystal was cleaned with ethanol, and a background spectrum (accumulation of 256 scans) of only the diamond crystal was performed before placing the freeze-dried cells. All measurements were conducted at room temperature, and data were collected from the accumulation of 256 scans from $1800\text{--}600\text{ cm}^{-1}$ with a spectral resolution of 1 cm^{-1} . ATR-FTIR spectra of the freeze-dried cells were taken for each control and NPL treatment group replicate. The averaged ATR-FTIR spectra (as triplicates) were baseline-corrected and normalized to the amide II band. Deconvolution of peak area at $\sim 990\text{ cm}^{-1}$ and $\sim 1656\text{ cm}^{-1}$ (representative of polysaccharides and proteins, respectively) was determined using the Peak Deconvolution app within the OriginPro software (OriginLab Corporation, Northampton, Massachusetts, USA). Peak area ratios at $\sim 990\text{ cm}^{-1}$ and $\sim 1656\text{ cm}^{-1}$ were calculated between the peak area of the NPL-exposed samples and the peak area of the control sample.

Complementary information about cell surfaces under NPL exposure was obtained using XPS. XPS Analyses were conducted on the freeze-dried cell biomass using a Thermo Scientific K-Alpha photoelectron spectrometer with monochromatized Al K α X-ray radiation (nominal 400 μm spot, 10 μm area of analysis). XPS analysis was also done on freeze-dried PS NPLs. Individual peak and survey analyses were obtained for each sample (as duplicates) with a pass energy of 150 eV and 0.2 eV step. High-resolution spectra of carbon (C1s), oxygen (O1s), and nitrogen (N1s) were measured using a pass energy of 50 eV and 0.1 eV step. To fix the C1s component to C–C bonds, 284.8 eV was set as the binding energy scale. In addition, the O1s components were adjusted to exclude the sodium auger peak present due to the washing procedure. XPS technique collects information only from the outermost molecular layers of the surface (2 to 5 nm) due to inelastic scattering of electrons in the sample.



Avantage software was used to fit the XPS spectra peaks. The atomic concentration ratios were calculated based on the fitting of the peak areas normalized to the basis of the acquisition parameters and sensitivity factors provided by the manufacturer. Refer to ESI† for details on the deconvolution of carbon, nitrogen, and oxygen peaks and their functional group assignments.

The surface composition of cell walls can be presented based on three classes of essential constituents: polysaccharides, hydrocarbon-like compounds (lipids), and peptides (proteins). Compounds such as peptidoglycans are considered a combination of peptides and polysaccharides.³⁴ This corresponds to $C_6H_{10}O_5$ for polysaccharides and CH_2 for hydrocarbon-like compounds. The weight fraction ratios of protein (C_{Pr}/C), polysaccharide (C_{PS}/C), and hydrocarbon-like compounds (C_{HC}/C) to total carbon were estimated from the XPS analyses according to the following equations:³⁹

$$[N/C]_{obs} = 0.279(C_{Pr}/C) \quad (1)$$

$$[O/C]_{obs} = 0.325(C_{Pr}/C) + 0.833(C_{PS}/C) \quad (2)$$

$$1 = (C_{Pr}/C) + (C_{PS}/C) + (C_{HC}/C) \quad (3)$$

EPS measurements and visualization

Cell suspensions were preserved in 4% buffered paraformaldehyde (containing 1% sodium dodecyl sulfate) in 1× PBS (pH = 7.4) and stored in −20 °C before EPS extraction and visualization. Cell suspensions were transferred to 1.5 mL microcentrifuge tubes and centrifuged at $11\,648 \times g$ for 10 min to remove the preservative and growth medium. Afterward, the cell pellets were washed three times with 1× PBS.

EPS was extracted according to Khatiwada *et al.*⁴⁰ Briefly, 1.5 mL of 0.9% (wt/vol) NaCl was added to the cell pellets; the samples were then incubated in an 80 °C water bath for 30 min with moderate shaking conditions. After cooling the samples to room temperature for 30 min, they were centrifuged at $7455 \times g$ for 10 min, and the supernatant was collected. Total polysaccharide content was determined using the phenol-sulfuric acid method.⁴¹ 50 μL of the supernatant was added to a 96-well plate, followed by 150 μL of concentrated sulfuric acid and 30 μL of 5% (v/v) phenol. The microplate was placed in a water bath at 90 °C for 5 min and then cooled down at room temperature for 5 min. The microplate was reheated and cooled to room temperature. Once dried, the absorbance of the solution was recorded at 490 nm spectrophotometrically. Blank samples were prepared using DI water instead of the supernatant. Calibration curves were constructed using different glucose concentrations (mg L^{−1}) as a standard. Measurements were performed with three measurements per replicate.

Calcofluor white was used to stain (exo)polysaccharides within EPS and cells, specifically β-D-glucopyranose

polysaccharides.⁴² After preserving and washing the cells with 1× PBS, the samples were stained with 100 μL of 300 mg L^{−1} calcofluor white (dissolved in water) and incubated at room temperature for 30 min in the dark. The stained cells were washed two times using 1× PBS to remove any excess stains before preparing glass slides. The samples were observed under LSCM (Leica Stellaris, Wetzlar, Germany). The excitation/emission wavelengths of cells and calcofluor white were 570/680 nm (red) and 405/435 nm (blue), respectively. Images were taken using a 63×/1.40 (numerical aperture) oil lens with a pinhole of 95.5 μm.

Blue fluorescence represents polysaccharides, and red fluorescence represents chlorophyll. Within the overlay images, a purple fluorescence signal indicates a higher fluorescence intensity of polysaccharides than the chlorophyll fluorescence signal, likely EPS coating the cells. A pink fluorescence signal represents a lower concentration of EPS coating cell surfaces.

Statistical analysis

Standard deviations and standard errors for the three independent replicates were calculated using the OriginPro 2024b software. The normality of the data was tested using the Shapiro–Wilk test. Statistical analysis was performed using one-way analysis of variance (ANOVA) followed by *post hoc* Tukey's honest significant difference (HSD) test. The homogeneity of variance was tested using the Brown–Forsythe test. Statistical significance was defined as $p < 0.05$.

Results and discussion

Characterization of PS NPLs under freshwater and marine conditions

TEM images showed PS NPLs with an average particle diameter of 549.5 ± 25.6 nm (Fig. 1a and S2†), a significant discrepancy from the manufacturer's reported diameter. A significant difference in average ZP and HD values of PS NPLs was observed in ASN-III, Zarrouk, and DI water (Fig. 1b and c). The higher HD values indicate aggregation compared to the nominal size of PS NPLs (549.5 ± 25.6 nm).

The positive trend in ZPs and higher HDs can be attributed to the solution's ionic strength; the ionic strength of ASN-III is higher than the ionic strength of Zarrouk and DI water. Based on the Derjaguin–Landau–Verwey–Overbeek (DLVO) theory, a higher ionic strength reduces the stability of individual particles and decreases electrostatic repulsion between negatively charged PS NPLs due to electrostatic charge screening of the NPLs' surfaces and compression of the electrical double layer.⁴³ Charge neutralization by cations produces a less negative ZP due to the screening of negatively charged surfaces.⁹ Thus, the more positive ZP of 20 mg L^{−1} NPLs in ASN-III (-14.31 ± 0.96 mV) compared to the ZPs in Zarrouk (-29.39 ± 1.69 mV) and DI water (-48.09 ± 1.63 mV) at the same NPL concentration (Fig. 1b) illustrates the influence of electrostatic charge screening.






Fig. 1 Visualization and measurements of PS NPLs in growth media and DI water. **a.** TEM image of 10 mg L⁻¹ PS NPLs in DI water. Average **b.** ZPs and **c.** HDs of the three PS NPLs concentrations in ASN-III (MM), Zarrouk (FM), and DI water. ZPs and HDs of PS NPLs in DI water were measured as a control. Values are shown as mean $\pm$ standard deviation (SD), $n = 9$. Different letters indicate significant differences between treatments ($p < 0.05$).

The compression of the electrical double layer promotes the aggregation and sedimentation of PS NPLs, thereby increasing HD *via* van der Waals interactions.¹ Indeed, we observed significantly higher HDs for 2, 10, and 20 mg L⁻¹ PS NPLs in ASN-III (1626.59 ± 116.43 nm, 806.75 ± 17.55 nm, and 614.51 ± 12.70 nm, respectively) compared to Zarrouk and DI water (Fig. 1c; $p < 0.05$). Likewise, in Schmidtman *et al.*,⁴⁴ the aggregation and sedimentation of 1 μ m PS MPs in 10 mM NaCl were reported despite highly negative ZPs. Additionally, the HDs of PS NPLs in DI water were similar to the nominal size of PS NPLs. Individual particles remain stable due to electrostatic repulsion at high positive or negative ZPs: ZPs higher than 30 mV or lower than -30 mV.³² This scenario applies to

environments with low ionic strength, *i.e.*, freshwater environments.⁴⁵ Therefore, we expect PS NPLs to persist mainly as individual particles in freshwater environments, whereas, in marine environments, we expect PS NPLs to form agglomerates. Furthermore, our findings suggest that the formation and sedimentation of agglomerates potentially mitigate the effects of PS NPLs on aquatic biota more in marine than in freshwater environments. Combined with other environmental factors such as dissolved organic matter, this has implications for the fate, stability, and transport of NPLs in aquatic environments and the extent of their impacts on cyanobacterial cell surfaces. Since agglomerates of PS NPLs will eventually settle to the sediment, this removal of PS NPLs in the water



Environ. Sci.: Nano, 2025, 12, 3139–3157 | 3145




Fig. 2 Morphology of PCC8806 and *Spirulina* exposed to PS NPLs. SEM images of PCC8806 and *Spirulina* cells at the exponential phase: a and b. control, c and d. with 2 mg L^{-1} PS NPLs, e and f. with 10 mg L^{-1} PS NPLs, g and h. with 20 mg L^{-1} PS NPLs. Surface interactions between PS NPLs and cyanobacterial cells were observed. The arrows indicate the presence of PS NPLs and EPS.

cycling.^{25,46} In particular, any inhibition in the uptake of trace and heavy metals by negatively and positively charged binding sites by bacterial surfaces, especially of dominant cyanobacterial species like *Synechococcus*, will have drastic consequences in primary productivity and growth of higher trophic-level organisms.^{56,57}

Moreover, the compression of the electrical double layer and sorption of opposite charge ions, particularly under high ionic strength, can reduce ZPs by promoting more NPL-cell surface interactions.¹¹ Similar hypotheses have been proposed to explain the drop in the ZPs of Gram-negative bacteria *Klebsiella pneumoniae* and green algae *Scenedesmus*

exposed to negatively charged NPLs.^{17,32} Furthermore, the ZPs of PCC8806 significantly decreased as PS NPL concentrations increased. A ZP dependence on PS NPL concentrations lets us assume that increasing particle concentration would enable more sorption and blockage of positively charged functional groups on cell surfaces.

Interestingly, the ZPs of *Spirulina* were found to be exclusively altered at lower pH values (pH 4–6) by PS NPLs (Fig. 3b; $p < 0.05$), contrary to the significant decrease in the ZPs of PCC8806 by the presence of PS NPLs at pH 4–10. Our results indicate a considerable pH dependence on the impacts of NPLs on the surface properties of *Spirulina*. This






Fig. 3 Surface charge of PCC8806 and *Spirulina* under PS NPI exposure. Effects of PS NPIs on the ZPs of a. PCC8806 and b. *Spirulina* cells at the exponential phase in 0.1 M NaNO_3 at pH 4–10. Values are shown as mean $\pm$ SD, $n = 15$. Statistical significance is defined as $p < 0.05$.

pH dependence for cell–NPI interactions carries profound implications for cyanobacterial adaptation to plastic pollution in the face of globally changing aquatic environments. The pH of natural waters ranges from pH 2–12, depending on the buffering capacities of water bodies and dissolved organic matter; fluctuations in pH values can occur during periods of high primary productivity and acidification.⁵⁸ The phenomenon of pH homeostasis, which regulates differences in intracellular and extracellular pH, is primarily maintained by channeling cations like sodium (Na^+) and Ca^{2+} through cell membranes and begins with the sorption of cations onto cell surfaces.^{46,59} Sorption of NPIs would block channel proteins, hindering pH homeostasis

and resulting in more susceptibility to cell lysis. Further consequences of the presence of NPIs include impacts on the uptake and transport of metals and nutrients, *i.e.*, phosphorus (P) and cyanobacterial-induced CaCO_3 formation, as both are linked to cell surfaces.^{34,46} We hypothesize that surface-driven cyanobacterial CaCO_3 precipitation will also be affected by NPIs.

The ATR-FTIR spectra of PCC8806 exhibited strong peaks associated with polysaccharides, including peaks at ~ 990 and $\sim 1040\text{ cm}^{-1}$ relating to C–O and C–C stretching and at $\sim 1150\text{ cm}^{-1}$ for C–O stretching in glycosidic bridges (Fig. 4a). Peaks associated with proteins were also observed, including peaks at 1536 cm^{-1} (N–H bending, C–N stretching, and C=O




Fig. 4 ATR-FTIR spectra of PCC8806 and *Spirulina* under NPI exposure. ATR-FTIR spectra of a. PCC8806 and b. *Spirulina* under control and NPI-exposed conditions. See Tables S1 and S2† for specific assignments of the ATR-FTIR peaks for PCC8806 and *Spirulina*.



stretching) and $\sim 1656\text{ cm}^{-1}$ ($\text{C}=\text{O}$ stretching) of amides in proteins (Table S2†). Similar findings have been reported in Paulo *et al.*,³⁴ peaks associated with polysaccharides (*i.e.*, at $\sim 1000\text{ cm}^{-1}$) and proteins (*i.e.*, at $\sim 1650\text{ cm}^{-1}$) were observed within the ATR-FTIR spectra of PCC8806 grown under different P concentrations. The ATR-FTIR spectra of *Spirulina* revealed strong peaks relating to proteins (Fig. 4b). This includes peaks at $\sim 1650\text{ cm}^{-1}$ and $\sim 1540\text{ cm}^{-1}$ corresponding to amide I and II, respectively. Broad bands were detected within the polysaccharide region, such as at ~ 990 and $\sim 1150\text{ cm}^{-1}$, corresponding to C–O stretching and C–C of pyranose (Table S3†). The obtained ATR-FTIR spectra of *Spirulina* are consistent with a previous study by Kavitha *et al.*,⁶⁰ where strong peaks associated with amide I and II bands and C–O stretching (1070 cm^{-1}) of glycogen were observed. Minor nucleic acid and lipids peaks were also discovered in PCC8806 and *Spirulina* ATR-FTIR spectra.

Variations in the macromolecular composition of cell surfaces are expected in the presence of stressors, such as PS NPLs. For instance, Déniel *et al.*⁶¹ reported increased protein content in the FTIR spectra of microalgae *Chlamydomonas reinhardtii* exposed to 1 mg L^{-1} PS NPLs. Principal component analysis revealed slight variations in polysaccharide and protein bands, indicating a reaction between PS NPLs and *C. reinhardtii*.⁶¹

Under PS NPL exposure, polysaccharide bands at ~ 990 and $\sim 1040\text{ cm}^{-1}$ shifted towards higher wavenumber for PCC8806 and *Spirulina* (Tables S2 and S3†). Furthermore, an increase in the peak area ratio at $\sim 990\text{ cm}^{-1}$ was detected for PCC8806 exposed to 2 and 20 mg L^{-1} PS NPLs and *Spirulina* under all tested PS NPL concentrations compared to the control (Fig. S10a†). Both observations indicate that PS NPLs react with cyanobacterial surfaces, producing more polysaccharides. This contrasts with decreased polysaccharide-related peak intensities in PS NPL-exposed microalgae *Chlorella vulgaris*.¹⁸

Moreover, there were spectral variations within the PCC8806 and *Spirulina* ATR-FTIR spectra towards lower and higher wavenumbers at peaks relating to the protonation and deprotonation of carboxylic acid or ester functional groups in proteins and lipids (at $\sim 1340\text{ cm}^{-1}$, $\sim 1399\text{ cm}^{-1}$, and $\sim 1739\text{ cm}^{-1}$). The peak area ratio at $\sim 1656\text{ cm}^{-1}$ increased and decreased for PCC8806 and *Spirulina* by PS NPL exposure, respectively (Fig. S10b†). The increased PCC8806 protein content is in agreement with previous studies.^{61,62} A decreased protein content in *Spirulina* further confirms that PS NPLs interacted with cyanobacterial surfaces. In Hazeem *et al.*,¹⁸ an increase in the amide I band was observed in conjunction with a decrease in the amide III band ($\sim 1450\text{ cm}^{-1}$). The authors concluded that PS NPLs interacted with proteins on the cell surfaces of *C. vulgaris*, resulting in lower total protein content being detected. *C. vulgaris* exposed to 20 mg L^{-1} PS NPLs revealed a decline in FTIR peak intensities associated with polysaccharides and proteins (amide I, II, and III bands) arising from the observed sorption and destruction of cell membranes. Interestingly, a higher peak intensity at $\sim 701\text{ cm}^{-1}$ was observed within the *Spirulina* FTIR spectra under NPL exposure; this band represents C–H out-of-plane bending vibrations within the PS structure.³² This confirms the presence of PS NPLs on the cell surfaces of *Spirulina*.

Complementary to ATR-FTIR, XPS is a surface characterization method with a penetration depth of a few nanometers.³⁹ In agreement with the findings by ATR-FTIR, the XPS spectra of PCC8806 showed that the carbon component at $\sim 284.8\text{ eV}$ (C1s), which corresponded to carbon bonded solely to carbon and hydrogen,⁶³ was lower under NPL exposure (Table 2). Total carbon mass fraction decreased under 10 and 20 mg L^{-1} NPLs. Qian *et al.*⁶⁴ reported similar findings; exposure to PS NPLs decreased C1s concentrations within EPS. The decrease in C1s concentrations and lower total carbon mass fraction imply

Table 2 Binding energies (eV), mass fractions (%), and high-resolution XPS spectral band assignments of PCC8806 at the exponential phase^{34,39,63}


Element/ratio	Peak energy (eV)	PCC8806				Assignment
		Mass fraction (%)				
		Control	2 mg L ⁻¹ NPLs	10 mg L ⁻¹ NPLs	20 mg L ⁻¹ NPLs	
Total C	284.76 ± 0.11	68.4	68.4	55.3	62.1	
Total O	399.61 ± 0.04	23.6	24.6	38.5	30.5	
Total N	531.63 ± 0.14	8.0	7.0	6.2	7.4	
O/C		0.3	0.4	0.7	0.5	
N/C		0.1	0.1	0.1	0.1	
C1s	284.63 ± 0.08	53.6	61.2	54.4	54.4	C-(C,H)
C1s A	286.01 ± 0.09	29.2	21.2	30.3	29.0	C-(O,N)
C1s B	287.57 ± 0.13	12.6	12.2	10.4	12.9	C=O; O-C-O
C1s C	288.42 ± 0.13	4.6	5.4	4.9	3.7	O=C-OH/C=O
O1s	530.68 ± 0.09	17.0	31.9	16.0	19.2	O=C; P=O, P-O-ring
O1s A	531.64 ± 0.16	48.9	40.2	50.6	49.2	C-OH; C-O-C; P-OH
O1s B	532.78 ± 0.10	34.0	28.0	33.3	31.6	HO-C
N1s	399.61 ± 0.04	91.7	100	100	98.9	Unprotonated amine or amide functional groups
N1s A	401.25 ± 0.37	8.3			1.1	Protonated amine or amide functional groups



The dominant macromolecules of PCC8806 and *Spirulina* cell envelopes were proteins and polysaccharides, respectively. Under PS NPL exposure, we observed an increased relative proportion of polysaccharides by up to 35% for PCC8806 exposed to 10 and 20 mg L⁻¹ PS NPLs (Fig. 5a); a decrease in the weight fraction of lipids accompanied this increase. The relative proportion of proteins remained relatively stable. Lower C1s concentrations, spectral variations within ATR-FTIR spectra, and the increased ATR-FTIR peak area ratios of polysaccharides and proteins (Fig. S10a and Table S2†) reinforce these observations. A decrease in hydrocarbon content could be attributed to favorable interactions of PS NPLs with lipids. PS NPLs could interact with the headgroups of membrane lipids or fatty acid residues within the lipid

Both nitrogen components at ~ 399.5 eV (N1s) and ~ 400.5 eV (N1s A) were detected in PCC8806 and *Spirulina*. [N/C] ratios of PCC8806 and *Spirulina* were similar under NPI exposure (Tables 2 and 3), corresponding to no significant changes in the abundance of nitrogenated compounds, such as amino acids and proteins. These observations differ from our ATR-FTIR results. In particular, the peak area ratio associated with proteins increased for PCC8806 in the presence of PS NPIs (Fig. S10b†). This can be linked to the penetration depth of X-rays during XPS analysis of a few

Element/ratio	Peak energy (eV)	<i>Spirulina</i>				Assignment	
		Mass fraction (%)					
		Control	2 mg L ⁻¹ NPLs	10 mg L ⁻¹ NPLs	20 mg L ⁻¹ NPLs		
Total C	284.65 ± 0.05	55.9	61.0	59.9	68.3		
Total O	399.43 ± 0.35	41.1	33.2	36.5	26.7		
Total N	531.11 ± 0.49	3.0	5.8	3.6	5.0		
O/C		0.7	0.5	0.6	0.4		
N/C		0.1	0.1	0.1	0.1		
C1s	284.61 ± 0.01	52.3	45.0	51.5	50.9	C-(C,H)	
C1s A	285.98 ± 0.11	18.0	33.0	22.6	18.8	C-(O,N)	
C1s B	287.31 ± 0.55	9.7	14.5	8.9	8.5	C=O; O-C-O	
C1s C	288.58 ± 0.42	20.1	7.6	17.0	21.8	O=C-OH/C=O	
O1s	530.70 ± 0.11	66.5	33.2	74.2	35.2	O=C; P=O, P-O-ring	
O1s A	531.74 ± 0.34	19.9	23.0	18.3	24.9	C-OH; C-O-C; P-OH	
O1s B	532.72 ± 0.20	13.6	43.7	7.6	39.9	HO-C	
N1s	399.61 ± 0.06	100	87.4	100	94.8	Unprotonated amine or amide functional groups	
N1s A	399.58 ± 1.36		12.6		5.2	Protonated amine or amide functional groups	



Furthermore, our TEM observations documented that cell wall thickness for PCC8806 and *Spirulina* increased under NPI exposure (Fig. S8 and S9†). The cell wall thickness of PCC8806 rose significantly in the presence of PS NPIs (34.52 ± 3.32 nm vs. 38.27 ± 3.03 nm, 49.51 ± 2.18 nm, and 50.56 ± 3.02 nm under 2, 10, and 20 mg L⁻¹ PS NPIs, respectively); the cell wall thickness of *Spirulina* increased under 10 and 20 mg L⁻¹ PS NPIs (Fig. S9; Table S1;† $p < 0.05$). Similar observations have been revealed in *Chlorella pyrenoidosa* exposed to PS MPs.²⁰ Our results indicate that though the thickness of cell envelopes increased, there was no uniform increase in the outer, peptidoglycan, and inner membrane layers. The outer and peptidoglycan layers may have increased for PCC8806, as both layers are rich in




Fig. 7 EPS secretion of PCC8806 by PS NPLs. LSCM images of PCC8806 cells at the exponential phase. a–c. Control, d–f. with 2 mg L⁻¹ PS NPLs, g–i. with 10 mg L⁻¹ PS NPLs, j–l. with 20 mg L⁻¹ PS NPLs. Scale bar is 5 μm. Cells were stained with calcofluor white (CW) for polysaccharides. Images (c, f, i and l) represent CW staining and chlorophyll fluorescence overlay. Purple color indicates a higher presence of polysaccharides than chlorophyll fluorescence.

NPLs from being in direct contact with, overcoming, and translocating across cell membranes. The contributions from secreted EPS could influence the presence of certain bonds and surface macromolecules detected. In particular, EPS has been shown to reorganize its contents in the presence of external stressors.¹⁸ Thus, besides increased cell wall thickness, EPS could explain the spectral shifts, increased polysaccharide and protein peak area ratios, and increased relative proportion of polysaccharides and lipids on the cell surfaces of NPL-exposed cells (Fig. 4 and 5 and S8†). Moreover, EPS and humic acids secreted as a physiological response to NPLs can facilitate eco-corona formation on NPLs, altering cell surface reactivity and reducing their affinity for

cyanobacterial surfaces.^{80,81} The encapsulation and eco-corona formation of NPLs with humic acids were hypothesized to explain the alleviation of the adverse effects on cell growth and photosynthetic activity of *M. aeruginosa*.⁸² However, each EPS strand has a unique electrostatic potential and role depending on its composition.^{79,83} Thus, EPS may influence ZP, particularly for the cyanobacterial cell surfaces of NPL-exposed cells. This has implications for the sorption and uptake of nutrients and metals dependent on cell surfaces and maintaining pH homeostasis. Furthermore, the increased EPS production will influence carbon-to-nutrient ratios in aquatic environments, potentially increasing the abundance of higher trophic-level organisms. Interestingly,



EPS could be a biological strategy for removing NPLs in WWTPs, as conventional methods cannot be used due to size limitations and costs.⁸⁴ Currently, MPs and NPLs are removed mainly by screens, gravity separation, or flocculation with coagulants such as aluminum and ferric salts.⁸⁵ However, residual MPs and NPLs not removed initially will be retained within the solid phase (sludge); this requires costly and careful sludge disposal practices or additional treatment. Therefore, cyanobacteria can serve as a bio-alternative to help prevent the transport and leakage of NPLs to aquatic systems through EPS secretion and biodegradation.

Conclusion

Global plastic production and plastic leakage to aquatic environments are expected to increase drastically over the next few decades, leading to higher levels of NPLs. Hence, further studies into the ecological risks of PS NPLs to cyanobacteria are justified. We observed no significant impact on the specific cell growth and morphology of PCC8806 and *Spirulina* under environmentally relevant PS NPL concentration despite cell-NPL interactions. We attributed these observations to the aggregation of NPLs, elevated EPS secretion, and increased cell wall thickness, thereby reducing the impacts of NPLs. Increased cell wall thickness and EPS secretion minimized the severity and bioavailability of PS NPLs to cell surfaces. On the other hand, our experiments clearly showed that the sorption of PS NPLs altered cyanobacterial cell surface properties since polysaccharide and lipid contents increased in the cell walls of PCC8806 and *Spirulina*, respectively; polysaccharide content in the cell walls of PCC8806 increased up to 35% whereas lipid content increased up to 37% for *Spirulina*. In addition, the presence of PS NPLs on the cell surfaces of *Spirulina* was also detected with ATR-FTIR.

Our findings showcase the coping mechanisms utilized by cyanobacteria exposed to PS NPLs, such as EPS secretion, changes in cell surface composition, and increased cell wall thickness. Specifically for cyanobacterial cell walls, the increases in polysaccharide and lipid contents highlight different strategies elicited by freshwater and marine cyanobacteria involving cell walls. Higher pectin and cellulose levels and increased thickness of outer and peptidoglycan layers may prevent the bioaccumulation of PS NPLs while maintaining growth and morphology for cyanobacteria like PCC8806. Conversely, cyanobacteria like *Spirulina* may increase lipid content and the thickness of the inner membrane as a coping mechanism. Further investigation is required regarding the molecular mechanisms involved in the presence of NPLs. Changes in gene expression within PCC8806 and *Spirulina* may explain the modifications in cell surface properties and the lack of significant changes in cell growth and morphology. Transcriptomic analysis needs to be conducted in future research to understand the mechanisms at play in the presence of PS NPLs.

Studying the impacts of NPLs on cyanobacterial cell surface properties can provide insights into the environmental consequences of NPLs, especially on the processes dependent on cyanobacterial cell surfaces in an increasingly plastic world. NPLs are complex and heterogeneous contaminants, so their risks to aquatic systems and cyanobacteria merit further examination. Though overall cyanobacterial physiology and ecological resilience were confirmed at environmentally relevant NPL concentrations, the effectiveness of the coping mechanisms under higher NPL concentrations remains unclear. Increasing cell wall thickness and EPS production consumes valuable resources, so an insufficient response to NPLs will severely harm cyanobacterial physiology and surface properties. As cyanobacterial species like *Synechococcus* and *Spirulina* are expected to increase in abundance, they will have an expanded role in aquatic systems as primary producers and players in biogeochemical processes (*i.e.*, CaCO₃ precipitation). Thus, any reduction in cyanobacterial populations will have detrimental effects on nutrient cycling and growth of higher-level trophic organisms and our combat against climate change and acidification. It is imperative to understand the ecological risks of NPLs at environmentally relevant concentrations to mitigate any severe effects on cyanobacterial cell surfaces that could be exacerbated under future trends of plastic pollution.

Data availability

Data supporting this article has been included as part of the ESI.† Data will be available on request.

Author contributions

NK – conceptualization, investigation, methodology, formal analysis, visualization, and writing – original draft, and writing – review and editing. BR – investigation and methodology – culturing. MD – conceptualization, investigation, writing – review and editing, supervision, and funding acquisition.

Conflicts of interest

There are no conflicts of interest.

Acknowledgements

MD thank the NSERC fund “Source-specific identification, characterization, and control of microplastics across a remote, rural, and urban gradient” and Canada Foundation for Innovation, Leaders Opportunity Fund for the PhD support for NK. The authors are grateful to Dr. DiLoreto, Dr. Avetisyan, and D. Aceituno-Cacedo for their guidance. We also thank Prof. Ruby Sullan (Nano(materials)-Bio Interactions Lab; University of Toronto, Scarborough) for access to the NanoBrook Omni and F. Perez for training. We wish to acknowledge S. Boccia and P. Brodersen (Open Centre for the Characterization of Advanced Materials (OCCAM); University



References

- ## References
1. A. Pradel, C. Catrouillet and J. Gigault, The environmental fate of nanoplastics: What we know and what we need to know about aggregation, *NanoImpact*, 2023, **29**, 100453.
 2. N. Kokilathasan and M. Dittrich, Nanoplastics: Detection and impacts in aquatic environments – A review, *Sci. Total Environ.*, 2022, **849**, 157852.
 3. A. Cózar, F. Echevarría, J. I. González-Gordillo, X. Irigoien, B. Úbeda, S. Hernández-León, Á. T. Palma, S. Navarro, J. García-de-Lomas, A. Ruiz, M. L. Fernández-de-Puelles and C. M. Duarte, Plastic debris in the open ocean, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 10239–10244.
 4. A. L. Andrady, Microplastics in the marine environment, *Mar. Pollut. Bull.*, 2011, **62**, 1596–1605.
 5. J. Gigault, H. El Hadri, B. Nguyen, B. Grassl, L. Rowenczyk, N. Tufenkji, S. Feng and M. Wiesner, Nanoplastics are neither microplastics nor engineered nanoparticles, *Nat. Nanotechnol.*, 2021, **16**, 501–507.
 6. J. Duan, N. Bolan, Y. Li, S. Ding, T. Atugoda, M. Vithanage, B. Sarkar, D. C. W. Tsang and M. B. Kirkham, Weathering of microplastics and interaction with other coexisting constituents in terrestrial and aquatic environments, *Water Res.*, 2021, **196**, 117011.
 7. L. Tian, Q. Chen, W. Jiang, L. Wang, H. Xie, N. Kalogerakis, Y. Ma and R. Ji, A carbon-14 radiotracer-based study on the phototransformation of polystyrene nanoplastics in water versus in air, *Environ. Sci.: Nano*, 2019, **6**, 2907–2917.
 8. G. Hul, H. Okutan, P. Le Coustumer, S. Ramseier Gentile, S. Zimmermann, P. Ramaciotti, P. Perdaems and S. Stoll, Influence of Concentration, Surface Charge, and Natural Water Components on the Transport and Adsorption of Polystyrene Nanoplastics in Sand Columns, *Nanomaterials*, 2024, **14**, 529.
 9. J. Wu, R. Jiang, W. Lin and G. Ouyang, Effect of salinity and humic acid on the aggregation and toxicity of polystyrene nanoplastics with different functional groups and charges, *Environ. Pollut.*, 2019, **245**, 836–843.
 10. M. Dittrich and S. Sibler, Cell surface groups of two picocyanobacteria strains studied by zeta potential investigations, potentiometric titration, and infrared spectroscopy, *J. Colloid Interface Sci.*, 2005, **286**, 487–495.
 11. T. M. Nolte, N. B. Hartmann, J. M. Kleijn, J. Garnæs, D. van de Meent, A. J. Hendriks and A. Baun, The toxicity of plastic nanoparticles to green algae as influenced by surface modification, medium hardness and cellular adsorption, *Aquat. Toxicol.*, 2017, **183**, 11–20.
 12. M. Tavafoghi, S. Garg, A. Korenevski and M. Dittrich, Environmentally friendly antibiofilm strategy based on cationized phytylglycogen nanoparticles, *Colloids Surf., B*, 2021, **207**, 111975–111986.
 13. T. T. S. de Oliveira, I. Andreu, M. C. Machado, G. Vimbela, A. Tripathi and A. Bose, Interaction of Cyanobacteria with Nanometer and Micron Sized Polystyrene Particles in Marine and Fresh Water, *Langmuir*, 2020, **36**, 3963–3969.
 14. T. Zhu and M. Dittrich, Carbonate Precipitation through Microbial Activities in Natural Environment, and Their Potential in Biotechnology: A Review, *Front. Bioeng. Biotechnol.*, 2016, **4**, 4.
 15. Q. Xiang, Y. Zhou and C. Tan, Toxicity Effects of Polystyrene Nanoplastics with Different Sizes on Freshwater Microalgae *Chlorella vulgaris*, *Molecules*, 2023, **28**, 3958.
 16. E. Yousif and R. Haddad, Photodegradation and Photostabilization of Polymers, especially Polystyrene: Review, *Springerplus*, 2013, **2**, 398.
 17. P. Bhattacharya, S. Lin, J. P. Turner and P. C. Ke, Physical Adsorption of Charged Plastic Nanoparticles Affects Algal Photosynthesis, *J. Phys. Chem. C*, 2010, **114**, 16556–16561.
 18. L. J. Hazeem, G. Yesilay, M. Bououdina, S. Perna, D. Cetin, Z. Suludere, A. Barras and R. Boukherroub, Investigation of the toxic effects of different polystyrene micro- and nanoplastics on microalgae *Chlorella vulgaris* by analysis of cell viability, pigment content, oxidative stress and ultrastructural changes, *Mar. Pollut. Bull.*, 2020, **156**, 111278.
 19. F. C. Camini, C. C. da Silva Caetano, L. T. Almeida and C. L. de Brito Magalhães, Implications of oxidative stress on viral pathogenesis, *Arch. Virol.*, 2017, **162**, 907–917.
 20. Y. Mao, H. Ai, Y. Chen, Z. Zhang, P. Zeng, L. Kang, W. Li, W. Gu, Q. He and H. Li, Phytoplankton response to polystyrene microplastics: Perspective from an entire growth period, *Chemosphere*, 2018, **208**, 59–68.
 21. S. Śliwińska-Wilczewska, J. Maculewicz, A. Barreiro Felpeto and A. Latala, Allelopathic and Bloom-Forming Picocyanobacteria in a Changing World, *Toxins*, 2018, **10**, 48.
 22. B. Raoof, B. D. Kaushik and R. Prasanna, Formulation of a low-cost medium for mass production of *Spirulina*, *Biomass Bioenergy*, 2006, **30**, 537–542.
 23. P. Flombaum, J. L. Gallegos, R. A. Gordillo, J. Rincón, L. L. Zabala, N. Jiao, D. M. Karl, W. K. W. Li, M. W. Lomas, D. Veneziano, C. S. Vera, J. A. Vrugt and A. C. Martiny, Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 9824–9829.
 24. A. Liang, C. Paulo, Y. Zhu and M. Dittrich, CaCO₃ biomineralization on cyanobacterial surfaces: Insights from experiments with three *Synechococcus* strains, *Colloids Surf., B*, 2013, **111**, 600–608.
 25. R. Ramanan, K. Kannan, A. Deshkar, R. Yadav and T. Chakrabarti, Enhanced algal CO₂ sequestration through calcite deposition by *Chlorella* sp. and *Spirulina platensis* in a mini-raceway pond, *Bioresour. Technol.*, 2010, **101**, 2616–2622.
 26. M. Chang, P. Sun, L. Zhang, Y. Liu, L. Chen, H. Ren and B. Wu, Changes in characteristics and risk of freshwater

- microplastics under global warming, *Water Res.*, 2024, **260**, 121960.
- 27 L. Peng, D. Fu, H. Qi, C. Q. Lan, H. Yu and C. Ge, Micro- and nano-plastics in marine environment: Source, distribution and threats — A review, *Sci. Total Environ.*, 2020, **698**, 134254.
 - 28 O. Pencik, M. Durdakova, K. Molnarova, A. Kucsera, D. Klofac, M. Kolackova, V. Adam and D. Huska, Microplastics and nanoplastics toxicity assays: A revision towards to environmental-relevance in water environment, *J. Hazard. Mater.*, 2023, **454**, 131476.
 - 29 S. Jiang, H. Lu, Y. Xie, T. Zhou, Z. Dai, R. Sun, L. He and C. Li, Toxicity of microplastics and nano-plastics to coral-symbiotic alga (Dinophyceae Symbiodinium): Evidence from alga physiology, ultrastructure, OJIP kinetics and multi-omics, *Water Res.*, 2025, **273**, 123002.
 - 30 R. Yang, J. Qu, H. Li, W. Meng, X. Xu, J. Guo and F. Fang, Effects of unmodified and amine-functionalized polystyrene nanoplastics on nitrogen removal by *Pseudomonas stutzeri*: strain characteristics, extracellular polymers, and transcriptomics, *Environ. Sci.: Nano*, 2025, **12**, 1531–1544.
 - 31 G. L. Sullivan, J. D. Gallardo, E. W. Jones, P. J. Holliman, T. M. Watson and S. Sarp, Detection of trace sub-micron (nano) plastics in water samples using pyrolysis-gas chromatography time of flight mass spectrometry (PY-GCToF), *Chemosphere*, 2020, **249**, 126179.
 - 32 M. Zając, J. Kotyńska, G. Zambrowski, J. Breczko, P. Deptula, M. Cieśluk, M. Zambrzycka, I. Święcicka, R. Bucki and M. Naumowicz, Exposure to polystyrene nanoparticles leads to changes in the zeta potential of bacterial cells, *Sci. Rep.*, 2023, **13**, 9552.
 - 33 R. Cavicchioli, W. J. Ripple, K. N. Timmis, F. Azam, L. R. Bakken, M. Baylis, M. J. Behrenfeld, A. Boetius, P. H. Boyd, A. T. Classen, T. W. Crowther, R. Danovaro, C. M. Foreman, J. Huisman, D. A. Hutchins, J. K. Jansson, D. M. Karl, B. Koskella, D. B. Mark Welch, J. B. H. Martiny, M. A. Moran, V. J. Orphan, D. S. Reay, J. V. Remais, V. I. Rich, B. K. Singh, L. Y. Stein, F. J. Stewart, M. B. Sullivan, M. J. H. Van Oppen, S. C. Weaver, E. A. Webb and N. S. Webster, Scientists' warning to humanity: microorganisms and climate change, *Nat. Rev. Microbiol.*, 2019, **17**, 569–586.
 - 34 C. Paulo, J. Kenney, P. Persson and M. Dittrich, Effects of Phosphorus in Growth Media on Biomineralization and Cell Surface Properties of Marine Cyanobacteria *Synechococcus*, *Geosciences*, 2018, **8**, 471.
 - 35 T. Zhu, Y. Lin, X. Lu and M. Dittrich, Assessment of cyanobacterial species for carbonate precipitation on mortar surface under different conditions, *Ecol. Eng.*, 2018, **120**, 154–163.
 - 36 M. Seoane, C. González-Fernández, P. Soudant, A. Huvet, M. Esperanza, Á. Cid and I. Paul-Pont, Polystyrene microbeads modulate the energy metabolism of the marine diatom *Chaetoceros neogracile*, *Environ. Pollut.*, 2019, **251**, 363–371.
 - 37 T. Zhu, C. Paulo, M. L. Merroun and M. Dittrich, Potential application of biomineralization by *Synechococcus* PCC8806 for concrete restoration, *Ecol. Eng.*, 2015, **82**, 459–468.
 - 38 I. A. Bundeleva, L. S. Shirokova, P. Bénézet, O. S. Pokrovsky, E. I. Kompantseva and S. Balor, Zeta potential of anoxygenic phototrophic bacteria and Ca adsorption at the cell surface: Possible implications for cell protection from CaCO₃ precipitation in alkaline solutions, *J. Colloid Interface Sci.*, 2011, **360**, 100–109.
 - 39 Y. F. Dufrêne, A. Van Der Wal, W. Norde and P. G. Rouxhet, X-Ray Photoelectron Spectroscopy Analysis of Whole Cells and Isolated Cell Walls of Gram-Positive Bacteria: Comparison with Biochemical Analysis, *J. Bacteriol.*, 1997, **179**, 1023–1028.
 - 40 J. R. Khatiwada, C. Madsen, C. Warwick, S. Shrestha, C. Chio and W. Qin, Interaction between polyethylene terephthalate (PET) microplastic and microalgae (*Scenedesmus* spp.): Effect on the growth, chlorophyll content, and hetero-aggregation, *Environ. Adv.*, 2023, **13**, 100399.
 - 41 T. Masuko, A. Minami, N. Iwasaki, T. Majima, S.-I. Nishimura and Y. C. Lee, Carbohydrate analysis by a phenol-sulfuric acid method in microplate format, *Anal. Biochem.*, 2005, **339**, 69–72.
 - 42 M.-Y. Chen, D.-J. Lee, Z. Yang, X. F. Peng and J. Y. Lai, Fluorescent Staining for Study of Extracellular Polymeric Substances in Membrane Biofouling Layers, *Environ. Sci. Technol.*, 2006, **40**, 6642–6646.
 - 43 G. V. Lowry, R. J. Hill, S. Harper, A. F. Rawle, C. O. Hendren, F. Klaessig, U. Nobbmann, P. Sayre and J. Rumble, Guidance to improve the scientific value of zeta-potential measurements in nanoEHS, *Environ. Sci.: Nano*, 2016, **3**, 953–965.
 - 44 J. Schmidtman, H. Elagami, B. S. Gilfedder, J. H. Fleckenstein, G. Papastavrou, U. Mansfeld and S. Peiffer, Heteroaggregation of PS microplastic with ferrihydrite leads to rapid removal of microplastic particles from the water column, *Environ. Sci.: Processes Impacts*, 2022, **24**, 1782–1789.
 - 45 H. Zhu, X. Fan, H. Zou, R.-B. Guo and S.-F. Fu, Effects of size and surface charge on the sedimentation of nanoplastics in freshwater, *Chemosphere*, 2023, **336**, 139194.
 - 46 M. Dittrich and S. Sibling, Influence of H⁺ and Calcium Ions on Surface Functional Groups of *Synechococcus* PCC 7942 Cells, *Langmuir*, 2006, **22**, 5435–5442.
 - 47 Y. Guo, A. M. O'Brien, T. F. Lins, R. S. Shahmohammadloo, X. O. Almirall, C. M. Rochman and D. Sinton, Effects of Hydrogen Peroxide on *Cyanobacterium Microcystis aeruginosa* in the Presence of Nanoplastics, *ACS ES&T Water*, 2021, **1**, 1596–1607.
 - 48 Y. Song, B. Zhang, M. Si, Z. Chen, J. Geng, F. Liang, M. Xi, X. Liu and R. Wang, Roles of extracellular polymeric substances on *Microcystis aeruginosa* exposed to different sizes of polystyrene microplastics, *Chemosphere*, 2023, **312**, 137225.
 - 49 X. You, X. Cao, X. Zhang, J. Guo and W. Sun, Unraveling individual and combined toxicity of nano/microplastics and ciprofloxacin to *Synechocystis* sp. at the cellular and molecular levels, *Environ. Int.*, 2021, **157**, 106842.
 - 50 X. Zheng, Y. Yuan, Y. Li, X. Liu, X. Wang and Z. Fan, Polystyrene nanoplastics affect growth and microcystin



- production of *Microcystis aeruginosa*, *Environ. Sci. Pollut. Res.*, 2021, **28**, 13394–13403.
- 51 M. C. Machado, G. V. Vimbela, T. T. Silva-Oliveira, A. Bose, A. Tripathi and W.-C. Chin, The response of *Synechococcus* sp. PCC 7002 to micro-/nano polyethylene particles - Investigation of a key anthropogenic stressor, *PLoS One*, 2020, **15**, e0232745.
 - 52 J. Karimi, A. Asgharpour, S. Mohsenzadeh and S. Abbasi, The impact of polystyrene nanoplastics (PSNPs) on physiological and biochemical parameters of the microalgae *Spirulina platensis*, *J. Hazard. Mater.*, 2024, **474**, 134644.
 - 53 S. Stoll and E. Pfefferkorn, Monte Carlo Simulation of Controlled Colloid Growth by Homo- and Heterocoagulation in Two Dimensions, *J. Colloid Interface Sci.*, 1996, **177**, 192–197.
 - 54 R. Li, B. Wang, F. Nan, J. Lv, X. Liu, Q. Liu, J. Feng and S. Xie, Effects of polystyrene nanoplastics on the physiological and biochemical characteristics of microalga *Scenedesmus quadricauda*, *Environ. Pollut.*, 2023, **319**, 120987.
 - 55 E. Bergami, S. Pugnali, M. L. Vannuccini, L. Manfra, C. Faleri, F. Savorelli, K. A. Dawson and I. Corsi, Long-term toxicity of surface-charged polystyrene nanoplastics to marine planktonic species *Dunaliella tertiolecta* and *Artemia franciscana*, *Aquat. Toxicol.*, 2017, **189**, 159–169.
 - 56 Y. Liu, D. S. Alessi, G. W. Owttrim, D. A. Petrash, A. M. Mloszewska, S. V. Lalonde, R. E. Martinez, Q. Zhou and K. O. Konhauser, Cell surface reactivity of *Synechococcus* sp. PCC 7002: Implications for metal sorption from seawater, *Geochim. Cosmochim. Acta*, 2015, **169**, 30–44.
 - 57 J. C. Brown, Q. Mackay, Q. Yu and J. B. Fein, The role of cell surface sulfhydryl and amine binding sites in the removal of Cr(VI) from solution by *Bacillus subtilis* bacterial cells, *Chem. Geol.*, 2024, **664**, 122299–122310.
 - 58 R. G. Wetzel, *Limnology: Lake and River Ecosystems*, Academic Press, San Diego, California, 3rd edn, 2001.
 - 59 N. Giraldez-Ruiz, P. Mateo, I. Bonilla and F. Fernandez-Pinas, The relationship between intracellular pH, growth characteristics and calcium in the cyanobacterium *Anabaena* sp. strain PCC7120 exposed to low pH, *New Phytol.*, 1997, **137**, 599–605.
 - 60 E. Kavitha, L. Devaraj Stephen, B. Fatema Hossain and S. Karthikeyan, Two-trace two-dimensional (2T2D) correlation infrared spectral analysis of *Spirulina platensis* and its commercial food products coupled with chemometric analysis, *J. Mol. Struct.*, 2021, **1244**, 130964.
 - 61 M. Dénél, F. Lagarde, A. Caruso and N. Errien, Infrared spectroscopy as a tool to monitor interactions between nanoplastics and microalgae, *Anal. Bioanal. Chem.*, 2020, **412**, 4413–4422.
 - 62 M. Dénél, N. Errien, F. Lagarde, M. Zanella and A. Caruso, Interactions between polystyrene nanoparticles and *Chlamydomonas reinhardtii* monitored by infrared spectroscopy combined with molecular biology, *Environ. Pollut.*, 2020, **266**, 115227.
 - 63 J. J. Ojeda, M. E. Romero-González, R. T. Bachmann, R. G. J. Edyvean and S. A. Banwart, Characterization of the Cell Surface and Cell Wall Chemistry of Drinking Water Bacteria by Combining XPS, FTIR Spectroscopy, Modeling, and Potentiometric Titrations, *Langmuir*, 2008, **24**, 4032–4040.
 - 64 J. Qian, X. He, P. Wang, B. Xu, K. Li, B. Lu, W. Jin and S. Tang, Effects of polystyrene nanoplastics on extracellular polymeric substance composition of activated sludge: The role of surface functional groups, *Environ. Pollut.*, 2021, **279**, 116904.
 - 65 I. A. Guschina, A. J. Hayes and S. J. Ormerod, Polystyrene microplastics decrease accumulation of essential fatty acids in common freshwater algae, *Environ. Pollut.*, 2020, **263**, 114425.
 - 66 S. Zhang, Z. Sun, T. Zheng, C. He and D. Lin, Nanoplastics increase algal absorption and toxicity of Cd through alterations in cell wall structure and composition, *Water Res.*, 2024, **254**, 121394.
 - 67 A. P. Dean, D. C. Sigeo, B. Estrada and J. K. Pittman, Using FTIR spectroscopy for rapid determination of lipid accumulation in response to nitrogen limitation in freshwater microalgae, *Bioresour. Technol.*, 2010, **101**, 4499–4507.
 - 68 O. Hollóczki and S. Gehrke, Nanoplastics can change the secondary structure of proteins, *Sci. Rep.*, 2019, **9**, 16013.
 - 69 S. Das, V. Thiagarajan, N. Chandrasekaran, B. Ravindran and A. Mukherjee, Nanoplastics enhance the toxic effects of titanium dioxide nanoparticle in freshwater algae *Scenedesmus obliquus*, *Comp. Biochem. Physiol., Part C: Toxicol. Pharmacol.*, 2022, **256**, 109305.
 - 70 S. C. Singh, R. P. Sinha and D. P. Häder, Role of lipids and fatty acids in stress tolerance in cyanobacteria, *Acta Protozool.*, 2002, **41**, 297–308.
 - 71 S. Dai, R. Ye, J. Huang, B. Wang, Z. Xie, X. Ou, N. Yu, C. Huang, Y. Hua, R. Zhou and B. Tian, Distinct lipid membrane interaction and uptake of differentially charged nanoplastics in bacteria, *J. Nanobiotechnol.*, 2022, **20**, 191.
 - 72 W. Chen, J. Xu, Q. Yu, Z. Yuan, X. Kong, Y. Sun, Z. Wang, X. Zhuang, Y. Zhang and Y. Guo, Structural insights reveal the effective *Spirulina platensis* cell wall dissociation methods for multi-output recovery, *Bioresour. Technol.*, 2020, **300**, 122628.
 - 73 J. Li and M. Dittrich, Dynamic polyphosphate metabolism in cyanobacteria responding to phosphorus availability, *Environ. Microbiol.*, 2019, **21**, 572–583.
 - 74 H. Nigam, R. Jain, A. Malik and V. Singh, Effect of different polystyrene nano-plastic concentrations on *Chlorella pyrenoidosa*, *Algal Res.*, 2022, **67**, 102835.
 - 75 Q. Wang, W. Liu, A. Zeb, Y. Lian, R. Shi, J. Li and Z. Zheng, Toxicity effects of polystyrene nanoplastics and arsenite on *Microcystis aeruginosa*, *Sci. Total Environ.*, 2023, **874**, 162496.
 - 76 S. Pereira, A. Zille, E. Micheletti, P. Moradas-Ferreira, R. De Philippis and P. Tamagnini, Complexity of cyanobacterial exopolysaccharides: composition, structures, inducing factors and putative genes involved in their biosynthesis and assembly, *FEMS Microbiol. Rev.*, 2009, **33**, 917–941.



- 77 B. Zhang, Z. Wang, D. Li, L. Li, Y. Zhao, X. Tang and Y. Zhao, Reactive oxygen species mediated extracellular polymeric substances production assisting the recovery of *Thalassiosira pseudonana* from polystyrene micro and nanoplastics exposure, *Environ. Pollut.*, 2024, **348**, 123850.
- 78 W. Yang, P. Gao, Z. Ye, F. Chen and L. Zhu, Micro/nanoplastics and microalgae in aquatic environment: Influence factor, interaction, and molecular mechanisms, *Sci. Total Environ.*, 2024, **934**, 173218.
- 79 K. Ikuma, A. W. Decho and B. L. T. Lau, When nanoparticles meet biofilms—interactions guiding the environmental fate and accumulation of nanoparticles, *Front. Microbiol.*, 2015, **6**, 591.
- 80 J. Saavedra, S. Stoll and V. I. Slaveykova, Influence of nanoplastic surface charge on eco-corona formation, aggregation and toxicity to freshwater zooplankton, *Environ. Pollut.*, 2019, **252**, 715–722.
- 81 Y. Jiao, Y. Zhu, M. Chen, L. Wan, Y. Zhao, J. Gao, M. Liao and X. Tian, The humic acid-like substances released from *Microcystis aeruginosa* contribute to defending against smaller-sized microplastics, *Chemosphere*, 2022, **303**, 135034.
- 82 G. Liu, R. Jiang, J. You, D. C. G. Muir and E. Y. Zeng, Microplastic Impacts on Microalgae Growth: Effects of Size and Humic Acid, *Environ. Sci. Technol.*, 2020, **54**, 1782–1789.
- 83 L. Cui, L. Fan, Z. Li, J. Wang, R. Chen, Y. Zhang, J. Cheng, X. Wu, J. Li, H. Yin, W. Zeng and L. Shen, Characterization of extracellular polymeric substances from *Synechocystis* sp. PCC6803 under Cd (II), Pb (II) and Cr (VI) stress, *J. Environ. Chem. Eng.*, 2021, **9**, 105347.
- 84 L.-J. Feng, J.-J. Wang, S.-C. Liu, X.-D. Sun, X.-Z. Yuan and S.-G. Wang, Role of extracellular polymeric substances in the acute inhibition of activated sludge by polystyrene nanoparticles, *Environ. Pollut.*, 2018, **238**, 859–865.
- 85 M. Keerthana Devi, N. Karmegam, S. Manikandan, R. Subbaiya, H. Song, E. E. Kwon, B. Sarkar, W. Kim, J. Rinklebe and M. Govarthan, Removal of nanoplastics in water treatment processes: A review, *Sci. Total Environ.*, 2022, **845**, 157168–157180.

