

# Chemical Science

Volume 16  
Number 24  
28 June 2025  
Pages 10625–11166

rsc.li/chemical-science



ISSN 2041-6539

**EDGE ARTICLE**

Qingyun Wan, Chi-Ming Che *et al.*

<sup>3</sup>MMLCT excited states of luminescent binuclear Pd<sup>II</sup> complexes: excited state inner-sphere electron-transfer reactions and application

Cite this: *Chem. Sci.*, 2025, 16, 10701

All publication charges for this article have been paid for by the Royal Society of Chemistry

## <sup>3</sup>MMLCT excited states of luminescent binuclear Pd<sup>II</sup> complexes: excited state inner-sphere electron-transfer reactions and application†

Minying Xue,<sup>ab</sup> Wai-Pong To,<sup>id</sup> Gang Cheng,<sup>id</sup> Yuzhen Zhang,<sup>id</sup> Zhou Tang,<sup>a</sup> Lili Du,<sup>e</sup> Kam-Hung Low,<sup>id</sup> Qingyun Wan<sup>\*f</sup> and Chi-Ming Che<sup>id</sup>\*

Compared with Pt<sup>II</sup> analogues that exhibit unique stimulus-induced switching luminescence properties and novel material applications, the properties and reactivity of the <sup>3</sup>MMLCT excited state of Pd<sup>II</sup> complexes in solutions are under-developed. Here, we prepared a series of binuclear cyclometalated Pd<sup>II</sup> complexes with short intramolecular Pd–Pd distances of 2.79–2.89 Å and luminescent <sup>3</sup>MMLCT excited states in solutions at 298 K (emission quantum yield and radiative decay rate constant up to 0.70 and  $2 \times 10^5 \text{ s}^{-1}$ , respectively). Their photophysical properties have been examined by femtosecond time-resolved absorption spectroscopy, and the 1e oxidation products of binuclear Pd<sup>II</sup> complexes have been studied by electron paramagnetic resonance spectroscopy and computational studies. Density functional theory (DFT) and time-dependent DFT (TDDFT) calculations show that changing the C-deprotonated aryl pyridine (C<sup>AN</sup>) ligand to the strong  $\sigma$ -donor aryl N-heterocyclic carbene (C<sup>AC</sup>\*) ligand significantly increases the energy level of the metal centered (<sup>3</sup>dd) excited state. The binuclear Pd<sup>II</sup> complex with a redox-active formamidinate bridging ligand reacts with benzyl bromide to immediately generate Pd<sup>II</sup>–Pd<sup>III</sup>–Br species upon light irradiation. Quenching and time-resolved absorption experiments show that the Pd<sup>II</sup>–<sup>3</sup>MMLCT excited state reacts with alkyl bromides *via* an inner-sphere electron transfer pathway. These binuclear Pd<sup>II</sup> complexes were examined as organic light-emitting diode (OLED) emitters and photocatalysts for C–C bond formation reactions.

Received 21st December 2024

Accepted 4th May 2025

DOI: 10.1039/d4sc08612k

rsc.li/chemical-science

## Introduction

The metal–metal-to-ligand charge transfer [also defined as MMLCT or ( $d\sigma^* \rightarrow \pi^*$ )] excited state of d<sup>8</sup> metal complexes, taking the Pt<sup>II</sup>–<sup>3</sup>MMLCT excited state as an example, exhibits unique photophysical properties and reactivity.<sup>1</sup> This type of excited state usually exhibits luminescence properties that can be reversibly switched by external stimulus,<sup>2</sup> as well as a large radiative decay rate constant ( $k_r \approx 10^5\text{--}10^6 \text{ s}^{-1}$ ) and low emission energy.<sup>3</sup> In addition, due to the presence of axial vacant

coordination sites, the <sup>3</sup>MMLCT excited state is reactive towards the activation of C–X bonds through the inner-sphere atom transfer or electron transfer mechanism.<sup>4</sup> We are interested in developing the photochemistry of binuclear Pd<sup>II</sup> complexes through the <sup>3</sup>MMLCT excited state, which reacts with C–X bonds to produce reactive Pd<sup>III</sup> species (Pd<sup>III</sup>–Pd<sup>II</sup>–X or X–Pd<sup>III</sup>–Pd<sup>II</sup>–X). Compared to Pt analogues, these Pd<sup>III</sup>–Pd<sup>II</sup>–X or X–Pd<sup>III</sup>–Pd<sup>II</sup>–X species are rarely studied and reported. Pd<sup>III</sup> species are generally reactive and unstable, and can be easily reduced back to Pd<sup>II</sup>,<sup>5</sup> suggesting that luminescent binuclear Pd<sup>II</sup> complexes with long-lived <sup>3</sup>MMLCT excited states can be more reactive photocatalysts. In the literature, only a few binuclear Pd<sup>II</sup> complexes with long-lived luminescent <sup>3</sup>MMLCT excited states in solution have been reported.<sup>6</sup> The small ligand-field splitting of the Pd<sup>II</sup> ion results in a low-lying metal-centered <sup>3</sup>dd excited state that is more thermally accessible than in the Pt<sup>II</sup> ion, thus allowing efficient non-radiative decay through electron population in the antibonding  $4d_{x^2-y^2}$  orbital.<sup>7</sup> To destabilize the dd excited state, researchers have employed strong-field C-deprotonated tridentate or tetradentate cyclometalated ligands to form luminescent Pd<sup>II</sup> complexes,<sup>8</sup> or used  $\sigma$ -donating diarylacetylide bridging ligands to construct discrete binuclear Pd<sup>II</sup> complexes.<sup>6a</sup>

<sup>a</sup>State Key Laboratory of Synthetic Chemistry, Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong, P. R. China. E-mail: cmche@hku.hk

<sup>b</sup>Hong Kong Quantum AI Lab Limited, Hong Kong Science Park, Units 909–915, Building 17W, Science Park West Avenue, Pak Shek Kok, Hong Kong, P. R. China

<sup>c</sup>HKU Shenzhen Institute of Research and Innovation, Shenzhen, Guangdong, P. R. China

<sup>d</sup>School of Chemistry and Chemical Engineering, Guangxi Minzu University, No. 158, Daxue West Road, Nanning, Guangxi, P. R. China

<sup>e</sup>School of Life Science, Jiangsu University, Zhejiang, P. R. China

<sup>f</sup>Department of Chemistry, The Chinese University of Hong Kong, Shatin, Hong Kong, P. R. China. E-mail: qingyunwan@cuhk.edu.hk

† Electronic supplementary information (ESI) available. CCDC 1976305, 2018758, 2053114, 2353065, 2353067 and 2353099. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4sc08612k>



We describe here the synthesis, photophysical and photochemical properties of a series of binuclear Pd<sup>II</sup> complexes (**1–7**) and two Pt<sup>II</sup> counterparts (**Ref-Pt1** and **Ref-Pt5**<sup>3c</sup>). Complexes **1** and **3–7** have short intramolecular Pd–Pd distances of 2.79–2.89 Å. Binuclear Pd<sup>II</sup> complexes with cyclometalated [C<sup>∧</sup>C\*] (aryl N-heterocyclic carbene) ligands exhibit strong <sup>3</sup>MMLCT emissions in dilute solution, solid state and PMMA thin films. Based on DFT/TDDFT calculations, two factors affecting the photophysical properties of binuclear Pd<sup>II</sup> complexes include (1) the relative energy of the <sup>3</sup>dd excited state compared to the lowest <sup>3</sup>MMLCT state, which is closely related to the non-radiative decay process ( $k_{nr}$ ); (2) radiative decay process, affected by the spin-orbit coupling constant of Pd atoms. Upon photoirradiation, the binuclear Pd<sup>II</sup> complex with DPF bridging ligand (DPF = *N,N*-diphenylformamidinate) activates alkyl bromides through the inner-sphere type electron transfer pathway, to produce Pd<sup>II</sup>–Pd<sup>III</sup>–Br species. A binuclear Pd<sup>II</sup> complex in this work was also shown to be a photosensitizer or photocatalyst for C–C bond coupling reactions. OLEDs fabricated using binuclear Pd<sup>II</sup> complexes exhibit high maximum luminance and external quantum efficiency (EQE) up to 104 000 cd m<sup>-2</sup> and 22.9%, respectively.

## Results and discussion

### Synthesis and characterization

The structures of the binuclear Pd<sup>II</sup> complexes (**1–7**) and Pt<sup>II</sup> analogues (**Ref-Pt1** and **Ref-Pt5**) are shown in Fig. 1. Their syntheses and characterization data are given in the ESI.† Complexes **1** and **2** were prepared by the reaction of [Pd(C<sup>∧</sup>N)(μ-OAc)]<sub>2</sub> (C<sup>∧</sup>N = 2-[1,1'-biphenyl]-3-yl-4-phenylpyridine) with pyrazole (Pz) ligands and NaOMe in THF. The Pt<sup>II</sup> analogue **Ref-Pt1** was similarly prepared from [Pt(C<sup>∧</sup>N)(μ-Cl)]<sub>2</sub>. These complexes were purified by crystallization to give a mixture of *cis* and *trans* isomers (Fig. 1a), as shown by <sup>1</sup>H NMR (see the ESI†). Complex **3** was prepared by treating [Pd(C<sup>∧</sup>N)(μ-Cl)]<sub>2</sub> with DPF ligand and KO<sup>t</sup>Bu in *N,N*-dimethylformamide (DMF), while **4–7** and **Ref-Pt5** were synthesized by a one-pot reaction; the [M(C<sup>∧</sup>C\*)(μ-Cl)]<sub>2</sub> (M

= Pd or Pt) was generated *in situ* through transmetalation between silver carbene intermediate and Pd(COD)Cl<sub>2</sub> or Pt(COD)Cl<sub>2</sub>. The products were purified by column chromatography to give complexes **3–7** and **Ref-Pt5** as pure *trans* isomers in moderate yields (15–40%). All binuclear Pd<sup>II</sup> complexes were obtained as air-stable light green to orange solids, and **Ref-Pt1** and **Ref-Pt5** are red solids. In the absence of light, complex **5** shows higher stability than complex **1** in aerated CH<sub>2</sub>Cl<sub>2</sub> and toluene; the former is stable for at least a week while the latter decomposed after one day (Fig. S7 and S8†).

The crystal structures of complexes **1–5** and **Ref-Pt1** are shown in Fig. 1b. Complex **2** shows a boat-shaped conformation, similar to that reported for [(ppy)Pd(μ-Pz)]<sub>2</sub> (Hppy = 2-phenylpyridine) with the same unsubstituted Pz ligand.<sup>9</sup> The other complexes adopt a “double-decker” conformation in which the bridging ligand is orthogonal to the plane containing the Pd/Pt atom and the cyclometalated ligand. From their crystal structures, **2** and **Ref-Pt1** exhibit the *cis* configuration. On the other hand, complex **1** shows the *trans* configuration in its crystal structure. As the steric bulk of the bridging Pz ligand increases, the Pd–N(Pz)–N(Pz) angle decreases from 119.8° in **2** to 108.5° in **1**, resulting in a shortening of the Pd–Pd distance from 3.40 Å in **2** to 2.79 Å in **1**. A similar finding was reported by Ma *et al.* in analogous binuclear Pt<sup>II</sup> complexes.<sup>10</sup> **Ref-Pt1**, which has the same ligand scaffold as **1**, also exhibits a close Pt–Pt distance of 2.90 Å. Complexes **3–5** with bridging DPF ligands have short intramolecular Pd–Pd distances of 2.86–2.89 Å and are in the *trans* configuration.

### Absorption and emission spectroscopy

The electronic absorption spectra of binuclear Pd<sup>II</sup> complexes, **Ref-Pt1**, and **Ref-Pt5** are shown in Fig. 2a, b and S5.† The spectral data are summarized in Table 1. In CH<sub>2</sub>Cl<sub>2</sub>, complexes **1**, **2**, and **Ref-Pt1** show an intense absorption band with λ<sub>max</sub> at 268–279 nm ( $\epsilon > 10^4$  M<sup>-1</sup> cm<sup>-1</sup>) and a less intense absorption band at 356–385 nm ( $\epsilon = 5.8 - 7.3 \times 10^3$  M<sup>-1</sup> cm<sup>-1</sup>). Both bands



Fig. 1 (a) Chemical structures of binuclear Pd<sup>II</sup> and Pt<sup>II</sup> complexes in this work. (b) Crystal structures of complexes **1–5** and **Ref-Pt1** (hydrogen atoms are omitted for clarity).





Fig. 2 Absorption spectra of (a) complexes **1**, **2**, and Ref-Pt**1**, (b) complexes **4**–**7** and Ref-Pt**5** in  $\text{CH}_2\text{Cl}_2$  at RT. Emission spectra of (c) complexes **1** and Ref-Pt**1**, (d) complexes **5**–**7** and Ref-Pt**5** in deoxygenated  $\text{CH}_2\text{Cl}_2$  ( $2 \times 10^{-5} \text{ M}$ ) at RT. Emission spectra of (e) complexes **1** and (f) **5** in various solvents. Asterisk (\*) indicates Raman scattering of excitation light.

were assigned to ( $\pi$ - $\pi^*$ ) singlet intraligand ( $^1\text{IL}$ ) transitions localized on the cyclometalated  $\text{C}^*\text{N}$  ligand. For **1**, there is an additional weak absorption band with  $\lambda_{\text{max}}$  at 433 nm ( $\epsilon = 2.8 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ ), and this band is blue-shifted relative to Ref-Pt**1** ( $\lambda_{\text{max}} = 518 \text{ nm}$ ). These low-energy absorption bands are attributed to the  $[\text{d}\sigma^*(\text{Pd}_2/\text{Pt}_2) \rightarrow \pi^*(\text{C}^*\text{N})]$   $^1\text{MMLCT}$  transition, as shown by the DFT calculations (*vide infra*). For **3**–**7**, the intense absorption bands at 242–358 nm ( $\epsilon > 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ ) were assigned to ( $\pi$ - $\pi^*$ )  $^1\text{IL}$  transition of the  $\text{C}^*\text{N}/\text{C}^*\text{C}^*$  or DPF ligands.

Complexes **3** and **5**–**7** show a broad shoulder band tailing to 500 nm ( $\epsilon = 2.0 - 5.5 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ ), which was assigned to a mixed  $[\pi(\text{DPF}) \rightarrow \pi^*(\text{C}^*\text{N}/\text{C}^*\text{C}^*)]$   $^1\text{LLCT}$  (major) and  $[\text{d}\sigma^*(\text{Pd}_2) \rightarrow \pi^*(\text{C}^*\text{N}/\text{C}^*\text{C}^*)]$   $^1\text{MMLCT}$  (minor) transitions (see DFT calculations). The energy of these low-energy absorption bands follows the order  $7 > 5 > 6$ , which is related to the energy level of the  $\pi^*$  orbitals of the  $\text{C}^*\text{C}^*$  ligand. The high energy level of  $\pi^*$  orbitals of the non-conjugated  $\text{C}^*\text{C}^*$  ligand of **4** causes its  $^1\text{LLCT}/^1\text{MMLCT}$  transitions to occur in the region  $< 400 \text{ nm}$ . The low-energy absorption bands ( $\lambda_{\text{max}} > 400 \text{ nm}$ ) of **1** and **5** show

Table 1 UV-visible absorption and emission data of complexes **1**–**7**, Ref-Pt**1**, and Ref-Pt**5** in  $\text{CH}_2\text{Cl}_2$ , solid state and 2 wt% PMMA thin films

	UV-vis absorption <sup>a</sup>	Emission at 298 K		
		$\lambda_{\text{em}}$ [nm] ( $\tau^d$ [ $\mu\text{s}$ ]; $\Phi_{\text{em}}$ [%]; $k_r^i$ [ $10^3 \text{ s}^{-1}$ ]; $k_{\text{nr}}^j$ [ $10^4 \text{ s}^{-1}$ ])		
	$\lambda_{\text{max}}$ [nm] ( $\epsilon$ [ $10^3 \text{ M}^{-1} \text{ cm}^{-1}$ ])	In deoxygenated $\text{CH}_2\text{Cl}_2$ <sup>b</sup>	Solid state <sup>c</sup>	2 wt% in PMMA <sup>c</sup>
<b>1</b>	269 (102.1), 365 (sh, 5.9), 433 (br, 2.8)	609 (0.30; 0.016 <sup>e</sup> ; 0.53; 333)	612 (4.8; 30; 63; 15)	592 (7.1; 3.0; 4.2; 14), 736 (sh)
<b>2</b>	269 (102.2), 356 (sh, 5.8)	Non-emissive	497 (sh), 521 (0.11; <0.1; n.d.; n.d.)	479, 506 (0.29; <0.1; n.d.; n.d.)
<b>3</b>	268 (119.7), 358 (24.9), 463 (br, 2.0)	595 (0.88; 2.0 <sup>f</sup> ; 23; 111)	585 (6.3; 38; 60; 9.8)	583 (9.8; 2.0; 2.0; 10)
<b>4</b>	242 (sh, 75.5), 293 (38.9), 332 (50.7)	Non-emissive	589 (63.9; 70; 11; 0.47)	579 (30.8; 29; 9.4; 2.3)
<b>5</b>	286 (55.7), 338 (40.2), 410 (br, 5.0)	532 (0.45; 3.0 <sup>f</sup> ; 67; 216)	550 (6.2; 45; 73; 8.9)	519 (7.4; 47; 64; 7.2)
<b>6</b>	286 (sh, 50.5), 307 (59.0), 335 (sh, 38.8), 450 (br, 4.7)	576 (5.9; 61 <sup>f</sup> ; 103; 6.6)	566 (0.25; 5.0; 200; 380)	559 (6.7; 67; 100; 4.9)
<b>7</b>	281 (50.1), 337 (47.8), 400 (5.5)	555 (0.062; 0.15 <sup>e</sup> ; 24; 1610)	510 (3.8 <sup>h</sup> ; 27; 71; 19)	541 (12.9; 58; 45; 3.3)
Ref-Pt <b>1</b>	279 (86.9), 385 (7.3), 460 (4.9), 518 (4.5)	806 (0.14; 8.7 <sup>g</sup> ; 621; 652)	661 (0.88; 46; 523; 61)	663 (0.76; 31; 408; 91)
Ref-Pt <b>5</b>	288 (58.6), 328 (sh, 43.6), 472 (br, 8.1)	619 (1.4; 62 <sup>f</sup> ; 443; 27)	628 (1.4; 52; 371; 34)	607 (1.7; 99; 582; 0.59)

<sup>a</sup> At a concentration of  $1 \times 10^{-5} \text{ M}$ . <sup>b</sup> At a concentration of  $2 \times 10^{-5} \text{ M}$ . <sup>c</sup> Measured in air. <sup>d</sup> Emission lifetime. <sup>e</sup> Emission quantum yield was calculated using  $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$  as reference (bpy = 2,2'-bipyridyl, in MeCN,  $\Phi_{\text{em}}$ : 0.062,  $\lambda_{\text{em}}$  = 619 nm). <sup>f</sup> Emission quantum yield was obtained by integrating sphere. <sup>g</sup> Emission quantum yield was calculated using  $[\text{Pt}(\text{tptbp})]$  as reference ( $\text{H}_2\text{tptbp}$  = tetraphenyltetraazaporphyrin, in toluene,  $\Phi_{\text{em}}$ : 0.51,  $\lambda_{\text{em}}$  = 770 nm). <sup>h</sup> Weighted average lifetime for biexponential decay. <sup>i</sup> Radiative decay rate constant estimated from the equation  $k_r = \Phi_{\text{em}}/\tau$ . <sup>j</sup> Non-radiative decay rate constant estimated from the equation  $k_{\text{nr}} = (1 - \Phi_{\text{em}})/\tau$ .



hypsochromic shifts of up to 13 nm with increasing solvent polarity (Fig. S6 and Table S7<sup>†</sup>), further indicating the charge transfer nature of these absorption bands.

All complexes, except 2 and 4, show emission in deoxygenated CH<sub>2</sub>Cl<sub>2</sub> at room temperature (RT) (Fig. 2c, d and Table 1). For the binuclear Pd<sup>II</sup> complex, an unstructured emission band is observed with  $\lambda_{\text{max}}$  at 532–609 nm and  $\tau$  in the sub-microsecond regime. The  $\lambda_{\text{max}}$  of **1** (609 nm) undergoes a hypsochromic shift from its Pt<sup>II</sup> counterpart **Ref-Pt1** (806 nm), consistent with the spectral assignment of the emission of **1** to  $[\text{d}\sigma^*(\text{Pd}_2) \rightarrow \pi^*(\text{C}^{\wedge}\text{N})]$  <sup>3</sup>MMLCT excited state. The  $k_r$  of **1** with  $\tau$  of 0.30  $\mu\text{s}$  and  $\Phi_{\text{em}}$  of <0.01 is small, about  $10^2 \text{ s}^{-1}$ . Replacing the bridging Pz ligands to DPF ligands increases the  $\Phi_{\text{em}}$  and  $k_r$ , as observed in **3** ( $\Phi_{\text{em}}$ : 0.02;  $k_r$ :  $2.3 \times 10^4 \text{ s}^{-1}$ ). On the other hand, the  $\Phi_{\text{em}}$  varies greatly (0.0015–0.61) for binuclear Pd<sup>II</sup> complexes with Pd–NHC bonds (5–7). Both complexes **5** and **7** have lower  $\Phi_{\text{em}}$  than **6**, which is mainly due to significant non-radiative decay in solution at RT ( $k_{\text{nr}}$ :  $2.2 \times 10^6 \text{ s}^{-1}$  [**5**],  $1.6 \times 10^7 \text{ s}^{-1}$  [**7**] vs.  $6.6 \times 10^4 \text{ s}^{-1}$  [**6**]). The emissions of complexes **3** and **5–7** were assigned to a mixed  $[\text{d}\sigma^*(\text{Pd}_2) \rightarrow \pi^*(\text{C}^{\wedge}\text{N}/\text{C}^{\wedge}\text{C}^*)]$  <sup>3</sup>MMLCT and  $[\pi(\text{DPF}) \rightarrow \pi^*(\text{C}^{\wedge}\text{N}/\text{C}^{\wedge}\text{C}^*)]$  <sup>3</sup>LLCT excited state. As the solvent polarity increases from toluene, through CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub> and EA to MeCN, the emission  $\lambda_{\text{max}}$  of **1** shows a hypsochromic shift of up to 11 nm (Fig. 2e). In contrast, the emission  $\lambda_{\text{max}}$  of **5** is bathochromic shifted by up to 7 nm (Fig. 2f). These shifts suggest that the differences in dipole changes between the ground and excited states of **1** and **5** are opposite. The emission of complex **1** is too weak to be detected in MeCN, and the  $\Phi_{\text{em}}$  of complex **5** drops from 0.043 in toluene to 0.007 in MeCN (Table S7<sup>†</sup>).

Complexes **1** and **3–7** show broad emission bands with  $\lambda_{\text{max}}$  at 510–612 nm in 2 wt% PMMA thin films and in the solid state at RT (Fig. S9 and S10<sup>†</sup>). In contrast, the emission band of **2** shows a vibronic structure with the first vibronic peak at 479–497 nm attributed to the  $[\pi-\pi^*]$  <sup>3</sup>IL excited state. Complexes **1** and **3–7** have  $\Phi_{\text{em}}$  spanning from 0.05 to 0.70, and  $\tau$  in the microsecond regime, revealing large  $k_r$  from  $10^4 \text{ s}^{-1}$  to  $2 \times 10^5 \text{ s}^{-1}$ , all of which are significantly larger than the  $k_r$  of **2** ( $\leq 10^3 \text{ s}^{-1}$ ) and other reported Pd<sup>II</sup> complexes with ligand-centered excited states.<sup>11</sup>

### Variable-temperature emission decay lifetime

The binuclear Pd<sup>II</sup> complexes (**1**, **3**, **5**, and **7**) exhibit weak emissions in dilute solutions at 298 K with low  $\Phi_{\text{em}}$  ( $\leq 0.03$ ) and large  $k_{\text{nr}}$  values ( $1.1 - 16.1 \times 10^6 \text{ s}^{-1}$ ). In this regard, the variation of  $\tau_{\text{obs}}$  over a large temperature range provides information on the radiationless decay process of the emitting excited state(s) and could be modelled by eqn (1):<sup>12</sup>

$$k_{\text{obs}} = k_r + k_{\text{nr}} + k_{\text{nr}}(T) = k_0 + A \exp\left(\frac{-E_a}{k_B T}\right) \quad (1)$$

where  $k_0$  is the temperature-independent decay rate of the decay process,  $A$  is the pre-exponential factor of the Arrhenius term that contributes to the non-radiative process,  $E_a$  is the activation energy, and  $k_B$  is the Boltzmann constant. To gain insight into the non-radiative processes of the binuclear Pd complexes, the




Fig. 3 Plots of the temperature dependence of the excited-state decay rate constant ( $k_{\text{obs}}$ ) of (a) complexes **1** ( $5 \times 10^{-5} \text{ M}$ ), **3**, **5**, and **7** ( $2 \times 10^{-5} \text{ M}$ ) and (b) **6** ( $2 \times 10^{-5} \text{ M}$ ) in deoxygenated toluene.

Table 2 Kinetic parameters for excited-state deactivation

Complex	$k_0$ [ $\text{s}^{-1}$ ]	$A_1$ [ $\text{s}^{-1}$ ]	$E_a$ [ $\text{cm}^{-1}$ ]
<b>1</b>	144 852	$1.08 \times 10^{12}$	2605
<b>3</b>	37 601	$2.05 \times 10^{12}$	2989
<b>5</b>	96 000	$2.81 \times 10^{10}$	2088
<b>6</b>	125 769	$1.73 \times 10^8$	1994
<b>7</b>	406 039	$1.56 \times 10^{10}$	1520

temperature-dependent emission lifetimes of complexes **1**, **3**, and **5–7** were measured in deoxygenated toluene. As shown in Fig. 3, using eqn (1), the  $k_{\text{obs}}$  values fit well with  $1/T$ . Complexes **1** and **3** have similarly large  $A$  values, about  $10^{12} \text{ s}^{-1}$  (Table 2), but the  $E_a$  value of **3** (2989  $\text{cm}^{-1}$ ) is slightly larger than that of **1** (2605  $\text{cm}^{-1}$ ). This is consistent with the higher  $\Phi_{\text{em}}$  and smaller  $k_{\text{nr}}$  of **3** compared to **1** in dilute solutions (Table 1). In particular, the  $A$  values for **1** and **3** are approximately  $10^{11}$ – $10^{14} \text{ s}^{-1}$ , suggesting the existence of an activated and rate-determining surface crossing mechanism to reach higher-lying dark electronic state based on previous studies on luminescent d<sup>6</sup> and d<sup>8</sup> metal complexes.<sup>13</sup> Although the  $A$  values of **5** ( $2.81 \times 10^{10} \text{ s}^{-1}$ ) and **7** ( $1.56 \times 10^{10} \text{ s}^{-1}$ ) are about two orders of magnitude smaller than those of **1** and **3**, the  $E_a$  values of **5** (2088  $\text{cm}^{-1}$ ) and **7** (1520  $\text{cm}^{-1}$ ) make the thermal deactivation process non-negligible. Despite having a similar  $E_a$  value to complex **5**, the pre-exponential term of complex **6** is significantly smaller ( $A = 1.73 \times 10^8 \text{ s}^{-1}$ ), which allows moderately efficient emission to occur at RT. The kinetic parameters of complexes **1**, **3**, **5**, and **7** (Table 2) reveal an overwhelming contribution (92–97%) of the temperature-dependent non-radiative decay rate  $k_{\text{nr}}(T)$  to  $k_{\text{obs}}$  at 298 K. For **6**, only 8.2% of the  $k_{\text{obs}}$  come from the  $k_{\text{nr}}(T)$  term. This is in sharp contrast to complexes **1**, **3**, **5**, and **7**, whose excited states undergo significant thermal populated deactivation processes, resulting in these four complexes having lower  $\Phi_{\text{em}}$  and large  $k_{\text{nr}}$  in dilute solutions at 298 K.

### Time-resolved absorption and emission spectroscopy

Complexes **1** and **5** were selected as representative examples to study the excited-state dynamics of binuclear Pd<sup>II</sup> complexes by femtosecond/nanosecond time-resolved absorption difference (fs/ns-TA) and femtosecond time-resolved fluorescence (fs-TRF) spectroscopy. In toluene, different excited state kinetics were observed for the two complexes. The initial TA ( $\sim 0.8 \text{ ps}$ ) of **1** has






Fig. 4 (a) and (b) fs-TA spectra of **5** at selected time delays; (c) fs-TRF spectra and (d) its kinetic trace of **5** in toluene.

two excited-state absorption (ESA) signals at 509 and 610 nm that decay throughout the delay time (0.8 ps–2.5 ns; Fig. S11a†). The kinetics at 509 nm showed two ps decay time constants of 1.1 and 8.8 ps (Fig. S11b†); the former is consistent with TRF decay ( $\tau_{\text{TRF}}$ : 1.0 ps; Fig. S12b†). For **5**, its initial TA ( $\sim 0.9$  ps) is characterized by two ESA signals at 462 and 625 nm (Fig. 4a). The spectral evolution of **5** occurs in three stages. The early phase occurs with a time constant of 2.0 ps and involves an ESA growth at 462 nm. This 2.0 ps time constant is consistent with the TRF decay ( $\tau_{\text{TRF}}$ : 2.5 ps; Fig. 4c and d). In the second stage, with a time constant of 13.8 ps, the TA rises at 462 and 560 nm, with the former undergoing an 18 nm redshift (Fig. 4b). After the second stage of TA evolution, the fs-TA signals undergo a long-lived decay process on the ns timescale to give a TA spectrum consistent with the respective ns-TA spectrum (Fig. S13b†), which can be attributed to the formation of  $T_1$  state. Based on these time-resolved spectroscopic measurements, the excited state cascades of complexes **1** and **5** were constructed (Fig. S14†). Upon excitation, the emission  $\lambda_{\text{max}}$  of the  $S_1$  excited state is at 521 nm for **1** (Fig. S12a†) and 486 nm for **5** (Fig. 4c), both of which are depleted by  $S_1$ -to-triplet ( $T_n$  or  $T_1$ ) intersystem crossing (ISC) with an ultrafast time constant (1.1 ps for **1**; 2.0 ps for **5**). The formed triplet excited state undergoes an internal conversion from  $T_n$  to  $T_1$  or vibrational cooling of  $T_1$  with time constants of 8.8 ps for **1** and 13.8 ps for **5**. This is followed by radiative decay to the ground state in a sub-microsecond timescale (Table 1).

### Electrochemistry

The electrochemical properties of complexes **1–7**, **Ref-Pt1**, and **Ref-Pt5** were investigated by cyclic voltammetry (CV). To avoid undesirable side reactions between oxidized species and coordinating solvent molecules, the non-coordinating solvent  $\text{CH}_2\text{Cl}_2$  was used for anodic scanning.<sup>14</sup> Cathodic scanning was performed in DMF due to its wide electrochemical window for reduction. The CV curves and electrochemical data are shown in Fig. 5, S15† and Table 3, respectively. Complex **1** exhibits a reversible oxidation couple with  $E_{1/2}$  at 0.86 V vs. SCE, while **2** shows an irreversible oxidation wave with  $E_{\text{pa}}$  at 1.22 V vs. SCE.




Fig. 5 Cyclic voltammograms of selected complexes in  $\text{CH}_2\text{Cl}_2$  for anodic sweeps and in DMF for cathodic sweeps with 0.1 M  $\text{NBu}_4\text{PF}_6$  as electrolyte.  $E_{1/2}$  ( $\text{Cp}_2\text{Fe}^{+/0}$ ) was recorded in the range of 0.43–0.52 V vs. SCE in  $\text{CH}_2\text{Cl}_2$  or DMF.

This is consistent with the HOMO of **1** being destabilized by metal–metal interactions. Complexes **3–7** show reversible/quasi-reversible oxidation couple with  $E_{1/2}$  at 0.72–0.87 V vs. SCE. Compared to their  $\text{Pt}^{\text{II}}$  analogues **Ref-Pt1** and **Ref-Pt5**, the  $E_{1/2}$  values of **1** and **5** are anodically shifted by 0.42 and 0.29 V, respectively (Fig. 5). Based on the DFT calculations (*vide infra*), the oxidation process of complex **1** is assigned to be metal-centered, whereas the oxidation process of complex **5** occurs on the DPF ligand. For the reduction process, complexes **1**, **3**, **6**, and **Ref-Pt1** exhibit two reversible reduction couples with  $E_{1/2}$  ranging from  $-1.68$  to  $-1.79$  and  $-1.82$  to  $-2.04$  V vs. SCE, respectively, while **2** exhibits one reversible reduction couple with  $E_{1/2}$  at  $-1.70$  V vs. SCE. The reduction process of complexes **4**, **5**, **7**, and **Ref-Pt5** is irreversible with  $E_{\text{pc}}$  spanning from  $-2.11$  to  $-2.73$  V vs. SCE. The reduction potentials of binuclear  $\text{Pd}^{\text{II}}$  complexes **1–7** vary with the cyclometalated ligands. Also pairs of binuclear complexes with the same cyclometalated and bridging ligands have similar reduction potentials, such as complex **1** ( $E_{1/2} = -1.72$  vs. SCE) and **Ref-Pt1** ( $E_{1/2} = -1.68$  vs. SCE), or complex **5** ( $E_{\text{pc}} = -2.17$  V vs. SCE) and **Ref-Pt5** ( $E_{1/2} = -2.11$  V vs. SCE). This result shows that the reduction of the binuclear  $\text{Pd}^{\text{II}}$  complex in this study is mainly ligand centered.

### DFT/TDDFT calculations

DFT/TDDFT calculations were performed on the binuclear  $\text{Pd}^{\text{II}}$  complex **1** and its  $\text{Pt}^{\text{II}}$  counterpart **Ref-Pt1** to study their electronic structures in the ground and excited states. Because both *trans* and *cis* isomers were observed in the  $^1\text{H}$  NMR spectra of **1** and **Ref-Pt1**, calculations were performed for both isomers. A small energy difference (ranging from 0.04 to 0.06 eV) was observed between the *cis* and *trans* isomers (Table S8†). Since calculations for the *cis* and *trans* isomers show no significant differences, only the complexes with the *trans* geometry are described here. The optimized geometries of **1** and **Ref-Pt1** match well with their crystal structures, as shown in Fig. 6a,



Table 3 Electrochemical data<sup>a</sup> and excited-state redox properties

Complex	$E(M^+/M^0)$ [V]	$E(M^0/M^-, M^-/M^{2-})$ [V]	$E_{0-0}$ <sup>e</sup> [V]	$E(M^+/*M^0)$ <sup>f</sup> [V]	$E(*M^0/M^-)$ <sup>f</sup> [V]
<b>1</b>	0.86 <sup>b</sup>	-1.72, -1.84 <sup>b</sup>	2.26	-1.40	0.42
<b>2</b>	1.22 <sup>c</sup>	-1.70 <sup>b</sup>	—	—	—
<b>3</b>	0.86 <sup>b</sup>	-1.79, -2.04 <sup>b</sup>	2.40	-1.54	0.67
<b>4</b>	0.72 <sup>b</sup>	-2.73 <sup>d</sup>	—	—	—
<b>5</b>	0.81 <sup>b</sup>	-2.17 <sup>d</sup>	2.69	-1.88	0.52
<b>6</b>	0.87 <sup>b</sup>	-1.68, -1.82 <sup>b</sup>	2.45	-1.58	0.80
<b>7</b>	0.78 <sup>b</sup>	-2.27 <sup>d</sup>	2.67	-1.89	0.40
<b>Ref-Pt1</b>	0.44 <sup>b</sup>	-1.68, -1.83 <sup>b</sup>	1.83	-1.39	0.15
<b>Ref-Pt5</b>	0.52 <sup>b</sup>	-2.11 <sup>b</sup>	2.25	-1.73	0.10

<sup>a</sup> Measured in CH<sub>2</sub>Cl<sub>2</sub> for anodic sweeps and DMF for cathodic sweeps with 0.1 M NBu<sub>4</sub>PF<sub>6</sub> as supporting electrolyte at a scan rate of 100 mV s<sup>-1</sup>;  $E_{1/2}$  (Cp<sub>2</sub>Fe<sup>+0</sup>) is recorded at the range of 0.43–0.52 V vs. SCE in CH<sub>2</sub>Cl<sub>2</sub> or DMF. <sup>b</sup> Values refer to  $E_{1/2}$  versus SCE. <sup>c</sup> Value refers to the anodic peak versus SCE. <sup>d</sup> Value refers to the cathodic peak versus SCE. <sup>e</sup> Approximate zero-zero excitation energy,  $E_{0-0}$ , was estimated from the emission onset at 298 K in CH<sub>2</sub>Cl<sub>2</sub>. <sup>f</sup> Estimation of approximate excited-state redox potentials:  $E(M^+/*M^0) = E(M^+/M^0) - E_{0-0}$  (V vs. SCE),  $E(*M^0/M^-) = E(M^0/M^-) + E_{0-0}$  (V vs. SCE).

where the intramolecular Pd–Pd and Pt–Pt distances are calculated to be 2.83 Å and 2.87 Å, respectively. In the calculated absorption spectrum (Fig. 6b), the lowest transition band is located at 462 nm for **1** and 530 nm for **Ref-Pt1**, both close to the experimental values ( $\lambda_{\max}$ : 440 nm for **1**, 518 nm for **Ref-Pt1**). As the distance between two metal atoms is less than the sum of their van der Waals radii, the overlap between the two valence  $nd_{z^2}$  ( $n = 4$  for Pd and 5 for Pt) orbitals will result in the formation of a bonding ( $nd\sigma$ ) and an antibonding ( $nd\sigma^*$ ) orbital. In the MO diagrams shown in Fig. 6c, the HOMOs of both **1** and **Ref-Pt1** consist of the  $nd\sigma^*$  orbital between two

metals, while the LUMO is mainly located at the C<sup>N</sup> ligand. The calculated S<sub>1</sub> states of **1** and **Ref-Pt1** are derived from the HOMO ( $nd\sigma^*$ ) to LUMO ( $\pi^*$ ) transition. The 4d $\sigma^*$  orbital energy level of complex **1** is lower than the 5d $\sigma^*$  orbital of **Ref-Pt1**, which causes the HOMO–LUMO gap of complex **1** to increase and the lowest absorption band to blue-shift. According to the calculation results, the lowest absorption bands (S<sub>0</sub> → S<sub>1</sub>) of **1** and **Ref-Pt1** are attributed to the <sup>1</sup>MMLCT transition, and the calculated oscillator strengths are 0.02 and 0.04, respectively.

The triplet excited state T<sub>1</sub> of **1** and **Ref-Pt1** was also calculated. Calculations show that the M–M distance in the T<sub>1</sub> state



Fig. 6 (a) Optimized structures of **1** and **Ref-Pt1** at S<sub>0</sub> and T<sub>1</sub> state. (b) Calculated absorption spectra of **1** and **Ref-Pt1** based on the optimized S<sub>0</sub> structure. (c) Calculated MO diagrams of **1** and **Ref-Pt1** at T<sub>1</sub> state. (d) Calculated ZFS and SOC coupling of **1** and **Ref-Pt1** at T<sub>1</sub> state.



of **1** is shortened to 2.61 Å and that of **Ref-Pt1** is shortened to 2.65 Å (Fig. 6a). This is due to the M–M bonding interaction formed by excitation of electrons from the antibonding  $d\sigma^*$  orbital to the ligand  $\pi^*$  orbital. The calculated emission wavelength of the  $^3\text{MMLCT}$  excited state of **1** is 632 nm, which is consistent with the experimentally observed emission  $\lambda_{\text{max}}$  at 609 nm.

The phosphorescent radiative decay process of **1** and **Ref-Pt1** was evaluated by calculating the values of zero-field splitting (ZFS) and SOC ( $\langle S_m | H_{\text{SO}} | T_1 \rangle$ ). As shown in Fig. 6d, the calculated ZFS of **Ref-Pt1** ( $39 \text{ cm}^{-1}$ ) is larger than that of **1** ( $3.8 \text{ cm}^{-1}$ ). The ZFS value is closely related to the SOC constant, which is 1412 and  $4000 \text{ cm}^{-1}$  for Pd and Pt, respectively. The contribution of metal orbitals in the singlet and triplet excited states also affects ZFS. Compared to **Ref-Pt1**, complex **1** exhibits a reduced contribution of Pd<sup>II</sup>-4d orbitals in the MMLCT excited state due to the low-lying Pd<sup>II</sup>-4d orbitals, resulting in a smaller ZFS in the Pd<sup>II</sup> complex. For **Ref-Pt1**, the SOC between  $S_2$  and  $T_{1z}$ ,  $S_4$  and  $T_{1y}$ ,  $S_6$  and  $T_{1z}$  states mainly contributes to the  $T_1 \rightarrow S_0$  transition (the  $T_{1x}$ ,  $T_{1y}$ , and  $T_{1z}$  refer to three  $T_1$  substates), with calculated values (SOC) of 404, 467 and  $952 \text{ cm}^{-1}$ , respectively (Fig. 6d). For complex **1**, the  $T_1 \rightarrow S_0$  transition is mainly contributed by the SOC between  $S_6$  and  $T_{1x}$  states, with a value of  $270 \text{ cm}^{-1}$ , which is smaller than that of **Ref-Pt1**.

DFT/TDDFT calculations were also performed on **5** to examine the effect of DPF ligands on the excited state. In the calculated absorption spectrum (Fig. 7a), the  $S_0 \rightarrow S_1$  transition of **5** is located at 418 nm, which is consistent with the experimental observation ( $\lambda_{\text{max}}$ : 410 nm). As shown in the MO diagram (Fig. 7c), the HOMO of **5** consists of  $\pi$  orbitals of the DPF ligand (82%) and the LUMO is mainly located at the C<sup>∗</sup>C<sup>∗</sup> ligand. The  $4d\sigma^*$  orbital is located in the H-2 orbital. The  $S_1$  state of **5** is mainly contributed by the HOMO  $\rightarrow$  LUMO transition, and the H-2  $\rightarrow$  LUMO transition contributes moderately.

Therefore, the lowest absorption band of **5** was assigned to a mixture of  $^1\text{LLCT}/^1\text{MMLCT}$  transitions. In the  $T_1$  excited state, the Pd–Pd distance is significantly shortened (Fig. 7b), which lifts the H-2 orbital ( $4d\sigma^*$ ) to the HOMO. The  $T_1 \rightarrow S_0$  transition of **5** is mainly contributed by the HOMO  $\rightarrow$  LUMO transition. The HOMO of complex **5** at  $T_1$  state contains contributions from both the ligand and the metal, so the  $T_1$  state was assigned as a mixed  $^3\text{MMLCT}/^3\text{LLCT}$  excited state.

Since the ligand field splitting of Pd<sup>II</sup> ions is smaller relative to Pt<sup>II</sup> ions, the thermal population of the metal centered ( $^3\text{dd}$ ) excited state provides a quenching pathway for the emissive  $^3\text{MMLCT}$  excited state of the binuclear Pd<sup>II</sup> complex. In this regard, the relative energies between the  $^3\text{MMLCT}$  and  $^3\text{dd}$  excited states of complexes **1** and **Ref-Pt1** were calculated (Fig. 7d, S18 and S19†). For complex **1**, the  $T_1$  state consists of 90%  $^3\text{MMLCT}$  and 10%  $^3\text{dd}$  characters. The  $^3\text{dd}$  character also has a major contribution to the  $T_2$  state and the  $T_1$ – $T_2$  energy difference is calculated to be 0.27 eV. In comparison, the  $^3\text{dd}$  character does not contribute to the  $T_1$  state of **Ref-Pt1**, but has a major contribution to the  $T_8$  state, which has an energy 1.4 eV higher than the  $T_1$  state. These results indicate that the excited-state deactivation process of **1** through the  $^3\text{dd}$  excited state is thermally more feasible relative to **Ref-Pt1**, leading to a faster non-radiative decay process of **1**. For complexes **5** and **6**, the excited states with dominant  $^3\text{dd}$  character are the  $T_5$  states at 2.92 and 2.77 eV, respectively (Fig. S20 and S21†). These energies are significantly higher than the  $T_2$  state energy of complex **1**, showing that the incorporation of strong  $\sigma$ -donating NHC ligands can destabilize the  $^3\text{dd}$  excited state.<sup>15</sup> The calculated  $T_1$ – $T_5$  energy differences of complexes **5** and **6** are 0.66 eV and 0.69 eV, respectively (Fig. 7d), indicating the deactivation pathway through the  $^3\text{dd}$  state of complexes **5** and **6** is less feasible than for complex **1**.



Fig. 7 (a) Calculated absorption spectrum of **5** (red line) based on the optimized structure, and the experimental absorption spectrum (black line). (b) Optimized structures of complex **5** at ground state ( $S_0$ ) and triplet excited state ( $T_1$ ). (c) Calculated MOs diagram of **5** at  $S_0$  (left) and  $T_1$  (right). (d) Calculated  $^3\text{MMLCT}$  and  $^3\text{dd}$  excited state energy of complexes **1**, **5**, **6**, and **Ref-Pt1**. Note that due to mixing, the state labels may not strictly apply.









Fig. 10 (a) X-band EPR spectrum and simulation of the reaction mixture of **5** and benzyl bromide in frozen benzene at 100 K after light irradiation for 15 minutes. (b) Optimized structure (left) and spin density population (right) of complex  $5^+$ . (c) Optimized structure (left) and spin density population (right) of complex  $5\text{-Br}$ .

by chemical oxidation of **5** with (TBPA)SbCl<sub>6</sub> (Fig. 8d). The Pd–Pd distance shrinks from 2.89 Å in **5** to 2.79 Å in  $5\text{-Br}$  (Fig. 10c), showing Pd–Pd bonding interactions (though weak) due to the oxidation of Pd atoms. The spin density distribution of  $5\text{-Br}$  is mainly localized on Pd and Br atoms, and the values of Pd1, Pd2 and Br atom are 0.14, 0.4 and 0.29, respectively. This significant difference in the spin density population between  $5\text{-Br}$  and  $5^+$  is consistent with the different EPR patterns of  $5\text{-Br}$  and  $5^+$ .

There are three possible reaction mechanisms for the photoreaction of binuclear Pd<sup>II</sup> complex **5** with alkyl bromides, as shown in Scheme 1:<sup>26</sup> (1) in the outer-sphere electron transfer mechanism, the alkyl bromide R–Br receives electrons from the excited state of **5** and undergoes C–Br bond cleavage to generate the bromide anion, which is subsequently added to the metal center; (2) the inner-sphere electron transfer reaction involves the formation of caged ion pairs as intermediates prior to the electron transfer process; (3) alternatively, the excited state of **5** can react with alkyl bromides through a halogen atom abstraction mechanism. The linear correlation between the log  $k_q$  values of **5** and the reduction potentials of the alkyl bromides is inconsistent with the halogen atom abstraction hypothesis, which predicts that the reaction rate depends on the BDE of the alkyl bromide. No formation of  $5^+$  was observed in laser flash


photolysis experiments of **5** with benzyl bromide, whereas the direct observation of  $5\text{-Br}$  on the nanosecond time scale suggests that the reaction most likely follows an inner-sphere electron transfer mechanism.

## Photocatalysis


The bimolecular photoreaction between **5** and alkyl halides prompts us to study the application of **5** as a photocatalyst in the photo-induced intramolecular cyclization of indole/pyrrole (Scheme 2).<sup>27</sup> For all substrates examined, 100% substrate conversion and high yields of cyclized products (68–95%) were achieved (Scheme 2). Both electron-rich indoles and electron-deficient indoles perform well in photo-induced cyclization reactions. Notably, the C3 electron-rich indole gave the highest product yield (95%). The conversion efficiency of 1-(3-iodopropyl)-1H-indole to the cyclized product was less satisfactory (68%), probably due to the higher ring strain of the five-membered fused ring than the six-membered fused ring. This photoredox reaction also worked for pyrrole, furnishing the cyclized product in 72% yield.

With  $E(\text{DIPEA}^+/\text{DIPEA}^0) = 0.65 \text{ V vs. SCE}$  and  $E_{\text{red}}$  of 1-(4-iodobutyl)-1H-indole =  $-1.68 \text{ V vs. SCE}$ , the oxidative quenching of **5** by 1-(4-iodobutyl)-1H-indole is thermodynamically favored by 0.20 V and the reductive quenching of **5** by DIPEA is thermodynamically uphill by 0.13 V. Stern–Volmer quenching experiments showed that 1-(4-iodobutyl)-1H-indole quenches the emission of **5** with a  $k_q$  of  $1.65 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ , while the quenching of the emission of **5** by DIPEA is negligible (Fig. S24<sup>†</sup>). In addition, we observed a long-lived species ( $\tau > 10 \mu\text{s}$ ) in the ns-TA spectrum of a deoxygenated toluene solution

### Photo-induced indoles/pyrroles functionalization by intramolecular cyclization<sup>a</sup>





### Photo-induced aerobic C–C coupling reactions of N-phenyl-1,2,3,4-tetrahydroisoquinoline with acetone<sup>a</sup>





Scheme 2 Photo-induced indoles/pyrroles functionalization by intramolecular cyclization (upper), and aerobic C–C coupling reactions of N-phenyl-1,2,3,4-tetrahydroisoquinoline with acetone (bottom). (a) Reaction conditions: indoles/pyrroles (0.05 mmol), DIPEA (0.10 mmol), complex **5** (2.5 mol%) in 0.56 mL deoxygenated benzene-*d*<sub>6</sub>, light (LED lamp,  $\lambda_{\text{max}} = 400 \text{ nm}$ , 24 W), irradiation time: 2 h. (b) Reaction conditions: N-phenyl-1,2,3,4-tetrahydroisoquinoline (0.128 mmol), acetone (0.4 mL), L-proline (0.128 mmol), PC (1 mol%), MeCN (0.4 mL), MeOH (0.8 mL), oxygen, light (LED lamp,  $\lambda_{\text{max}} = 455 \text{ nm}$ , 24 W), irradiation time: 6 h. Product yields and substrate conversion were calculated by <sup>1</sup>H NMR using 1,3,5-trimethoxybenzene as an internal standard.


### 1. Outer-Sphere Electron Transfer



### 2. Inner-Sphere Electron Transfer



### 3. Halogen Atom Abstraction



Scheme 1 Possible reaction mechanisms for the photoreaction of complex **5** with alkyl bromides.





**Fig. 11** (a) ns-TA spectra at 0.01 and 4  $\mu\text{s}$  after flashing a deoxygenated toluene solution of **5** ( $2 \times 10^{-5}$  M). (b) ns-TA spectra at 0.01 and 4  $\mu\text{s}$  after flashing a deoxygenated toluene solution containing **5** ( $2 \times 10^{-5}$  M) and 1-(4-iodobutyl)-1*H*-indole (0.1 M). (c) Photoreaction of 1-(4-iodobutyl)-1*H*-indole in the presence of TEMPO. Product yield was calculated by  $^1\text{H}$  NMR using 1,3,5-trimethoxybenzene as an internal standard.


containing **5** and 1-(4-iodobutyl)-1*H*-indole (Fig. 11b). This shows that the photoreaction is initiated by oxidative quenching of  $^*5$  with indole. Photoreaction of 1-(4-iodobutyl)-1*H*-indole in the presence of TEMPO (2,2,6,6-tetramethyl-1-piperidyl)oxy as a scavenger was conducted; the TEMPO-coupling product was obtained in 90% yield (Fig. 11c). In view of the above results, a plausible reaction mechanism was proposed and shown in Fig. S25.† The mixed  $^3\text{MMLCT}/^3\text{LLCT}$  excited state of **5** can induce reductive dehalogenation of *N*-alkylated indole/pyrrole to form an alkyl radical, which subsequently undergoes cyclization reaction to give a benzylic radical. Oxidation of the benzylic radical followed by deprotonation affords the desired cyclized product. DIPEA was used as a sacrificial electron donor to regenerate **5**.

The feasibility of these binuclear  $\text{Pd}^{\text{II}}$  complexes as photosensitizers (PC) was also examined in the photochemical generation of singlet oxygen ( $^1\text{O}_2$ ). As shown in Fig. S26,† the emission of  $^1\text{O}_2$  was detected upon photoexcitation of binuclear  $\text{Pd}^{\text{II}}$  complexes (**3**, **5**, and **7**) in aerated  $\text{CHCl}_3$ . Referring to the emission intensity of  $^1\text{O}_2$  produced with  $\text{H}_2\text{tpp}$  as the photosensitizer ( $\Phi_{\text{so}} = 0.55$  in  $\text{CHCl}_3$ ) at  $\lambda_{\text{max}} = 1270$  nm,<sup>28</sup> it was found that the  $\Phi_{\text{so}}$  values of **3**, **5**, and **7** were 0.19, 0.15 and 0.08, respectively. Subsequently, we examined the photo-induced aerobic C–C coupling reaction of *N*-phenyl-1,2,3,4-tetrahydroisoquinoline with acetone using these complexes as photosensitizers. This reaction is known to be initiated by the oxidation of amines to iminium ion intermediates *via* photochemically generated  $^1\text{O}_2$ .<sup>29</sup> As shown in Scheme 2, complete substrate conversion (100%) and high product yields (81–88%) were achieved. Control experiments showed that in the absence of binuclear  $\text{Pd}^{\text{II}}$  complexes, *L*-proline,  $\text{O}_2$  or light irradiation, only trace amounts of coupling products were detected.

## Electroluminescence

The high emission quantum yields of **5** and **6** in PMMA thin films prompted us to investigate their potential applications as

OLED emitters. Devices with architecture consisting of ITO/HAT-CN (5 nm)/TAPC (40 nm)/CCP (10 nm)/**5**: PPF (10 nm)/PPF (10 nm)/TmPyPb (40 nm)/LiF (1.2 nm)/Al (100 nm) were fabricated to examine the electroluminescent (EL) properties of **5**. In these devices, di-[4-(*N,N*-ditolyl-amino)-phenyl]cyclohexane (TAPC) and 1,3,5-tri(*m*-pyrid-3-yl-phenyl)benzene (TmPyPb) were used as the hole-transporting layer (HTL) and electron-transporting layer (ETL), respectively, and 1,4,5,8,9,11-hexaazatriphenylene hexacarbonitrile (HAT-CN) was used as the hole-injecting layer to facilitate the hole injection from ITO to TAPC. Complex **5** was doped in 2,8 bis(diphenylphosphoryl) dibenzo[*b,d*]furan (PPF) at concentrations ranging from 4 to 16 wt% to form an emissive layer (EML). PPF and CCP (9-phenyl-3,9'-bicarbazole), with triplet energies of 3.0 and 3.1 eV, were used as an exciton-blocking layer to confine the excitons within the EML.<sup>30</sup> When the doping concentration is 4 wt%, the EL maximum of **5** is located at 521 nm (Fig. 12a). When the doping concentration is increased to 8 wt%, the EL spectrum of **5** is almost unchanged and is slightly red-shifted to 525 nm at 16 wt%. The device with 4 wt% of **5** achieved a high max. EQE of 22.9% (Fig. 12b), corresponding to a current efficiency (CE) of 71.2  $\text{cd A}^{-1}$  (Table 5). At high luminance of 1000  $\text{cd m}^{-2}$ , the EQE and CE of this device slightly dropped to 19.6% and 60.9  $\text{cd A}^{-1}$ , respectively. In devices with 8 wt% and 16 wt% doping concentrations of **5**, max. EQE values dropped slightly to 21.8% and 20.7%, respectively. Although the PLQY (0.92–0.93) of **5** in PPF is similar at different doping concentrations (Fig. S27†), the luminance and EQE at 1000  $\text{cd m}^{-2}$  of devices with higher concentrations of **5** are lower than those with 4 wt% of **5**. The inferior device performance of **5**-based devices at higher doping concentrations can be attributed to the strong trapping effect of **5**, as shown in Fig. S28.† The HOMO energy level (−5.2 eV) of **5** is 1.5 eV higher than the host PPF (−6.7 eV), causing holes injected from the HTL deeply trapped.



**Fig. 12** (a) Normalized EL spectra and (b) EQE-luminance characteristics of OLEDs with **5** at various doping concentrations. (c) Normalized EL spectra and (d) EQE-luminance characteristics of OLEDs with **6** at various doping concentrations.



Table 5 Key performances of OLEDs with **5** and **6** at various concentrations<sup>a</sup>

	$L_{\max}^b$ [cd m <sup>-2</sup> ]	Max. EQE <sup>c</sup> [%]	EQE <sup>c</sup> at 1000 cd m <sup>-2</sup> [%]	Max. CE <sup>d</sup> [cd A <sup>-1</sup> ]	Max. PE <sup>e</sup> [lm W <sup>-1</sup> ]	CIE coordinates <sup>f</sup> (x, y)
PPF: <b>5</b> (4 wt%)	18 200	22.9	19.6	71.2	85.2	0.29, 0.57
PPF: <b>5</b> (8 wt%)	12 750	21.8	18.4	70.2	70.0	0.30, 0.57
PPF: <b>5</b> (16 wt%)	11 200	20.7	18.2	64.1	39.5	0.31, 0.58
mCBP: <b>6</b> (4 wt%)	94 100	21.8	18.9	76.5	61.9	0.41, 0.56
mCBP: <b>6</b> (8 wt%)	99 100	21.2	17.9	73.8	62.3	0.42, 0.56
mCBP: <b>6</b> (12 wt%)	104 000	21.5	18.4	73.4	65.9	0.43, 0.55

<sup>a</sup> Fabricated by vacuum deposition. <sup>b</sup> Max. luminance. <sup>c</sup> External quantum efficiency. <sup>d</sup> Current efficiency. <sup>e</sup> Power efficiency. <sup>f</sup> CIE coordinates at 1000 cd m<sup>-2</sup>.

For **6**-based OLEDs, EML is formed by doping **6** in mCBP (3,3'-di(9*H*-carbazol-9-yl)-1,1'-biphenyl) at a concentration of 4 to 12 wt%. As shown in Fig. 12c, the EL spectrum of the **6**-based device is unstructured with a peak maximum at 554–560 nm. The max. luminance and max. power efficiency (PE) increase as the doping concentration increases from 4 wt% to 12 wt%, up to 104 000 cd m<sup>-2</sup> and 69.5 lm W<sup>-1</sup>, respectively. The max. EQE values (21.2–21.8%) are almost insensitive to the doping concentration (Fig. 12d). The OLED performance of complex **5** is slightly inferior to that of the Pt<sup>II</sup> analogue; the max. EQE of the latter is reported to be over 25%.<sup>31</sup> Nevertheless, the max. EQE and max. luminance of the **5**-based OLEDs are up to 22.9% and 104 000 cd m<sup>-2</sup>, respectively; these two values are the highest among OLEDs doped with binuclear Pd<sup>II</sup> emitters.<sup>6</sup> At high luminance of 1000 cd m<sup>-2</sup>, the EQE values drop slightly to 17.9–18.9%, corresponding to an efficiency roll-off of 13.3–15.6%.

## Conclusion

In this work, binuclear Pd<sup>II</sup> complexes with short intramolecular Pd–Pd distances of 2.79–2.89 Å and <sup>3</sup>MMLCT excited state with high  $\Phi_{\text{em}}$  (up to 0.7) and large  $k_r$  (up to  $2 \times 10^5$  s<sup>-1</sup>) have been achieved. These binuclear Pd<sup>II</sup> complexes show potential applications as light-emitting dopants for OLEDs and as photocatalysts or photosensitizers for C–C coupling reactions. According to DFT/TDDFT calculations, the energy difference between the <sup>3</sup>MMLCT and <sup>3</sup>dd excited states of complex **1** is smaller than that of **Ref-Pt1**. Together with the smaller SOC, complex **1** shows faster non-radiative decay and slower radiative decay processes. The strongly  $\sigma$ -donating (C<sup>^</sup>C\*) ligands of complexes **5** and **6** can significantly destabilize the <sup>3</sup>dd excited state. Complex **5** reacts with alkyl bromide through the inner-sphere electron transfer pathway under photo-irradiation, and the Pd-containing product is [(Pd<sup>II</sup>–Pd<sup>III</sup>–Br)(DPF<sup>-</sup>)<sub>2</sub>]<sup>2+</sup> (**5-Br**). Overall, this work provides a comprehensive study of the <sup>3</sup>MMLCT excited states of binuclear Pd<sup>II</sup> complexes from basic research to applications.

## Data availability

The data supporting this article have been included as part of the ESI.†

## Author contributions

Chi-Ming Che designed and initiated this research project. Mingyong Xue synthesized and characterized all complexes in this work, performed the spectroscopic and electrochemical measurements and carried out the photochemical studies. Wai-Pong To provided experimental guidance for this research project. Chi-Ming Che, Qingyun Wan and Mingyong Xue wrote and revised the manuscript. Yuzhen Zhang grew the single crystals of complex **1** and **Ref-Pt1**. Gang Cheng performed OLED fabrication and electroluminescence measurements. Zhou Tang carried out the EPR simulations. Lili Du carried out the femtosecond time-resolved absorption and emission spectroscopic measurements. Kam-Hung Low was responsible for X-ray crystal structure determinations. Qingyun Wan was responsible for all the DFT/TDDFT calculations in this work.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We acknowledge the financial support by the Major Program of Guangdong Basic and Applied Research (2019B030302009), the Science, Technology, and Innovation Commission of Shenzhen Municipality (JCYJ20200109150414471 and JCYJ20180508162429786), the Research Grants Council (17309823) of Hong Kong, and the Hong Kong Quantum AI Lab Limited, AIR @ InnoHK of Hong Kong Government. This work was conducted, in part, using the research computing facilities and advisory services offered by Information Technology Services, The University of Hong Kong.

## References

- (a) M. Yoshida and M. Kato, *Coord. Chem. Rev.*, 2018, **355**, 101–115; (b) S.-Y. Yang, Y. Chen, R. T. K. Kwok, J. W. Y. Lam and B. Z. Tang, *Chem. Soc. Rev.*, 2024, **53**, 5366.
- (a) M. Han, Y. Tian, Z. Yuan, L. Zhu and B. Ma, *Angew. Chem.*, 2014, **126**, 11088; *Angew. Chem., Int. Ed.*, 2014, **53**, 10908; (b) X.-P. Zhang, J.-F. Mei, J.-C. Lai, C.-H. Li and X.-Z. You, *J. Mater. Chem. C*, 2015, **3**, 2350; (c) J. Ni, Y. Zhang, S. Liu



- and J. Zhang, *J. Organomet. Chem.*, 2024, **1004**, 122948; (d) J. Ni, Q. Zhu, X. Wang, S. Liu and J. Zhang, *J. Organomet. Chem.*, 2024, **1015**, 123233.
- 3 (a) K. T. Ly, R.-W. Chen-Cheng, H.-W. Lin, Y.-J. Shiau, S.-H. Liu, P.-T. Chou, C.-S. Tsao, Y.-C. Huang and Y. Chi, *Nat. Photonics*, 2017, **11**, 63; (b) K.-H. Kim, J.-L. Liao, S. W. Lee, B. Sim, C.-K. Moon, G.-H. Lee, H. J. Kim, Y. Chi and J.-J. Kim, *Adv. Mater.*, 2016, **28**, 2526; (c) P. Pinter, J. Soellner and T. Strassner, *Eur. J. Inorg. Chem.*, 2021, 3104; (d) H. Leopold, M. Tenne, A. Tronnier, S. Metz, I. Münster, G. Wagenblast and T. Strassner, *Angew. Chem.*, 2016, **128**, 16011; *Angew. Chem., Int. Ed.*, 2016, **55**, 15779; (e) M. Xue, T.-L. Lam, G. Cheng, W. Liu, K.-H. Low, L. Du, S. Xu, F.-F. Hung, D. L. Phillips and C.-M. Che, *Adv. Opt. Mater.*, 2022, **10**, 2200741.
- 4 (a) V. Sicilia, L. Arnal, S. Fuertes, A. Martín and M. Baya, *Inorg. Chem.*, 2020, **59**, 12586; (b) V. Sicilia, P. Borja and A. Martín, *Inorganics*, 2014, **2**, 508; (c) V. Sicilia, M. Baya, P. Borja and A. Martín, *Inorg. Chem.*, 2015, **54**, 7316.
- 5 (a) J. J. Topczewski and M. S. Sanford, *Chem. Sci.*, 2015, **6**, 70; (b) D. C. Powers and T. Ritter, *Acc. Chem. Res.*, 2012, **45**, 840; (c) D. C. Powers, M. A. L. Geibel, J. E. M. N. Klein and T. Ritter, *J. Am. Chem. Soc.*, 2009, **131**, 17050; (d) D. C. Powers and T. Ritter, *Nat. Chem.*, 2009, **1**, 302.
- 6 (a) J. Lin, C. Zou, X. Zhang, Q. Gao, S. Suo, Q. Zhuo, X. Chang, M. Xie and W. Lu, *Dalton Trans.*, 2019, **48**, 10417; (b) L. Qiao, X. Kong, K. Li, L. Yuan, Y. Shen, Y. Zhang and L. Zhou, *Adv. Sci.*, 2024, 2404621.
- 7 P. K. Chow, C. Ma, W.-P. To, G. S. M. Tong, S.-L. Lai, S. C. F. Kui, W.-M. Kwok and C.-M. Che, *Angew. Chem., Int. Ed.*, 2013, **52**, 11775.
- 8 (a) Q. Wan, W.-P. To, C. Yang and C.-M. Che, *Angew. Chem.*, 2018, **130**, 3143; *Angew. Chem., Int. Ed.*, 2018, **57**, 3089; (b) C. Zou, J. Lin, S. Suo, M. Xie, X. Chang and W. Lu, *Chem. Commun.*, 2018, **54**, 5319; (c) L. Cao, K. Klimes, Y. Ji, T. Fleetham and J. Li, *Nat. Photonics*, 2019, **15**, 185; (d) L. Ameri, L. Cao, X. Tan and J. Li, *Adv. Mater.*, 2023, **35**, 2208361.
- 9 J. Pérez, A. Espinosa, J. M. Galiana, E. Pérez, J. L. Serrano, M. A. G. Aranda and M. Insausti, *Dalton Trans.*, 2009, 9625.
- 10 B. Ma, J. Li, P. I. Djurovich, M. Yousufuddin, R. Bau and M. E. Thompson, *J. Am. Chem. Soc.*, 2005, **127**, 28.
- 11 (a) P.-K. Chow, G. Cheng, G. S. M. Tong, C. Ma, W.-M. Kwok, W.-H. Ang, C. Y.-S. Chung, C. Yang, F. Wang and C.-M. Che, *Chem. Sci.*, 2016, **7**, 6083; (b) G. Li, J. Zheng, X. Fang, K. Xu, Y.-F. Yang, J. Wu, L. Cao, J. Li and Y. She, *Organometallics*, 2021, **40**, 472; (c) G. Li, H. Guo, X. Fang, Y.-F. Yang, Y. Sun, W. Lou, Q. Zhang and Y. She, *Chin. J. Chem.*, 2022, **40**, 223.
- 12 J. Liu, T.-L. Lam, M.-K. Sit, Q. Wan, C. Yang, G. Cheng and C.-M. Che, *J. Mater. Chem. C*, 2022, **10**, 10271.
- 13 P. S. Wagenknecht and P. C. Ford, *Coord. Chem. Rev.*, 2011, **225**, 591.
- 14 (a) M. Chaaban, Y.-C. Chi, M. Worku, C. Zhou, H. Lin, S. Lee, A. Ben-Akacha, X. Lin, C. Huang and B. Ma, *Inorg. Chem.*, 2020, **59**, 13109; (b) S. F. Wang, L. W. Fu, Y.-C. Wei, S.-H. Liu, J.-A. Lin, G.-H. Lee, P.-T. Chou, J.-Z. Huang, C.-I. Wu, Y. Yuan, C.-S. Lee and Y. Chi, *Inorg. Chem.*, 2019, **58**, 13892.
- 15 H. Amouri, *Chem. Rev.*, 2023, **123**, 230.
- 16 C.-L. Yao, L.-P. He, J. D. Korp and J. L. Bear, *Inorg. Chem.*, 1988, **27**, 4389.
- 17 J. F. Berry, E. Bill, E. Bothe, F. A. Cotton, N. S. Dalal, S. A. Ibragimov, N. Kaur, C. Y. Liu, C. A. Murillo, S. Nellutla, J. M. North and D. Villagrán, *J. Am. Chem. Soc.*, 2007, **129**, 1393.
- 18 C.-H. Lam, W. K. Tang and V. W.-W. Yam, *Inorg. Chem.*, 2023, **62**, 1942.
- 19 N. Bayliss, A. Cole and B. Green, *Aust. J. Chem.*, 1948, **1**, 472.
- 20 M. Hermet, L. Bakás, S. R. Morcelle and D. L. Bernik, *Spectrochim. Acta, Part A*, 2019, **223**, 117266.
- 21 S. Yamashita, K. Iwamatsu, Y. Maehashi, M. Taguchi, K. Hata, Y. Muroya and Y. Katsumura, *RSC Adv.*, 2015, **5**, 25877.
- 22 M. Dey, S. S. Dhar and M. Kalita, *Synth. Commun.*, 2013, **43**, 1734.
- 23 (a) J. P. Mittal and E. Hayon, *Nat. Phys. Sci.*, 1972, **240**, 20; (b) N. A. McAskill and D. F. Sangster, *Aust. J. Chem.*, 1977, **30**, 2107.
- 24 S. H. Eitel, M. Bauer, D. Schweinfurth, N. Deibel, B. Sarkar, H. Kelm, H.-J. Kruger, W. Frey and R. Peters, *J. Am. Chem. Soc.*, 2012, **134**, 4683.
- 25 J. Harmer, C. Finazzo, R. Piskorski, C. Bauer, B. Jaun, E. C. Duin, M. Goenrich, R. K. Thauer, S. V. Doorslaer and A. Schweiger, *J. Am. Chem. Soc.*, 2005, **127**, 17744.
- 26 (a) D. C. Smith and H. B. Gray, *Coord. Chem. Rev.*, 1990, **100**, 169; (b) S. J. Atherton and D. M. Roundhill, *Inorg. Chem.*, 1986, **25**, 4072.
- 27 S. J. Kaldas, A. Cannillo, T. McCallum and L. Barriault, *Org. Lett.*, 2015, **17**, 2864.
- 28 F. Wilkinson, W. P. Helman and A. B. Ross, *J. Phys. Chem. Ref. Data*, 1993, **22**, 113.
- 29 W.-P. To, Y. Liu, T.-C. Lau and C.-M. Che, *Chem.-Eur. J.*, 2013, **19**, 5654.
- 30 (a) M. Numata, T. Yasuda and C. Adachi, *Chem. Commun.*, 2015, **51**, 9443; (b) I. S. Park, K. Matsuo, N. Aizawa and T. Yasuda, *Adv. Funct. Mater.*, 2018, **28**, 1802031.
- 31 (a) T.-L. Lam, H. Li, K. Tan, Z. Chen, Y.-K. Tang, J. Yang, G. Cheng, L. Dai and C.-M. Che, *Small*, 2024, **20**, 2307393; (b) L. Wang, J. Miao, Y. Zhang, C. Wu, H. Huang, X. Wang and C. Yang, *Adv. Mater.*, 2023, **35**, 2303066.

