Nanoscale



PAPER

View Article Online
View Journal | View Issue



Cite this: Nanoscale, 2024, 16, 4095

Oxygen self-supplying small size magnetic nanoenzymes for synergistic photodynamic and catalytic therapy of breast cancer†

In recent years, tumor catalytic therapy based on nanozymes has attracted widespread attention. However, its application is limited by the tumor hypoxic microenvironment (TME). In this study, we developed oxygen-supplying magnetic bead nanozymes that integrate hemoglobin and encapsulate the photosensitizer curcumin, demonstrating reactive oxygen species (ROS)-induced synergistic breast cancer therapy. Fe₃O₄ magnetic bead-mediated catalytic dynamic therapy (CDT) generates hydroxyl radicals ('OH) through the Fenton reaction in the tumor microenvironment. The Hb-encapsulated Fe₃O₄ magnetic beads can be co-loaded with the photosensitizer/chemotherapeutic agent curcumin (cur), resulting in Fe₃O₄-Hb@cur. Under hypoxic conditions, oxygen molecules are released from Fe₃O₄-Hb@cur to overcome the TME hypoxia, resulting in comprehensive effects favoring anti-tumor responses. Upon near-infrared (NIR) irradiation, Fe₃O₄-Hb@cur activates the surrounding molecular oxygen to generate a certain amount of singlet oxygen ⁽¹O₂), which is utilized for photodynamic therapy (PDT) in cancer treatment. Meanwhile, we validated that the O₂ carried by Hb significantly enhances the intracellular ROS level, intensifying the catalytic therapy mediated by Fe₃O₄ magnetic beads and inflicting lethal damage to cancer cells, effectively inhibiting tumor growth. Therefore, significant *in vivo* synergistic therapeutic effects can be achieved through catalytic-photodynamic combination therapy.

Received 20th October 2023, Accepted 15th January 2024 DOI: 10.1039/d3nr05289c

rsc.li/nanoscale

1. Introduction

Nanocatalytic therapy is widely used in cancer treatment due to its enormous advantages such as high selectivity, low toxicity, high efficiency, and tunability. Efficient tumor catalysts, such as hydrogen peroxide enzymes, metal nanoparticles, and porphyrin compounds, can induce the generation of reactive oxygen species (ROS), including singlet oxygen (lO₂), hydroxyl radicals (OH), and superoxide anions

('O^{2–}).^{8,9} By harnessing the power of these natural ROS, it is possible to effectively induce oxidative damage to cancer cells and shrink tumors, which has been adopted and further explored for *in vivo* tumor therapy.¹⁰ Despite these significant advancements, there are still challenging factors that limit the potential efficacy of catalytic therapy.^{11,12} One of the major challenges is the inherently hypoxic tumor microenvironment (TME) within solid tumors, which significantly restricts the effectiveness of nanocatalytic therapy in clinical applications.^{13,14}

Chemodynamic therapy (CDT), which enhances intracellular ROS through the utilization of the Fenton reaction, has garnered significant attention in cancer treatment. 15,16 Considering the overexpression of hydrogen peroxide (H_2O_2) and mild acidic conditions in the TME, extensive exploration has been conducted on advanced nanocatalysts to capture H_2O_2 as a substrate. $^{17-19}$ However, achieving sufficient and satisfactory therapeutic efficacy remains challenging due to the limited endogenous oxygen (O_2) content. 20 Hemoglobin (Hb) is a composite protein composed of four subunits. Each Hb molecule consists of two α subunits and two β subunits, with each subunit containing a heme molecule capable of binding oxygen. The structure and function of hemoglobin enable it to

^aDongguan Key Laboratory of Screening and Research of Anti-inflammatory
Ingredients in Chinese Medicine, School of Pharmacy, Guangdong Medical
University, Dongguan 523808, China. E-mail: wsyt132@163.com

^bGuangdong Key Laboratory of Nanomedicine, Shenzhen Institute of Advanced
Technology, Chinese Academy of Sciences, Shenzhen 518055, China

^cDepartment of Gastroenterology, the Tenth Affiliated Hospital of Southern Medical
University (Dongguan People's Hospital), Dongguan 523000, China

^dInstitute of Nano Biomedicine and Engineering, School of Sensing Science and

^aInstitute of Nano Biomedicine and Engineering, School of Sensing Science and Engineering, School of Electronic Information and Electrical Engineering, Shanghai Jiao Tong University, 800 Dongchuan RD, Shanghai 200240, China

^eCancer Centre, Faculty of Health Sciences, University of Macau, Macau SAR, China †Electronic supplementary information (ESI) available. See DOI: https://doi.org/ 10.1039/d3nr05289c

[‡]These authors contributed equally to this work.

Paper

anti-tumor effects. 25,26

encapsulate and transport oxygen. ²¹ Using Hb to elevate the oxygen concentration within tumors, this combination therapy approach enhances the effectiveness of anti-tumor treatments. ^{22,23} However, few researchers have explored the simultaneous elevation of O_2 levels and increased ROS within cancer cells to improve therapeutic outcomes. Therefore, there is an urgent need to develop novel combination therapies to enhance the efficacy of catalytic treatments. Curcumin (Cur), a diketone compound extracted from the rhizomes of turmeric, exhibits diverse pharmacological activities such as anti-inflammatory and anti-tumor effects. ²⁴ Cur, being a natural photo-

sensitizer, can mediate the generation of singlet oxygen in

photodynamic therapy (PDT) to kill tumor cells. Additionally,

as a chemotherapeutic agent, it synergizes with PDT to exert

In this study, we constructed a composite magnetic nanosystem, utilizing a Hb-coloaded Fe_3O_4 based nanoenzyme and Cur (referred to as Fe_3O_4 -Hb@cur), for enhancing ROS and designing catalytic reactions in cancer nanotherapy (Fig. 1). This strategy possesses the following characteristics: (1) Nanozymes exhibit inherent peroxidase-like activity, decomposing H_2O_2 into OH to induce tumor cell damage. (2) Oxygen




Fig. 1 A scheme of the preparation process (A) of the oxygen-supplying magnetic nanoparticle enzyme (Fe $_3$ O $_4$ -Hb@cur) and its mechanism for achieving photodynamic and catalytic therapy of tumor (B). In a hypoxic environment, oxygen molecules are released from Fe $_3$ O $_4$ -Hb@cur to overcome the oxygen-deficient TME. Hb carrying O $_2$ is co-loaded with the photosensitizer/chemotherapeutic agent cur, and upon NIR irradiation, it activates the surrounding molecular oxygen to generate a certain amount of 1 O $_2$ for PDT. Simultaneously, O $_2$ -enhanced Fe $_3$ O $_4$ mediates CDT by producing 'OH through the Fenton reaction. Therefore, Fe $_3$ O $_4$ -Hb@cur achieves significant *in vivo* synergistic therapeutic effects to accomplish the mechanism of catalytic-photodynamic therapy for breast cancer.

self-supply can be achieved through the encapsulation of Hb, and the increased $\rm O_2$ can be activated by Cur to generate $^{\rm 1}{\rm O_2}$ for tumor cell death, thereby enhancing the catalytic performance of the nanozymes. Based on this therapeutic approach, we report the preparation of a small-sized magnetic nanozyme with intrinsic oxygen supply (Fe₃O₄-Hb@cur), which demonstrates synergistic effects on tumor damage through combined photodynamic and catalytic therapy. In summary, our work highlights the tremendous potential of modulating the TME using intelligent nanosystems to enhance the effectiveness of various treatment modalities, potentially inhibiting tumor growth and achieving comprehensive effects on cancer. This has significant implications in the field of nanocatalytic medicine.

2. Experimental section

2.1. Materials

Curcuminoids (Cur) and hemoglobin (Hb) were purchased from Macklin Biochemical Co., Ltd (Shanghai, China). 3,3',5,5'-Tetramethylbenzidine (TMB), 2',7'-dichlorofluorescein diacetate (DCFH-DA), 4',6-diamidino-2-phenylindole (DAPI) and 1,3-diphenylisobenzofuran (DPBF) were purchased from Aladdin Reagent Co., Ltd (Shanghai, China). Cy5.5 was purchased from Xi'an ruixi Biological Technology Co., Ltd. The cell counting kit was purchased from ZETA life. All reagents were of analytical grade and used without any purification.

2.2. Synthesis of Fe₃O₄

Monodisperse Fe_3O_4 nanoparticles with average diameters of around 15 nm were synthesized using a previously published protocol. Afterwards, the product was precipitated with ethanol and redispersed in chloroform. The obtained Fe_3O_4 nanoparticles were then transferred into water by a polymer coating process, which was carried out by a published method. Finally, Fe_3O_4 nanoparticles were stored in water at 4 °C for further use.

2.3. Synthesis of Fe₃O₄-Hb@cur

Under magnetic stirring, 2 mL of Fe_3O_4 (2 μM) was mixed with a 4 mL solution of Hb (dissolved in PBS at a concentration of 1 mg mL⁻¹) overnight. The mixture was then slowly added dropwise to 2 mL of cur (300 μM) and further blitzed overnight on a rotary mixer. The final product was concentrated using an ultrafiltration tube to 4 mL and stored at 4 °C.

2.4. Characterization of Fe₃O₄-Hb@cur

The morphological features of Fe₃O₄ and Fe₃O₄-Hb were analyzed using Transmission Electron Microscopy (TEM, JEOL JEM 2100F from Japan Electron). UV-vis absorption spectra of Fe₃O₄, cur, Hb, and Fe₃O₄-Hb@cur were recorded using an ultraviolet-visible spectrophotometer (Lambda25, PerkinElmer, USA). A Dynamic Light Scattering (DLS) analyzer (Malvern Instruments, UK) was used to measure the particle size of Fe₃O₄-Hb@cur. Electron Paramagnetic Resonance (EPR,

Nanoscale Paper

BRUKER EMXPLU) was used to measure the signals of reactive oxygen species. The EDS (Energy Dispersive Spectrometer) used for detecting the elements, composition information, and concentration of the samples was Oxford Xplore 30. *In vitro* magnetic resonance (MR) images using T_2 -weighted sequences were acquired using a 3.0 T clinical MR scanner (Erlangen, GER). The signal intensity in the region of interest (ROI) was measured for all T_2 -weighted MR images. To calculate the relaxation rates $(r_2, 1/T_2)$, a range of concentrations of the Fe₃O₄-Hb@cur complex were measured. FL images were analyzed with the IVIS imaging system (CALIPER IVIS SPECTRUM). The collection parameters were Em = 700 nm and Ex = 640 nm. Fluorescence spectral scanning was performed using an FL970 fluorescence spectrophotometer (Tianmei, China).

2.5. Hemolysis assay

0.5 mL of blood sample was added to 1 mL of PBS solution and centrifuged at 4500 rpm for 5 min, and the red blood cells were separated from the serum. Then, the red blood cells were washed 5 times with PBS and resuspended in 5 mL of PBS. Next, 0.3 mL of the above-mentioned red blood cells was mixed with 1.2 mL of Fe₃O₄-Hb@cur at different concentrations (0–100 μ M, prepared in PBS). The solution was gently mixed and allowed to stand at 25 °C for 4 h, followed by centrifugation to separate the red blood cells. The dissolved red blood cells could not be separated. The absorbance of the supernatant was then measured at a wavelength of 541 nm (mean \pm SD, n = 3). Red blood cells treated with PBS and deionized water were used as negative and positive control groups, respectively. The percentage of hemolysis was calculated using the equation:

Hemolysis(%) = ((OD sample - OD negative control)/(OD positive control – OD negative control)) × 100%.

2.6. H₂O₂-dependent catalysis of Fe₃O₄-Hb@cur

Briefly, 10 μ L TMB (1 mM) and 150 μ L acetic acid buffer solution (pH 5.5) were added to acetic acid buffer solution (H₂O₂ + Fe₃O₄@cur, H₂O₂ + Fe₃O₄-Hb@cur, and 150 μ L H₂O₂ (1 mM)), respectively. Degradation of TMB was monitored with UV-Visible absorbance at 652 nm 1 h after the reaction. The degradation of TMB was monitored by observing the change in UV-Visible absorbance to evaluate the peroxidase activity of Fe₃O₄. Specifically, different concentrations of Fe₃O₄-Hb@cur (3, 6, 9, 12, and 15 μ M) were added to a mixture containing acetic acid buffer solution (pH 5.5) and H₂O₂ (1 mM), following the addition of 10 μ L TMB (1 mM). The absorbance at 652 nm was immediately detected and the data were recorded continuously for 8 min.

2.7. Light-triggered ROS generation

Diphenylisobenzofuran (DPBF) was used in this experiment to trap the 1O_2 produced by Fe $_3O_4$ @cur under light conditions. We prepared Fe $_3O_4$ @cur (Fe $_3O_4$ (2 μ M, 15 μ L) + cur (300 μ M, 150 μ L) + 300 μ L PBS) and Fe $_3O_4$ -Hb@cur (Fe $_3O_4$ (2 μ M, 15 μ L)

+ cur (300 μ M, 150 μ L) + Hb (1 mg mL⁻¹, 50 μ L) + 250 μ L PBS). 100 μ L of the above samples were taken and mixed with 10 μ L DPBF (20 mM) at 25 °C. Fe₃O₄ @cur and Fe₃O₄ -Hb@cur (300 μ M as cur) and DPBF (20 mM) were irradiated with a 660 nm laser (0.5 W cm⁻²) for 1, 2, 3, 4, or 5 min, respectively. Subsequently, the absorbance was recorded at a wavelength of 405 nm using a microplate reader.

2.8. Cell culture

4T1 mouse mammary adenocarcinoma cells were procured from Beina Biotech Co., Ltd and maintained in culture medium (DMEM) supplemented with 10% (v/v) fetal bovine serum and a 1% penicillin–streptomycin mixture. The cells were cultured under standard conditions (5% CO_2 , 37 °C).

2.9. Cell uptake of Fe₃O₄-Hb@cur

4T1 cells were inoculated in confocal culture dishes at a density of 1×10^5 cells per dish and cultured in 5% CO $_2$ at 37 °C or 24 h. In 300 μM Fe $_3O_4$ -Hb@cur after replacing medium, the cells were subjected to further different incubation times (1 h, 5 h, 9 h). Subsequently, the culture medium was aspirated, and the cells were washed three times with PBS. Paraformaldehyde was then added for an 8-minute incubation period to fix the cells. After removing the fixative, DAPI was added and incubated for 20 min. The cell nuclei were stained with DAPI in the co-localization experiments. The cells were observed using a confocal laser scanning microscope (CLSM) with an excitation wavelength of 471 nm for cur and 405 nm for DAPI.

2.10. Cell viability

The viability of 4T1 cells treated with different concentrations of PBS (as control), Fe $_3$ O $_4$, Fe $_3$ O $_4$ @cur, Fe $_3$ O $_4$ -Hb@cur, Fe $_3$ O $_4$ @cur + laser, and Fe $_3$ O $_4$ -Hb@cur + laser was determined. The 4T1 cells were cultured in 96-well plates at an initial density of 10^4 cells per well and with overnight incubation at 37 °C. After that, gradient concentrations of Fe $_3$ O $_4$, Fe $_3$ O $_4$ @cur and Fe $_3$ O $_4$ -Hb@cur were introduced and co-incubated with the cells for 24 h. A 100 μ L medium solution containing 10 μ L of CCK-8 was added to each well and incubated with the cells for 0.5 h. The optical density (OD) at 450 nm was measured using a microplate reader. Cell viability was calculated using the following formula:

 $\begin{aligned} & \text{Cell viability}(\%) = (\text{OD}(\text{sample}) - \text{OD}(\text{blank group})) / \\ & (\text{OD}(\text{control}) - \text{OD}(\text{blank group})) \times 100. \end{aligned}$

2.11. Intracellular 'OH detection

The intracellular 'OH was measured *via* CLSM, and DCF served as a 'OH indicator. The 4T1 cells were plated in 6-well plates and incubated at 37 °C overnight. Then, the cells were treated with different formulations (PBS (as a control), Fe₃O₄@cur, Fe₃O₄-Hb@cur, Fe₃O₄@cur + laser (Fe₃O₄@cur (+)) and Fe₃O₄-Hb@cur + laser (Fe₃O₄-Hb@cur (+)) (300 μ M as cur)) for 12 h. Next, after washing the cells with PBS several times, the DCF solution (10 μ M) was added and incubated

Paper Nanoscale

with the cells for 30 min. Finally, the cells were observed by CLSM.

2.12. Animal model

Female BALB/c mice aged 6 weeks and weighing 18-22 g were procured from Zhuhai Baishitong Biotechnology Co., Ltd. Subsequently, 4T1 cells which were suspended in 100 µL of phosphate-buffered saline (PBS) at a concentration of 3×10^6 were inoculated into the subcutaneous site of the right hind limb of each BALB/c mouse. The tumor volume was estimated every other day according to the following formula: (length × width²)/2. All animal procedures were performed in accordance with the Guidelines for Care and Use of Laboratory Animals of Guangdong Medical University and approved by the Animal Ethics Committee of Guangdong Medical University (GDY2204043).

2.13. Monitoring the distribution of drugs in tumor-bearing mice

The distribution of nanoparticles was studied using the 4T1 mouse breast cancer model. Upon reaching a tumor volume of 150 mm³, mice bearing tumors were randomly assigned into groups (n = 3) and received intravenous injections of either free Cy5.5 (2 mg kg⁻¹) or Cy5.5@ Fe₃O₄-Hb@cur (2 mg kg⁻¹ of free Cy5.5). At specified time points after injection (1, 2, 4, 8, 12, 24, and 48 h), fluorescence biodistribution was analyzed using the IVIS in vivo imaging system. The collection parameters were Em = 700 nm and Ex = 640 nm.

2.14. MRI evaluation

In order to evaluate the tumor accumulation of nanoparticles, 4T1 tumor-bearing mice were subjected to MRI imaging at different time intervals following intravenous administration of either PBS or Fe₃O₄-Hb@cur. The mice were intravenous injection with 100 µL of PBS or Fe₃O₄-Hb@cur (containing 2 μM Fe₃O₄). MRI intensity imaging and semi-quantitative analysis were conducted using a 3.0 T clinical MR scanner (Erlangen, GER) at two time points: immediately after injection and 3 hours post-injection.

2.15. In vivo synergistic photodynamic and catalytic antitumor therapy

A 4T1 xenograft tumor model was established as mentioned above for evaluation of antitumor efficacy. The mice were then randomly separated into five groups (n = 6 mice per group): saline (as a control group), Fe₃O₄, Fe₃O₄@cur, Fe₃O₄-Hb@cur, Fe₃O₄@cur + laser (Fe₃O₄@cur (+)) and Fe₃O₄-Hb@cur + laser (Fe₃O₄-Hb@cur (+)). Therapeutic measures were commenced when the tumors reached a volume of approximately 150 cubic millimeters. On day 0, mice received combined therapy via tail vein injection. The tumor size and body weight were documented for each mouse from day 0 to day 16, and the tumor volume V was computed as $V = W^2 \times L/2$ (W and L are the shortest and longest diameters of the tumor, respectively). After 16 days of treatment, all mice were euthanized, and the major organs and tumor tissues were collected for further analysis.

2.16. Statistical analysis

All statistical analyses were analyzed using GraphPad Prism version 9 software. Experimental data were collected three times or more. The results were expressed as means \pm standard deviation. Statistical analysis of the differences between two groups was conducted using a two-tailed Student's t-test. Difference between the two groups was considered statistically significant when the two-sided P-value was less than 0.05, where all significant values were indicated as follows: *P < 0.05, **P < 0.01, and ***P < 0.001.

Results and discussion 3.

3.1. Construction and characterization of the Fe₃O₄-Hb@cur nanoplatform

In our study, homogeneous Fe₃O₄ nanoparticles with the average size of 15 nm were prepared by the thermal decomposition method, and after phase transfer into aqueous solution by polymer coating, a large number of carboxyl groups on their surface can then physically adsorb proteins (such as Hb) through electrostatic interactions. Small-sized magnetic nanozymes (Fe₃O₄-Hb) were further created by loading the Hb shells on the surface of Fe₃O₄ nanoparticles. The cavities of Hb can easily be loaded with many small molecule drugs (such as cur). Then, a photosensitizer/chemotherapeutic drug, cur, is loaded into the Fe₃O₄-Hb structure to obtain Fe₃O₄-Hb@cur. Therefore, the interaction during the formation of Fe₃O₄-Hb@cur is through physical absorption. Finally, oxygen is introduced into the solution to obtain Fe_3O_4 -Hb@cur (O_2) . Representative transmission electron microscopy (TEM) images show that the prepared nanozyme in this design have a single spherical morphology with an average diameter of about 15 nm (Fig. 2A). Notably, the successful encapsulation of Hb can be evidenced by the appearance of white halos on the surface of Fe₃O₄-Hb after Hb encapsulation of Fe₃O₄ (Fig. 2B). The dispersion of Fe₃O₄-Hb@cur was evaluated by dynamic light scattering (DLS). The DLS measurements indicate that Fe₃O₄-Hb@cur is homogeneous and well dispersed, with an average particle size of approximately 30 nm (Fig. S1†). Small-sized nanoparticles facilitate drug uptake and penetration within the tumor owing to elevated permeability and retention (EPR) effects. 29,30 As shown in the UV-vis spectra, due to the interaction between the carrier during cur loading, a red shift occurs in the peak position. As shown in Fig. 2C, the UV absorption peak of Fe3O4-Hb@cur matches the peak shape of cur. This absorption red shift demonstrates the successful encapsulation of cur in Fe₃O₄-Hb@cur. Moreover, the spectra changed significantly after Hb encapsulation with a strong absorption at 410 nm, indicating the formation of the Hb layer. The Fe₃O₄-Hb@cur sample shows both Hb and Fe₃O₄ absorption peaks after cur and Hb-bound Fe₃O₄. The successful loading of Cur was indicated by the change in the zeta potential, which was -15 mV for Fe₃O₄, -0.15 mV for cur, and -5.5 mV for the synthesized Fe₃O₄-Hb (Fig. S2†). Subsequently, we tested the zeta potential of the final product,

Nanoscale Paper




Fig. 2 Structural and compositional characterization of the Fe₃O₄-Hb@cur magnetic nanozyme. TEM images of (A) Fe₃O₄, scale bar: 25 nm, and (B) Fe₃O₄-Hb, scale bar: 50 nm. (C) UV-vis spectra of Fe₃O₄, cur, Hb, and Fe₃O₄-Hb@cur NPs. (D) Dark-field TEM image and the corresponding elemental mapping of C, O, Fe, N, and S of Fe₃O₄-Hb@cur. Scale bar: 30 nm. (E) Longitudinal relaxation rates $(1/T_2)$ of Fe₃O₄-Hb@cur solution treated with pH 7.4 as a function of Fe concentration. The following images show T_2 -weighted MR images of Fe₃O₄-Hb@cur after treatment at a pH of 7.4. (F) FL intensity of Fe₃O₄-Hb@cur solution at 673 nm as a function of Fe concentration. The inset shows the corresponding FL images of Fe₃O₄-Hb@cur solution.

Fe₃O₄-Hb@cur, and the result was -1.5 mV. The loading of cur successfully neutralized the electronegativity of the final product. Moreover, we compared the fluorescence spectra of cur, Fe₃O₄-Hb@cur, and Fe₃O₄-Hb in the range of 450-700 nm under the excitation wavelength of cur (420 nm) (Fig. S3†). The results revealed that cur and Fe₃O₄-Hb@cur exhibited identical fluorescence peak positions, providing evidence for the successful loading of cur onto Fe₃O₄-Hb@cur. However, Fe₃O₄-Hb did not exhibit the same fluorescence peak. In addition, elemental mapping was performed to reveal the distribution of the elements C, N, O, S, and Fe in Fe₃O₄-Hb@cur (Fig. 2D). Fe₃O₄-Hb@cur nanoenzymes contain elements C, O, Fe, N and S. The successful synthesis of Fe₃O₄-Hb@cur was further demonstrated by the combination of UV-vis and zeta potential. The atomic percentages of the elements in the obtained Fe₃O₄-Hb@cur solution were as follows: C 86.41%, N 0.08%, O 13.24%, S 0.01%, and Fe 0.25%. The zeta potential, changes in DLS size, TEM imaging, and absorption peaks indicate the successful preparation of Fe₃O₄-Hb@cur. Finally, we performed hemolysis experiments to demonstrate that Fe₃O₄-Hb@cur has good biocompatibility and safety (Fig. S4†).

Magnetic resonance imaging (MRI) and fluorescence intensity (FL) experiments were performed in order to reveal the bioimaging capabilities of Fe₃O₄-Hb@cur. As shown in Fig. 2E, MRI images of Fe₃O₄-Hb@cur at different concentrations were acquired using a clinical magnetic resonance scanner at room temperature. The transverse relaxivity (r_2) measurement for iron concentration was found to be 17.04 mM⁻¹ s⁻¹, indicating a linear relationship. Fe₃O₄-Hb@cur exhibited a decrease in

signal intensity (shortening of T_2 relaxation time) in T_2 weighted imaging with increasing concentration. 31,32 The fluorescence properties of the nanoparticles were subsequently investigated by attaching Cy5.5 to Fe₃O₄-Hb@cur. We compared the fluorescence spectra of Cy5.5, Fe₃O₄-Hb@cur-Cy5.5, and Fe₃O₄-Hb@cur in the range of 500-1000 nm at an excitation wavelength of Cy5.5 (673 nm). The results revealed that Cy5.5 and Fe₃O₄-Hb@cur-Cy5.5 exhibited identical fluorescence peak positions, providing evidence for the successful loading of Cy5.5 onto Fe₃O₄-Hb@cur (Fig. S5†). Cy5.5 has a specific fluorescence emission upon excitation at 673 nm, which can be further used for in vivo imaging. The FL signal intensity in aqueous solution under 673 nm laser excitation was linearly correlated with the concentration of Fe (Fig. 2F). Overall, Fe₃O₄-Hb@cur combined with Cy5.5 showed excellent MRI and FL imaging properties, which could be used as an ideal MRI contrast and fluorescence indicator for therapeutic diagnostic applications.

3.2. H₂O₂-dependent catalysis of Fe₃O₄-Hb@cur

In the presence of acetic acid buffer solution (pH 5.5), Fe₃O₄ can exhibit peroxidase-like activity that catalyzes the degradation of H2O2 to generate highly toxic 'OH for chemodynamic cancer therapy. 33-35 3,3',5,5'-Tetramethylbenzidine (TMB), oxidized by 'OH, changes the color of the mixture from colorless to blue-green with a typical absorbance peak at 652 nm. Therefore, it can be used to detect the generated 'OH. As shown in Fig. 3A, H₂O₂ in the absence of Fe₃O₄ based nanozymes cannot induce the Fenton reaction without obvious absorption at 652 nm. When H2O2 is mixed with Fe3O4-Hb@cur or Fe₃O₄@cur, an evident absorption at 652 nm from TMB was detected and the color changes of the solution was recorded, indicating the peroxidase-like activity of Fe₃O₄. The




Fig. 3 Characterization of the enzyme-like activity of the Fe₃O₄-Hb@cur nanozyme. (A) UV-vis absorption spectra of TMB in different groups with pH 5.5. The inset shows the corresponding visual color changes. Generation of ¹O₂ signals (B) and [•]OH signals (E) recorded by ESR spectra at different conditioned stimuli, respectively. (C) H₂O₂-concentration dependence of UV-vis absorbance changes employing a TMB probe at 652 nm. (D) Changes in DPBF absorbance at 405 nm were observed after treating it with Fe₃O₄@cur and Fe₃O₄-Hb@cur NPs under a 660 nm laser for different irradiation times. (F) Schematic representation of possible mechanisms of oxygen enhancement and tumorexpressed ROS under Fe₃O₄-Hb@cur conditions.

Paper Nanoscale

generated 'OH can be used for chemokinetic therapy and combined with chemotherapy to achieve multi-mode cancer therapy. When Hb was loaded, the peak absorbance increased, indicating that O2 carried on Hb enhanced the peroxidase activity of Fe₃O₄ and promoted the generation of 'OH. According to reports, the catalytic activity is associated with the generation capacity of 'OH.36 Therefore, the detection of 'OH production using electron spin resonance (ESR) is consistent. Compared to Fe₃O₄@cur and Fe₃O₄-Hb@cur groups, the 'OH signal was detected in both Fe₃O₄@cur and Fe₃O₄-Hb@cur groups containing H2O2. Furthermore, a more pronounced signal was detected in the Fe₃O₄-Hb@cur group containing H_2O_2 (Fig. 3B).

To systematically evaluate the catalytic performance of Fe₃O₄-Hb@cur, We investigated the ability of the nanoenzyme to generate 'OH under different pH conditions and found that our studied material only produced a significant amount of 'OH at pH 5.5. In contrast, very little 'OH was generated at pH 6.5 or pH 7.4. This indicates that a certain amount of H⁺ is required for Fe₃O₄-Hb@cur to exhibit positive feedback, which is beneficial for the effective functioning of the nanoenzyme after entering tumor cells (Fig. S6†). Moreover, different concentrations (3, 6, 9, 12, and 15 µM) of H₂O₂ were added to Fe₃O₄-Hb@cur in the presence of TMB (1 mM). The absorbance was then monitored as a function of time and H₂O₂ concentration to observe the dependence (Fig. 3C). Therefore, Fe₃O₄-Hb@cur based enzyme can be used as an efficientmimicking enzyme for cancer therapy.

3.3. O₂-enhanced PDT effect in vitro

Oxygen is an essential element in the photodynamic therapy process, yet the hypoxic TME greatly reduces the efficacy of PDT for solid tumors. Therefore, an adequate supply of oxygen at the tumor site is very important for generating ${}^{1}O_{2}$ to treat tumors.^{37,38} In this study, the absorption of DPBF at 405 nm was irreversibly quenched by 1O2, and thus was used as a probe for monitoring the production of ¹O₂ by Fe₃O₄ @cur and Fe₃O₄-Hb@cur under 660 nm light irradiation.^{39,40} At various time points, the absorption values of the samples at 405 nm were continuously recorded using an UV-Vis spectrophotometer. As shown in Fig. 3D, at the fifth minute, there was a significant decrease in absorbance values for both groups. However, the group containing encapsulated O₂ in Fe₃O₄-Hb@cur exhibited a stronger signal. In particular, the Fe₃O₄-Hb@cur group showed a lower absorbance value, around 50%, indicating an increased generation of ¹O₂. This observation suggests that the presence of Fe₃O₄-Hb@cur (O₂) enhances the production of ¹O₂. In summary, these results indicate that the release of O2 is crucial for the antitumor effect of PDT. The loading of O2 by Hb enhances the PDT efficacy of Fe₃O₄@cur, which has the potential to become a more efficient photocatalyst.

To further demonstrate the applicability of Fe₃O₄-Hb@cur in PDT, the ESR technique was employed to investigate the generation of ¹O₂ under near-infrared laser irradiation. As shown in Fig. 3E, in the absence of light and the photosensiti-

zer cur, no signal of 1O2 was detected. Compared to the other groups, the Fe₃O₄-Hb@cur group carrying O₂ exhibited a stronger signal of ¹O₂, indicating that O₂ effectively enhanced the PDT effect. This finding highlights the crucial role of O₂ in improving PDT within a hypoxic TME. These results emphasize the potential of Fe₃O₄-Hb@cur(O₂) in enhancing the efficacy of PDT, particularly under hypoxic conditions, and provide promising prospects for cancer treatment. The aforementioned principle and mechanism are presented using a schematic diagram labeled in Fig. 3F.

3.4. Cellular uptake of Fe₃O₄-Hb@cur

The cellular uptake behavior of Fe₃O₄-Hb@cur was studied by Confocal Laser Scanning Microscopy (CLSM). As shown in Fig. 4A, intense fluorescence of cur was observed in the cytoplasm, indicating the successful uptake of Fe₃O₄-Hb@cur by 4T1 cells. With the prolongation of the incubation time, cur was significantly accumulated in the nucleus, which indicated that after the cellular uptake of Fe₃O₄-Hb@cur magnetic nanoenzymes, the Fe₃O₄ nanocarriers were ruptured in acidic lysogeny and the intracellular cur was gradually released. Semiquantitative analysis of CLSM images in Fig. 4B showed that the uptake of Fe₃O₄-Hb@cur by 4T1 tumor cells increased with prolonged incubation times (1, 5, and 9 h).

3.5. Laser enhances cytotoxicity and cell apoptosis

Next, we investigated the synergistic photodynamic and catalytic effects of Fe₃O₄-Hb@cur as a nanoenzyme on tumor photodynamic and catalysis in vitro. We firstly tested the cytotoxicity of Fe₃O₄-Hb@cur in the dark. The release of cur and the generation of 'OH could significantly enhance the synergistic photodynamic/catalytic treatment of cancer with the help of enhanced O2 concentration. The synergistic anticancer effect of the drugs was studied using cell viability assay kit (CCK-8) analysis. When electronically coupled reagents are




Fig. 4 Cellular uptake (A) and relative fluorescence intensity (B) of Fe₃O₄-Hb@cur after 1, 5, and 9 h of co-incubation with 4T1 cells. Scale bar: 8 nm. Cytotoxicity at different concentrations of Fe₃O₄, Fe₃O₄@cur, and Fe₃O₄-Hb@cur against 4T1 cells in the dark (C) and after 30 seconds of laser irradiation (D). Data are presented as means \pm SD (n = 3 per group). ***P < 0.001 and ****P < 0.0001. Detection of ROS levels (E) with the relative fluorescence intensity (F) after different treatments in DCFH-DA-stained 4T1 cells for 12 h. Scale bar: 10 nm.

Nanoscale Paper

present, WST-8 (2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-dithiobenzyl)-2H-tetrazole monosodium salt) undergoes reduction by mitochondrial dehydrogenase, resulting in the formation of a water-soluble, orange-yellow product (formazan) that produces methane. The optical density (OD) of the formazan product can be measured at 450 nm using an enzyme marker. It can indirectly reflect the number of living cells. As shown in Fig. 4C, Fe₃O₄-Hb@cur showed stronger cell damage than Fe₃O₄ @cur or free Fe₃O₄ at each concentration, which was attributed to the much stronger effect of synergistic photodynamic therapy/catalytic treatment than PDT or catalytic treatment alone. Furthermore, the synergistic killing effect of O₂ for both treatments was confirmed.

To evaluate the photodynamic conversion efficiency of cur, the cells were treated with a 660 nm laser (0.5 W cm⁻², 2 min). The relative survival was determined after 24 h using a colorimetric assay with tetramethyl azole salts. As shown in Fig. 4D, both types of irradiation with 660 nm excitation light exhibited enhanced drug cytotoxicity. Compared with the unenhanced light irradiation (Fe₃O₄-Hb@cur), the combined Fe₃O₄-Hb@cur under laser treatment showed the best killing effect on cancer cells at different concentrations (Fig. S7†). Therefore, the high antitumor effect of Fe₃O₄-Hb@cur in these treatments was mainly attributed to the enhanced catalytic therapy induced by the self-contained O2 of the magnetic nanoenzymes as well as PDT.

3.6. Light-triggered ROS generation

Fe₃O₄@cur can react with local H₂O₂ to produce highly toxic OH in the acidic tumor microenvironment. In this study, 2',7'dichlorofluorescein diacetate (DCFH-DA) was used to detect the intracellular ROS levels to confirm whether light irradiation increased intracellular ROS levels. Intracellular ROS can catalyze the generation of fluorescent 2,7-dichlorofluorescent diacetate (DCF) from non-fluorescent DCFH, allowing it to be used as a ROS probe to dynamically monitor ROS by fluorescent signals. 41,42 As shown in Fig. 4E, little fluorescence was observed in 4T1 cells without any treatment, regardless of whether light was increased, indicating basal intracellular ROS levels. In contrast, ROS levels were significantly up-regulated after 4T1 cells co-cultured with Fe₃O₄@cur or Fe₃O₄-Hb@cur, which was shown as the intense green fluorescence in CLSM images (Fig. S8†). Thus, Fe₃O₄@cur contributed to the improvement of ROS production. Furthermore, a higher fluorescence intensity was detected in Fe₃O₄ -Hb@cur treated cells, proving that O2 in Fe3O4-Hb@cur was successfully involved in ¹O₂ generation. This further confirms that the PDT toxicity of Fe₃O₄ @cur can be enhanced by the increase of O₂.

3.7. Biodistribution and tumor-targeting activity of Fe₃O₄ -Hb@cur

The biological distribution and anticancer effect of Fe₃O₄-Hb@cur in vivo was further investigated in 4T1 Tumor-bearing Balb/c Mice. With its long excitation and emission wavelengths, Cy5.5 exhibits the capability of penetrating deep tissues, making it an ideal probe for the detection of tumor accumulation. To visually observe the biological distribution of Fe₃O₄-Hb@cur in vivo, intravenous Fe₃O₄-Hb@cur was labeled with Cy5.5 (denoted as Fe₃O₄-Hb@cur-Cy5.5). Imaging and quantification of the whole body and major organs were recorded and presented using an IVIS spectrum small animal imaging system at different time intervals after intravenous injection (Fig. 5A and B). Due to the enhanced permeability and retention (EPR) effect of tumor, the material accumulates at the tumor site. 43-45 As time progressed, the fluorescence intensity of the Fe₃O₄-Hb@cur-Cy5.5 group remained consistently stronger than that of the Cy5.5 group. Free Cy5.5 exhibited a shorter circulation half-life in mice, while the Fe₃O₄-Hb@cur-Cy5.5 group demonstrated a prolonged circulation time. At 12 and 48 h post-injection, the in vivo fluorescence intensity of the Fe₃O₄-Hb@cur-Cy5.5 group was 1.76-fold and 3.05-fold higher, respectively, compared to free Cy5.5. This indicates effective accumulation of Fe₃O₄-Hb@cur-Cy5.5 within the tumor (Fig. 5C). Moreover, in vitro imaging and quantitative analysis of major organs in Fig. 5D showed that there was still a strong fluorescence signal in Fe₃O₄-Hb@cur-




Fig. 5 In vivo imaging study of Fe₃O₄-Hb@cur. (A and B) In vivo fluorescence images of 4T1 tumor-bearing mice dissected organs, and tumors after intravenous injected with free Cy5.5 or Fe₃O₄-Hb@cur-Cy5.5 for different times. (C) Quantitative mean fluorescence intensity analysis of tumor tissues at different time intervals (n = 5 per group). ***P < 0.001 and ****P < 0.0001. (D) Fluorescence intensities of the organs and tumors collected at different times. (E) Over a period of time following the intravenous administration of Fe₃O₄-Hb@cur nanoparticles, in vivo magnetic resonance (MR) imaging was performed in the tumor region to monitor changes. (F) T2-Weighted magnetic resonance imaging of mouse tumors at 0 h and 3 h after tail vein injection of Fe₃O₄-Hb@cur nanoparticles.

Paper Nanoscale

Cy5.5 in the tumor and liver 48 h after injection, further confirming the long cycle and EPR effect of Fe₃O₄-Hb@cur.

Furthermore, MRI was employed to evaluate the accumulation of Fe₃O₄-Hb@cur. To evaluate the accumulation of Fe₃O₄-Hb@cur in vivo, T₂-weighted MRI was performed by utilizing 4T1 breast cancer xenografts after intravenous administration of Fe₃O₄-Hb@cur. Fig. 5E shows the observed MRI signal, which increased over time. Furthermore, further analysis was conducted on the statistically semi-quantitative T_2 signal intensity. The signal intensity of Fe₃O₄-Hb@cur was the lowest at 3 h, while the signal intensity of the PBS group showed no difference at 3 h, consistent with the imaging results (Fig. 5F). This further confirms the strong tumor accumulation ability of Fe₃O₄-Hb@cur through prolonged circulation and the EPR effect. Therefore, Fe₃O₄-Hb@cur nanoparticles can effectively target tumor sites.

3.8. In vivo synergistic photodynamic and catalytic antitumor effects of Fe₃O₄-Hb@cur

The combined therapeutic effect of Fe₃O₄-Hb@cur, which exhibits effective tumor targeting and PDT effects, was evaluated by inoculating 4T1 cells in BALB/c mice. As shown in Fig. 6A, the mice were randomly divided into 5 groups (6 mice per group), and each group was injected with 100 µL saline (as a control), Fe₃O₄, Fe₃O₄@cur, Fe₃O₄-Hb@cur, Fe₃O₄@cur + laser (Fe₃O₄@cur (+)) and Fe₃O₄-Hb@cur + laser (Fe₃O₄-




Fig. 6 (A) Schematic illustration showing the time line of the efficacy study. (B) Tumor weights in each group at the end of 16 days of treatment. (C) Tumor volume curves of 4T1 tumor-bearing mice with different treatments at 16 days. Obvious inhibition occurred in the Fe_3O_4 -Hb@cur (+) group (***P < 0.001 and ****P < 0.001). (D) Representative images of the 4T1-bearing mice at day 0 and day 16 after different treatments. (E) Tumor histological images of H&E staining and Ki67 staining of tumors after different treatments. Scale bar: 100 μm. TUNEL staining for apoptosis in tumors extracted from mice with different treatments (green, TUNEL; blue, and DAPI). Scale bar: 50 µm.

Hb@cur (+)) through the caudal vein (an equivalent dose of 25 mg kg⁻¹ cur or 0.75 mg kg⁻¹ Fe₃O₄). When the tumor volume reached approximately 150 mm³, the treatments were initiated. Irradiation was scheduled at 12 and 24 h after drug injection based on the results of biodistribution of nanoparticles. Subsequently, the tumors were exposed to a 660 nm laser with a power density of 0.5 W cm⁻² for a duration of 5 minutes. Fig. 6B-D illustrates that only the Fe₃O₄-Hb@cur(+) group exhibited a significant inhibition of tumor volume and tumor load, following the completion of the treatments, whereas the other four groups displayed only slight tumor growth suppression. In contrast, the Fe₃O₄-Hb@cur treatment group observed a strong inhibitory effect on the tumor, which was completely suppressed within 16 days of tumor growth. The therapeutic effect was clearly observed in post-treatment mouse tumor photographs (Fig. 6B). It is worth noting that the Fe₃O₄-Hb@cur nanoparticle group also demonstrated a moderate level of antitumor activity, further confirming the important role of O₂ carried by Hb in enhanced catalytic therapy and PDT. Additionally, Fe₃O₄@cur also exhibited certain therapeutic efficacy, which may be attributed to the ability of Fe₃O₄@cur to effectively accumulate tumors and administer chemotherapy. Overall, there were few significant fluctuations in the body weights of the mice after the various treatments, suggesting that Fe₃O₄ nanoparticles can have negligible side effects in the tumor treatment at the doses used (an equivalent dose of 25 mg kg⁻¹ cur or 0.75 mg kg⁻¹ Fe₃O₄) (Fig. S9 \dagger).

At the end of the treatment, the mice were euthanized, and tumor tissues were dissected and stained for histological analysis (Fig. 6E). Hematoxylin and eosin (H&E) staining of the tumor tissues revealed typical characteristics: enlarged and darkly stained cell nuclei with minimal cytoplasm. Compared to the saline group, the treatment group with Fe₃O₄-Hb@cur (+) showed severe cell necrosis in the tumor tissue, while other groups exhibited varying degrees of tumor necrosis induction. Additionally, the TUNEL assay and Ki67 immunohistochemistry were performed to further support the above findings, indicating that Fe₃O₄-Hb@cur had the strongest positive reaction and the most significant therapeutic effect. Therefore, Fe₃O₄-Hb@cur demonstrated effective tumor targeting and synergistic anti-tumor activity.

3.9. Biosafety evaluation

To investigate the in vivo toxicity of the nanomedicine, on the 16th day of treatment, the mice were killed by cervical disand tumors were isolated and weighed. Comprehensive analysis of blood, physiological, and biochemical parameters of the mice was conducted. H&E staining was performed on major organ tissue samples (Fig. 7A). No significant organ damage or inflammatory injury was observed in any of the treatment groups compared with the physiological saline group, indicating that photodynamic and catalytic synergistic therapies had no significant impact on the organ function of mice. To assess liver and kidney damage, blood levels of AST, ALT, BUN and CRE were measured and analyzed. No significant differences occurred and remained within the

Nanoscale **Paper**




Fig. 7 In vivo biosafety of Fe₃O₄-Hb@cur. (A) H&E-stained images of major organs (heart, liver, spleen, lungs, and kidneys) harvested from 4T1 tumor-bearing mice. Scale bar, 100 nm. Each mouse was treated with a single dose of saline (as a control), Fe₃O₄, Fe₃O₄@cur, Fe₃O₄-Hb@cur, Fe_3O_4 @cur + laser (Fe_3O_4 @cur (+)) and Fe_3O_4 -Hb@cur + laser (Fe₃O₄-Hb@cur (+)) through the caudal vein ((an equivalent dose of 25 mg kg⁻¹ cur or 0.75 mg kg⁻¹ Fe₃O₄)) through i.v. administration. Mice were sacrificed 16 days after drug administration. (B-D) The serology panel for biomarkers of body function and hepatic function of mice in the control and various Fe containing groups (an equivalent dose of 25 mg kg⁻¹ cur or 0.75 mg kg⁻¹ Fe₃O₄) after treatment for 16 days. n =6 mice per group. AST: aspartate aminotransferase; ALT: alanine transaminase; BUN: blood urea nitrogen; and CRE: creatinine.

normal physiological range with regard to all these biomarkers (Fig. 7B-D). Therefore, all the measured parameters and pathological assessments indicated that Fe₃O₄, Fe₃O₄@cur, Fe₃O₄-Hb@cur, Fe₃O₄ @cur + laser and Fe₃O₄-Hb@cur + laser (at an equivalent dose of 25 mg kg⁻¹ cur or 0.75 mg kg⁻¹ Fe₃O₄) did not exhibit significant toxicity.

4. Conclusions

In summary, we reported a unique type of O2 enhanced multifunctional nano-sonosensitizer platform using Fe₃O₄ and cur, encapsulated by Hb (Fe₃O₄-Hb@cur) successfully. Fe₃O₄ can catalyze the conversion of H2O2 to 'OH through the Fenton reaction, while Hb can self-supply O2, overcoming the hypoxia barrier of tumor cells. Working as a photosensitizer, cur was effectively stimulated by low-intensity laser irradiation at 660 nm and this excited the surrounding O2 to generate a certain amount of ¹O₂ for tumor inhibition. Therefore, Fe₃O₄-Hb@cur exhibited a synergistic CDT/PDT anticancer ability in vitro and in vivo. Additionally, Fe₃O₄-Hb@cur remarkably enhanced T₂-weighted MR imaging. Overall, Fe₃O₄-Hb@cur

with MR imaging-guided synergistic cancer therapy has the potential for bio-application in the bio-nanomedicine field.

Author contributions

X.Y.C. directed and conceived this project with input from T.Y. and Q.Z. synthesized the materials, performed most of the experimental work, and wrote the draft of the manuscript. T.T. X., R.D. and D.Z. performed the experimental data analysis. G. Q.C. and W.C.Z. performed the theoretical study. H.P. and J.J. H. designed the project. All the authors discussed the results and contributed to the writing and revision of the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (81901864 and 82272821), the Natural Science Foundation of Guangdong Province (2022A1515011337), the Guangdong Province Universities and Colleges Characteristic Innovation (2021KTSCX036), the Traditional Chinese Medicine Research Project of Guangdong Province Traditional Chinese Medicine Bureau (20221206), the Natural Science Foundation of Shanghai (22ZR1467600), the Special Funds for Scientific Technological Innovation of Undergraduates in Guangdong Province (pdjh2022b0227), the Funds for Ph.D. researchers of Guangdong Medical University in 2023, and the Discipline Construction Project of Guangdong Medical University (4SG24015G, 4SG22009G).

References

- 1 S. Yao, M. Zheng, S. Wang, T. Huang, Z. Wang, Y. Zhao, W. Yuan, Z. Li, Z. L. Wang and L. Li, Adv. Funct. Mater., 2022, 32, 2209142.
- 2 S. Yao, X. Zhao, X. Wang, T. Huang, Y. Ding, J. Zhang, Z. Zhang, Z. L. Wang and L. Li, Adv. Mater., 2022, 34, 2109568.
- 3 Y. Nie, W. Chen, Y. Kang, X. Yuan, Y. Li, J. Zhou, W. Tao and X. Ji, Biomaterials, 2023, 295, 122031.
- 4 L. Hou, F. Gong, Z. Han, Y. Wang, Y. Yang, S. Cheng, N. Yang, Z. Liu and L. Cheng, Angew. Chem., Int. Ed., 2022, 61, e202208849.
- 5 X. Wang, Q. Chen, Y. Zhu, K. Wang, Y. Chang, X. Wu, W. Bao, T. Cao, H. Chen, Y. Zhang and H. Qin, Signal Transduction Targeted Ther., 2023, 8, 277.
- 6 L. Chen, Z. Mao, Y. Wang, Y. Kang, Y. Wang, L. Mei and X. Ji, Sci. Adv., 2022, 8, eabo7372.
- 7 C. Zhang, J. Huang, Z. Zeng, S. He, P. Cheng, J. Li and K. Pu, Nat. Commun., 2022, 13, 3468.

Paper

8 C. Cao, N. Yang, X. Wang, J. Shao, X. Song, C. Liang, W. Wang and X. Dong, Coord. Chem. Rev., 2023, 491,

215245. 9 X. Song, H. Huang, L. Xia, W. Jia, S. Yang, C. Wang and

Y. Chen, Adv. Sci., 2023, 2301279.

- 10 Z. Li, C. Wang, C. Dai, R. Hu, L. Ding, W. Feng, H. Huang, Y. Wang, J. Bai and Y. Chen, *Biomaterials*, 2022, 287, 121668.
- 11 H. Yu, Y. Cheng, C. Wen, Y.-Q. Sun and X.-B. Yin, *Biomaterials*, 2022, **280**, 121308.
- 12 Y. Zhu, R. Zhao, L. Feng, C. Wang, S. Dong, M. V. Zyuzin, A. Timin, N. Hu, B. Liu and P. Yang, ACS Nano, 2023, 17, 6833–6848.
- 13 C. Zhang, W.-J. Qin, X.-F. Bai and X.-Z. Zhang, *Nano Today*, 2020, 35, 100960.
- 14 C. Liao, X. Liu, C. Zhang and Q. Zhang, Semin. Cancer Biol., 2023, 88, 172–186.
- 15 S. Wang, Z. Wang, Z. Li, J. Xu, X. Meng, Z. Zhao and Y. Hou, *Adv. Healthc. Mater.*, 2022, **11**, e2201240.
- 16 S. L. Li, P. Jiang, F. L. Jiang and Y. Liu, Adv. Funct. Mater., 2021, 31, 2100243.
- 17 Y. Rao, T. Fan, L. Zhou, K. Fang, Y. Sun, X. Hu, A. Wang, R. Li, Z. Zhu and C. Dong, *J. Controlled Release*, 2023, 354, 701–712.
- 18 Y. He, S. Guo, Y. Zhang, Y. Liu and H. Ju, *Biomaterials*, 2021, 275, 120962.
- 19 C. Zhang, J. Huang, X. Guo, X. Da, Z. Dai, M. Hassan, Y. Yu, X. Wang and Q. Zhou, *Nano Today*, 2023, **50**, 101824.
- 20 P. Lin, J. Shi, L. Ming, Y. Sheng, L. Song, M. Hong and Y. Zhang, *Chem. Eng. J.*, 2022, 442, 135638.
- 21 Q. Yu, J. Zhou, J. Song, H. Zhou, B. Kang, H. Y. Chen and J. J. Xu, *Small*, 2023, **19**, 2206592.
- 22 M. Yuan, S. Liang, Y. Zhou, X. Xiao, B. Liu, C. Yang, P. a. Ma, Z. Cheng and J. Lin, *Nano Lett.*, 2021, 21, 6042–6050.
- 23 S. Y. Wu, Y. X. Ye, Q. Zhang, Q. J. Kang, Z. M. Xu, S. Z. Ren, F. Lin, Y. T. Duan, H. J. Xu, Z. Y. Hu, S. S. Yang, H. L. Zhu, M. J. Zou and Z. C. Wang, *Adv. Sci.*, 2023, 10, e2203742.
- 24 R. Li, Z. Lin, Q. Zhang, Y. Zhang, Y. Liu, Y. Lyu, X. Li, C. Zhou, G. Wu, N. Ao and L. Li, ACS Appl. Mater. Interfaces, 2020, 12, 17936–17948.
- 25 R. Li, Z. Lin, Q. Zhang, Y. Zhang, Y. Liu, Y. Lyu, X. Li, C. Zhou, G. Wu and N. Ao, ACS Appl. Mater. Interfaces, 2020, 12, 17936–17948.
- 26 Z. Zhang, R. Wang, X. Huang, R. Luo, J. Xue, J. Gao, W. Liu, F. Liu, F. Feng and W. Qu, ACS Appl. Mater. Interfaces, 2020, 12, 5680–5694.
- 27 Q. Zhang, W. Lai, T. Yin, C. Zhang, C. Yue, J. Cheng, K. Wang, Y. Yang, D. Cui and W. J. Parak, *Bioconjugate Chem.*, 2018, 29, 2120–2125.

- 28 Q. Zhang, T. Yin, G. Gao, J. G. Shapter, W. Lai, P. Huang, W. Qi, J. Song and D. Cui, ACS Appl. Mater. Interfaces, 2017, 9, 17777–17785.
- 29 Y. Matsumura and H. Maeda, Cancer Res., 1986, 46, 6387–6392.
- 30 H. Maeda, T. Sawa and T. Konno, J. Controlled Release, 2001, 74, 47-61.
- 31 X. Huang, B. Sheng, H. Tian, Q. Chen, Y. Yang, B. Bui, J. Pi, H. Cai, S. Chen and J. Zhang, *Acta Pharm. Sin. B*, 2023, 13, 1303–1317.
- 32 W. Yang, C. Deng, X. Shi, Y. Xu, C. Dai, H. Wang, K. Bian, T. Cui and B. Zhang, ACS Nano, 2023, 17, 4009–4022.
- 33 Y. Wang, D. Wang, Y. Zhang, H. Xu, L. Shen, J. Cheng, X. Xu, H. Tan, X. Chen and J. Li, *Bioact. Mater.*, 2023, 22, 239–253.
- 34 J. Li, H. Tian, F. Zhu, S. Jiang, M. He, Y. Li, Q. Luo, W. Sun, X. Liu and P. Wang, *Adv. Healthcare Mater.*, 2022, **11**, 2201986.
- 35 M. Tang, Y. Shi, L. Lu, J. Li, Z. Zhang, J. Ni, W. Wang, Y. Zhang, T. Sun and Z. Wu, *Chem. Eng. J.*, 2022, 449, 137847.
- 36 S. Gao, H. Lin, H. Zhang, H. Yao, Y. Chen and J. Shi, *Adv. Sci.*, 2019, **6**, 1801733.
- 37 L. H. Liu, Y. H. Zhang, W. X. Qiu, L. Zhang, F. Gao, B. Li, L. Xu, J. X. Fan, Z. H. Li and X. Z. Zhang, *Small*, 2017, 13, 1701621.
- 38 C. P. Liu, T. H. Wu, C. Y. Liu, K. C. Chen, Y. X. Chen, G. S. Chen and S. Y. Lin, *Small*, 2017, 13, 1700278.
- 39 Z. Zhang, R. Wang, X. Huang, R. Luo, J. Xue, J. Gao, W. Liu, F. Liu, F. Feng and W. Qu, ACS Appl. Mater. Interfaces, 2020, 12, 5680–5694.
- 40 Q. You, Q. Sun, M. Yu, J. Wang, S. Wang, L. Liu, Y. Cheng, Y. Wang, Y. Song, F. Tan and N. Li, ACS Appl. Mater. Interfaces, 2017, 9, 40017–40030.
- 41 Y. Ma, Z. Su, L. Zhou, L. He, Z. Hou, J. Zou, Y. Cai, D. Chang, J. Xie, C. Zhu, W. Fan, X. Chen and S. Ju, Adv. Mater., 2022, 34, e2107560.
- 42 T. Yin, H. Chen, A. Ma, H. Pan, Z. Chen, X. Tang, G. Huang, J. Liao, B. Zhang, M. Zheng and L. Cai, *Biomaterials*, 2023, 293, 121992.
- 43 S. Wilhelm, A. J. Tavares, Q. Dai, S. Ohta, J. Audet, H. F. Dvorak and W. C. W. Chan, *Nat. Rev. Mater.*, 2016, 1, 16014.
- 44 B. Ouyang, W. Poon, Y. N. Zhang, Z. P. Lin, B. R. Kingston, A. J. Tavares, Y. Zhang, J. Chen, M. S. Valic, A. M. Syed, P. MacMillan, J. Couture-Senécal, G. Zheng and W. C. W. Chan, *Nat. Mater.*, 2020, 19, 1362–1371.
- 45 Y. Matsumoto, J. W. Nichols, K. Toh, T. Nomoto, H. Cabral, Y. Miura, R. J. Christie, N. Yamada, T. Ogura, M. R. Kano, Y. Matsumura, N. Nishiyama, T. Yamasoba, Y. H. Bae and K. Kataoka, *Nat. Nanotechnol.*, 2016, 11, 533–538.