



 Cite this: *RSC Adv.*, 2024, 14, 39436

Construction, structural modification, and bioactivity evaluation of pentacyclic triterpenoid privileged scaffolds in active natural products

 Ling Rong Zeng,[†] Bo Wen Pan,[‡]  [†] Juan Cai,[†] Li Juan Liu, Zhang Chao Dong, Ying Zhou,^{*} Ting Ting Feng^{*} and Yang Shi^{*}

Pentacyclic triterpenoids, as important representatives of natural products, have garnered widespread attention due to their diverse biological activities, including anti-inflammatory, antiviral, and antitumor effects. Oleanolic acid (OA), betulinic acid (BA), ursolic acid (UA), triptolide, and glycyrrhetic acid (GA) are typical examples of pentacyclic triterpenoids. Despite their significant biological activities, their poor water solubility and low bioavailability have limited further development and application. In recent years, researchers have developed a series of derivatives with enhanced biological activities and improved drug properties through structural modifications of these compounds, particularly achieving notable progress in the field of antitumor therapy. This review summarizes recent advances in the structural modification of pentacyclic triterpenoids and explores their promising applications in the development of antitumor, antiviral, and other therapeutic agents.

 Received 24th October 2024
 Accepted 9th December 2024

DOI: 10.1039/d4ra07602h

rsc.li/rsc-advances

Terpenoids are a class of compounds and their derivatives with a molecular scaffold based on isoprene units, which are widely distributed in nature. They are major components of secondary metabolites such as plant essential oils, resins, and pigments. Terpenoids are classified into monoterpenes, sesquiterpenes, diterpenes, sesterterpenes, triterpenes, tetraterpenes, and polyterpenes based on the number of isoprene units in their molecular structure. Triterpenes, composed of a basic skeleton of 30 carbon atoms, are important plant secondary metabolites found in nature either in free form or as glycosides or esters bound to sugars. They exhibit various pharmacological activities, including anti-cancer, antiviral, and cholesterol-lowering effects. Triterpenoids can be further classified into monocyclic, bicyclic, tricyclic, tetracyclic, and pentacyclic triterpenes, based on the number of rings in their structure, with tetracyclic and pentacyclic triterpenes being the most common.

Pentacyclic triterpenoids possess various modification types in their carbon skeletons, resulting in complex and diverse structures, which have garnered significant attention due to their wide range of functions. Based on their aglycone types, pentacyclic triterpenoids are divided into four main categories: oleanane-type [e.g., OA, GA, and hawthorn acid (HA)], lupane-type (e.g., BA, lupeol, and betulin), ursane-type [e.g., UA and asiatic acid (AA)], and friedelane-type (e.g., *Tripterygium wilfordii* polyglycosides). Oleanane-type triterpenoids exhibit extensive pharmacological activity; OA is commonly used clinically for its

hepatoprotective effects,^{1–3} while glycyrrhizin-based formulations have been marketed for the treatment of hepatitis.⁴ The oleanane-type, also known as the β -amyrane-type, includes compounds such as β -amyrin, germanicol, OA, and GA.

Lupane-type compounds mainly include lupeol, betulin, BA, and their derivatives. Lupeol is widely distributed in nature and serves as the core skeleton for lupane-type triterpenoids, showing significant anti-inflammatory⁵ and anti-cancer⁶ activities. Betulinic acid, formed through three consecutive oxidation steps at the C-28 position of lupeol, not only demonstrates potent anti-HIV activity,⁷ but also selectively inhibits melanoma cell proliferation,⁸ making it a promising candidate for anti-HIV and anti-melanoma drug development.^{9,10} BA is predominantly found in the bark of birch trees.

The ursane-type, also known as the α -amyrane-type, includes compounds such as α -amyrin, UA, and taraxasterol. α -Amyrin is a precursor for ursane-type triterpenoids, whose structural diversity can be enhanced through glycosylation, acylation, and redox reactions. UA, also known as ursane acid or UA, is an oxidation product of α -amyrin at the C-28 position and exhibits anti-cancer, anti-diabetic, anti-ulcer, and lipid-lowering effects.^{11–13} UA is found in various plants, including fruit trees, aromatic herbs, and traditional Chinese medicinal herbs.

1 Construction of pentacyclic triterpenoid privileged scaffolds

The synthesis of pentacyclic triterpenoid scaffolds primarily occurs through two pathways:¹⁴ the methylerythritol phosphate (MEP) pathway in plastids and the mevalonate (MVA) pathway.

College of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang, 550025, China. E-mail: xiaoyewater@163.com; ft0809@163.com; yingzhou71@126.com

[†] These authors contributed equally to this work.






Fig. 1 Synthesis of the pentacyclic triterpenoid frameworks OA, BA, and UA.

In plastids, isopentenyl diphosphate (IPP) is synthesized from pyruvate and glyceraldehyde-3-phosphate *via* seven enzymatic reactions, after which IPP is transported into the cytoplasm to participate in terpenoid synthesis. Under the action of IPP isomerase, IPP and dimethylallyl pyrophosphate (DMAPP) interconvert. In the cytoplasmic MVA pathway, acetyl-CoA acts as the initial donor to produce IPP. IPP and DMAPP are then converted into farnesyl diphosphate under the action of farnesyl diphosphate synthase, which is subsequently converted into squalene by squalene synthase. Squalene is further oxidized into 2,3-oxidosqualene *via* squalene epoxidase (SE).¹⁵ The cyclization of 2,3-oxidosqualene, catalyzed by oxidosqualene cyclase (OSC), produces various tetracyclic and pentacyclic triterpenoid scaffolds. These scaffolds undergo chemical modifications, such as oxidation, substitution, and glycosylation, through the action of cytochrome P450 monooxygenases and glycosyltransferases, with uridine diphosphate (UDP) as the

glycosyl donor, ultimately forming various pentacyclic triterpenoid compounds. The synthesis pathways of **OA**, **BA**, and **UA** scaffolds are shown in Fig. 1. OSC, a key enzyme in this process, catalyzes the cyclization of 2,3-oxidosqualene to produce sterol and triterpenoid precursors, representing the critical step in generating the diversity of triterpenoid products.

2 Structural modification of pentacyclic triterpenoid compounds

In the structure of pentacyclic triterpenoids, the modifications of **OA**, **BA**, and **UA** primarily occur at the hydroxyl group on the A-ring, the carboxyl group at the C-28 position, and the double bond on the C-ring. Rational structural modifications of these active groups can significantly enhance the antitumor activity of these compounds.



2.1 Salt formation modification

Based on the structural characteristics of pentacyclic triterpenoid compounds, there are three direct methods of salt formation: carboxyl groups forming salts with alkali metal ions; introduction of hydrophilic groups followed by salt formation with alkali metal ions; and the introduction of quaternary ammonium salts. Zhao *et al.*¹⁶ and Zhai *et al.*¹⁷ demonstrated that converting **OA** into its sodium salt significantly increased its solubility in water compared to **OA**. After administering sodium oleanolate to rats, Li *et al.*¹⁸ observed a marked improvement in subacute liver injury, with the hepatoprotective effect being significantly stronger than that of **OA**. This indicates that salt formation significantly enhances the water solubility, bioavailability, and bioactivity of **OA**.

2.2 Amino acid conjugation

Amino acids, as essential nutrients for the human body, are commonly used in the treatment of liver diseases, neurological disorders, and psychiatric conditions. Due to the presence of both carboxyl and hydroxyl polar groups in their structure, coupling amino acids with pentacyclic triterpenoid compounds can enhance the water solubility of the drug and improve its bioavailability. Zhang *et al.*¹⁹ introduced amino acids at the C-28 position of **OA**, and the resulting **OA** derivatives exhibited significantly higher antitumor activity compared to the parent compound.

2.3 Glycosylation

Glycosides are polyhydroxy compounds widely distributed in the roots, stems, and leaves of plants, characterized by high solubility and strong polarity. Currently, numerous studies have focused on using **OA** as a starting material to design and synthesize **OA** glycoside derivatives with enhanced biological activities. These **OA** derivatives have shown significantly improved pharmacological activities. Meng *et al.*²⁰ synthesized 10 **OA** derivatives through a series of reactions, using natural **OA** as the lead compound. Pharmacological activity analysis of two glycoside derivatives revealed that one of the derivatives exhibited significantly better antitumor activity than **OA**, while the other showed even stronger antitumor activity, comparable to the antitumor drug gefitinib.

2.4 Introduction of ester and amide bonds

Based on the structural characteristics of **UA**, **OA**, and **BA**, the hydroxyl group at the C-3 position can undergo esterification reactions with compounds containing carboxyl groups to form ester bonds. Additionally, the carboxyl group at the C-28 position can react with amino or hydroxyl groups through dehydration to form either ester or amide bonds, thereby improving their water solubility and enhancing bioavailability. Zhang *et al.*²¹ applied the principle of molecular hybridization in drug synthesis, coupling oxo-oleanolic acid with aminophosphonic acid aryl esters through amide bonds. The resulting novel **OA** derivatives exhibited better water solubility and higher bioactivity compared to the parent compound.

3 Structural modification and bioactivity of oleanolic acid

OA, also known as Cenoside, is a pentacyclic triterpenoid compound of the oleanane type. It is widely distributed in nature and can be found in various plants such as the fruit of *Ligustrum lucidum* and the whole herb of *Rabdosia rubescens*, existing either in its free form or as glycosides. **OA** exhibits a wide range of biological activities, common to pentacyclic triterpenoids, including antitumor activity,^{22–24} antiviral properties,²⁵ antihyperlipidemic effects,^{26,27} anti-inflammatory actions,^{28,29} and hypoglycemic effects. **OA** is considered an ideal drug for the treatment of chronic viral hepatitis and acute jaundice hepatitis³⁰ due to its low toxicity and side effects. The anticancer activity of **OA** and its derivatives is particularly broad, showing inhibitory effects on various human tumor cells. Yang *et al.*³¹ reported that **OA** and its derivatives are well-known for their potent antitumor activity, including inhibiting tumor cell proliferation, inducing cell cycle arrest, apoptosis, and differentiation. These effects have been observed in various cancer cell lines, such as A549, SGC7901, MCF-7, HT1080, and MKN-45 cells. Jannus *et al.*³² further highlighted that the anticancer properties of **OA** and its derivatives have been demonstrated in numerous *in vitro* and *in vivo* models, showing significant anticancer effects. In cancer, the homeostatic balance between cell proliferation and death is disrupted, and apoptosis, a physiological process of programmed cell death, is one of the key mechanisms for controlling this balance.

However, the major limitation of **OA** and its derivatives as anticancer agents is their poor water solubility, which leads to low bioavailability *in vivo*. Therefore, it is necessary to introduce polar groups and perform rational structural modifications to enhance solubility and bioavailability, thus improving the clinical applicability of these compounds.

3.1 Structural modifications of oleanolic acid and antitumor activity

3.1.1 Modification of hydroxyl group at C-3 or carboxyl group at C-28 of oleanolic acid.

Spivak *et al.*³³ designed and synthesized derivatives of **OA** with a C-28 carboxyl group connected to a guanidine group through various linkers to investigate the influence of the guanidine moiety on the anticancer properties of **OA**. The synthesized guanidine-containing **OA** derivatives (compound 3, Fig. 2) exhibited significantly higher cytotoxicity in Jurkat cells compared to the original **OA**. Most of the guanidine derivatives tested demonstrated higher IC₅₀ values than the amine counterparts, while showing lower toxicity towards human fibroblasts.

Chouaib *et al.*³⁴ synthesized two series of **OA** 1,2,3-triazole derivatives (compounds 4 and 5, Fig. 3) using CuAAC or RuAAC as catalysts under microwave conditions. The introduction of the 1,2,3-triazole moiety enhanced their activity against mouse mammary cancer (EMT-6) and human colon cancer (SW480) cells.

Khusnutdinova *et al.*³⁵ synthesized alkynyl derivatives 6 and 7 (Fig. 4) through amidation and the Mannich reaction. These






Fig. 2 Synthesis of OA derivative 3.




Fig. 3 Synthesis of OA derivatives 4 and 5.




Fig. 4 Synthesis of OA derivatives 6 and 7.

derivatives exhibited *in vitro* anticancer activity against leukemia cells and human colon cancer, acting through their pro-apoptotic activity, stimulating the activity of apoptosis-related caspases 3 and 8, enhancing the activity of NF-κB transcription factors, influencing BAX protein expression, and inhibiting DNA polymerase.

3.1.2 Modification of C-2, ring A, and other positions of oleanolic acid. Liu *et al.*³⁶ synthesized a variety of OA derivatives (compounds 8 and 9, Fig. 5) involving the condensation of formaldehyde at the C-2 position, oxidation of the C-3 hydroxyl group to a carbonyl, and the attachment of ethylenediamine at C-28. Compounds 8 and 9 demonstrated inhibitory effects on

various types of cancer stem cells (CSCs) involved in tumorigenesis, tumor recurrence, invasion, metastasis, and resistance.

Meng *et al.*³⁷ modified the A ring and C-28 position of OA to synthesize ten OA derivatives (compounds 12–21, Fig. 6). The inhibitory activities of these derivatives were evaluated against the SGC7901 and A549 cell lines, confirmed through the MTT assay. The experimental results indicated that all compounds exhibited certain antitumor activities against the SGC-7901 and A-549 cell lines. Among them, the OA derivatives 15 and 21 demonstrated superior antitumor activity against both SGC7901 and A549 cells compared to the positive drug gefitinib.







Fig. 5 The structures of OA derivatives 8 and 9.



12. $R_1 = \text{CH}_2\text{C}_6\text{H}_5$; 13. $R_1 = \text{CH}_2\text{CH}_3$; 14. $R_1 = \text{CH}(\text{CH}_3)_2$; 15. $R_1 = \text{CH}_2\text{CH}(\text{CH}_3)_2$; 16. $R_1 = (\text{CH}_2)_5\text{CH}_3$;
 17. $R_2 = \text{CH}_2\text{CH}_3$; 18. $R_2 = \text{CH}(\text{CH}_3)_2$; 19. $R_2 = (\text{CH}_2)_3\text{CH}_3$; 20. $R_2 = \text{C}_2\text{H}_4\text{CH}(\text{CH}_3)_2$; 21. $R_2 = (\text{CH}_2)_5\text{CH}_3$

Reagents and conditions: (a) acetone, Jones Reagent, rt; (b) DMF, anhydrous K_2CO_3 , RBr, rt; (c) *t*-BuOH, *t*-BuOK, 50 °C; (d) anhydrous ethanol, $\text{C}_2\text{H}_8\text{N}_2$, Magnesium sulfate anhydrous, 80 °C; (e) 1, 2-Benzenediamine, anhydrous ethanol, reflux, 8 h.


Fig. 6 Synthesis of OA derivatives 12 to 21.

Nitric oxide (NO) is associated with several carcinogenic signaling pathways. Ling's group synthesized several hybrid compounds of OA and ivy saponin NO donors³⁸ (compounds 24–26, Fig. 7) and evaluated their cytotoxicity against various cancer cell lines. The most effective derivative, compound 25, significantly inhibited the proliferation of five cancer cell lines ($\text{IC}_{50} = 4.6\text{--}5.2 \mu\text{mol L}^{-1}$). Furthermore, compound 25 exhibited a stronger inhibitory effect on EGFR-LTC kinase activity ($\text{IC}_{50} = 0.01 \mu\text{mol L}^{-1}$) compared to ivy saponin ($\text{IC}_{50} > 20 \mu\text{mol L}^{-1}$) and inhibited the proliferation of gefitinib-resistant H1975 ($\text{IC}_{50} = 8.1 \mu\text{mol L}^{-1}$) and osimertinib-resistant H1975-LTC ($\text{IC}_{50} = 7.6 \mu\text{mol L}^{-1}$) non-small cell lung cancer (NSCLC) cells. This suggests that NO may synergistically act with compound 15

(Fig. 6) to induce the maximum production of NO in H1975 cells.

In 2018, Raghuvanshi *et al.*³⁹ synthesized a series of OA-based chromene derivatives and evaluated their anticancer activity. Among the derivatives with promising activity (compounds 27 and 28, Fig. 8), they further analyzed their ability to trigger apoptosis in A549 and MDA-MB-231 cells and found that these compounds caused cell cycle arrest in the G2/M phase. The most effective compound, 27 ($\text{IC}_{50} = 3.6 \mu\text{mol L}^{-1}$), exhibited anti-cancer activity through microtubule destabilization, and it was found to be non-toxic to human red blood cells.

Froelich *et al.*⁴⁰ introduced a lactam system into the pentacyclic triterpene structure of OA, facilitating the Beckmann



(a) Ac_2O , pyridine, rt, 12 h; (b) $\text{Br}(\text{CH}_2)_n\text{Br}$, K_2CO_3 , DMF, rt, 8 h; (c) AgNO_3 , CH_3CN , light-protected, reflux 3 h.

Fig. 7 Synthesis of OA derivatives 24 to 26.






Fig. 8 The structures of OA derivatives 27 and 28.




Fig. 9 Synthesis of OA derivatives 32 to 34.

rearrangement of the C-ring lactone to form OA derivatives (oxime 32, nitrile 33, and lactam 34, Fig. 9). Zaprutko *et al.*⁴¹ conducted tests on these derivatives as percutaneous permeation enhancers, finding that their activity was comparable to that of the positive control drug Azone.

3.1.3 Modifications of C-3, C-4 of the A ring and C-17 of oleanolic acid. Xu *et al.*⁴² discovered that methoxy substitutions in triterpene structures exhibit better anticancer activity compared to hydroxyl substitutions. Based on this finding, they modified the pentacyclic triterpene acid through esterification, using compound 35 as the ester moiety to synthesize a series of triterpene derivatives (compounds 36 and 37, Fig. 10), with modifications at the C-3 and C-4 positions of the A ring and the C-17 position of the five-membered ring. Activity assays indicated that most derivatives showed enhanced antifibrotic and antiproliferative effects. The increased lipophilicity improved the ability of the drugs to penetrate cell membranes, demonstrating that the introduction of ester moieties can enhance the anticancer activity of triterpenes. Furthermore, studies have shown that introducing amide bonds at the C-28 position and carbonyl groups at the C-11 position can also enhance the antitumor activity of triterpene acids.⁴³ Heise *et al.*⁴⁴ modified OA derivatives to obtain two structurally unique lactone

triterpenes (derivatives 38–40, Fig. 10) and a tri-hydroxy OA derivative (compound 40). Among these, derivative 38 exhibited no cytotoxicity against all tested cell lines, while derivative 39 showed a degree of selectivity against MCF-7 and FaDu cancer cells. The tri-hydroxy compound 40 demonstrated the strongest cytotoxic effect on MCF-7 cells, with toxicity approximately 5.5 times greater than that observed in non-malignant fibroblasts. This unique structural modification provides new avenues for enhancing the anticancer activity and selectivity of triterpenes.

3.1.4 Improvements and antitumor activity of CDDO based on oleanolic acid derivatives. Honda *et al.*⁴⁵ synthesized a derivative of OA over two decades ago, known as 2-cyano-3,12-dioxo-olean-1,9(11)-dien-28-oic acid (CDDO, Fig. 11). Their research revealed that CDDO not only prevents somatic cancers but also shows increasing evidence of efficacy against gender-specific cancers. CDDO and its derivatives are relatively non-toxic and serve as promising candidates for testing therapies for various diseases, demonstrating effectiveness against multiple types of cancer. The CDDO compound provides protection against ROS-induced cell death in normal cells by upregulating Nrf2, while simultaneously inducing death in cancerous or abnormal cells. In ovarian cancer models, CDDO directly targets and inhibits hsp90, reducing cell proliferation;






Fig. 10 The structures of triterpenic acid derivatives 35 to 40.




Fig. 11 Synthesis of OA derivative CDDO.

it also upregulates NQO1, which contributes to PARP1-mediated programmed necrosis in breast cancer, offering a potential therapeutic approach for this malignancy.

The most common modification site for CDDO is the C-28 carboxyl group. For instance, the introduction of a methyl ester at C-28 yields CDDO-Me (Fig. 12), which exhibits potent anti-inflammatory and anticancer activity.⁴⁶ CDDO-Me has shown inhibitory effects on prostate cancer cells, ovarian cancer cells, acute myeloid leukemia, pancreatic cancer cells, esophageal cancer cell lines, and triple-negative breast cancer cells. In ovarian cancer, CDDO-Me induces apoptosis by inhibiting AKT, NF- κ B, and mTOR signaling pathways, without affecting PDK1 kinase or PP2A activity. It induces prostate cancer cell death by inhibiting Akt, and apoptosis in acute myeloid leukemia by inhibiting ERK 1/2 phosphorylation through activation of p38/MAPK. CDDO-Me can also downregulate telomerase activity (hTERT) and induce cell death in pancreatic cancer cell lines.⁴⁷ Wong *et al.*⁴⁸ synthesized a novel CDDO-Me analog (compound

41, Fig. 12), which can irreversibly react with model thiols, aiding in the identification of pharmacologically relevant target interaction sites. Compound 41 exhibited cytotoxicity in rat liver cancer cells and comparable efficacy to CDDO-Me in activating Nrf2. Molecular modeling supported the stable modification of specific cysteine residues within Keap1 by compound 41. Gao *et al.*⁴⁹ explored whether CDDO-NFM (Fig. 12) has potential antitumor effects for cancer treatment, and they found that CDDO-NFM effectively inhibits OS cell growth, and this inhibitory effect is independent of apoptosis-related and cell cycle-related proteins.

Zhang *et al.*⁵⁰ designed and synthesized CDDO derivatives (compounds 42–44, Fig. 13). Their anticancer activity was further evaluated by measuring IC₅₀ values using the MTT assay. These compounds demonstrated potent inhibitory activity against both A549/Taxol (IC₅₀ = 0.349–1.087 $\mu\text{mol L}^{-1}$) and A549 (IC₅₀ = 1.066–1.438 $\mu\text{mol L}^{-1}$) cells, showing greater antiproliferative efficacy than CDDO-Me (IC₅₀ = 1.703 and 2.313





Fig. 12 The structures of OA derivatives CDDO-Me, CDDO-NFM, and 41.





(i) $(\text{COCl})_2$, anhydrous CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{r.t.}$, 12 h; (ii) anhydrous piperidine, TEA, anhydrous CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{r.t.}$, 12 h; (iii) Na_2CO_3 , DMF
(iv) N-Boc-protected L-amino acid, EDCI, DMAP, anhydrous, CH_2Cl_2 , 25°C ; (v) $\text{BF}_3 \cdot \text{Et}_2\text{O}$, anhydrous CH_2Cl_2 , 25°C .

Fig. 13 Synthesis of OA derivatives 42 to 44.

$\mu\text{mol L}^{-1}$) and JS-K ($\text{IC}_{50} = 1.987 \mu\text{mol L}^{-1}$ and $2.313 \mu\text{mol L}^{-1}$). The most active compound, **44**, induced higher levels of NO and reactive oxygen species (ROS) in paclitaxel-resistant A549/Taxol cells with overexpression of glutathione *S*-transferase π ($\text{GST}\pi$), significantly inhibiting cell proliferation ($\text{IC}_{50} = 0.349 \pm 0.051 \mu\text{mol L}^{-1}$). Moreover, compound **44** showed stronger inhibition of Lon protease expression, as well as more pronounced induction of apoptosis and cell cycle arrest in A549/Taxol cells, compared to CDDO-Me.

3.2 Structural modifications of oleanolic acid and antiviral activity

Yan *et al.*⁵¹ synthesized a series of OA derivatives by introducing conjugated dienes and utilizing photosensitized oxidation to form epoxides. The cytotoxicity of these derivatives was

evaluated in HepG 2.2.15 cells using the standard MTS assay. Most of the derivatives exhibited activity in inhibiting HBV antigen secretion in HepG 2.2.15 cells. Among the compounds tested, the most active derivative (compound **46**, Fig. 14) demonstrated significant activity in inhibiting the secretion of HBsAg, HBeAg, and the replication of HBV DNA. It was more effective than lamivudine in preventing the rebound of viral replication.

Li *et al.*⁵² introduced amino conjugates at the C-28 position of OA (compounds **48–52**, Fig. 15) and evaluated their antiviral activity against influenza A virus A/WSN/33 (H1N1) in Madin-Darby Canine Kidney (MDCK) cells. Based on the biological results, the structure–activity relationship of the compounds was discussed. Compound **52**, which contains a six-carbon chain and a terminal hydroxyl group, exhibited the strongest anti-influenza activity, with an IC_{50} of $2.98 \mu\text{mol L}^{-1}$. Hemagglutination




Fig. 14 Synthesis of OA derivative 46.




Fig. 15 Synthesis of OA derivatives 48 to 52.






Fig. 16 Synthesis of OA derivative 53 and derivatives 54a to 54c.

inhibition and surface plasmon resonance (SPR) analysis suggested that compound 52 may interfere with viral invasion by interacting with the HA protein of the influenza virus.

In 2019, Yang's team⁵³ designed and synthesized 32 derivatives by coupling various amino acids with the 28-COOH of OA. The most active derivative (compound 53, Fig. 16) demonstrated robust efficacy and a broad antiviral spectrum against four different influenza strains. Hemagglutination inhibition (HI) assays and docking studies suggested that these derivatives likely share the same mechanism as their parent compound, blocking viral entry by binding to hemagglutinin (HA) and preventing the interaction between the viral HA protein and the sialic acid receptors on host cells. In the same year, Medina-O'Donnell *et al.*⁵⁴ esterified one or two amino acids at the C-28 position and substituted phthaloyl groups at the C-3 position to obtain three 3-phthaloyl derivatives (compounds 54a–54c, Fig. 16), which showed enhanced inhibition of HIV-1 protease activity.

4 Structural modifications and biological activities of glycyrrhetic acid

Glycyrrhiza, derived from the dried roots and rhizomes of *Glycyrrhiza uralensis*, *Glycyrrhiza inflata*, or *Glycyrrhiza glabra*, is a valuable traditional Chinese medicine. The active constituents found in its dried roots and rhizomes primarily include flavonoids such as glycyrrhizin, GA, liquiritin, and isoliquiritin. Current research on the structural modifications of GA mainly focuses on the hydroxyl group at C-3, the carboxyl group at C-30, and the carbonyl group at C-11. Through modifications of the GA skeleton, a series of derivatives with physiological activities have been developed.

For instance, after replacing the hydroxyl group at the C-3 position of GA, the anti-inflammatory and anti-ulcer activities were significantly enhanced. When acyl groups, glycosides, or alkoxy and glycosyl groups were introduced at the C-3 hydroxyl or C-30 carboxyl positions, the anti-ulcer and anti-inflammatory activities were improved. Introducing substituents at both the C-3 hydroxyl and C-30 carboxyl positions further increased anti-inflammatory and antiviral activities. Additionally, the removal of the hydroxyl group at the C-11 position of GA has been shown to reduce or eliminate side effects and enhance pharmacological activity.

4.1 Structural modifications of the C-30 carboxyl group of glycyrrhetic acid

Wang *et al.*^{55,56} utilized addition and elimination reactions to modify substituted benzaldehyde, yielding substituted methylamine intermediates. These intermediates were then reacted with GA and deoxyglycyrrhetic acid (DGA), respectively, to produce GA and DGA derivatives at the C-30 position with amide bonds, forming heterocyclic derivatives (compound 55, Fig. 17).

The activity testing results for these compounds indicated that they exhibited high analgesic, anti-inflammatory, and antitussive activities, along with increased safety. The findings also demonstrated that 1-(3-methoxyphenyl)-5-aminomethyl-2-isoxazole DGA possesses certain hepatoprotective effects, effectively reducing the mortality rate of mice infected with the H1N1 influenza virus at a dose of 100 mg kg⁻¹. Lee *et al.*⁵⁷ synthesized 11 GA derivatives by introducing dehydrogingerol at the C-30 position. The activity tests revealed that these GA derivatives significantly inhibited tumor cell replication and exhibited anticancer effects. Some derivatives displayed strong cytotoxic activity; however, substituting the methoxy groups in compounds 56 and 57 with ethoxy groups resulted in a reduction in the cytotoxic activity of compounds 60 and 61 (Fig. 18).

Zhang Yan *et al.*⁵⁸ synthesized GA methyl ester and GA ethyl ester through esterification reactions between GA and methanol or ethanol. GA and its derivatives reacted with peracetic acid, resulting in the oxidation of the C-3 hydroxyl group to a carbonyl group, yielding oxidized GA and its methyl and ethyl ester derivatives. The catalytic asymmetric epoxidation activity of these compounds was tested, revealing the following activity order: oxidized glycyrrhetic acid (OGA) > oxidized





Fig. 17 The structure of isoxazole heterocyclic derivatives 55 of glycyrrhetic acid and dehydroglycyrrhetic acid.





- | | |
|--|---|
| 56. 4-OGA R ₁ =3-OCH ₃ R ₂ =CH ₃ | 62. 2-OGA R ₁ =3-F R ₂ =CH ₃ |
| 57. 2-OGA R ₁ =3-OCH ₃ R ₂ =CH ₃ | 63. 4-OGA R ₁ =3-OCH ₃ R ₂ =Ph |
| 58. 3-OGA R ₁ =4-OCH ₃ R ₂ =CH ₃ | 64. 2-OGA R ₁ =3-OCH ₃ R ₂ =Ph |
| 59. 2-OGA R ₁ =3-OCH ₃ R ₂ =CH ₃ | 65. 3-OGA R ₁ =4-OCH ₃ R ₂ =Ph |
| 60. 4-OGA R ₁ =3-OC ₂ H ₅ R ₂ =CH ₃ | 66. 2-OGA R ₁ =4-OCH ₃ R ₂ =Ph |
| 61. 2-OGA R ₁ =3-OC ₂ H ₅ R ₂ =CH ₃ | |

Fig. 18 The structures of dehydrogingdiolide derivatives of glycyrrhizic acid **56** to **66**.

glycyrrhetic acid derivatives > GA derivatives. Furthermore, GA derivatives were found to be ineffective in catalyzing the reactions of sterically hindered alkenes.

4.2 Structural modifications of C-3 hydroxyl and C-30 carboxyl groups of glycyrrhetic acid

Zhang Na *et al.*⁵⁹ introduced an acetyl group at the C-3 position through an acetylation reaction to synthesize compound **67**. Subsequently, under the reaction conditions of (COCl)₂,

compounds **56** and **57** (Fig. 18) were reacted to obtain compounds **68** and **69**. Using berberine, 18 α -GA, and methotrexate as controls, it was found that compounds a and b exhibited significant inhibitory effects on the liver cancer cell line SMMC-7721, with IC₅₀ values of 85.7 $\mu\text{mol L}^{-1}$ and 178.2 $\mu\text{mol L}^{-1}$, respectively. Animal experiments indicated that these compounds were almost non-toxic to normal liver cells (Fig. 19).

Csuk *et al.*⁶⁰ modified GA methyl ester (GA-Me) by reacting it with chloroacetyl chloride, followed by a condensation reaction with various diamines to introduce straight-chain amino groups at the C-3 hydroxyl position, resulting in a series of GA derivatives **70**. The results of antitumor activity testing demonstrated that the introduction of straight-chain amino groups at the C-3 position enhanced the activity of GA, with the highest activity observed when the carbon chain length $n = 6$, yielding an IC₅₀ range of 0.6–3.0 $\mu\text{mol L}^{-1}$ (Fig. 20).

Schwarz *et al.*⁶¹ synthesized GA derivatives by introducing methoxy, ethoxy, and benzyloxy groups at the C-30 position through esterification reactions. They also introduced different configurations of alanine at the C-3 position, resulting in a series of C-3 and C-30 GA derivatives **71**. The activity and selectivity of these derivatives against tumor cells were found to be superior to those of the control group, which consisted of GA (Fig. 21).

Gao *et al.*⁶² simultaneously modified the C-30 carboxyl and C-3 hydroxyl groups of GA through oxidation, esterification, substitution, and hydrolysis reactions to obtain compound **72**.




Fig. 19 Synthesis of 3-acetyl glycyrrhizic acid lipid derivatives **68** to **69**.




Fig. 20 Synthesis of 3-O-(linear aminocarbonyl) glycyrrhizic acid lipid derivative **70**.






Fig. 21 Synthesis of 3-alanine type glycyrrhizic acid lipid derivative 71.

Compound 72 underwent nucleophilic substitution to introduce various *N*-substituents at the carboxyl group, yielding amide derivatives 73a–73c. Compound 2 was synthesized by forming a ring between C-3 and C-2, followed by nucleophilic substitution to obtain compounds 74a–74d. Subsequently, ring-opening elimination and reduction led to various structural derivatives of oxidized GA amides, including 75a–75d and 76a–76d. Activity studies indicated that these compounds exhibited strong inhibitory effects on the growth of HL-60 leukemic cells (Fig. 22).

4.3 Structural modification of the C-11 carbonyl group of glycyrrhetic acid

Meng Yanqiu *et al.*⁶³ reduced the C-11 carbonyl group of GA to obtain DGA (compound 77) using zinc–mercury amalgam.

Subsequently, they introduced an ester bond at the C-30 position through an esterification reaction. The reaction with *p*-toluenesulfonyl chloride in a dichloromethane and triethylamine medium at low temperatures facilitated the introduction of a *p*-toluenesulfonyl group at the C-3 hydroxyl. Finally, refluxing with sodium azide in tetrahydrofuran yielded compounds 80a–80g.

In vitro antitumor activity tests, using paclitaxel and gefitinib as controls, revealed that compounds 80a, 80c, and 80e exhibited significant inhibitory effects on human breast cancer cells (MCF-7) and lung cancer cells (A549). At a concentration of $10 \mu\text{mol L}^{-1}$, the inhibition rates against MCF-7 were 71.57%, 14.53%, and 22.02%, respectively. Notably, compound 80a demonstrated a significantly enhanced inhibitory capacity against breast cancer cells compared to glycyrrhetic acid, with




Fig. 22 Synthetic route of oxidized glycyrrhizic acid amide derivatives.







Fig. 23 Synthesis route of 3-azido glycyrrhizic acid derivatives.

Fig. 24 The synthesis route of 11-position modified glycyrrhizic acid derivatives **81** and **82**.


an IC₅₀ of 4.29 ± 0.07 μmol L⁻¹. The inhibition rates against A549 cells at 10 μmol L⁻¹ were 4.91%, 26.60%, and 19.97%, all exceeding that of glycyrrhetic acid. The activity testing results of compounds **80a**, **80c**, and **80e** further indicated that the different ester bonds at the C-30 position significantly influenced the inhibition of MCF-7 cells (Fig. 23).

Su *et al.*⁶⁴ modified the carbonyl group at the C-11 position and the carboxyl group at the C-30 position of GA, yielding two GA derivatives, **81** and **82**. These derivatives exhibited strong inhibitory activity against 11β-HSD1, with IC₅₀ values of 0.4 μmol L⁻¹ and 1.1 μmol L⁻¹, respectively. Notably, the presence of the carboxyl group at the C-30 position contributed to the inhibition of 11β-HSD1 activity (Fig. 24).

5 Betulinic acid: structural modifications and biological activity

5.1 Structural modifications of betulinic acid and antitumor activity

5.1.1 Structural modifications at C-3 and C-28 of betulinic acid. Research has indicated that derivatives with modifications at the C-3 and C-28 positions show promising potential. The introduction of amino acid conjugates at the C-28 position enhances solubility and cytotoxicity, while the incorporation of

Fig. 25 Synthesis of BA derivatives **85** and **86**.




Fig. 26 Synthesis of BA derivative 89.

polar groups such as acids, amines, or hydroxyls increases biological activity.⁶⁵ Hydroxylation at the C-3 position has yielded encouraging results against murine melanoma cells. Additionally, another chemical modification at C-3, namely dimethylsuccinoyl BA, has transformed BA from a proteasome activator to a proteasome inhibitor. BA derivatives substituted with new carbamate derivatives at C-3 and C-28 (compounds 85 and 86, Fig. 25) exhibit strong cytotoxicity and can induce apoptosis in lung tumor cell lines. Furthermore, compounds 85 and 86 demonstrate higher selectivity for tumor cells compared to normal cells.⁶⁶

Spivak *et al.*³³ designed and synthesized guanidine-containing BA derivatives (compound 89, Fig. 26), which

exhibited higher cytotoxicity in Jurkat cells compared to the original triterpenic acid. Most of the tested guanidine derivatives showed higher IC₅₀ values than amines, while demonstrating lower toxicity to human fibroblasts.

5.1.2 Structural modifications of betulinic acid in the A ring. Chatterjee *et al.*⁶⁷ investigated the metabolic conversion of BA using *Bacillus subtilis* to biosynthesize derivative 90 (Fig. 27). Compared to the activity of BA (ED₅₀ ranging from 2.1 μmol L⁻¹ to 7.0 μmol L⁻¹), compound 90 exhibited cytotoxic activity against human melanoma cell lines (Mel-1 and Mel-2) in the micromolar range (ED₅₀ < 40 μmol L⁻¹).

In the experiment by Mukherjee *et al.*,⁶⁸ BA was oxidized with Jones reagent to form its 3-oxo derivative, BA ketone (compound




Fig. 27 Synthesis of BA derivative 90.




Fig. 28 Synthesis of BA derivatives 91 to 93.



91, Fig. 28), which then reacted with phenylhydrazine and sodium acetate to yield the 3-phenylhydrazone (compound **92**, Fig. 28). The results from anti-angiogenesis screening indicated that derivative **91** exhibited lower cytotoxicity compared to **BA** (with an IC_{50} of $4.4 \mu\text{mol L}^{-1}$ against the ECV304 endothelial cell line) and comparable low endothelial cell specificity (with an EC_{50} of $1.9\text{--}4.2 \mu\text{mol L}^{-1}$ across four cancer cell lines). Compound **91** was then reacted with isopropenyl acetate and *p*-toluenesulfonyl chloride under reflux for 9 hours, resulting in the enol acetate (compound **93**, Fig. 28). Compound **93** was screened against a panel of five cancer cell lines and demonstrated low cytotoxic activity (IC_{50} ranging from $40 \mu\text{mol L}^{-1}$ to $100 \mu\text{mol L}^{-1}$).

Li *et al.*⁶⁹ synthesized difluorinated derivatives (compounds **96** and **97**, Fig. 29) starting from dihydrobetulinic acid (DHBA) through a five-step or six-step reaction pathway. They investigated the cytotoxic activity of all compounds on the CCRF-CEM cell line. Compared to DHBA, the difluorinated derivatives **96** and **97** exhibited higher cytotoxicity, with IC_{50} values of $2.2 \mu\text{mol L}^{-1}$ and $4.0 \mu\text{mol L}^{-1}$, respectively.

5.1.3 Structural modifications of betulinic acid at the C-3 and C-28 positions with phosphorus. Tsepaveva *et al.*⁷⁰ used **BA** as the starting material and, in the presence of K_2CO_3 , reacted it with phosphoric acid esters in DMF solvent to obtain derivative **98** (Fig. 30). The compounds containing triphenylphosphonium cations are promising ionic molecules for the targeted delivery of neutral bioactive compounds to the mitochondria of cancer

cells, including anticancer agents that possess pro-oxidant properties and destabilize the mitochondrial membrane.

In the study by Spivak *et al.*,⁷¹ **BA** derivatives (compounds **99**–**104**, Fig. 31) containing triphenylphosphonium cations featured various functional groups at the C-3 and C-28 positions ($3\beta\text{-OAc}$, $3\alpha\text{-OAc}$, $3\beta\text{-OH}$, $3\alpha\text{-OH}$, $3\beta\text{-O-phthalate}$, 28-COOMe , 28-COOH). Despite the differences in their chemical structures, the cytotoxic effects exhibited against EAC cells and mastocytoma P-815 cells were not significantly different, with IC_{50} values in Ehrlich cells ranging from 1.2 to $1.9 \mu\text{mol L}^{-1}$ and in P-815 cells from 0.9 to $1.1 \mu\text{mol L}^{-1}$. However, the benzyl ester of DHBA (compound **99**) demonstrated reduced antitumor activity against both cell types, with IC_{50} values of $4.80 \mu\text{mol L}^{-1}$ for Ehrlich cells and $4.89 \mu\text{mol L}^{-1}$ for P-815 cells.

5.2 Structural modifications of betulinic acid and their antiviral activity against HIV-1

Chen *et al.*⁷² designed and synthesized a series of C-28 amine derivatives of **BA** (**107**) based on C-3 benzoyl-substituted **BA** derivatives (**105**, **106**) as HIV-1 inhibitors. Compared to the C-28 amide series, the C-28 amine derivatives (**107**) showed enhanced inhibitory activity against HIV-1, specifically targeting Gag polyprotein polymorphisms. Their activity was further improved in the presence of human serum, even though basic amines generally exhibit lower plasma exposure after oral administration in rats. Among the C-28 amine series, the




Fig. 29 Synthesis of BA derivatives **96** and **97**.




Fig. 30 Synthesis of BA derivative **98**.






Fig. 31 The structures of BA derivatives 99 to 104.




Fig. 32 Synthesis of BA derivative 108.

derivative with the best antiviral activity and acceptable pharmacokinetic properties (compound **108**, Fig. 32) demonstrated a 2- to 4-fold improvement in antiviral potency compared to the C-28 amide. It exhibited low EC_{50} shifts in the presence of human serum or human serum albumin against the V370A and DV370 viral strains and showed promising efficacy against polymorphic variants T371A and V362L.

Reutrakul *et al.*⁷³ isolated two partially acetylated 2 α -hydroxy BA derivatives (compounds **109** and **110**, Fig. 33) from the leaves, branches, and resin of *Garcinia* species. Both acetylated derivatives, **109** and **110**, exhibited anti-HIV-1 activity in HIV-1 reverse transcriptase and syncytium formation assays. In the syncytium formation assay, compounds **109** and **110**




Fig. 33 Structure of BA derivatives 109 and 110.

demonstrated efficacy with IC_{50} values of $30.9 \mu mol L^{-1}$ and $38.6 \mu mol L^{-1}$, respectively, which is 2–3 times more potent than the activity of BA.

6 Structural modifications of ursolic acid and its biological activities

Ursolic acid (UA), also known as UA or ursolic acid, is a representative compound of the ursane-type pentacyclic triterpenoids. It is widely distributed in many plants, including medicinal herbs and edible plants such as the leaves of Rosaceae plants like loquat and species from the Ericaceae family. Significant progress has been made in recent years in studying the antitumor activities of UA. Li *et al.*⁷⁴ found that UA can exert anticancer effects by inhibiting cancer initiation and progression as well as DNA mutations. Young *et al.*⁷⁵ reported that UA can induce the differentiation of teratoma stem cells, demonstrating its anti-carcinogenic properties, and has been recognized as one of the most promising natural compounds for cancer prevention. Additionally, UA shows potential therapeutic applications for prostate cancer by controlling the proliferation and apoptosis of PC-3 and LNCaP cells.⁷⁶ UA exerts its



anticancer activity by inhibiting kinases involved in cancer cell signaling, showing antitumor effects in various cell types.

6.1 Structural modifications of ursolic acid and antitumor activity

6.1.1 Structural modifications of ursolic acid at the C-3 and C-28 positions.

Zhang Dajun *et al.*⁷⁷ synthesized benzyl phosphonic acid diethyl ester acetamide derivatives of UA (compounds **111–115**, Fig. 34) through a substitution reaction between UA and various substituted diethyl benzyl phosphonic acids. The *in vitro* anti-hepatocellular activity of the target compounds was evaluated using the MTT assay. All target compounds exhibited low cytotoxicity against normal human liver cells (L02), while compound **115** demonstrated significant antiproliferative activity against HepG2 cells, with an IC_{50} value of $2.69 \mu\text{mol L}^{-1}$, showing good selectivity between tumor cells and normal cells.

Spivak *et al.*³³ designed and synthesized guanidine-containing UA derivatives (compound **118**, Fig. 35). Compared to the original UA, these guanidine-containing derivatives exhibited lower toxicity toward human fibroblasts while demonstrating higher cytotoxicity in Jurkat cells. Most of the guanidine derivatives tested showed higher IC_{50} values compared to their amine counterparts.

Tian *et al.*⁷⁸ designed and synthesized a series of novel nitrogen-containing UA derivatives by introducing diamine fragments at the C-3 and C-28 positions. The derivatives with the highest activity (compounds **120** and **122**, Fig. 36) were tested for their cytotoxicity against MCF-7, HeLa, and A549 cell

lines. UA-4a and UA-8a showed significant inhibitory activity against all three human cancer cell lines, with IC_{50} values of less than $10 \mu\text{mol L}^{-1}$. Analyzing the antitumor activity of the synthesized compounds revealed no significant difference in activity between the introduction of diamine compounds at the C-3 and C-28 positions, both displaying strong antitumor properties.

Furthermore, analysis of the synthesized series suggested that the carbon chain length of the diamine compounds had little impact on antitumor activity, and no linear correlation was observed. However, a slight decrease in activity was noted when the carbon chain length reached six carbons. Considering factors such as ease of isolation, purification, yield, and activity during synthesis, the diamine compounds with a two-carbon chain were deemed the most valuable for further research. These compounds were characterized by low synthesis cost, easy purification, simple synthetic steps, and high cytotoxicity against cancer cell lines, making them promising candidates for further study as novel antitumor drugs. Additionally, they provide the potential for further salt formation modifications.

Bai *et al.*^{79–81} conducted comprehensive studies on modifications at the C-3 and C-28 positions of UA. The introduction of hydrophilic polyethylene glycol (PEG) significantly enhanced the solubility of the drug. Conjugating PEG at the C-28 position of the UA scaffold notably improved the *in vitro* antitumor activity. For example, compound **123** (Fig. 37) exhibited a solubility between 0.01 g mL^{-1} and 0.033 g mL^{-1} in water and demonstrated potent anticancer activity with IC_{50} values below $10 \mu\text{mol L}^{-1}$ against five tumor cell lines: HepG2, BGC-823, AGS, HT-29, and PC-3.





Fig. 34 Synthesis of UA derivatives **111–115**.



- (a) 1. (COCl)₂, CH₂Cl₂; 2. Tris(hydroxymethyl)aminomethane, DMAP, Py, DCM;
(b) 1,3-Di-Boc-2-(trifluoromethylsulfonyl)guanidine; Et₃N, CHCl₃, reflux;
(c) 50% TFA, CH₂Cl₂, 2–4 h, rt; (d) 5M HCl, MeOH.

Fig. 35 Synthesis of UA derivative **118**.






Fig. 36 Synthesis of UA derivatives 120 and 122.




Fig. 37 Structure of UA derivatives 123 to 125.

Furthermore, the attachment of groups with varying electro-negativities to UA effectively improved its water solubility, particularly the positively charged UA derivatives, which exhibited significantly enhanced anticancer activity. Compound 124 (Fig. 37) displayed a 70-fold increase in water solubility and an IC_{50} value below $10 \mu\text{mol L}^{-1}$ against PC-3 cancer cells. Additionally, it was found that introducing amino acid derivatives with primary amino groups at the C-28 position of UA substantially enhanced its anticancer activity compared to UA alone, with the mechanism of action involving the induction of apoptosis.

Zhao *et al.*⁸² carried out structural modifications on the A and C rings of UA, as exemplified by compound 125 (Fig. 37). These modifications involved multiple oxygen substitutions on the A ring and oxidation at the 11th position of the C ring. The resulting compounds exhibited cytotoxicity against cancer cells such as KB and PC-3, comparable to cisplatin, and demonstrated an inhibitory effect on α -glucosidase that was an order of magnitude stronger than that of acarbose.

6.1.2 Structural modifications of ursolic acid at the C-2 and C-3 positions of the A ring. Research on the structural modification of UA at the C-2 and C-3 positions of the A ring has provided insights into enhancing its biological activity.


Modifications in these positions have been shown to impact the compound's pharmacological properties, particularly its anti-cancer effects. Wu *et al.*⁸³ designed and synthesized a series of UA derivatives containing an aminoguanidine structure and evaluated their activity as HIF-1 α inhibitors and anticancer agents. Using a luciferase reporter gene assay based on Hep3B cells, most compounds exhibited good HIF-1 α inhibitory activity. Among them, the most active derivative (compound 128, Fig. 38) demonstrated the strongest HIF-1 α inhibition under hypoxic conditions, with an IC_{50} value of $4.0 \mu\text{mol L}^{-1}$, and showed no significant cytotoxicity in the tested cells. The mechanism of action may involve the downregulation of HIF-1 α protein expression through inhibition of HIF-1 α protein synthesis, as indicated by western blotting, RT-PCR assays, and cell aggregation experiments. This reduction in HIF-1 α expression subsequently leads to decreased production of vascular endothelial growth factor (VEGF), thereby inhibiting cancer cell proliferation.

Wang *et al.*⁸⁴ synthesized derivatives of UA with a furan moiety at the C-2 and C-3 positions based on the method of Khan (compound 130, Fig. 31). These derivatives were further modified by conjugation with ethyl-piperazine to obtain piperidine-containing derivatives (compound 131, Fig. 39). The anti-proliferative activities of the target compounds against the HeLa and MKN45 cancer cell lines were evaluated using the MTT assay, with cisplatin and UA used as positive controls. Derivative 130 exhibited significantly higher activity than UA, with an IC_{50} value below $10 \mu\text{mol L}^{-1}$, while derivative 131 displayed moderate to low activity against both cancer cell lines. Structure–activity relationship studies indicated that the introduction of thiazole on the A ring and the incorporation of triazole or tetrazole structures at C-28 had minimal impact on enhancing antitumor activity. However, the inclusion of piperazine or higher piperazine derivatives significantly improved the antitumor activity.

6.2 Structural modifications of ursolic acid and antiviral activity


The structural modifications of UA have been investigated to enhance its antiviral activity. Various derivatives have been





(a) Jones Reagent, acetone, 0 °C; (b) Formaldehyde, 5%NaOH, Anhydrous Ethanol, rt, 2 h;
(c) ConcHCl, Anhydrous Ethanol, reflux, 8 h.

Fig. 38 Synthesis of UA derivative 128.



(a) Isoamyl nitrite, t-BuOK, t-BuOH, rt, 2.5 h; (b) Hydroxylamine hydrochloride, Pyridine, 110 °C, 2 h; (c) EtOH, NaOH, NaOCl, rt, 2 min;
(d) 2-chloroethanol, K₂CO₃, DMF, 60 °C, 3 h (e) MsCl, Pyridine, 10 °C~rt, 3 h; (f) Piperazine, K₂CO₃, DMF, 60 °C, 30 min.

Fig. 39 Synthesis of UA derivatives 130 and 131.

synthesized to explore their potential effects against different viral strains. Min *et al.*⁸⁵ reported that the 3-*O*-acyl derivatives of ursolic acid (compounds 132–142, Fig. 40) exhibited significant inhibitory effects on HIV replication in human T lymphocytes (H9 cells), with IC₅₀ values of 0.31 and 2.1 μmol L⁻¹, respectively. Compared to the lead compound ursolic acid (IC₅₀ value of 4.4 μmol L⁻¹), these two compounds demonstrated improved anti-HIV activity and safety profiles. They further reported that a series of anhydrides were introduced at the C-3 position of UA to form corresponding dicarboxylic acid

monoesters, evaluating the impact of the length of the anhydride chain on anti-HIV activity. Analysis of the IC₅₀ values revealed that, when introducing anhydride chains containing up to six carbons, the anti-HIV activity of the compounds increased with the number of carbon atoms. However, when two methylene units were added, a slight decrease in activity was observed. Additionally, when the carboxyl groups of the anhydride formed carboxylic acid methyl esters, the anti-HIV activity of the compounds decreased, and in some cases, was completely lost.




Fig. 40 Structure of UA derivatives 132 to 142.






Fig. 41 Synthesis of ursolic acid 3- β ester derivatives (143a–143o).

6.3 Ursolic acid 3- β ester derivatives and corresponding activity evaluation

UA has garnered attention for its versatile pharmacological properties, and the development of 3- β ester derivatives is a promising approach to enhance its biological activity. Various studies have explored the synthesis of these ester derivatives and evaluated their potential in different therapeutic areas.

For instance, Chen *et al.*⁸⁶ synthesized a series of UA 3- β ester derivatives (143a–143o, Fig. 41). Activity assays indicated that compounds 143a and 143b, which contain phthalate esters or isophthalate esters, exhibited moderate inhibitory effects on cholesterol ester transfer protein (CETP), while the inhibitory effect of derivative 143c, containing terephthalate esters, was completely lost. Further comparison of derivatives 143b, 143d, and 143e revealed that the carboxyl group on the backbone chain is essential for enhancing inhibitory activity. In contrast to derivative 143g, the derivatives containing succinate esters (143j and 143k) lost their inhibitory effects on CETP, possibly due to steric hindrance from the methyl group. Derivative 143l, which contains 1,2-cyclopropanedicarboxylate esters, exhibited the strongest CETP inhibitory activity, with an IC_{50} of 2.4 $\mu\text{mol L}^{-1}$. The presence of a carboxyl group at the end of the backbone chain and the length of the backbone chain are critical factors for maintaining or enhancing CETP inhibitory activity. These types of 3- β ester triterpenoid derivatives (*e.g.*, derivative 143l) hold promise as lead compounds with potent CETP enzyme inhibitory activity. Xu *et al.*⁸⁷ found that glycosylated corosolic acid derivatives possess certain α -glucosidase inhibitory activity and improved water solubility, which is associated with the presence of hydroxyl groups at the C-2 and C-3 positions. This activity is also closely related to the substituents at the C-28 position.^{88,89}

7 Structural modifications and pharmacological activities of celastrol

Celastrol, also known as tripterine, is a triterpenoid compound belonging to the oleanane-type pentacyclic triterpenoids. It was

first extracted from the traditional Chinese medicinal herb *Tripterygium wilfordii* by the renowned organic chemist Zhao Chenggu in 1936.⁹⁰ As the first active compound isolated from the root bark of *Tripterygium wilfordii*, celastrol is the most abundant bioactive component in this plant and has been demonstrated to exhibit potent anticancer activity.⁹¹ Compared to some conventional chemotherapeutic agents, celastrol shows similar or even superior anticancer effects, particularly against multidrug-resistant cancer cells. In 2007, it was listed by *Cell* magazine as one of the five natural products most likely to be developed into modern pharmaceuticals.⁹² Despite extensive research into its potential anticancer properties, its further application remains limited due to challenges such as poor water solubility, low bioavailability, and high toxicity.⁹³

7.1 Structural modifications of celastrol at the C-29 position and its anticancer activity

Feng *et al.*⁹⁴ modified the hydroxyl group at C-3 and the carboxyl group at C-29 of celastrol to synthesize a series of 1,2,3-triazole derivatives of celastrol (compound 145, Fig. 42). They evaluated the anti-glioma activity of these derivatives *in vitro* against four human glioma cell lines (A172, LN229, U87, and U251). Structure–activity relationship studies indicated that the modifications at C-3 and C-29 significantly enhanced the anti-proliferative activity compared to celastrol, with the C-29 carboxyl group derivatives exhibiting stronger anti-proliferative activity than the C-3 hydroxyl group derivatives across the four human glioma cell lines. Among these, compound 145 demonstrated the highest anti-proliferative activity in the U251 cell line, with an IC_{50} of 0.94 $\mu\text{mol L}^{-1}$. Further studies revealed that its effective anti-glioma activity is associated with the activation of the receptor–interacting protein 1/receptor–interacting protein 3/mixed lineage kinase domain-like protein (RIPI/RIP3/MLKL) pathway.

Ferulic acid (FA) is a type of phenolic acid that has been demonstrated to possess various pharmacological activities, including antioxidant, anti-diabetic, anti-inflammatory, and neuroprotective effects. FA and its derivatives have also shown






Fig. 42 Synthesis of celastrol derivative 145.

significant anticancer efficacy both *in vitro* and *in vivo*.⁹⁵ Coghi *et al.*⁹⁶ designed and synthesized a series of novel hybrids of celastrol and ferulic acid, evaluating their antitumor activity. Structure–activity relationship (SAR) studies indicated that different linkers influenced the pharmacological effects, with a two-carbon linker exhibiting stronger anticancer capabilities compared to a three-carbon linker. Moreover, when a methoxy group was substituted on the FA moiety, the hybrids displayed improved efficacy. Among these, the most active derivative (compound **146**, Fig. 43) exhibited the strongest inhibitory capacity. Mechanistic studies revealed that compound **146** could disrupt the heat shock protein Hsp90–Cdc37 complex while also inducing abnormal regulation of heat shock protein (Hsp90) receptors (*p*-Akt and Cdk4), leading to concentration-dependent cell cycle arrest in the G0/G1 phase.

Disrupting the interaction between Hsp90 and Cdc37 has emerged as a promising strategy for investigating the antitumor activity of celastrol.⁹⁷ To further explore the effects of celastrol C-

29 derivatives on the Hsp90–Cdc37 complex, Li *et al.*⁹⁸ introduced several lipophilic fragments into celastrol C-29, synthesizing 48 new celastrol derivatives and testing their effects on the Hsp90–Cdc37 complex and antitumor cell proliferation activity. SAR studies revealed that the length of the linker significantly influenced the activity of the compounds. Using two-carbon and three-carbon linkers resulted in much higher inhibition rates on tumor cells compared to four-carbon linkers, with 8-cyanide groups exhibiting better efficacy than α -hydroxyl groups. Among the three-amine substitutions, introducing electron-donating groups onto the phenyl ring enhanced the antiproliferative activity of the compounds. One particularly active derivative (compound **147**, Fig. 44) demonstrated excellent antiproliferative activity against relevant cancer cells (IC_{50} : 0.41–0.94 $\mu\text{mol L}^{-1}$). Further mechanistic studies indicated that compound **147** could reduce the expression of the Hsp90–Cdc37 complex in A549 cells and form a complex with Cdc37 in tumor cells, thereby disrupting the Hsp90–Cdc37 interaction.




Fig. 43 Synthesis of celastrol derivative 146.




Fig. 44 Synthesis of celastrol derivative 147.






Fig. 45 Synthesis of celastrol derivative 148.

CSCs are associated with cancer progression, tumor recurrence, metastasis, and chemotherapy resistance. To develop effective and selective anti-CSC agents, Li *et al.*⁹⁹ synthesized two series of celastrol derivatives with cinnamoylamine linkers, utilizing ethylenediamine and piperazine as connecting chains. Their antitumor activity against ovarian cancer was evaluated. SAR analysis revealed that most compounds exhibited stronger antiproliferative activity compared to celastrol. Additionally, no significant differences in activity were observed between the two series of compounds with different linker chains. However, the celastrol derivative bearing a 3,4,5-trimethoxy cinnamoylamine side chain (compound 148, Fig. 45) demonstrated the strongest antiproliferative effect against ovarian cancer cells ($IC_{50} = 0.6 \mu\text{mol L}^{-1}$). Further pharmacological tests indicated that compound 148 significantly inhibited the colony formation ability of tumor cells, reduced the number of tumor spheres, and decreased the percentage of ovarian CSCs.

Nitric oxide (NO) is a key signaling and effector molecule in tumor development. Generally, high concentrations of NO can induce apoptosis in tumor cells, while low concentrations typically offer cellular protection. Based on this, Tang *et al.*¹⁰⁰ synthesized a series of celastrol nitric oxide donor drugs and evaluated their NO release capabilities and antiproliferative activities. The results showed that the most active derivative (compound 149, Fig. 46) released the highest levels of NO *in vitro* and exhibited greater antiproliferative activity compared to other compounds (A549: $IC_{50} = 0.48 \mu\text{mol L}^{-1}$). SAR analysis indicated that the C-29 amide bond was beneficial for enhancing antiproliferative activity. Furthermore, furoxan nitrogen oxides demonstrated stronger antiproliferative activity as NO-releasing components than nitrate esters. Additional pharmacological studies revealed that compound 149 could induce dysregulation of the Hsp90 receptor, apoptosis, and cell cycle arrest at the G0/

G1 phase. These findings suggest that the inhibition of Hsp90 and NO release have a synergistic effect in tumor cells, providing new directions for the discovery of antitumor drugs.

7.2 A/B ring-modified celastrol derivatives and their anticancer activity

In the exploration of the anticancer activity of celastrol, the modification of its A/B rings has become an important strategy. By chemically modifying the A/B rings of celastrol, it is possible to enhance its biological activity and improve its drug properties. Tang *et al.*¹⁰¹ synthesized a series of derivatives of celastrol with urea moieties as linkers at the C-29 position and modifications at the A/B rings, followed by testing their antitumor cell proliferation activity. SAR analysis revealed that the C-29 carboxy urea derivatives exhibited enhanced cytotoxicity against tumor cell lines compared to celastrol. However, modifications to the A/B rings resulted in diminished antiproliferative activity of the derivatives. This suggests that the intact quinone methide structure of the A/B rings and the reactive Michael addition site at position 6 may be essential groups for the anticancer effects of celastrol. Among these, the most active derivative (compound 150, Fig. 47) demonstrated superior antiproliferative activity with greater selectivity. Preliminary studies on its mechanism of action indicated that compound 150 induces apoptosis through the activation of cysteine protease-8 (caspase-8) and cysteine protease-3 (caspase-3), as well as the cleavage of poly (ADP-ribose) polymerase (PARP) *via* an extrinsic pathway. Additionally, it induced downregulation of p53 in the SKOV-3 cell line, which is a p53-mutant cell line.

7.3 C-6 modified celastrol derivatives and anticancer activity

To explore the impact of A-ring and C-6 modifications on the antitumor activity of celastrol, Tang *et al.*¹⁰² synthesized a series




Fig. 46 Synthesis of celastrol derivative 149.






Fig. 47 Synthesis of celastrol derivative 150.




Fig. 48 Synthesis of celastrol derivative 152.

of C-6 sulfonated, thiolated, and carbon-modified celastrol derivatives *via* a Michael addition reaction. The cytotoxicity of these derivatives was evaluated in human cancer cell lines (BGC-823, H4, Bel7402, H522, Colo205, HepG2, and MDA-MB468). SAR analysis showed that C-6 sulfonated and thiolated derivatives exhibited stronger cytotoxicity compared to carbon-modified derivatives at the C-6 position, with 3-hydroxy acetylation proving more effective than propionylation. Among these derivatives, compound 152 (Fig. 48) demonstrated the most potent bioactivity in the Colo205 cell line ($IC_{50} = 0.06 \mu\text{mol L}^{-1}$). Further experimental results revealed that, in nude mice bearing human colorectal cancer xenografts, compound 152 effectively inhibited the growth of Colo205 xenografts *in vivo* and exhibited better safety profiles compared to celastrol.

8 Conclusion

Natural products, as an important source of lead compounds for drug development, play a crucial role in pharmaceutical research. However, their inherent limitations, such as poor water solubility, low bioavailability, and significant toxicity, have led researchers to focus increasingly on structural modifications to obtain more efficient and less toxic new drugs. Among these, pentacyclic triterpenes have garnered growing attention due to their diverse biological activities, with research on these compounds becoming progressively more in-depth in recent years. Despite the structural diversity of pentacyclic triterpenes, their limited content in plants, complex extraction processes, and our insufficient understanding of their biosynthetic pathways, effective chemical modifications offer promising opportunities for their development. Structural modifications, particularly aimed at their hydrophobic framework, are essential for further exploration.

OA, BA and UA are pentacyclic triterpenes widely distributed in nature, known for their anti-inflammatory, antiviral, and anticancer properties, showcasing significant potential for therapeutic applications. However, due to their poor water solubility and low bioavailability, structural modifications can effectively improve these properties by enhancing water solubility, increasing bioactivity, and reducing toxic side effects. Although numerous bioactive derivatives have been developed, few have advanced to clinical application. Research into the structural modification of OA and BA has yielded significant progress in the field of anticancer therapy, identifying key targets and mechanisms of action. SAR studies have provided insights that will guide the design of more potent pentacyclic triterpene derivatives in the future.^{103–106} Additionally, as research continues, UA has also become a focal point for anticancer drug development through structural modification. The continued exploration of OA, BA, and UA derivatives for the development of highly effective, low-toxicity anticancer drugs remains a promising area of research.

The structural modification of celastrol has mainly focused on the C-29 position, with relatively little research on the 3-OH group and A/B rings. However, the high reactivity of the A/B rings may be crucial to celastrol's pharmacological effects. Therefore, further studies on the modification of the A/B rings are essential, and even ring-opening reactions could be explored to generate celastrol derivatives with reduced toxicity and enhanced efficacy. This could also lead to a more comprehensive understanding of the SAR of celastrol. Moreover, biotransformation could provide novel approaches for the structural modification of celastrol, potentially yielding derivatives that are difficult to obtain through chemical methods, thus offering new directions for celastrol's modification.



GA is another pentacyclic triterpene with broad biological activities, particularly in the fields of anti-inflammatory and anticancer research, showcasing immense potential for therapeutic applications. Similar to OA, BA, and UA, GA's poor water solubility and high toxicity have limited its clinical application. Recent studies on GA's structural modifications have led to the discovery of derivatives with promising anticancer activities, along with initial insights into their mechanisms of action, laying the groundwork for the development of GA as an anti-cancer agent. Further optimization of GA's structural modifications to enhance bioactivity and reduce toxicity will facilitate the clinical translation of GA and its derivatives.

In conclusion, structural modifications of natural products hold great potential for the development of novel anticancer drugs. The modification studies on pentacyclic triterpenes, such as OA, BA, UA, and celastrol, not only enrich the theoretical foundation of chemistry and pharmacology but also provide new avenues for drug discovery.

Data availability

No new data were created or analyzed in this study. This article is a review of existing literature, and all data supporting the findings are derived from previously published sources, which are appropriately cited in the manuscript. Should any further clarification on data sources be needed, the corresponding author can provide additional information upon request.

Conflicts of interest

The authors declare that they have no competing interests.

Acknowledgements

This research was funded by Guiyang City Science and Technology Plan Project (No. [2024]2-37), the Guizhou Provincial Basic Research Program (Natural Science) (No. ZK[2023] general 404 and ZK [2024] general 370), the Research Project on Traditional Chinese Medicine and Ethnic Medicine Science and Technology of Guizhou Provincial Administration of Traditional Chinese Medicine (QZY-2024-130), High-level Innovative Talents of Guizhou Province (QianKeHe platform talents-GCC [2023] 047), Guizhou Province Innovation Talent Team Construction for the Efficient and Comprehensive Utilization of Unique Food and Medicinal Materials (Platform Talent-CXTD [2023]020), Natural Science Research Project of Guizhou Department of Education (Qianjiaoj [2023]069) and the Projects of Guizhou Province (Qian Jiao He KY Zi [2022]253, gzwkj2022-232 and gzwkj2022-467).

References

- 1 D. Wang, B. Wang, Y. Liu, *et al.*, Synthetic pathway and fermentation process optimization of oleanolic acid in yeast cell factory, *China J. Chin. Mater. Med.*, 2014, **39**(14), 2640–2645.
- 2 J. Pollier and A. Goossens, Oleanolic acid, *Phytochemistry*, 2012, **77**, 10–15.
- 3 J. Wei, H. Liu, M. Liu, *et al.*, Oleanolic acid potentiates the antitumor activity of 5-fluorouracil in pancreatic cancer cells, *Oncol. Rep.*, 2012, **28**, 1339–1345.
- 4 M. Zhu, C. Wang, W. Sun, *et al.*, Boosting 11-oxo-beta-amyrin and glycyrrhetic acid synthesis in *Saccharomyces cerevisiae* via pairing novel oxidation and reduction system from legume plants, *Metab. Eng.*, 2018, **45**, 43–50.
- 5 M. A. Fernández, B. I. Heras, M. D. García, *et al.*, New insights into the mechanism of action of the anti-inflammatory triterpene lupeol, *J. Pharm. Pharmacol.*, 2010, **53**, 1533–1539.
- 6 W. N. Setzer and M. C. Setzer, Plant-derived triterpenoids as potential antineoplastic agents, *Mini-Rev. Med. Chem.*, 2003, **3**(6), 540–556.
- 7 K. Qian, R.-Y. Kuo, C.-H. Chen, *et al.*, Anti-AIDS agents 81. design, synthesis, and structure-activity relationship study of betulinic acid and moronic acid derivatives as potent HIV maturation inhibitors, *J. Med. Chem.*, 2010, **53**(8), 3133–3141.
- 8 E. Pisha, H. Chai, I.-S. Lee, *et al.*, Discovery of betulinic acid as a selective inhibitor of human melanoma that functions by induction of apoptosis, *Nat. Med.*, 1995, **1**(10), 1046–1051.
- 9 E. Urano, S. D. Ablan, R. Mandt, *et al.*, Alkyl amine bevirimat derivatives are potent and broadly active HIV-1 maturation inhibitors, *Antimicrob. Agents Chemother.*, 2016, **60**(1), 190–197.
- 10 T. Lin, D. Wang, Z. Dai, *et al.*, Creation of a *Saccharomyces cerevisiae* cell factory for the fermentation production of lupeol, *China J. Chin. Mater. Med.*, 2016, **41**(6), 1008–1015.
- 11 C. Z. Lu, C. Zhang, F. Zhao, *et al.*, Biosynthesis of ursolic acid and oleanolic acid in *Saccharomyces cerevisiae*, *AIChE J.*, 2018, **64**, 3794–3802.
- 12 X. He and R. H. Liu, Triterpenoids isolated from apple peels have potent antiproliferative activity and may be partially responsible for apple's anticancer activity, *J. Agric. Food Chem.*, 2007, **55**, 4366–4370.
- 13 B. D. Abrosca, A. Fiorentino, P. Monaco, *et al.*, Annurcoic acid : A new antioxidant ursane triterpene from fruits of cv. Annurca apple, *Food Chem.*, 2006, **98**, 285–290.
- 14 E. Vranová, D. Coman and W. Grussem, Network analysis of the MVA and MEP pathways for isoprenoid synthesis, *Annu. Rev. Plant Biol.*, 2013, **64**, 665–700.
- 15 Z. Luo, K. Zhang, X. Ma, *et al.*, Research Progress in Synthetic Biology of Triterpenoid Saponins, *Chin. Tradit. Herb. Drugs*, 2016, **47**(10), 1806–1814.
- 16 Z. Zhao, L. Xing, Y. Zhou, *et al.*, Preparation of Oleanolic Acid Sodium Salt and Its Solubility, *Chin. Tradit. Pat. Med.*, 1991, **13**(8), 32.
- 17 G. Zhai, H. Lou, L. Zou, *et al.*, Research progress on drug-phospholipid complexes, *Chin. J. Pharm.*, 2001, **36**(12), 800–806.



- 18 Z. Li, Z. Sun and H. Liu, The protective effect of sodium oleanolate on chemical liver injury in rats, *Guizhou Med. J.*, 2004, **28**(7), 582–583.
- 19 Y. Zhang, J. X. Li and J. Zhao, *et al.*, Synthesis and Activity of Oleanolic Acid Derivatives, A Novel Class of Inhibitors of Osteoclast Formation, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 1629–1632.
- 20 Y. Meng, H. Nie, X. Wang, *et al.*, Synthesis and Anti-Tumor Activity of Oleanolic Acid Derivatives, *Guizhou Med. J.*, 2011, **28**(7), 582–583.
- 21 Y. Zhang, L. Gu, L. Su, *et al.*, Synthesis of Oleanolic Acid Derivatives Containing Amino Phosphonate and Their Anti-Tumor Activity, *J. Anhui Agric. Sci.*, 2012, **40**(5), 2737–2738.
- 22 W. Huang, J. Huang, D. Zhang, *et al.*, Study on the Anti-Invasion and Apoptosis-Inducing Effects of Pentacyclic Triterpenoids on Human Lung Cancer Cells, *Chin. J. Lung Cancer*, 2003, **6**(4), 254–257.
- 23 Z. Huang, H. Li, L. Sun, *et al.*, Study on the Anti-Metastatic Effect of Oleanolic Acid on Rat Transplantable Liver Cancer, *J. Anhui Agric. Sci.*, 2011, **39**(18), 10807–10808.
- 24 H. Sun, C. Hu, W. Fang, *et al.*, Synthesis of 3-Oxo-Oleanolic Acid Amino Acid Conjugates, Determination of Aqueous Solubility, and Research on Anticancer Activity, *Chin. J. Med. Chem.*, 2008, **18**(1), 11–15.
- 25 Q. Li, J. Gu, L. Li, *et al.*, Synthesis of Oleanolic Acid Derivatives, *Synth. Chem.*, 2010, **18**(3), 290–296.
- 26 J. Xi, Q. Chang, C. K. Chan, *et al.*, Formulation Development and Bioavailability Evaluation of a Self-nanoemulsified Drug Delivery System of Oleanolic Acid, *AAPS PharmSciTech*, 2009, **10**, 172–182.
- 27 Y. Zhang, J. X. Li, J. Zhao, *et al.*, Synthesis and activity of oleanolic acid derivatives, a novel class of inhibitors of osteoclast formation, *Bioorg. Med. Chem. Lett.*, 2005, **15**(6), 1629–1632.
- 28 T. Akihisa, S. Kamo, T. Uchiyama, *et al.*, Cytotoxic activity of *Perilla frutescens* var. *japonica* leaf extract is due to high concentrations of oleanolic and ursolic acids, *J. Nat. Med.*, 2006, **60**, 331–333.
- 29 W. Li, S. das, K. Ng, *et al.*, Formulation, biological and pharmacokinetic studies of sucrose ester-stabilized nanosuspensions of oleanolic acid, *Pharm. Res.*, 2011, **28**, 2020–2033.
- 30 L. Wang, G. Han, W. Liu, *et al.*, Chemical and Pharmacological Research on Oleanolic Acid, *J. Pharm. Pract.*, 2001, **19**(2), 104–107.
- 31 H. Yang, M. Deng, H. Jia, *et al.*, A review of structural modification and biological activities of oleanolic acid, *Chin. J. Nat. Med.*, 2024, **22**(1), 15–30.
- 32 F. Jannus, J. Sainz, F. J. Reyes-Zurita, *et al.*, Principal Bioactive Properties of Oleanolic Acid, Its Derivatives and Analogues, *Molecules*, 2024, **29**, 3291–3312.
- 33 A. Spivak, R. Khalitova, D. Nedopekina, *et al.*, Synthesis and evaluation of anticancer activities of novel c-28 guanidine-functionalized triterpene acid derivatives, *Molecules*, 2018, **23**, 3000–3022.
- 34 K. Chouaïb, A. Romdhane, S. Delemasure, *et al.*, Regiospecific synthesis by copper- and ruthenium-catalyzed azide–alkyne 1, 3-dipolar cycloaddition, anticancer and anti-inflammatory activities of oleanolic acid triazole derivatives, *Arabian J. Chem.*, 2019, **12**(8), 3732–3742.
- 35 E. F. Khusnutdinova, G. N. Apryshko and A. V. Petrova, The Synthesis and Selective Cytotoxicity of New Mannich Bases, Derivatives of 19- and 28-Alkynyltriterpenoids, *Russ. J. Bioorg. Chem.*, 2018, **44**, 123–127.
- 36 X. Liu, B. Li, Z. Zhang, *et al.*, Synthesis and discovery novel anti-cancer stem cells compounds derived from the natural triterpenic acids, *J. Med. Chem.*, 2018, **61**(23), 10814–10833.
- 37 Y.-Q. Meng, Z.-Y. Kuai, S.-W. Zhan, *et al.*, Synthesis and anti-tumor activity of oleanolic acid derivatives, *Acta Pharm. Sin.*, 2011, **46**(10), 1215–1220.
- 38 Z. Chen, K.-Y. Huang, Y. Ling, *et al.*, Discovery of an oleanolic acid/hederagenin–nitric oxide donor hybrid as an EGFR tyrosine kinase inhibitor for non-small-cell lung cancer, *J. Nat. Prod.*, 2019, **82**(11), 3065–3073.
- 39 D. S. Raghuvanshi, N. Verma, S. Singh, *et al.*, Design and synthesis of novel oleanolic acid based chromenes as anti-proliferative and anti-inflammatory agents, *New J. Chem.*, 2018, **42**(20), 16782–16794.
- 40 A. Froelich, B. Bednarczyk-Cwynar, L. Zaprutko, *et al.*, Beckmann rearrangement within the ring C of oleanolic acid lactone: Synthesis, structural study and reaction mechanism analysis, *J. Mol. Struct.*, 2017, **1136**, 173–181.
- 41 L. Zaprutko, D. Partyka and B. Bednarczyk-Cwynar, Triterpenoids. Part 21: Oleanolic acid azaderivatives as percutaneous transport promoters, *Bioorg. Med. Chem. Lett.*, 2004, **14**(18), 4723–4726.
- 42 J. Xu, X. Wang, H. Zhang, *et al.*, Synthesis of triterpenoid derivatives and their anti-tumor and anti-hepatic fibrosis activities, *Nat. Prod. Res.*, 2020, **34**(6), 766–772.
- 43 J. F. Li, R. Z. Huang, G. Y. Yao, *et al.*, Synthesis and biological evaluation of novel aniline-derived asiatic acid derivatives as potential anticancer agents, *Eur. J. Med. Chem.*, 2014, **86**, 175–188.
- 44 N. Heisi, B. Siewert, D. Strohl, *et al.*, A simple but unusual rearrangement of an oleanane to a taraxerane 28, 14 β -olide, *Steroids*, 2021, **172**, 108853–108858.
- 45 T. Honda, B. V. Rounds, G. W. Gribble, *et al.*, Design and synthesis of 2-cyano-3, 12-dioxoolean-1, 9-dien-28-oic acid, a novel and highly active inhibitor of nitric oxide production in mouse macrophages, *Bioorg. Med. Chem. Lett.*, 1998, **8**(19), 2711–2714.
- 46 R. Borella, L. Forti, L. Gibellini, *et al.*, Synthesis and anticancer activity of CDDO and CDDO-Me, two derivatives of natural triterpenoids, *Molecules*, 2019, **24**(22), 4097–4117.
- 47 Y.-Y. Wang, Y.-X. Yang, H. Zhe, *et al.*, Bardoxolone methyl (CDDO-Me) as a therapeutic agent: an update on its pharmacokinetic and pharmacodynamic properties, *Drug Des., Dev. Ther.*, 2014, 2075–2088.



- 48 M. H. L. Wong, I. K. Bryan, I. M. Copple, *et al.*, Design and synthesis of irreversible analogues of bardoxolone methyl for the identification of pharmacologically relevant targets and interaction sites, *J. Med. Chem.*, 2016, **59**(6), 2396–2409.
- 49 F. Gao, Q. Zuo, T. Jiang, *et al.*, A newly synthesized oleanolic acid derivative inhibits the growth of osteosarcoma cells in vitro and in vivo by decreasing c-MYC-dependent glycolysis, *J. Cell. Biochem.*, 2019, **120**(6), 9264–9276.
- 50 F. Kang, Y. Ai, Y. Zhang, *et al.*, Design and synthesis of new hybrids from 2-cyano-3, 12-dioxooleana-9-dien-28-oic acid and O2-(2, 4-dinitrophenyl) diazeniumdiolate for intervention of drug-resistant lung cancer, *Eur. J. Med. Chem.*, 2018, **149**, 269–306.
- 51 W. Yan, C. Zhang, B. Li, *et al.*, A Series of oleanolic acid derivatives as anti-hepatitis B virus agents: Design, synthesis, and in vitro and in vivo biological evaluation, *Molecules*, 2016, **21**(4), 402–413.
- 52 W. Li, F. Yang, L. Meng, *et al.*, Synthesis, structure activity relationship and anti-influenza A virus evaluation of oleanolic acid-linear amino derivatives, *Chem. Pharm. Bull.*, 2019, **67**(11), 1201–1207.
- 53 L. Meng, Y. Su, F. Yang, *et al.*, Design, synthesis and biological evaluation of amino acids-oleanolic acid conjugates as influenza virus inhibitors, *Bioorg. Med. Chem.*, 2019, **27**(23), 115147–115156.
- 54 M. Medina-O'Donnell, F. Rivas, F. J. Reyes-Zurita, *et al.*, Oleanolic acid derivatives as potential inhibitors of HIV-1 protease, *J. Nat. Prod.*, 2019, **82**(10), 2886–2896.
- 55 J. W. Wang, W. R. Xu, P. Yong, *et al.*, Glycyrrhetic acid-30 amide derivatives and their use, US2008214636(A1), 2008.
- 56 J. Yong, Research Progress on the Synthesis and Bioactivity of Glycyrrhizic Acid and Its Triterpenoid Saponin Derivatives, *Shizhen Guoyi Guoyao*, 2012, **23**(6), 1495–1498.
- 57 J. Tatsuzaki, M. Taniguchi, K. F. Bastow, *et al.*, Antitumor agents glycyrrhetic acid dehydrozingerone conjugates as cytotoxic agents, *Bioorg. Med. Chem.*, 2007, **15**, 6193–6199.
- 58 Y. Zhang, Research on Chiral Ketones of Galactose and Triterpenoid Saponins and Isoxazole and Benzimidazole Ring Isoxazolidinones, PhD Dissertation, Shandong University, 2007.
- 59 N. Zhang, X. Cui, X. Zhao, *et al.*, Synthesis and Anti-Liver Cancer Activity of Glycyrrhetic Acid Derivatives, *Chin. J. Exp. Tradit. Med. Formulae*, 2015, **10**, 37–41.
- 60 R. Csuk, S. Schwarz, R. Kluge, *et al.*, structure-Activity Relationships of Glycyrrhetic Acid Derivatives Modified at Position C-11, *Arch. Pharm.*, 2012, **345**(1), 28–32.
- 61 S. Schwarz, R. Csuk, *et al.*, synthesis and antitumor activity of glycyrrhetic acid derivatives, *Bioorg. Med. Chem.*, 2010, **18**(21), 7458–7474.
- 62 Y. Gao, X. Guo, L. Zhao, *et al.*, The synthesis of Glycyrrhetic Acid Derivatives Containing A Nitrogen Heterocycle and Their Antiproliferative Effects in Human Leukemia Cells, *Molecules*, 2010, **15**, 4439–4449.
- 63 Y. Meng, Z. Yang, M. Zhang, *et al.*, Synthesis of Glycyrrhetic Acid C3, C11, and C30 Derivatives and Their Anti-Tumor Activity In Vitro, *Drugs Clin.*, 2014, **29**(1), 6–10.
- 64 X. Su, N. Vicker, H. Lawrence, *et al.*, Inhibition of human and rat 11 β -hydroxysteroid dehydrogenase type 1 by 18 β -glycyrrhetic acid derivatives, *J. Steroid Biochem. Mol. Biol.*, 2007, **104**(3–5), 312–320.
- 65 R. C. Santos, J. A. R. Salvador, S. Marin, *et al.*, Novel semisynthetic derivatives of betulin and betulinic acid with cytotoxic activity, *Bioorg. Med. Chem.*, 2009, **17**(17), 6241–6250.
- 66 E. R. Trumbull, E. Bianchi, D. J. Eckert, *et al.*, inhibitory agents from *Vauquelinia corymbosa* (Rosaceae), *J. Pharm. Sci.*, 1976, **65**(9), 1407–1408.
- 67 P. Chatterjee, S. A. Kouzi, J. M. Pezzuto, *et al.*, Biotransformation of the antimelanoma agent betulinic acid by *Bacillus megaterium* ATCC 13368, *Appl. Environ. Microbiol.*, 2000, **66**(9), 3850–3855.
- 68 R. Mukherjee, M. Jaggi, P. Rajendran, *et al.*, Betulinic acid and its derivatives as anti-angiogenic agents, *Bioorg. Med. Chem. Lett.*, 2004, **14**(9), 2181–2184.
- 69 J. Li, M. Goto, X. Yang, *et al.*, Fluorinated betulinic acid derivatives and evaluation of their anti-HIV activity, *Bioorg. Med. Chem. Lett.*, 2016, **26**(1), 68–71.
- 70 O. V. Tsepaveva, A. V. Nemtarev, L. R. Grigor'eva, *et al.*, Synthesis of C (28)-linker derivatives of betulinic acid bearing phosphonate group, *Russ. Chem. Bull.*, 2021, **70**, 179–182.
- 71 A. Y. Spivak, D. A. Nedopekina, E. R. Shakurova, *et al.*, Synthesis of lupane triterpenoids with triphenylphosphonium substituents and studies of their antitumor activity, *Russ. Chem. Bull.*, 2013, **62**, 188–198.
- 72 Y. Chen, S. Y. Sit, J. Chen, *et al.*, The design, synthesis and structure-activity relationships associated with C28 amine-based betulinic acid derivatives as inhibitors of HIV-1 maturation, *Bioorg. Med. Chem. Lett.*, 2018, **28**, 1550–1557.
- 73 V. Reutrakul, N. Anantachoke, M. Pohmakotr, *et al.*, Anti-HIV-1 and anti-inflammatory lupanes from the leaves, twigs, and resin of *Garcinia hanburyi*, *Planta Med.*, 2010, **76**(4), 68–71.
- 74 J. Li, W. Guo and Q. Yang, Effects of ursolic acid and oleanolic acid on human colon carcinoma cell line HCT15, *World J. Gastroenterol.*, 2002, **8**(3), 493–495.
- 75 H. S. Young, H. Y. Chung, C. K. Lee, *et al.*, Ursolic acid inhibits aflatoxin B1-induced mutagenicity in a Salmonella assay system, *Biol. Pharm. Bull.*, 1994, **17**(7), 990–992.
- 76 E. Kassi, Z. Papoutsis, H. Pratsinis, *et al.*, Ursolic Acid, a Naturally Occurring Triterpenoid, Demonstrates Anticancer Activity on Human Prostate Cancer Cells, *J. Cancer Res. Clin. Oncol.*, 2007, **133**, 493–500.
- 77 D. Zhang and L. L. Zhao, Synthesis and Anti-hepatocellular Carcinoma In Vitro of Ursolic Acid Derivatives Bearing Benzylphosphonic Acid Diethyl Ester Acetamide Moieties, *Chin. J. Mod. Appl. Pharm.*, 2022, **39**(06), 738–744.
- 78 T. Tian, Synthesis Characterization and Anticancer Activity Research of Three Pentacyclic Triterpenoid Nitrogen-Containing Derivatives, PhD Dissertation, Liaoning Normal University, 2016.



- 79 K. Bai, F. Chen and Y. Guo, Synthesis and In Vitro Anticancer Activity of Polyethylene Glycol Conjugated Ursolic Acid Derivatives, *Chin. J. New Drugs*, 2012, **13**(21), 1536–1540.
- 80 K. Bai, F. Chen and Y. Guo, Synthesis and In Vitro Anticancer Activity of Ursolic Acid Derivatives on Prostate Cancer PC-3 Cells, *Chin. J. New Drugs*, 2012, **22**(21), 2667–2673.
- 81 K. Bai, F. Chen, Y. Zheng, *et al.*, Synthesis and In Vitro Anti-Gastric Cancer Activity of Amino Acid Modified Ursolic Acid Derivatives, *Chin. J. Pharm.*, 2012, **4**(47), 265–269.
- 82 Y. Zhao, A-ring multi-oxidized derivatives of pentacyclic triterpenes, their preparation methods, and applications, *J. Chin. Med. Mater.*, 2006, (05), 427–428.
- 83 J. Wu, Z. H. Zhang, L. H. Zhang, *et al.*, Design, synthesis, and screening of novel ursolic acid derivatives as potential anti-cancer agents that target the HIF-1 α pathway, *Bioorg. Med. Chem. Lett.*, 2019, **29**(6), 853–858.
- 84 W. Wang, L. Lei, Z. Liu, *et al.*, Design, synthesis, and biological evaluation of novel nitrogen heterocycle-containing ursolic acid analogs as antitumor agents, *Molecules*, 2019, **24**(5), 877–891.
- 85 Q. Min, Synthesis of 3,4-Fissured Pentacyclic Triterpenoid-3,4-Lactone Derivatives and Their Anti-Hepatitis B Virus Activity, PhD Dissertation, Southern Medical University, 2015.
- 86 C. Chen, R. Sun, Y. Sun, *et al.*, Synthesis, biological evaluation and SAR studies of ursolic acid 3 β -ester derivatives as novel CETP inhibitors, *Bioorg. Med. Chem. Lett.*, 2020, **30**(2), 126–824.
- 87 J. Xu, X. Nie, Y. Hong, *et al.*, Synthesis of water soluble glycosides of pentacyclic dihydroxytriterpene carboxylic acids as inhibitors of α -glucosidase, *Carbohydr. Res.*, 2016, **424**, 42–53.
- 88 Z. Zeng, X. Yin, X. Wang, *et al.*, Synthesis of water soluble pentacyclic dihydroxyterpene carboxylic acid derivatives coupled amino acids and their inhibition activities on α -glucosidase, *Bioorg. Chem.*, 2019, **86**, 277–287.
- 89 X. Liu, X. Zang, X. Yin, *et al.*, Semi-synthesis of C28-modified triterpene acid derivatives from maslinic acid or corosolic acid as potential α -glucosidase inhibitors, *Bioorg. Chem.*, 2020, **97**, 103694–103740.
- 90 A. R. Setty and L. H. Sigal, Herbal medications commonly used in the practice of rheumatology: mechanisms of action, efficacy, and side effects, *Semin. Arthritis Rheum.*, 2005, **34**(6), 773–784.
- 91 M. Jing, J. Yang, L. Zhang, *et al.*, Celastrol inhibits rheumatoid arthritis through the ROS-NF- κ B-NLRP3 inflammasome axis, *Int. Immunopharmacol.*, 2021, **98**, 107–879.
- 92 Z. Li, R. Liu, Z. Guo, *et al.*, Celastrol-based nanomedicine promotes corneal allograft survival, *J. Nanobiotechnol.*, 2021, **19**(1), 1–19.
- 93 J. Zhang, K. Zhou, X. Zhang, *et al.*, Celastrol ameliorates inflammation in human retinal pigment epithelial cells by suppressing NF- κ B signaling, *J. Ocul. Pharmacol. Ther.*, 2019, **35**(2), 116–123.
- 94 Y. Feng, W. Wang, Y. Zhang, *et al.*, Synthesis and biological evaluation of celastrol derivatives as potential anti-glioma agents by activating RIP1/RIP3/MLKL pathway to induce necroptosis, *Eur. J. Med. Chem.*, 2022, **229**, 114070–114086.
- 95 M. Xu, N. Li, Z. Zhao, *et al.*, Design, synthesis and antitumor evaluation of novel celastrol derivatives, *Eur. J. Med. Chem.*, 2019, **174**, 265–276.
- 96 P. Coghi, J. P. L. Ng, O. Kadioglu, *et al.*, Synthesis, computational docking and biological evaluation of celastrol derivatives as dual inhibitors of SERCA and P-glycoprotein in cancer therapy, *Eur. J. Med. Chem.*, 2021, **224**, 113676–113690.
- 97 N. Li, M. Xu, B. Wang, *et al.*, Discovery of Novel Celastrol Derivatives as Hsp90–Cdc37 Interaction Disruptors with Antitumor Activity, *J. Med. Chem.*, 2019, **62**(23), 10798–10815.
- 98 X. Li, J. Ding, N. Li, *et al.*, Synthesis and biological evaluation of celastrol derivatives as anti-ovarian cancer stem cell agents, *Eur. J. Med. Chem.*, 2019, **179**, 667–679.
- 99 N. Li, M. Xu, N. Bao, *et al.*, Discovery of novel NO-releasing celastrol derivatives with Hsp90 inhibition and cytotoxic activities, *Eur. J. Med. Chem.*, 2018, **160**, 1–8.
- 100 W. J. Tang, J. Wang, X. Tong, *et al.*, Design and synthesis of celastrol derivatives as anticancer agents, *Eur. J. Med. Chem.*, 2015, **95**, 166–173.
- 101 Y. Jing, G. Wang, Y. Ge, *et al.*, Synthesis, anti-tumor and anti-angiogenic activity evaluations of asiatic acid amino acid derivatives, *Molecules*, 2015, **20**(4), 7309–7324.
- 102 K. Tang, J. Huang, J. Pan, *et al.*, Design, synthesis and biological evaluation of C (6)-indole celastrol derivatives as potential antitumor agents, *RSC Adv.*, 2015, **5**(25), 19620–19623.
- 103 R. G. Brahmachari, N. C. Mandal, R. Roy, *et al.*, A new pentacyclic triterpene with potent antibacterial activity from *Limnophila indica* Linn.(Druce), *Fitoterapia*, 2013, **90**, 104–111.
- 104 A. A. Lasisi, B. W. Ayinde, A. O. Adeleye, *et al.*, New triterpene isovanniloyl and antibacterial activity of constituents from the roots of *Paullinia pinnata* Linn (Sapindaceae), *J. Saudi Chem. Soc.*, 2015, **19**(2), 117–122.
- 105 L. J. Shai, L. J. McGaw, M. A. Aderogba, *et al.*, Four pentacyclic triterpenoids with antifungal and antibacterial activity from *Curtisia dentata* (Burm.f) CA Sm. Leaves, *J. Ethnopharmacol.*, 2008, **119**(2), 238–244.
- 106 A. Parra, S. Martin-Fonseca, F. Rivas, *et al.*, Semi-synthesis of acylated triterpenes from olive-oil industry wastes for the development of anticancer and anti-HIV agents, *Eur. J. Med. Chem.*, 2014, **74**, 278–301.

