


Dalton Transactions

An international journal of inorganic chemistry

rsc.li/dalton

Volume 52
Number 23
21 June 2023
Pages 7757-8156



ISSN 1477-9226

PAPER

M. Concepción Gimeno *et al.*
Catalysis-free synthesis of thiazolidine-thiourea ligands for
metal coordination (Au and Ag) and preliminary cytotoxic
studies

PAPER

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *Dalton Trans.*, 2023, **52**, 7797

Catalysis-free synthesis of thiazolidine–thiourea ligands for metal coordination (Au and Ag) and preliminary cytotoxic studies†

Daniel Salvador-Gil,^a Raquel P. Herrera^b and M. Concepción Gimeno ^{*,a}

The reaction of propargylamines with isothiocyanates results in the selective formation of iminothiazolidines, aminothiazolines or mixed thiazolidine–thiourea compounds under mild conditions. It has been observed that secondary propargylamines lead to the selective formation of cyclic 2-amino-2-thiazoline derivatives, while primary propargylamines form iminothiazoline species. In addition, these cyclic thiazoline derivatives can further react with an excess of isothiocyanate to give rise to thiazolidine–thiourea compounds. These species can also be achieved by reaction of propargylamines with isothiocyanates in a molar ratio of 1 : 2. Coordination studies of these heterocyclic species towards silver and gold with different stoichiometries have been carried out and complexes of the type [ML(PPh₃)]OTf, [ML₂]OTf (M = Ag, Au) or [Au(C₆F₅)L] have been synthesised. Preliminary studies of the cytotoxic activity in lung cancer cells have also been performed in both ligands and complexes, showing that although the ligands do not exhibit anticancer activity, their coordination to metals, especially silver, greatly enhances the cytotoxic activity.

Received 9th January 2023,
Accepted 21st February 2023

DOI: 10.1039/d3dt00079f

rsc.li/dalton

Introduction

In recent years, there has been an increased interest in the discovery and development of novel biologically active compounds as more selective agents for the treatment of different diseases. Cancer is a common illness in our society, the second leading cause of death in the world, responsible for important health costs associated with the treatment of patients and with a high level of mortality. As a result, numerous chemotherapeutic agents have been developed to avoid the progression of tumours or to prevent their recurrence, as well as with the main aim of avoiding many of the side effects of conventional antitumor drugs.¹ In the continuous efforts to develop drugs with such anticancer properties, scientists have focused upon many different aspects of this illness. Among the antitumor organic compounds discovered in recent years, urea² and thiourea³ derivatives have exhibited potent anticancer activity and have been reported as prodrugs. These structures have also been reported as important biologically active compounds in a

variety of promising chemical prototypes for drug development. Additionally, thioureas could display different properties, such as anti-HIV,⁴ antiviral,⁵ antibacterial,⁶ antifungal,⁷ analgesic,⁸ or antithrombotic activity,⁹ among others (Fig. 1).

Heterocyclic compounds, such as thiazoline and thiazolidine scaffolds,¹⁰ have also been considered as privileged structural motifs in medicinal chemistry and of high importance in organic synthesis.¹¹ These heterocycles provide the final drugs with improved pharmacokinetic properties and facilitate the construction of a large variety of biologically active molecules,




Fig. 1 Biologically active thiourea and aminothiazole scaffolds.

^aDepartamento de Química Inorgánica, Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), CSIC-Universidad de Zaragoza, 50009 Zaragoza, Spain.
E-mail: gimeno@unizar.es

^bDepartamento de Química Orgánica, Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), CSIC-Universidad de Zaragoza, 50009 Zaragoza, Spain

† Electronic supplementary information (ESI) available. CCDC 2234522–2234528. For ESI and crystallographic data in CIF or other electronic format see DOI:

<https://doi.org/10.1039/d3dt00079f>

such as anti-tumour,¹² anti-bacterial,¹³ anti-viral,¹⁴ and anti-inflammatory compounds.¹⁵ Moreover, several aminothiazole derivatives are clinically approved drugs for the treatment of amyotrophic lateral sclerosis (Riluzole),¹⁶ or anti-fungals (Abafungin),¹⁷ or third generation cephalosporin antibiotics (Cefdinir) (Fig. 1).

Additionally, the development of new metallodrugs is currently becoming very important for cancer chemotherapy since they try to improve the activity of existing drugs such as cisplatin and to avoid its important side effects and resistance. In this field, improvements have been achieved with ruthenium¹⁸ and also with the design of gold and silver-based drugs,¹⁹ which present great potential as anticancer agents.

In this context, we have focused our efforts during the last decade on the search and development of new biologically active gold and silver²⁰ complexes with new biological ligands²¹ in a synergic way as an excellent approach for discovering novel bioactive metallodrugs.

The search for new biological compounds goes in parallel with the development of new organic ligands or their corresponding metallodrug based compounds, because of the interesting properties that both parts could exhibit, thiourea or aminothiazoline derivatives and gold or silver complexes. We envisioned that coordination of these organic molecules with the respective group 11 metal centres should increase the biological activity of the resulting complexes in comparison with each part alone.

The conversion of propargylamines and isothiocyanates to afford the thiazolidine or thiazole moiety has been known long time ago; however, usually harsh conditions, microwaves or metal catalysts are used to achieve these heterocycles.

Results and discussion

Considering that heterocycles play an important role in many natural products and drug development, with a wide range of biological properties, we consider studying the formation of thiazolidines and thiazolidine-thiourea species under mild conditions. Additionally, their coordination to gold and silver centres will be explored because to the best of our knowledge the coordination of thiazoline containing heterocycles to gold or silver metal for biological purpose has been overlooked in the literature so far.

To initiate the study, different model propargylamines **1a–c** and isothiocyanates **2a–c** were selected (Fig. 2).




Fig. 2 Propargylamines **1a–c** and isothiocyanates **2a–c**.


For this purpose, the conditions for the formation of thiourea species **3** or aminothiazoline derivative **4** were investigated as disclosed in Scheme 1.

Surprisingly, the normal reaction to afford thiourea **3** did not work between amine **1a** and isothiocyanate **2a** using EtOH as the solvent at room temperature. To our delight, after analysis of the ¹H NMR spectrum of the reaction, it was observed that the signals of the obtained product were in agreement with the cyclic compound **4**, the 2-aminothiazoline derivative.²² In fact, to obtain the desired thiourea **3**, the solvent of the process was changed to a less polar solvent such as acetonitrile (Scheme 1). The reaction was carried out under the same conditions of time and temperature and finally the thiourea **3** was obtained.

However, in most of the assays the thiourea could not be isolated in the pure form, and a mixture of both thiazoline and thiourea derivatives was obtained, even if the reaction was carried out in acetonitrile, whereas pure thiazoline compounds are achieved in ethanol. Therefore, we continued working with EtOH. It is worth mentioning that when 2-thiazolines have an amino group at position 2, a tautomeric equilibrium is observed between the thiazoline and thiazolidine forms, that is 2-amino-2-thiazoline and 2-iminothiazolidine. Interestingly, we have observed that the use of secondary propargylamines **1b,c** leads selectively to the formation of cyclic iminothiazolidine derivatives **5**, **6**, **8** and **9**, while primary propargylamine **1a** could form 2-aminothiazoline derivatives **4** and **7** (Scheme 2).


The easy formation of these thiazoline and thiazolidine derivatives was accomplished for all the alkynes and isothiocyanates assayed with very good yields. Thiazoline **4** crystallises with four independent molecules, with similar distances and bond angles in all the molecules (Fig. 3B).²³ Moreover, it is possible to observe the existence of a double bond between the N1–C1 atoms of 1.278(2) Å, which corroborates the formation of the 2-amino-2-thiazoline **4** and also the presence of hydrogen bonds among the molecules that form a chain polymer. The shortest bonds correspond to the N7...H–N6 unit with a distance of 2.047 Å and an angle of 174° (Fig. 3B). In the structure of compound **6**, the distances C5–N2 of 1.2759(16) and C5–N1 of 1.3590(15) support the formation of the iminothiazolidine (Fig. 3C). The crystal structure of compound **7** also supports the formation of the 2-amino-2-thiazoline derivative, with distances N1–C4 of 1.276(2) and N2–C4 of 1.357(2) Å (Fig. 3D).

Encouraged with this reactivity, we tested the scope of this approach by adding 2 equivalents of isothiocyanate **2a–c** to propargylamine **1a**. Interestingly, new thiazolidine-thiourea



Scheme 1 Formation of thiourea **3** or aminothiazoline **4**.





Scheme 2 Synthesis of 2-amino-2-thiazolines and 2-iminothiazolidines.





Fig. 3 Crystal molecular structures of 4 (A), 6 (C), and 7 (D), and the chain formed by hydrogen bonding in 4 (B) showing ellipsoids at the 50% probability level.

derivatives **10–12** were prepared, incorporating three molecules of reagents in the final product in an atom-economical process (Scheme 3).

For the formation of these thiazolidine–thiourea species, the phenyl (**2a**), 3-fluorophenyl (**2b**) and benzyl (**2c**) isocyanates have been used. The reactions were carried out in ethanol and



Scheme 3 Synthesis of thiazolidine–thiourea adducts **10–12**.

the compounds precipitated as a white solid, being obtained in the pure form and in very good yields. As a proof of the final structures of thiourea–thiazolidine derivatives **10** and **11**, crystal structures were also corroborated by X-ray diffraction and are presented in Fig. 4.²⁴ In both cases, we observe the presence of an intramolecular hydrogen bond between the amino group of the thiourea and the imino group of the iminothiazolidine. For compound **10** the hydrogen bond N1–H6...N3 has a donor...acceptor distance of 2.621 Å and an angle of 142.2°. In compound **11**, the N1–H1...N3 bond has a donor...acceptor distance of 2.602 Å and an angle of 146.1°.

Mechanism of the reaction

A plausible mechanism for the formation of thiazolines or thiazolidine–thiourea derivatives is proposed in Scheme 4. The first step would be the reaction of propargylamine **1a** with the isothiocyanate **2**, giving rise to the formation of the corresponding thiourea **I**. If two equivalents of isothiocyanate are used, thiourea **I** could be first generated. In this case, only thiourea **I** could be isolated in a pure form. The next step would be the generation of a more reactive allene intermediate **II** or **IV**, in equilibrium with its thiourea species, probably favoured by the use of a polar solvent.²⁵ The formation of these intermediates would receive a posterior intramolecular nucleophilic attack by the sulfur atom to the quaternary carbon of the allene, giving rise to the cyclisation of the compound and the formation of the corresponding 2-iminothiazolidines **V** and **VI**. At this point, we cannot discard the direct attack of the sulfur atom over the triple bond as previously proposed by other authors.²⁶ In the case of 2-iminothiazolidines





Fig. 4 Crystal molecular structures of compounds **10** (A) and **11** (B) showing ellipsoids at the 50% probability level.



Scheme 4 Proposed formation of the iminothiazolidine **V** and thiazolidine–thiourea **VI**.

V, the internal amino NH group could also produce a nucleophilic attack on another molecule of isothiocyanate giving rise to the same thiazolidine–thiourea VI. Alternatively, this route could also justify the obtainment of the same compound VI (Scheme 4).

Metal coordination studies


It is known that both thioureas and thiazolidines alone may have some biological activity as described above. Therefore, we envisioned that the tandem of the two units in the same molecule could improve their activity. We also foresaw that coordinating these types of ligands to group 11 metals, with proven biological activity, probably would result in an enhancement of the potential biological activity.²⁷ For this reason, the binding of thiazolidine–thiourea ligands to silver or gold fragments has been addressed in this section.

First, the coordination of silver to **10–12** was explored, using $[\text{Ag}(\text{OTf})(\text{PPh}_3)]$ or $\text{Ag}(\text{OTf})$ as the metallic precursor, to afford the heteroleptic **13–15** or the homoleptic **16–18** silver complexes (Scheme 5).


In both cases, coordination of silver to the sulfur atoms of the C=S group of the ligands is proposed. The resonances of the complexes in the spectra appear shifted with respect to those of the ligands. In all cases, the new complexes were obtained in very good yields.

Attempts were also made to synthesize the analogous cationic gold complexes, starting from $[\text{Au}(\text{PPh}_3)(\text{MeCN})]\text{SbF}_6$ or $[\text{Au}(\text{tht})_2]\text{OTf}$ precursors, or neutral derivatives with $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ as the starting material (Scheme 6). The heteroleptic complexes **19–21** with the thiazolidine–thiourea and phosphine ligands were obtained with a very small amount of the homoleptic $[\text{Au}(\text{PPh}_3)_2]\text{OTf}$ complex.

The crystal structure of complex **20** has been established by X-ray diffraction studies. In this complex, the gold atom is in a linear environment bonded to the sulfur atom of thiourea with a distance Au1–S1 of 2.336(2) Å and to the phosphorus atom with a distance of Au1–P1 2.274(2) Å (Fig. 5).²⁸ The angle around the gold centre is slightly distorted from linearity, P1–Au1–S1 175.34(9)°. The molecules associate in dimers through a short aurophilic interaction of Au1...Au1 3.0861(7) Å (Fig. 5).



Scheme 5 Preparation of silver complexes **13–18**.



Scheme 6 Preparation of gold complexes **19–27**.




Fig. 5 Molecular structure of complex **20** showing ellipsoids at the 30% probability level, and its association in dimers through aurophilic interactions. Hydrogen atoms are omitted for clarity.

In complexes **25–27**, the ^{19}F NMR spectra provide good information about the formation of the complexes, in which three different resonances are assigned to the pentafluorophenyl group, two multiplets for the *ortho*- and *meta*-fluoro and a triplet for the *para*-fluoro. In addition, for compound **26** two additional signals are observed, as expected, corresponding to the fluorine atoms of the thiazolidine–thiourea ligand.

The crystal structure of complex **27** has also been characterised by X-ray diffraction (Fig. 6).²⁹ The geometry around the gold centre is also linear with a C1–Au1–S1 angle of 175.51(6)°. The main bond distances are Au1–C1 2.019(2) and Au1–S1 2.3069(6) Å. There is an intramolecular hydrogen bond as occurred in the ligands, with a donor...acceptor N3–H3...N2 distance of 2.626 Å and an angle of 146.2°.






Fig. 6 Molecular structure of complex 27 showing ellipsoids at the 50% probability level.

Biological studies

Before the biological tests, analyses of solubility and stability were performed for our complexes. Hence, the ligands and the complexes were not soluble in DMSO and mixtures of DMSO/water were used to carry out the tests. Stability studies were performed in DMSO by ^1H NMR studies and all compounds maintained stable unless for a period of 48 h, as no changes were observed between the initial ^1H NMR spectra and the corresponding ones after 24 and 48 h, confirming the retention of the complexes in solution. Preliminary studies of the cytotoxic activity of the compounds were carried out using the MTT assay for the lung carcinoma cell line A549.³⁰ The IC_{50} values for the complexes after 24 h of incubation are shown in Table 1.

From these results some statements can be concluded. Interestingly, the thiazolidine-thiourea ligands **10–12** did not show any type of anticancer activity against this cancer line. In contrast, the silver complexes **13–15** that incorporate the tri-

phenylphosphine ligand in their structure show excellent activity with IC_{50} values ranging from 3.97 ± 0.63 to 6.26 ± 0.29 , respectively. Unfortunately, no comparison is possible with the equivalent gold species **19–21** because in these cases the complexes were not obtained completely pure and, consequently, the biological test has not been performed. It is possible to compare the cytotoxic activity of the silver phosphine complexes with other related complexes, such as $[\text{Ag}(\text{PPh}_3)(\text{PyCO-aaOMe})]\text{OTf}$ (aaOMe = amino acid ester), which exhibit much less activity in A549 cells (around $25 \mu\text{M}$),^{20f} or with thiourea silver species, which present similar activity in the same cells (around $7 \mu\text{M}$).^{27c} Additionally, those complexes bearing pentafluorophenyl groups, such as **25–27**, do not show in our experience cytotoxic activity and they have not been measured. The homoleptic silver and gold complexes exhibit low activity. The best one corresponds to the silver complex **17** with an IC_{50} of 25.16 ± 1.8 .

On comparison with the reference drug cisplatin, which exhibits a high value after 24 h (in water) and only a moderate value after 48 h, our silver derivatives present much higher activity at 24 h.

In summary, we can conclude that these types of organic ligands do not exhibit antiproliferative activity in cancer cells but their combination with metal compounds strongly enhance their activity, especially when a phosphine ancillary ligand is present.

Experimental

Experimental details

Mass spectra were recorded on a BRUKER ESQUIRE 3000 PLUS, with the electrospray (ESI) technique. ^1H , $^{13}\text{C}\{^1\text{H}\}$, $^{31}\text{P}\{^1\text{H}\}$ and ^{19}F NMR spectra were recorded at room temperature on a BRUKER AVANCE 400 spectrometer (^1H , 400 MHz, $^{13}\text{C}\{^1\text{H}\}$, 101 MHz, $^{31}\text{P}\{^1\text{H}\}$, 162 MHz and ^{19}F 376.5 MHz) with chemical shifts (δ , ppm) reported relative to the solvent peaks of the deuterated solvent.

Crystal structure determination

Crystals were mounted in inert oil on glass fibres and transferred to the cold gas stream of an Xcalibur Oxford diffraction or a Smart APEX CCD diffractometer equipped with a low-temperature attachment. Data were collected using monochromated $\text{MoK}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$). Scan type ω . Absorption corrections based on multiple scans were applied using SADABS³³ or spherical harmonics implemented in the SCALE3 ABSPACK scaling algorithm.³⁴ The structures were solved by direct methods and refined on F^2 using the program SHELXT-2016,³⁵ or by using Olex2 as the graphical interface.³⁶ All non-hydrogen atoms were refined anisotropically.

Materials and procedures

The starting materials $[\text{AuCl}(\text{PPh}_3)]$, $[\text{Ag}(\text{OTf})(\text{PPh}_3)]$, $[\text{Au}(\text{tht})_2]\text{OTf}$ and $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ were prepared according to published procedures.³⁷ All other reagents were commercially available

Table 1 IC_{50} values for complexes **10–18** and **22–24**, after 24 h of incubation with A549 cells

Compounds	IC_{50} (μM)
Cisplatin	29.21 ± 1.92 (48 h) ³¹ 114.2 ± 9.1 (24 h) ³²
Ligands	
10	>50
11	>50
12	>50
$[\text{Ag}(\text{L})(\text{PPh}_3)]\text{OTf}$	
13	3.97 ± 0.63
14	5.78 ± 0.36
15	6.26 ± 0.29
$[\text{Ag}(\text{L})_2]\text{OTf}$	
16	>50
17	25.16 ± 1.8
18	>50
$[\text{Au}(\text{L})_2]\text{OTf}$	
22	39.71 ± 1.32
23	42.36 ± 0.56
24	>50



and used without further purification. Solvents were dried with a SPS solvent purification system.

Cell culture

A549 (human lung carcinoma) cells were grown in DMEM medium supplemented with 5% fetal bovine serum (SFB), 100 mg mL⁻¹ pyruvate, 2.2 g L⁻¹ sodium bicarbonate, 5 mL non-essential amino acids (Invitrogen), 2 mM L-glutamine, and antibiotics. Cells were grown at 2×10^5 cells per mL, in 96-well flat-bottomed microplates and allowed to attach for 24 h prior to addition of compounds.

Antiproliferative studies: MTT assay

The complexes were dissolved in DMSO and added at different concentrations between 1 and 50 μ M in quadruplicate. Each complex was added in four different concentrations of the drugs obtained by different dilutions and with a column of negative controls (cells with a DMSO concentration similar to which contain the solutions of the complexes) and another column of blank (medium only).

To make these dilutions, between 2 and 3 mg of the compound were weighed and dissolved in DMSO to obtain a stock solution of 0.1 M. It was then diluted to obtain a solution of 0.001 M in medium (DMEM) and finally from this, dilutions were made to obtain the different concentrations that would be added in the 96-well plate.

Cells were incubated with our compounds for 24 h at 37 °C and with a 5% CO₂ atmosphere. Then, 10 μ L of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazole) of concentration 5 mg mL⁻¹ was added to each well and plates were incubated for 2 h under the same conditions as described above (37 °C and with a 5% CO₂ atmosphere). The crystals of blue formazan that formed are adhered to the bottom of the well, so before dissolving them it is necessary to remove the supernatant by the “flipping” technique and then, dissolve them in DMSO (100 μ L per well). The plates are shaken to help to dissolve the formazan precipitated. The optical density was measured at 550 nm using a 96-well multiscanner autoreader (ELISA). The IC₅₀ was calculated by nonlinear regression analysis.

Synthesis of thioureas 3

1-Phenyl-3-(prop-2-ynyl)thiourea (3). To a mixture of propargylamine **1a** (6.4 μ L, 0.1 mmol) in CH₃CN (10 mL), isothiocyanatobenzene **2a** (11.9 μ L, 0.1 mmol) was added. The mixture was stirred for 24 h. After that, the dark brown solid was filtered under vacuum, washed with cold CH₃CN (3×1 mL), and dried in air. Product **3** was obtained with 80% yield. ¹H NMR (ppm) (400 MHz, CDCl₃): δ = 8.10 (br s, 1H, Ph-NH-C=S), 7.45 (m, 2H, Ph_{ortho}), 7.33 (m, 1H, Ph_{para}), 7.23 (m, 1H, Ph_{meta}), 6.10 (br s, 1H, C=S-NH-CH₂), 4.46 (dd, 2H, ³J_{HH} = 5.0 Hz, ⁴J_{HH} = 2.6 Hz, NH-CH₂-C), 2.26 (t, 1H, ⁴J_{HH} = 2.6 Hz, CH). ¹³C{¹H} APT NMR (ppm) (400 MHz, CDCl₃): δ = 180.7 (s, 1C, C=S), 135.8 (s, 1C, Ph_{ipso}), 130.4 (s, 2C, Ph_{ortho}), 127.7 (s, 1C, Ph_{para}), 125.4 (s, 1C, Ph_{meta}), 77.4 (s, 1C, C-CH₂), 72.6 (s, 1C, CH), 35.2 (s, 1C, C-CH₂-NH).

Synthesis of thiazolidines 4–9

N-(5-Methylenethiazolidin-2-ylidene)aniline (4). To a solution of propargylamine **1a** (6.4 μ L, 0.1 mmol) in EtOH (5 mL), isothiocyanate **2a** (11.9 μ L, 0.1 mmol) was added. The mixture was stirred for 24 h. After that, a light brown solid was filtered under vacuum, washed with cold EtOH (3×1 mL), and dried in air. Product **4** was obtained with 88% yield. ¹H NMR (ppm) (300 MHz, CDCl₃): δ = 7.28 (m, 2H, H_{ortho}Ph), 7.17 (m, 2H, H_{meta}Ph), 7.07 (tt, 1H, ³J_{HH} = 7.3 Hz, ⁴J_{HH} = 2.4 Hz, H_{para}Ph), 5.20 (dd, 1H, ²J_{HH} = 3.9 Hz, ³J_{HH} = 2.2 Hz, C=CH₂), 5.12 (dd, 1H, ²J_{HH} = 4.3 Hz, ³J_{HH} = 2.6 Hz, C=CH₂), 4.58 (t, 2H, ⁴J_{HH} = 2.3 Hz, N-CH₂). ¹³C{¹H} APT NMR (ppm) (101 MHz, CDCl₃): δ = 158.4 (s, 1C, N-C=N), 145.1 (s, 1C, C_{ipso}Ph), 144.2 (s, 1C, C=CH₂), 129.1 (s, 2C, C_{ortho}Ph), 123.6 (s, 1C, C_{para}Ph), 121.0 (s, 2C, C_{meta}Ph), 104.2 (s, 1C, C=CH₂), 59.3 (s, 1C, N=CH₂).

N-(5-Methylene-3-(prop-2-ynyl)thiazolidin-2-ylidene)aniline (5). To a mixture of propargylamine **1b** (10.3 μ L, 0.1 mmol) in EtOH (10 mL), isothiocyanate **2a** (11.9 μ L, 0.1 mmol) was added. The mixture was stirred for 24 h. After evaporation of the solvent under vacuum, a yellow oil was obtained. Product **5** was obtained with >95% yield. ¹H NMR (ppm) (400 MHz, CDCl₃): δ = 7.30 (m, 2H, H_{ortho}Ph), 7.09 (m, 1H, H_{para}Ph), 6.95 (m, 2H, H_{meta}Ph), 5.28 (dd, 1H, ²J_{HH} = 3.8 Hz, ³J_{HH} = 2.1 Hz, C=CH₂), 5.12 (dd, 1H, ²J_{HH} = 4.0 Hz, ³J_{HH} = 2.4 Hz, C=CH₂), 4.38 (d, 2H, ⁴J_{HH} = 2.5 Hz, CH₂-C≡CH), 4.37 (t, 2H, ⁴J_{HH} = 2.3 Hz, N-CH₂), 2.35 (t, 1H, ⁴J_{HH} = 2.5 Hz, C≡CH). ¹³C{¹H} APT NMR (ppm) (101 MHz, CDCl₃): δ = 156.3 (s, 1C, N-C=N), 151.2 (s, 1C, C_{ipso}Ph), 137.0 (s, 1C, C=CH₂), 129.0 (s, 2C, C_{ortho}Ph), 123.6 (s, 1C, C_{para}Ph), 121.9 (s, 2C, C_{meta}Ph), 106.0 (s, 1C, C=CH₂), 77.4 (s, 1C, C≡CH), 72.9 (s, 1C, C≡CH), 55.1 (s, 1C, N-CH₂), 33.3 (s, 1C, CH₂-C≡CH).

N-(3-Methyl-5-methylenethiazolidin-2-ylidene)aniline (6). To a mixture of propargylamine **1c** (8.2 μ L, 0.1 mmol) in EtOH (5 mL), isothiocyanate **2a** (11.9 μ L, 0.1 mmol) was added. The mixture was stirred for 24 h. After evaporation of the solvent under vacuum, a yellow oil was obtained. Product **6** was obtained with >95% yield. ¹H NMR (ppm) (400 MHz, CDCl₃): δ = 7.27 (m, 2H, H_{ortho}Ph), 7.05 (m, 1H, H_{para}Ph), 6.93 (m, 2H, H_{meta}Ph), 5.20 (dd, 1H, ²J_{HH} = 3.7 Hz, ³J_{HH} = 2.1 Hz, C=CH₂), 5.07 (dd, 1H, ²J_{HH} = 4.0 Hz, ³J_{HH} = 2.4 Hz, C=CH₂), 4.28 (t, 2H, ⁴J_{HH} = 2.3 Hz, N-CH₂), 3.09 (s, 3H, CH₃). ¹³C{¹H} APT NMR (ppm) (101 MHz, CDCl₃): δ = 151.8 (s, 2C, C_{ipso}Ph, N-C=N), 137.7 (s, 1C, C=CH₂), 129.0 (s, 2C, C_{ortho}Ph), 123.4 (s, 1C, C_{para}Ph), 122.2 (s, 2C, C_{meta}Ph), 105.3 (s, 1C, C=CH₂), 58.1 (s, 1C, N-CH₂), 33.2 (s, 1C, CH₃).

3-Fluoro-N-(5-methylenethiazolidin-2-ylidene)aniline (7). To a mixture of propargylamine **1a** (8.2 μ L, 0.1 mmol) in EtOH (5 mL), isothiocyanate **2b** (12 μ L, 0.1 mmol) was added. The mixture was stirred for 24 h. After that, a white solid was filtered under vacuum, washed with cold EtOH (3×1 mL), and dried in air. Product **7** was obtained with 85% yield. ¹H NMR (ppm) (400 MHz, CDCl₃): δ = 7.23 (m, 1H, Ar), 6.98 (td, 1H, ⁴J_{HH} = 10.2 Hz, ⁵J_{HH} = 2.1 Hz, Ar), 6.87 (ddd, 1H, ³J_{HH} = 8.0 Hz, ⁴J_{HH} = 2.0 Hz, ⁴J_{HH} = 0.9 Hz, Ar), 6.75 (tdd, 1H, ³J_{HH} = 8.4 Hz, ⁴J_{HH} = 2.5 Hz, ⁴J_{HH} = 0.9 Hz, Ar), 5.23 (dd, 1H, ²J_{HH} = 4.1 Hz,



$^3J_{\text{HH}} = 2.3$ Hz, $\text{C}=\text{CH}_2$), 5.14 (dd, 1H, $^2J_{\text{HH}} = 4.4$ Hz, $^3J_{\text{HH}} = 2.6$ Hz, $\text{C}=\text{CH}_2$), 4.56 (t, 2H, $^3J_{\text{HH}} = 2.4$ Hz, $\text{N}-\text{CH}_2$). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, CDCl_3): $\delta = 163.3$ (d, 1C, C_{ipsoPh} , $^1J_{\text{CF}} = 243.7$ Hz), 162.1 (s, 1C, $\text{N}-\text{C}=\text{N}$), 149.8 (s, 1C), 130.2 (d, 1C, $^3J_{\text{CF}} = 9.6$ Hz), 116.4 (d, 1C, $^4J_{\text{CF}} = 2.5$ Hz), 110.2 (d, 1C, $^2J_{\text{CF}} = 21.12$ Hz), 108.2 (d, 1C, $^2J_{\text{CF}} = 23.8$ Hz), 104.8 (s, 1C, CH_2), 60.44 (s, 1C, CH_2-N). ^{19}F NMR (ppm) (376.5 MHz, CDCl_3): $\delta = -108.5$ (m, 1F, Ph-F).

3-Fluoro-N-(5-methylene-3-(prop-2-ynyl)thiazolidin-2-ylidene)aniline (8). To a mixture of propargylamine **1b** (10.3 μL , 0.1 mmol) in EtOH (5 mL), isothiocyanate **2b** (12 μL , 0.1 mmol) was added. The mixture was stirred for 24 h. After evaporation of the solvent under vacuum, a yellow oil was obtained. Product **8** was obtained with >95% yield. ^1H NMR (ppm) (400 MHz, CDCl_3): $\delta = 7.22$ (m, 1H, Ar), 6.72 (m, 3H, Ar), 5.27 (m, 1H, $\text{C}=\text{CH}_2$), 5.12 (m, 1H, $\text{C}=\text{CH}_2$), 4.38 (t, 2H, $^4J_{\text{HH}} = 2.3$ Hz, $\text{N}-\text{CH}_2$), 4.36 (d, 2H, $^4J_{\text{HH}} = 2.5$ Hz, $\text{CH}_2-\text{C}\equiv\text{CH}$), 2.33 (t, 1H, $^4J_{\text{HH}} = 2.5$ Hz, $\text{CH}_2-\text{C}\equiv\text{CH}$). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, CDCl_3): $\delta = 163.2$ (d, 1C, $^1J_{\text{CF}} = 243.7$ Hz, C_{FPh}), 156.7 (s, 1C, $\text{N}-\text{C}=\text{N}$), 151.9 (d, 1C, $^3J_{\text{CF}} = 9.6$ Hz, C_{ipsoPh}), 136.7 (s, 1C, $\text{C}=\text{CH}_2$), 130.1 (d, 1C, $^3J_{\text{CF}} = 9.6$ Hz, $C-\text{Ar}$), 117.7 (d, 1C, $^4J_{\text{CF}} = 2.7$ Hz, $C-\text{Ar}$), 110.3 (d, 1C, $^2J_{\text{CF}} = 21.1$ Hz, $C-\text{Ar}$), 109.3 (d, 1C, $^2J_{\text{CF}} = 22.0$ Hz, $C-\text{Ar}$), 106.3 (s, 1C, $\text{C}=\text{CH}_2$), 77.7 (s, 1C, $\text{C}=\text{CH}$), 73.1 (s, 1C, $\text{C}=\text{CH}$), 55.2 (s, 1C, $\text{N}-\text{CH}_2$), 33.3 (s, 1C, $\text{CH}_2-\text{C}\equiv\text{CH}$). $^{19}\text{F}\{^1\text{H}\}$ NMR (ppm) (376.5 MHz, CDCl_3): $\delta = -112.9$ (m, 1F, Ph-F).

3-Fluoro-N-(3-methyl-5-methylenethiazolidin-2-ylidene)aniline (9). To a mixture of propargylamine **1c** (8.2 μL , 0.1 mmol) in EtOH (5 mL), isothiocyanate **2b** (12 μL , 0.1 mmol) was added. The mixture was stirred for 24 h. After evaporation of the solvent under vacuum, a colourless oil was obtained. Product **9** was obtained with >95% yield. ^1H NMR (ppm) (400 MHz, CDCl_3): $\delta = 7.20$ (m, 1H, Ar), 6.74 (m, 1H, Ar), 6.71 (m, 1H, Ar), 6.65 (m, 1H, Ar), 5.21 (m, 1H, $\text{C}=\text{CH}_2$), 5.09 (m, 1H, $\text{C}=\text{CH}_2$), 4.28 (t, 2H, $^4J_{\text{HH}} = 2.3$ Hz, $\text{N}-\text{CH}_2$), 3.07 (s, 3H, CH_3). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, CDCl_3): $\delta = 163.8$ (d, 1C, $^1J_{\text{CF}} = 245.3$ Hz, C_{FPh}), 157.8 (s, 1C, $\text{N}-\text{C}=\text{N}$), 154.0 (d, 1C, $^3J_{\text{CF}} = 9.6$ Hz, C_{ipsoPh}), 137.6 (s, 1C, $\text{C}=\text{CH}_2$), 130.3 (d, 1C, $^3J_{\text{CF}} = 9.7$ Hz, $C-\text{Ar}$), 118.3 (d, 1C, $^4J_{\text{CF}} = 2.7$ Hz, $C-\text{Ar}$), 110.2 (d, 1C, $^2J_{\text{CF}} = 21.3$ Hz, $C-\text{Ar}$), 109.7 (d, 1C, $^2J_{\text{CF}} = 21.9$ Hz, $C-\text{Ar}$), 105.9 (s, 1C, $\text{C}=\text{CH}_2$), 57.9 (s, 1C, $\text{N}-\text{CH}_2$), 32.8 (s, 1C, CH_3). ^{19}F NMR (ppm) (376.5 MHz, CDCl_3): $\delta = -113.0$ (m, 1F, Ph-F).

Synthesis of thiazolidine-thioamide derivative 10. To a solution of propargylamine **1a** (12.8 μL , 0.2 mmol) in ethanol (20 mL), an excess of phenyl isothiocyanate **2a** (71.5 μL , 0.6 mmol) was added and the reaction was stirred for 72 h. A white precipitate was formed which was filtered under vacuum, washed with cold EtOH (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product **10** was obtained with 83% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.67$ (m, 2H, Ph), 7.42 (m, 4H, Ph), 7.23 (m, 2H, Ph), 7.13 (m, 2H, Ph), 5.52 (m, 1H, $\text{C}=\text{CH}_2$), 5.38 (t, 2H, $^4J_{\text{HH}} = 2.4$ Hz, $\text{N}-\text{CH}_2$), 5.24 (dt, 1H, $^2J_{\text{HH}} = 2.7$ Hz, $^4J_{\text{HH}} = 2.2$ Hz, $\text{C}=\text{CH}_2$). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 178.6$ (s, 1C, $\text{C}=\text{S}$), 158.6 (s, 1C, $\text{N}=\text{C}-\text{N}$), 149.2 (s, 1C, $C_{\text{ipsoPh-N}}$), 139.9 (s, 1C, $C_{\text{ipsoPh-NH}}$),

132.8 (s, 1C, $\text{CH}_2=\text{C}$), 130.3 (s, 2C, Ph), 129.4 (s, 2C, Ph), 126.8 (s, 2C, Ph), 126.3 (s, 1C, Ph), 125.2 (s, 1C, Ph), 122.3 (s, 2C, Ph), 107.9 (s, 1C, $\text{C}=\text{CH}_2$), 60.9 (s, 1C, $\text{N}-\text{CH}_2$).

Synthesis of thiazolidine-thioamide derivative 11. To a solution of propargylamine **1a** (12.8 μL , 0.2 mmol) in ethanol (20 mL), an excess of 3-fluorophenyl isothiocyanate **2b** (72.0 μL , 0.6 mmol) was added and the reaction was stirred for 72 h. A white precipitate was formed, which was filtered under vacuum, washed with cold EtOH (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product **11** was obtained with 81% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.79$ –6.97 (m, 8H, Ar), 5.54 (m, 1H, $\text{C}=\text{CH}_2$), 5.38 (t, 2H, $J_{\text{HH}} = 2.4$ Hz, $\text{N}-\text{CH}_2$), 5.28 (m, 1H, $\text{C}=\text{CH}_2$). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 165.5$ (s, 1C, $\text{C}=\text{S}$), 157.9 (s, 1C, $\text{N}=\text{C}-\text{N}$), 150.0 (s, 1C, C_{ipsoPh}), 140.9 (s, 1C, C_{ipsoPh}), 131.2 (s, 1C, $\text{CH}_2=\text{C}$), 131.9 (d, 1C, $^3J_{\text{CF}} = 9.5$ Hz, C_{metaPh}), 130.9 (d, 1C, $^3J_{\text{CF}} = 9.4$ Hz, C_{metaPh}), 120.8 (d, 1C, $^4J_{\text{CF}} = 3.0$ Hz, C_{paraPh}), 118.4 (d, 1C, $^4J_{\text{CF}} = 3.0$ Hz, C_{paraPh}), 113.3 (d, 1C, $^2J_{\text{CF}} = 21.4$ Hz, C_{orthoPh}), 112.9 (d, 1C, $^2J_{\text{CF}} = 21.3$ Hz, C_{orthoPh}), 112.0 (d, 1C, $^2J_{\text{CF}} = 25.6$ Hz, C_{orthoPh}), 109.8 (d, 1C, $^2J_{\text{CF}} = 23.3$ Hz, C_{orthoPh}), 108.3 (s, 1C, $\text{C}=\text{CH}_2$), 61.0 (s, 1C, $\text{N}-\text{CH}_2$). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = -111.0$ (m, 1F, Ph-F), -111.5 (m, 1F, Ph-F).

Synthesis of thiazolidine thioamide derivative 12. To a solution of propargylamine **1a** (12.8 μL , 0.2 mmol) in ethanol (20 mL), an excess of benzyl isothiocyanate **2c** (78 μL , 0.6 mmol) was added and the reaction was stirred for 72 h. A white precipitate was formed, which was filtered under vacuum, washed with cold EtOH (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product **12** was obtained with 84% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 12.71$ (br s, 1H, NH), 7.32–7.29 (m, 5H, Ph), 7.25–7.23 (m, 3H, Ph), 7.16–7.13 (m, 2H, Ph), 5.52 (m, 1H, $\text{C}=\text{CH}_2$), 5.30 (m, 1H, $\text{C}=\text{CH}_2$), 5.26 (t, 2H, $^4J_{\text{HH}} = 2.3$ Hz, $\text{N}-\text{CH}_2$), 4.81 (d, 2H, $^3J_{\text{HH}} = 4.9$ Hz, $\text{NH}-\text{CH}_2$), 4.41 (s, 2H, $=\text{N}-\text{CH}_2-\text{Ph}$). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 180.2$ (s, 1C, $\text{C}=\text{S}$), 156.9 (s, 1C, $\text{N}-\text{C}=\text{N}$), 139.5 (s, 1C, $C_{\text{ipsoPh-CH}_2-\text{N}}$), 138.2 (s, 1C, $C_{\text{ipsoPh-CH}_2-\text{NH}}$), 133.2 (s, 1C, $\text{C}=\text{CH}_2$), 129.5–127.8 (s, 10C, Ph), 107.8 (s, 1C, CH_2), 60.4 (s, 1C, CH_2-N), 59.0 (s, 1C, $\text{N}-\text{CH}_2-\text{Ph}$), 50.2 (s, 1C, $\text{NH}-\text{CH}_2$).

Synthesis of silver complex 13. To a solution of compound **10** (32.5 mg, 0.1 mmol) in CH_2Cl_2 (10 mL), $[\text{Ag}(\text{OTf})(\text{PPh}_3)]$ (51.9 mg, 0.1 mmol) was added and the reaction was stirred for 2 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 mL and then, hexane (10 mL) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product **13** was obtained with 72% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.54$ (m, 21H, Ph), 7.26 (m, 1H, Ph), 7.20 (m, 1H, Ph), 7.12 (m, 2H, Ph), 5.59 (br s, 1H, $\text{C}=\text{CH}_2$), 5.46 (br t, 2H, $\text{N}-\text{CH}_2$), 5.35 (br s, 1H, $\text{C}=\text{CH}_2$). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 164.8$ (s, 1C), 156.5 (s, 1C), 151.2 (s, 1C), 134.8 (d, 6C, $^3J_{\text{H-P}} = 15.6$ Hz, C_{orthoPPh_3}), 130.2 (d, 6C, $^3J_{\text{H-P}} = 9.6$ Hz, C_{metaPPh_3}), 132.2–130.4 (s, 14C, $C_{\text{paraPPh}_3} + \text{Ph}$), 129.1 (s, 1C, Ph), 126.9 (s, 1C, Ph), 126.5 (s, 1C, Ph), 122.1 (s, 1C, Ph), 109.0 (s, 1C, $\text{C}=\text{CH}_2$), 60.8 (s, 1C,



N-CH₂). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 12.0 (br s, 1P, *PPh*₃). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{35}\text{H}_{30}\text{AgN}_3\text{PS}_2]^+$ 694.0664, found 694.0647.

Synthesis of silver complex 14. To a solution of compound **11** (36.1 mg, 0.1 mmol) in CH_2Cl_2 (10 ml), $[\text{Ag}(\text{OTf})(\text{PPh}_3)]$ (51.9 mg, 0.1 mmol) was added and the reaction was stirred for 2 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **14** was obtained with 69% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.51–7.34 (m, 20H, *Ar* + *PPh*₃), 7.06–6.95 (m, 3H, *Ar*), 5.57 (s, 1H, *C=CH*₂), 5.46 (s, 2H, *C=CH*₂), 5.36 (m, 1H, *N-CH*₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 165.2 (s, 1C, *C=S*), 162.7 (s, 1C, *C_q*), 151.1 (s, 1C), 149.5 (s, 1C), 134.7 (d, 6C, $^2J_{\text{CP}}$ = 15.3 Hz, *C_{ortho}PPh*₃), 132.1 (s, 3C, *C_{para}PPh*₃), 131.4 (d, 3C, $^1J_{\text{CP}}$ = 35.1 Hz, *C_{ipso}PPh*₃), 130.2 (d, 6C, $^3J_{\text{CP}}$ = 8.6 Hz, *C_{meta}PPh*₃), 118.3–109.6 (m, 10C, *Ph-F*), 109.5 (s, 2C, *C=CH*₂), 61.0 (s, 1C, *N-CH*₂). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): δ = –79.9 (s, 1F, *OTf*), –113.8 (m, 1F, *Ph-F*), –114.3 (m, 1F, *Ph-F*). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 12.2 (s, 1P, *PPh*₃).

Synthesis of silver complex 15. To a solution of compound **12** (35.3 mg, 0.1 mmol) in CH_2Cl_2 (10 ml), $[\text{Ag}(\text{OTf})(\text{PPh}_3)]$ (51.9 mg, 0.1 mmol) was added and the reaction was stirred for 2 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **15** was obtained with 68% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.55–7.26 (m, 25H, *Ph*), 5.46 (m, 1H, *C=CH*₂), 5.38 (t, 2H, *N-CH*₂), 5.34 (m, 1H, *C=CH*₂), 4.95 (d, 2H, J_{HH} = 4.8 Hz, *NH-CH*₂), 4.44 (s, 2H, *=N-CH*₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 177.9 (s, 1C, *C=S*), 158.2 (s, 1C, *N-C=N*), 138.8 (s, 1C, *C_{ipso}Ph-CH*₂-N), 136.5 (s, 1C, *C_{ipso}Ph-CH*₂-NH), 133.2 (s, 1C, *C=CH*₂), 134.7 (d, 6C, $^2J_{\text{HP}}$ = 15.3 Hz, *C_{ortho}PPh*₃), 132.1 (s, 3C, *C_{para}PPh*₃), 131.6 (d, 3C, *C_{ipso}PPh*₃, $^1J_{\text{HP}}$ = 31.5 Hz), 130.2 (d, 6C, $^3J_{\text{HP}}$ = 8.4 Hz, *C_{meta}PPh*₃), 129.8–128.1 (m, 10C, *Ph*), 108.9 (s, 1C, *C=CH*₂), 60.6 (s, 1C, *N-CH*₂), 59.2 (s, 1C, *Ph-CH*₂-N=C), 51.3 (s, 1C, *Ph-CH*₂-NH). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 11.5 (s, 1P, *PPh*₃). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{37}\text{H}_{34}\text{AgN}_3\text{PS}_2]^+$ 724.0975, found 724.0977.

Synthesis of silver complex 16. To a solution of compound **10** (65.0 mg, 0.2 mmol) in CH_2Cl_2 (10 ml), AgOTf (25.7 mg, 0.1 mmol) was added and the reaction was stirred for 1 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **16** was obtained with 87% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.61 (m, 4H, *Ph*), 7.41 (m, 10H, *Ph*), 7.26 (br t, 2H, *Ph*), 7.11 (br d, 4H, *Ph*), 5.64 (br s, 2H, *C=CH*₂), 5.42 (br s,

4H, *N-CH*₂), 5.39 (br s, 2H, *C=CH*₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 178.8 (s, 2C, *C=S*), 160.1 (s, 2C, *C_q*), 148.1 (s, 2C, *C_{ipso}*), 137.5 (s, 2C, *C_{ipso}*), 132.0 (s, 2C, *C_{quaternary}*), 131.1–122.1 (s, 20C, *Ph*), 109.1 (s, 2C, *CH*₂), 60.7 (s, 2C, *N-CH*₂).

Synthesis of silver complex 17. To a solution of compound **11** (72.0 mg, 0.2 mmol) in CH_2Cl_2 (10 ml), AgOTf was added (25.7 mg, 0.1 mmol) and the solution was stirred for 1 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **17** was obtained with 90% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.61–6.92 (m, 16H, *Ar*), 5.64 (br s, 2H, *C=CH*₂), 5.44 (br s, 4H, *N-CH*₂), 5.42 (br s, 2H, *CH*₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 168.1 (s, 2C, *C=S*), 163.3 (s, 1C), 157.4 (s, 1C), 149.1 (s, 1C), 132.5 (d, 4C, $^3J_{\text{HF}}$ = 9.2 Hz, *Ar*), 132.2 (d, 4C, $^3J_{\text{HF}}$ = 9.3 Hz, *Ar*), 123.1 (s, 2C, *Ar*), 118.2 (s, 2C, *Ar*), 116.4 (d, 2C, $^2J_{\text{HF}}$ = 21.9 Hz, *Ar*), 114.6 (d, 2C, $^2J_{\text{HF}}$ = 24.8 Hz, *Ar*), 113.6 (d, 2C, $^2J_{\text{HF}}$ = 22.1 Hz, *Ar*), 109.7 (d, 2C, $^2J_{\text{HF}}$ = 23.9 Hz, *Ar*), 109.5 (s, 2C, *C=CH*₂), 60.8 (s, 2C, *N-CH*₂). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): δ = –79.9 (s, 1F, *OTf*), –112.2 (m, 2F, *Ph*), –114.2 (m, 2F, *Ph*). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{34}\text{H}_{26}\text{AgF}_4\text{N}_6\text{S}_4]^+$ 831.0079, found 831.0056.

Synthesis of silver complex 18. To a solution of compound **12** (72.0 mg, 0.2 mmol) in CH_2Cl_2 (10 ml), AgOTf (25.7 mg, 0.1 mmol) was added and the reaction was stirred for 1 h at room temperature with the exclusion of light. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **18** was obtained with 88% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.33 (m, 10H, *Ph*), 7.24 (m, 6H, *Ph*), 7.13 (m, 4H, *Ph*), 5.52 (m, 2H, *C=CH*₂), 5.39 (m, 4H, *N-CH*₂), 5.35 (m, 2H, *C=CH*₂), 4.98 (m, 4H, *NH-CH*₂), 4.43 (s, 4H, *=N-CH*₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 176.5 (s, 2C, *C=S*), 156.5 (s, 2C, *N-C=N*), 138.7 (s, 2C, *C_{ipso}Ph-CH*₂-N), 136.0 (s, 2C, *C_{ipso}Ph-CH*₂-NH), 132.0 (s, 2C, *C=CH*₂), 130.0–128.1 (s, 20C, *Ph*), 109.0 (s, 2C, *C=CH*₂), 60.5 (s, 2C, *N-CH*₂), 59.2 (s, 1C, *Ph-CH*₂-N=C), 51.5 (s, 1C, *Ph-CH*₂-NH). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{38}\text{H}_{38}\text{AgN}_6\text{S}_4]^+$ 815.1083, found 815.1099.

Synthesis of gold complexes 19. To a solution of $[\text{Au}(\text{PPh}_3)(\text{MeCN})]\text{SbF}_6$ (70.4 mg, 0.1 mmol) which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound **10** (32.5 mg, 0.1 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **19** was obtained with 80% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.62 (m, 9H, *Ph* + *H_{meta}PPh*₃ + *H_{para}PPh*₃), 7.43 (m, 8H, *Ph* + *H_{ortho}PPh*₃), 7.30 (m, 3H, *Ph*), 7.12 (m, 3H, *Ph*), 5.66 (m, 1H, *CH*₂), 5.52 (m, 2H, *CH*₂-N), 5.42 (m, 1H, *CH*₂).



$^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 177.4 (s, 1C, C=S), 160.8 (s, 1C, N-C=N), 147.7 (s, 1C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-N}$), 138.1 (s, 1C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-NH}$), 135.3 (s, 2C, $C_{\text{ortho}}\text{Ph}$), 135.2 (d, 6C, $C_{\text{ortho}}\text{PPh}_3$, $^2J_{\text{C-P}} = 13.2$ Hz), 133.4 (d, 3C, $C_{\text{para}}\text{PPh}_3$, $^4J_{\text{C-P}} = 2.7$ Hz), 131.6 (s, 1C, C=CH₂), 130.9 (s, 2C, $C_{\text{meta}}\text{Ph}$), 130.5 (d, 6C, $C_{\text{meta}}\text{PPh}_3$, $^3J_{\text{C-P}} = 11.8$ Hz), 130.5 (s, 2C, $C_{\text{ortho}}\text{Ph}$), 128.7 (d, 3C, $C_{\text{ipso}}\text{PPh}_3$, $^1J_{\text{C-P}} = 63.2$ Hz), 127.2 (s, 2C, $C_{\text{para}}\text{Ph}$), 122.1 (s, 2C, $C_{\text{meta}}\text{Ph}$), 109.5 (s, 2C, CH₂), 60.5 (s, 1C, CH₂-N). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 45.2 (s, 1P, $\text{Au}(\text{PPh}_3)_2$), 37.5 (s, 1P, PPh_3). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{35}\text{H}_{30}\text{AuN}_3\text{PS}_2]^+$ 784.1279, found 784.1277.

Synthesis of gold complexes 20. To a solution of $[\text{Au}(\text{PPh}_3)(\text{MeCN})]\text{SbF}_6$ (70.4 mg, 0.1 mmol) which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound 11 (36.1 mg, 0.1 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product 20 was obtained with 83% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.63–7.48 (m, 16H, PPh_3 + H_3), 7.33 (m, 3H, H_1 + H_4 + H_7), 7.06 (m, 1H, H_2), 6.98 (m, 1H, H_8), 6.90 (m, 2H, H_5 + H_6), 5.68 (m, 1H, CH₂), 5.53 (m, 2H, CH₂-N), 5.44 (m, 1H, CH₂). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 177.5 (s, 1C, C=S), 164.9 (d, 1C, $C_{\text{ipso}}\text{Ph-F-NH}$, $^1J_{\text{C-F}} = 247.7$ Hz), 162.4 (d, 1C, $C_{\text{ipso}}\text{Ph-F-N}$, $^1J_{\text{C-F}} = 245.8$ Hz), 161.4 (s, 1C, N-C=N), 149.2 (d, 1C, $C_{\text{ipso}}\text{Ph-F-N}$, $^3J_{\text{C-F}} = 9.2$ Hz), 139.5 (d, 1C, $C_{\text{ipso}}\text{Ph-F-NH}$, $^3J_{\text{C-F}} = 10.0$ Hz), 134.9 (d, 6C, $C_{\text{ortho}}\text{PPh}_3$, $^2J_{\text{C-P}} = 13.8$ Hz), 133.4 (d, 3C, $C_{\text{para}}\text{PPh}_3$, $^4J_{\text{C-P}} = 2.6$ Hz), 132.5 (d, 1C, C_7 , $^3J_{\text{C-F}} = 9.2$ Hz), 132.1 (d, 1C, C_3 , $^3J_{\text{C-F}} = 9.4$ Hz), 131.2 (s, 1C, C=CH₂), 130.4 (d, 6C, $C_{\text{meta}}\text{PPh}_3$, $^3J_{\text{C-P}} = 11.9$ Hz), 129.0 (d, 3C, $C_{\text{ipso}}\text{PPh}_3$, $^1J_{\text{C-P}} = 61.0$ Hz), 123.4 (s, 1C, C_8), 118.2 (d, 1C, C_4 , $^4J_{\text{C-F}} = 2.7$ Hz), 116.6 (d, 1C, C_6 , $^2J_{\text{C-F}} = 21.0$ Hz), 115.1 (d, 1C, C_1 , $^2J_{\text{C-F}} = 24.4$ Hz), 113.9 (d, 1C, C_2 , $^2J_{\text{C-F}} = 21.1$ Hz), 109.9 (s, 1C, CH₂), 109.5 (d, 1C, C_5 , $^2J_{\text{C-F}} = 23.5$ Hz), 60.7 (s, 1C, CH₂-N). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): δ = -118.8 (s, 1F, F), -113.9 (s, 1F, F), -127.5 (m, 6F, SbF_6). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 45.2 (s, 1P, $\text{Au}(\text{PPh}_3)_2$), 37.4 (s, 1P, PPh_3). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{35}\text{H}_{28}\text{AuF}_2\text{N}_3\text{PS}_2]^+$ 820.1091, found 820.1088.

Synthesis of gold complexes 21. To a solution of $[\text{Au}(\text{PPh}_3)(\text{MeCN})]\text{SbF}_6$ (70.4 mg, 0.1 mmol) which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound 12 (35.3 mg, 0.1 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product 21 was obtained with 79% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.73–7.17 (m, 26H, Ph + PPh_3 + NH), 5.58 (m, 1H, CH₂), 5.50 (t, 2H, CH₂-N, $^4J_{\text{H-H}} = 2.3$ Hz), 5.43 (m, 1H, CH₂), 5.13 (d, 2H, CH₂-NH, $^3J_{\text{H-H}} = 5.1$ Hz), 4.48 (s, 2H, $\text{Ph-CH}_2\text{-N}$). ^{13}C APT (ppm) (100 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 175.2 (s, 1C, C=S), 158.9 (s, 1C, N-C=N), 138.4 (s, 1C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-N}$), 136.0 (s, 1C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-NH}$), 135.0 (d, 6C, $C_{\text{ortho}}\text{PPh}_3$, $^2J_{\text{C-P}} = 13.8$ Hz); 133.4 (d, 3C, $C_{\text{para}}\text{PPh}_3$, $^4J_{\text{C-P}} = 2.7$ Hz), 131.7 (s, 1C,

C=CH₂), 130.6 (d, 6C, $C_{\text{meta}}\text{PPh}_3$, $^3J_{\text{C-P}} = 11.9$ Hz), 129.9 (s, 2C, $C_{\text{ortho}}\text{Ph}$), 129.2 (s, 2C, $C_{\text{meta}}\text{Ph}$), 129.0 (s, 2C, $C_{\text{para}}\text{Ph}$), 128.6 (s, 2C, $C_{\text{ortho}}\text{Ph}$), 128.4 (s, 2C, $C_{\text{meta}}\text{Ph}$), 128.2 (s, 2C, $C_{\text{para}}\text{Ph}$), 109.4 (s, 2C, CH₂), 60.6 (s, 1C, CH₂-N), 59.1 (s, 1C, $\text{Ph-CH}_2\text{-N}$), 51.8 (s, 1C, CH₂-NH). $^{31}\text{P}\{^1\text{H}\}$ NMR (ppm) (162 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 45.2 (s, 1P, $\text{Au}(\text{PPh}_3)_2$), 37.6 (s, 1P, PPh_3). HRMS (ESI+ μ -TOF): m/z (%) = $[\text{M}]^+$ calcd for $[\text{C}_{37}\text{H}_{34}\text{AuN}_3\text{PS}_2]^+$ 812.1592, found 812.1591.

Synthesis of gold complex 22. To a solution of $[\text{Au}(\text{tht})_2]\text{OTf}$ (52.2 mg, 0.1 mmol), which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound 10 (65.0 mg, 0.2 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product 22 was obtained with 68% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.59–7.11 (m, 20H, Ph), 5.67 (br s, 2H, C=CH₂), 5.43 (br s, 6H, N-CH₂ + C=CH₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 160.1 (s, 1C), 154.1 (s, 1C), 146.4 (s, 1C), 138.8–131.8 (s, 6C, Ph), 130.6–122.1 (s, 20C, Ph), 109.5 (s, 2C, C=CH₂), 60.2 (s, 2C, N-CH₂). $^{19}\text{F}\{^1\text{H}\}$ NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): δ = -80.1 (s, 3F, OTf).

Synthesis of gold complex 23. To a solution of $[\text{Au}(\text{tht})_2]\text{OTf}$ (52.2 mg, 0.1 mmol) which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound 11 (72.0 mg, 0.2 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product 23 was obtained with 68% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.64–6.92 (m, 16H, Ph), 5.70 (br d, 2H, C=CH₂), 5.47 (br d, 6H, N-CH₂ + C=CH₂). $^{13}\text{C}\{^1\text{H}\}$ APT NMR (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 166.2 (s, 2C, C=S), 162.8 (s, 1C, C_q), 132.1 (d, 4C, $^3J_{\text{CF}} = 9.3$ Hz, Ar), 132.1 (d, 4C, $^3J_{\text{CF}} = 9.4$ Hz, Ar), 118.3 (s, 4C, Ar), 113.4 (d, 4C, $^2J_{\text{CF}} = 21.2$ Hz, Ar), 109.7 (d, 4C, $^2J_{\text{CF}} = 23.5$ Hz, Ar), 109.2 (s, 2C, C=CH₂), 60.9 (s, 1C, N-CH₂). $^{19}\text{F}\{^1\text{H}\}$ NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): δ = -73.5 (s, 1F, OTf), -106.4 (m, 2F, Ph-F), -107.5 (m, 2F, Ph-F).

Synthesis of gold complex 24. To a solution of $[\text{Au}(\text{tht})_2]\text{OTf}$ (52.2 mg, 0.1 mmol) which has been prepared *in situ* in CH_2Cl_2 (10 ml), compound 12 (72.0 mg, 0.2 mmol) was added and the reaction was stirred for 3 h. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3 $\times$ 1 mL), and vacuum dried to obtain the product. Product 23 was obtained with 72% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 7.32–7.14 (m, 20H, Ph), 5.60 (br s, 2H, C=CH₂), 5.42 (m, 6H, N-CH₂, C=CH₂), 5.08 (m, 2H, NH-CH_2), 4.48 (s, 2H, N-CH₂). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): δ = 176.1 (s, 2C, C=S), 158.2 (s, 2C, N-C=N), 138.9 (s, 2C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-N}$), 136.9 (s, 2C, $C_{\text{ipso}}\text{Ph-CH}_2\text{-NH}$), 133.3 (s, 2C, CH₂=C), 129.7–128.1 (s, 20C, Ph), 108.9 (s, 2C, C=CH₂), 60.4 (s, 2C, CH₂), 59.1 (s, 2C, $\text{Ph-CH}_2\text{-N=C}$), 51.2 (s, 2C, $\text{Ph-CH}_2\text{-NH}$). ^{19}F



$\{^1\text{H}\}$ NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = -79.8$ (s, 1F, OTf).

Synthesis of gold complex 25. To a solution of compound **10** (32.5 mg, 0.1 mmol) in CH_2Cl_2 (10 ml), $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ (45.3 mg, 0.1 mmol) was added and the reaction was stirred for 2 h at room temperature. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **25** was obtained with 80% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.39$ – 7.12 (m, 10H, Ph), 5.67 (br s, 1H, CH_2), 5.55 (br t, 2H, CH_2 -N), 5.40 (br s, 1H, CH_2). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 160.6$ (s, 1C), 148.1 (s, 1C, $\text{C}_{\text{ipso}}\text{Ph}$), 138.2 (s, 1C, $\text{C}_{\text{ipso}}\text{Ph}$), 132.1 (s, 1C, $\text{CH}_2=\text{C}$), 130.4 (s, 4C, $\text{C}_{\text{ortho}}\text{Ph}$), 129.3 (s, 1C, $\text{C}_{\text{para}}\text{Ph}$), 127.4 (s, 2C, $\text{C}_{\text{meta}}\text{Ph}$), 127.0 (s, 1C, $\text{C}_{\text{para}}\text{Ph}$), 121.1 (s, 2C, $\text{C}_{\text{meta}}\text{Ph}$), 109.2 (s, 1C, CH_2), 60.5 (s, 1C, CH_2 -N). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = -116.4$ (m, 2F, $\text{F}_{\text{ortho}}\text{C}_6\text{F}_5$), -164.1 (t, 1F, $\text{F}_{\text{para}}\text{C}_6\text{F}_5$, $^3J_{\text{FF}} = 19.7$ Hz), -166.3 (m, 2F, $\text{F}_{\text{meta}}\text{C}_6\text{F}_5$).

Synthesis of gold complex 26. To a solution of compound **11** (36.1 mg, 0.1 mmol) in CH_2Cl_2 (10 ml), $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ (45.3 mg, 0.1 mmol) was added and the reaction was stirred for 2.5 h at room temperature. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **26** was obtained with 70% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.48$ (m, 2H, Ph-F), 7.25 (m, 2H, Ph-F), 6.99 (m, 4H, Ph-F), 5.70 (br s, 1H, CH_2), 5.55 (br t, 2H, CH_2 -N), 5.43 (br s, 1H, CH_2). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 165.1$ (d, 1C, $\text{C}_{\text{ipso}}\text{Ph-F}$, $^3J_{\text{CF}} = 15.3$ Hz), 162.7 (d, 1C, $\text{C}_{\text{ipso}}\text{Ph-F}$, $^3J_{\text{CF}} = 15.8$ Hz), 132.2 (d, 1C, $^3J_{\text{HF}} = 9.4$ Hz), 132.0 (d, 1C, $^3J_{\text{CF}} = 9.4$ Hz), 131.6 (s, 1C, $\text{CH}_2=\text{C}$), 123.8 (s, 1C), 118.2 (s, 1C), 116.2 (d, 1C, $^2J_{\text{CF}} = 21.3$ Hz), 115.3 (d, 1C, $^2J_{\text{CF}} = 28.5$ Hz), 113.6 (d, 1C, $^2J_{\text{CF}} = 21.3$ Hz), 109.7 (d, 1C, $^2J_{\text{CF}} = 23.7$ Hz), 109.6 (s, 1C, CH_2), 60.6 (s, 1C, CH_2 -N). ^{19}F NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = -114.0$ (m, 1F, Ph-F), -114.1 (m, 1F, Ph-F), -116.6 (m, 2F, $\text{F}_{\text{ortho}}\text{C}_6\text{F}_5$), -163.8 (t, 1F, $\text{F}_{\text{para}}\text{C}_6\text{F}_5$, $^3J_{\text{FF}} = 19.7$ Hz), -166.2 (m, 2F, $\text{F}_{\text{meta}}\text{C}_6\text{F}_5$).

Synthesis of gold complex 27. To a solution of compound **12** (35.3 mg, 0.1 mmol) in CH_2Cl_2 (10 ml), $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ (45.3 mg, 0.1 mmol) was added and the reaction was stirred for 2.5 h at room temperature. The solution was concentrated under reduced pressure to approximately 1 ml and hexane (10 ml) was added to precipitate a white solid, which was filtered under vacuum, washed with hexane (3×1 mL), and vacuum dried to obtain the product. Product **27** was obtained with 83% yield. ^1H NMR (ppm) (400 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 7.37$ – 7.11 (m, 10H, Ph), 5.67 (br s, 1H, CH_2), 5.57 (m, 2H, CH_2 -N), 5.45 (m, 1H, CH_2), 5.34 (m, 2H, NH- CH_2), 4.48 (s, 2H, N- CH_2). $^{13}\text{C}\{^1\text{H}\}$ APT (ppm) (101 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = 157.9$ (s, 1C, N-C=N), 138.5 (s, 1C, $\text{C}_{\text{ipso}}\text{Ph-F}$), 132.0 (s, 1C, $\text{CH}_2=\text{C}$), 129.8–128.1 (m, 10C, Ph), 109.3 (s, 1C, CH_2), 60.5 (s, 1C, CH_2 -N), 59.1 (s, 1C, Ph- CH_2 -N=C), 52.3 (s, 1C, Ph- CH_2 -NH). ^{19}F

NMR (ppm) (376.5 MHz, $(\text{CD}_3)_2\text{CO}$): $\delta = -117.7$ (m, 2F, $\text{F}_{\text{ortho}}\text{C}_6\text{F}_5$), -163.4 (t, 1F, $\text{F}_{\text{para}}\text{C}_6\text{F}_5$, $^3J_{\text{FF}} = 19.6$ Hz), -166.7 (m, 2F, $\text{F}_{\text{meta}}\text{C}_6\text{F}_5$).

Conclusions

Several thiazole-based heterocycles have been achieved by the reaction of propargylamines with isothiocyanates. The selective formation of thioureas, iminothiazolidines, aminothiazolines or mixed thiazolidine–thiourea compounds has been accomplished depending upon the reaction conditions. The use of secondary propargylamines leads selectively to the formation of cyclic iminothiazolidine derivatives, while primary propargylamines form iminothiazoline species. Further reaction of the cyclic thiazoline derivatives with isothiocyanate or the reaction of the starting compounds in a 1 : 2 molar ratio affords the thiazolidine–thiourea derivatives. The proposed mechanism for the formation of these heterocycles starts with the formation of the thiourea species following two possible pathways. The first one is the formation of the thiourea, cyclisation to form the iminothiazolidine and further attack of the isothiocyanate to produce the thiazolidine–thiourea. Alternatively, the generation of bis(thiourea) and cyclisation produce the same compounds. Coordination studies of these heterocyclic species towards silver and gold with different stoichiometries have been carried out to afford cationic $[\text{ML}(\text{PPh}_3)]\text{OTf}$, $[\text{ML}_2]\text{OTf}$ ($\text{M} = \text{Ag}, \text{Au}$) or neutral $[\text{Au}(\text{C}_6\text{F}_5)\text{L}]$ complexes. Preliminary studies of the cytotoxic activity in lung cancer cells have been carried out in both ligand and complexes, showing that although the ligands do not exhibit anti-cancer activity, the coordination of the metals, especially silver, greatly enhances the cytotoxic activity.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors also thank Agencia Estatal de Investigación (PID2019-104379RB-C21 and PID2020-117455GB-I00/AEI/10.13039/501100011033; and RED2018-102471-T (MCIN/AEI/10.13039/501100011033)) and Gobierno de Aragón-Fondo Social Europeo (research group E07_20R) for financial support of our research.

References

- 1 N. Aydemir and R. Bilaloğlu, *Mutat. Res.*, 2003, **537**, 43–51.
- 2 For selected examples, see: (a) C. Sanmartín, M. Echeverría, B. Mendivil, L. Cordeu, E. Cubedo, J. García-Foncillas, M. Font and J. A. Palop, *Bioorg. Med. Chem.*, 2005, **13**, 2031–2044; (b) E. Moreau, S. Fortin, M. Desjardins,



- J. L. C. Rousseau, E. Petitclerc and R. C. -Gaudreault, *Bioorg. Med. Chem.*, 2005, **13**, 6703–6712; (c) G. Hallur, A. Jimeno, S. Dalrymple, T. Zhu, M. K. Jung, M. Hidalgo, J. T. Isaacs, S. Sukumar, E. Hamel and S. R. Khan, *J. Med. Chem.*, 2006, **49**, 2357–2360; (d) H.-Q. Li, P.-C. Lv, T. Yan and H.-L. Zhu, *Anti-Cancer Agents Med. Chem.*, 2009, **9**, 471–480; (e) H.-Q. Li, T.-T. Zhu, T. Yan, Y. Luo and H.-L. Zhu, *Eur. J. Med. Chem.*, 2009, **44**, 453–459; (f) L. Garuti, M. Roberti, G. Bottegoni and M. Ferraro, *Curr. Med. Chem.*, 2016, **23**, 1528–1548; (g) F. Auria-Luna, E. Marqués-López, E. Romanos, V. Fernández-Moreira, M. C. Gimeno, I. Marzo and R. P. Herrera, *Bioorg. Chem.*, 2020, **105**, 104364.
- 3 For selected examples, see: (a) A. Esteves-Souza, K. Pissinate, M. G. Nascimento, N. F. Grynberg and E. Aurea, *Bioorg. Med. Chem.*, 2006, **14**, 492–499; (b) S. N. Manjula, N. M. Noolvi, K. V. Parihar, S. A. M. Reddy, V. Ramani, A. K. Gadad, G. Singh, N. G. Kutty and C. M. Rao, *Eur. J. Med. Chem.*, 2009, **44**, 2923–2929; (c) S. Saeed, N. Rashid, P. G. Jones, M. Ali and R. Hussain, *Eur. J. Med. Chem.*, 2010, **45**, 1323–1331; (d) J. Yao, J. Chen, Z. He, W. Sun and W. Xu, *Bioorg. Med. Chem.*, 2012, **20**, 2923–2929; (e) C. Ning, Y. Bi, Y. He, W. Y. Huang, L. Liu, Y. Li, S. Zhang, X. Liu and N. Yu, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 6432–6435.
- 4 (a) T. K. Venkatachalam, C. Mao and F. M. Uckun, *Bioorg. Med. Chem.*, 2004, **12**, 4275–4284; (b) R. Hunter, Y. Younis, C. I. Muhanji, T.-I. Curtin, K. J. Naidoo, M. Petersen, C. M. Bailey, A. Basavapathruni and K. S. Anderson, *Bioorg. Med. Chem.*, 2008, **16**, 10270–10280; (c) K. Chen, Z. Tan, M. He, J. Li, S. Tang, I. Hewlett, F. Yu, Y. Jin and M. Yang, *Chem. Biol. Drug Des.*, 2010, **76**, 25–33.
- 5 J. D. Bloom, R. G. Dushin, K. J. Curran, F. Donahue, E. B. Norton, E. Terefenko, T. R. Jones, A. A. Ross, B. Feld, S. A. Lang and M. J. DiGrandi, *Bioorg. Med. Chem.*, 2004, **14**, 3401–3406.
- 6 S. A. Khan, N. Singh and K. Saleem, *Eur. J. Med. Chem.*, 2008, **43**, 2272–2277.
- 7 (a) J. Stefanska, D. Szulczyk, A. E. Koziol, B. Mirosław, E. Kedzierska, S. Fidecka, B. Busonera, G. Sanna, G. Giliberti, P. La Colla and M. Struga, *Eur. J. Med. Chem.*, 2012, **55**, 205–213; (b) B. Kocyigit-Kaymakcioglu, A. O. Celen, N. Tabanca, A. Ali, S. I. Khan, I. A. Khan and D. E. Wedge, *Molecules*, 2013, **18**, 3562–3567.
- 8 L. dos Santos, L. A. Lima, V. Cechinel-Filho, R. Corrêa, F. C. Buzzzi and R. J. Nunes, *Bioorg. Med. Chem.*, 2008, **16**, 8526–8534.
- 9 A. L. Lourenço, M. S. Saito, L. E. G. Dorneles, G. M. Viana, P. C. Sathler, L. C. S. Aguiar, M. de Pádula, T. F. S. Domingos, A. G. M. Fraga, C. R. Rodrigues, V. P. de Sousa, H. C. Castro and L. M. Cabral, *Molecules*, 2015, **20**, 7174–7200.
- 10 (a) N. Sahiba, A. Sethiya, J. Soni, D. K. Agarwal and S. Agarwal, *Top. Curr. Chem.*, 2020, **378**, 34; (b) X. Just-Baringo, F. Albericio and M. Alvarez, *Curr. Top. Med. Chem.*, 2014, **14**, 1244–1256; (c) A.-C. Gaumont, M. Gulea and J. Levillain, *Chem. Rev.*, 2009, **109**, 1371–1401.
- 11 (a) D. Das, P. Sikdar and M. Baigari, *Eur. J. Med. Chem.*, 2016, **109**, 89–98; (b) R. P. Singh, M. N. Aziz, D. Gout, W. Fayad, M. A. El-Manawaty and C. J. Lovely, *Bioorg. Med. Chem.*, 2019, **27**, 115047; (c) S. A. Shehzadi, I. Khan, A. Saeed, F. A. Larik, P. A. Channar, M. Hassan, H. Raza, Q. Abbas and S.-Y. Seo, *Bioorg. Chem.*, 2019, **84**, 518–528; (d) A. Singh, D. Malhotra, K. Singh, R. Chadha, P. Mohinder and S. Bedi, *J. Mol. Chem.*, 2022, **1266**, 133479.
- 12 (a) S. R. Alizadeh and S. M. Hashemi, *Med. Chem. Res.*, 2021, **30**, 771–806; (b) Y. Wan, J. Long, H. Gao and Z. Tang, *Eur. J. Med. Chem.*, 2021, **210**, 112953.
- 13 (a) S. K. Bharti, G. Nath, R. Tilak and S. K. Singh, *Eur. J. Med. Chem.*, 2010, **45**, 651–660; (b) A.-M. Borcea, I. Ionuț, O. Crișan and O. Oniga, *Molecules*, 2021, **26**, 624.
- 14 I. P. Singh, S. Gupta and S. Kumar, *Med. Chem.*, 2020, **16**, 4–23.
- 15 K. Liaras, M. Fesatidou and A. Geronikaki, *Molecules*, 2018, **23**, 685.
- 16 R. G. Miller, J. D. Mitchell and D. H. Moore, *Cochrane Database Syst. Rev.*, 2012, **3**, CD001447.
- 17 C. Borelli, M. Schaller, M. Niewerth, K. Nocker, B. Baasner, D. Berg, R. Tiemann, K. Tietjen, B. Fugmann, S. Lang-Fugmann and H. C. Korting, *Chemotherapy*, 2008, **54**, 245–259.
- 18 A. Bergamo and G. Sava, *Dalton Trans.*, 2011, **40**, 7817–7823.
- 19 (a) I. Ott, *Coord. Chem. Rev.*, 2009, **153**, 1670–1681; (b) S. Nobili, E. Mini, I. Landini, C. Gabbiani, A. Casini and L. Messori, *Med. Res. Rev.*, 2010, **30**, 550–580; (c) C.-M. Che and R. W.-Y. Sun, *Chem. Commun.*, 2011, **47**, 9554–9560; (d) A. Casini and L. Messori, *Curr. Top. Med. Chem.*, 2011, **11**, 2647–2660; (e) C. N. Banti and S. K. Hadjikakou, *Metallomics*, 2013, **5**, 569–596; (f) B. Bertrand and A. Casini, *Dalton Trans.*, 2014, **43**, 4209–4219; (g) S. Medici, M. Peana, V. M. Nurchi, J. I. Lachowicz, G. Crisponi and M. A. Zoroddu, *Coord. Chem. Rev.*, 2015, **284**, 329–350; (h) T. Zou, C. T. Lum, C.-N. Lok, J.-J. Zhang and C.-M. Che, *Chem. Soc. Rev.*, 2015, **44**, 8786–8801; (i) C. I. Yeo, K. K. Ooi and E. R. T. Tiekink, *Molecules*, 2018, **23**, 1410; (j) V. Fernández-Moreira, R. P. Herrera and M. C. Gimeno, *Pure Appl. Chem.*, 2019, **91**, 247.
- 20 For selected examples, see: (a) M. C. Gimeno, H. Goitia, A. Laguna, M. E. Luque, M. D. Villacampa, C. Sepúlveda and M. Meireles, *J. Inorg. Biochem.*, 2011, **105**, 1373–1382; (b) H. Goitia, Y. Nieto, M. D. Villacampa, C. Kasper, A. Laguna and M. C. Gimeno, *Organometallics*, 2013, **32**, 6069–6078; (c) A. Gutiérrez, L. Gracia-Fleta, I. Marzo, C. Cativiela, A. Laguna and M. C. Gimeno, *Dalton Trans.*, 2014, **43**, 17054–17066; (d) L. Ortego, F. Cardoso, S. Martins, M. F. Fillat, A. Laguna, M. Meireles, M. D. Villacampa and M. C. Gimeno, *J. Inorg. Biochem.*, 2014, **130**, 32–37; (e) A. Gutiérrez, I. Marzo, C. Cativiela, A. Laguna and M. C. Gimeno, *Chem. – Eur. J.*, 2015, **21**, 11088–11095; (f) L. Ortego, M. Meireles, C. Kasper, A. Laguna, M. D. Villacampa and M. C. Gimeno, *J. Inorg. Biochem.*, 2016, **156**, 133–144; (g) D. Salvador-Gil, L. Ortego,



- R. P. Herrera, I. Marzo and M. C. Gimeno, *Dalton Trans.*, 2017, **46**, 13745–13755; (h) V. Fernández-Moreira, C. Val-Campillo, I. Ospino, R. P. Herrera, I. Marzo, A. Laguna and M. C. Gimeno, *Dalton Trans.*, 2019, **48**, 3098–3108.
- 21 (a) I. Mármol, S. Montanel-Perez, J. C. Royo, M. C. Gimeno, M. D. Villacampa, M. J. Rodríguez-Yoldi and E. Cerrada, *Inorg. Chem.*, 2020, **59**, 17732–17745; (b) A. Luengo, I. Marzo, M. Reback, I. M. Daubitz, V. Fernández-Moreira, N. Metzler-Nolte and M. C. Gimeno, *Chem. – Eur. J.*, 2020, **26**, 12158–12167; (c) I. Mármol, P. Castellnou, R. Alvarez, M. C. Gimeno, M. J. Rodríguez-Yoldi and E. Cerrada, *Eur. J. Med. Chem.*, 2019, **183**, 111661; (d) H. Goitia, A. Laguna and M. C. Gimeno, *Inorg. Chim. Acta*, 2018, **24**, 53–58.
- 22 For a few precedents of this type of reaction under other conditions, see: (a) H. M.-F. Viart, T. S. Larsen, C. Tassone, T. L. Andresen and M. H. Clausen, *Chem. Commun.*, 2014, **50**, 7800–7802; (b) N. Scalacci, C. Pellajo, M. Radi and D. Castagnolo, *Synlett*, 2016, **27**, 1883–1887; (c) R. P. Singh, D. Gout and C. J. Lovely, *Eur. J. Org. Chem.*, 2019, 1726–1740.
- 23 CCDC 2234522 (4), 2234523 (6) and 2234524 (7)[†] contain the supplementary crystallographic data for this paper.
- 24 CCDC 2234525 (10) and 2234526 (11)[†] contain the supplementary crystallographic data for this paper.
- 25 A. Johnson, A. Laguna and M. C. Gimeno, *J. Am. Chem. Soc.*, 2014, **136**, 12812–12815.
- 26 (a) A. Ranjan, R. Yerande, P. B. Wakchaure, S. G. Yerande and D. H. Dethe, *Org. Lett.*, 2014, **16**, 5788–5791; (b) A. Ranjan, A. S. Deore, S. G. Yerande and D. H. Dethe, *Eur. J. Org. Chem.*, 2017, 4130–4139.
- 27 (a) K. Yan, C.-N. Lok, K. Bierla and C.-M. Che, *Chem. Commun.*, 2010, **46**, 7691–7693; (b) S. Adhikari, O. Hussain, R. M. Phillips, W. Kaminsky and M. R. Kollipara, *Appl. Organomet. Chem.*, 2018, **32**, e4476G; (c) G. Canudo-Barreras, L. Ortego, A. Izaga, I. Marzo, R. P. Herrera and M. C. Gimeno, *Molecules*, 2021, **26**, 6891.
- 28 CCDC 2234527 (20)[†] contains the supplementary crystallographic data for this paper.
- 29 CCDC 2234528 (27)[†] contains the supplementary crystallographic data for this paper.
- 30 J. van Meerloo, G. J. Kaspers and J. Cloos, *Methods Mol. Biol.*, 2011, **731**, 237–245.
- 31 C. Marzano, V. Gandin, M. Pellei, D. Colavito, G. Papini, G. G. Lobbia, E. Del Giudice, M. Porchia, F. Tisato and C. Santi, *J. Med. Chem.*, 2008, **51**, 798–808.
- 32 M. Frik, J. Fernández-Gallardo, O. Gonzalo, V. Mangas-Sanjuan, M. González-Álvarez, A. Serrano del Valle, C. Hu, I. González-Álvarez, M. Bermejo, I. Marzo and M. Contel, *J. Med. Chem.*, 2015, **58**, 5825–5841.
- 33 G. M. Sheldrick, *SADABS, Program for adsorption correction*, University of Göttingen, Göttingen, Germany, 1996.
- 34 CysAlisPro, Version 1.171.35.11; Agilent Technologies. Multiscan absorption correction with SCALE3 ABSPACK scaling algorithm.
- 35 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Adv.*, 2015, **71**, 3–8.
- 36 (a) O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, **42**, 339–341; (b) L. J. Bourhis, O. V. Dolomanov, R. J. Gildea, J. A. K. Howard and H. Puschmann, *Acta Crystallogr., Sect. A: Found. Adv.*, 2015, **71**, 59–71.
- 37 (a) R. Uson, A. Laguna, M. Laguna, D. A. Briggs, H. H. Murray and J. P. Fackler, in *Inorg. Synth*, John Wiley & Sons, Inc., 1989, pp. 85–91; (b) L. Malatesta, L. Naldini, G. Simonetta and F. Cariati, *Coord. Chem. Rev.*, 1966, **1**, 255–262.

