



Cite this: *Dalton Trans.*, 2024, **53**, 13709

Received 12th July 2024,
Accepted 2nd August 2024

DOI: 10.1039/d4dt02018a

rs.c.li/dalton

Lessons from recent theoretical treatments of Al–M bonds (M = Fe, Cu, Ag, Au) that capture CO₂

S. M. Supundrika Subasinghe  and Neal P. Mankad *

Complexes with Al–M bonds (M = transition metal) have emerged as platforms for discovering new reaction chemistry either through cooperative bond activation behaviour of the heterobinuclear unit or by modifying the properties of the M site through its interaction with the Al centre. Therefore, elucidating the nature of Al–M bonding is critical to advancing this research area and typically involves careful theoretical modelling. This Frontier article reviews selected recent case studies that included theoretical treatments of Al–M bonds, specifically highlighting complexes capable of cooperative CO₂ activation and focusing on extracting lessons particular to the Al–M sub-field that will inform future studies with theoretical/computational components.

Introduction

As the area of heterobimetallic chemistry continues to attract attention,^{1–8} it is of importance to consider heterobinuclear complexes that feature earth's most abundant metal, aluminium. Indeed, studying complexes with direct Al–M bonds has become a frontier area of research with applications in areas such as catalysis^{9–12} and small molecule activation.^{13–20} As these applications continue to emerge, it becomes increasingly important to understand the nature of Al–M bonding itself. Canonically, heterobinuclear complexes pairing d-block

elements with aluminium are viewed as featuring electropositive aluminium centres serving as donor atoms (aluminumyl “metalloligands”) towards transition metal sites. According to this model, the unique reactivity of the transition metal site is imparted by this unconventional bond polarity *vis à vis* traditional coordination complexes with electronegative donor atoms. This bonding motif is certainly valid in some cases but may not be universal to all Al–M complexes.

In this context, elucidating the electronic structures of Al–M bonds is critical. Because gaining this understanding will necessarily involve theoretical analysis, it is important to identify aspects of Al–M bonding that require special treatment theoretically.²¹ Lessons learnt from such case studies will inform future research in this frontier area. In this Frontier article, we briefly review selected recent cases to summarise

Department of Chemistry, University of Illinois Chicago, Chicago, IL 60607, USA.
E-mail: npm@uic.edu



**S. M. Supundrika
Subasinghe**

Supundrika Subasinghe is currently a PhD candidate at the University of Illinois Chicago. In 2017, she earned a Bachelor of Science degree from the University of Colombo Sri Lanka, specialising in Computational Chemistry. Her research interests lie at the intersection of organometallic chemistry and theoretical chemistry. Supundrika has gained valuable experience working as a research assistant, contributing to various projects and expanding her expertise in the field of Chemistry.



Neal P. Mankad

Neal Mankad is currently Professor and Associate Head of Chemistry at the University of Illinois Chicago as well as Associate Editor of Dalton Transactions. His group pursues projects spanning bioinorganic chemistry, organometallic chemistry, and electrochemistry that are generally motivated by impacting global energy and sustainability.

some important lessons that have emerged. We have chosen to use Al–M complexes that capture CO₂ as a venue. In the domain of anthropogenic greenhouse gases, carbon dioxide (CO₂) stands out as the primary contributor to global warming.²² As atmospheric CO₂ levels continue to rise, there is an increasing urgency for developing innovative strategies to capture and valorise CO₂ to mitigate its environmental impact.^{23,24} Transition metal complexes have long been known to present promising pathways for CO₂ capture due, in part to their ability to access multiple oxidation states during small-molecule activation.^{25,26} In addition to mononuclear and homobinuclear compounds, CO₂ activation by heterobinuclear complexes, particularly those combining main group and transition metal systems, continues to be an area of focus dating back to seminal observations by Floriani.^{27,28} As discussed herein, our group and several others have recently studied CO₂ activation with heterobinuclear complexes pairing d-block elements (e.g. Fe, Cu, Ag, Au) with aluminium.

Nucleophilic Al–M (M = coinage metal) complexes: covalent vs. polarised bonding


Initial discovery & assignment of nucleophilic Au(–I)

In an influential 2019 report,²⁹ Hicks *et al.* synthesised (NON) Al–AuP^tBu₃ (**1**) by reacting the Al(I) nucleophile, K[Al(NON)], with the Au(I) complex, ^tBu₃PAuI (Scheme 1). The resulting Al–Au bond was found to be the shortest on record (2.40 Å). The Al–N and Al–O bond distances in **1** being shorter than in the K[Al(NON)] precursor was attributed by the authors to 2e[–] oxidation of Al(I) to Al(III) during the formation of **1**. However, contributions of molecular charge (*i.e.*, transforming an anion to a neutral species) to these bond contractions were not considered at the time. The authors' preliminary model for the short Al–Au interaction in **1** was that of a polarised Al^{δ+}–Au^{δ–} bond, *i.e.* resembling a strong [(NON)Al]⁺ ← [AuP^tBu₃][–] dative interaction, based on the dramatically higher electronegativity of atomic Au (2.54) compared to atomic Al (1.61) on the Pauling scale. The implication that complex **1** contains a [AuP^tBu₃][–] fragment, and thus a gold centre with formally Au(–I) (auride) character, was provocative since most sub-valent

transition metal centres require stabilisation by strong π-acids like CO.³⁰

The nature of the putatively Al^{δ+}–Au^{δ–} bond was investigated computationally in the same study using density functional theory (DFT), quantum theory of atoms in molecules (QTAIM), and natural orbitals for chemical valence (NOCV) calculations. QTAIM is a conceptual framework used in theoretical chemistry to understand the electronic structure of molecules through analysis of electron density distribution and topology.^{31,32} In this case, partial atomic charges were assigned from QTAIM calculations based on the integration of electron density over each atomic basin found within the overall density map. NOCV is a method that identifies the natural bond orbitals (NBOs,³³ *i.e.*, localised wavefunctions resembling canonical Lewis structures) predominantly involved in chemical bonding.³⁴ Like in QTAIM, the NOCV orbitals are derived from analysis of the total electron density of the molecule rather than from arbitrary basis functions sometimes used in standard DFT and NBO calculations.

Comparison of QTAIM partial atomic charges for **1** to those of its synthetic precursors indicated transfer of ~1.6e[–] from [Al(NON)][–] to [AuP^tBu₃]⁺ during Al–Au bond formation.²⁹ The most significant Al/Au NOCV interaction in **1** compared to its synthetic precursor fragments was calculated to be a σ-bond assigned by the authors as Al-to-Au lone pair donation. These two observations were used to justify the formal 2e[–] oxidation of Al(I) to Al(III) and formal 2e[–] reduction of Au(I) to Au(–I) during formation of **1**. However, one can argue that these analyses were biased because they relied on comparison of **1** to starting fragments defined by the theorists. In other words, the authors based their interpretations of NOCV and QTAIM calculations on comparison of **1** to its ionised fragments, [Al(NON)][–] and [AuP^tBu₃]⁺. However, DFT calculations disclosed in the same study instead indicated that the lowest energy pathway (by 37–62 kcal mol^{–1})³⁵ for Al–Au cleavage is not heterolytic but homolytic, *i.e.* to [Al(NON)][•] and [AuP^tBu₃][•] fragments.²⁹ Thus, in contrast to the theoretical analyses described above and to use the authors' own words written in the article's ESI, “the bonding interaction should rather be considered as a conventional covalent bond instead of a coordination or some other dative bond”. The further refinement of the bonding model for complex **1** as involving a highly covalent bond between formally Al(II) and Au(0) centres would emerge in later studies (see below).



Scheme 1 Synthesis and reactivity of a nucleophilic Al–Au bond originally reported by Hicks *et al.*²⁹ dipp = 2,6-di-iso-propylphenyl.

Based on the observations above and the fact that the QTAIM charge at the Au centre in **1** was calculated to be quite negative (-0.82 au), the authors hypothesised that complex **1** would be nucleophilic at gold, in stark contrast to the typical electrophilic chemistry of molecular gold complexes. Addition of CO_2 gas to **1** resulted in metallacarboxylate species **2** in which the Au centre is C -bound to CO_2 (Scheme 1). This reaction outcome was used by the authors as further evidence for the presence of a nucleophilic centre with $\text{Au}(-\text{I})$ character in **1** that had added to the electrophilic carbon centre of CO_2 . However, soon afterwards, the same team as well as Liu *et al.* showed that CO_2 inserts readily into corresponding $\text{Al}-\text{Ag}$, $\text{Al}-\text{Cu}$, and $\text{Al}-\text{Zn}$ bonds (Fig. 1).^{36–39} These lighter analogues were calculated to feature significantly less $\text{Al}-\text{M}$ polarisation and less accumulation of negative charge on the d -block metal; yet, they reacted as (or more) efficiently with CO_2 as **1**. In the most extreme case, an $\text{Al}-\text{Zn}$ bond was calculated to accumulate *positive* charge of 1.26 au on Zn and still react with CO_2 to form a metallacarboxylate species with a C -bound Zn centre.³⁹ In other words, efficient CO_2 activation to form $\text{AlO}_2\text{C}-\text{M}$ bonds does not correlate with nor require negative charge accumulation at M, thus negating the only experimental evidence for the presence of a nucleophilic $\text{Au}(-\text{I})$ -like ion in **1**. Indeed, as later pointed out by Sorbelli *et al.*,³⁵ even homobinuclear complexes (*i.e.*, inherently apolar metal–metal bonds) efficiently insert CO_2 to form analogous metallacarboxylate structures.

Re-interpretation as a covalent, nucleophilic Al–Au bond

In 2021, Sorbelli *et al.* revisited the computational analysis of **1** and its CO_2 reactivity using DFT, extended transition state natural orbitals for chemical valence (ETS-NOCV), and charge displacement (CD) calculations.³⁵ ETS-NOCV is a theoretical method that combines aspects of both transition state theory

and NOCV analysis, essentially calculating changes in NOCV distributions occurring along a reaction coordinate.⁴⁰ CD further partitions this data into partial atomic charge contributions from individual components of the molecule(s).⁴¹ Relatedly, the CD-NOCV method breaks down partial atomic charge density contributions that, together, comprise a covalent interaction.⁴²

In their report,³⁵ the Sorbelli *et al.* disfavoured the nucleophilic auride model for **1**, instead proposing that **1** contains a highly covalent $\text{Al}-\text{Au}$ bond that, itself, serves as the nucleophilic electron pair towards CO_2 . The previous use of Pauling electronegativity values to formulate a polarised $\text{Al}^{\delta+}-\text{Au}^{\delta-}$ bond was contradicted by noting that the ionisation energies of the $[\text{Al}(\text{NON})]^\cdot$ and $[\text{AuP}^t\text{Bu}_3]^\cdot$ fragments are calculated to be nearly identical (2.53 and 2.56 eV, respectively). The previous use of partial atomic charge calculations to support the $\text{Al}^{\delta+}-\text{Au}^{\delta-}$ proposal was also called into question by noting that the partial atomic charge at Au in **1** varies from -0.83 to $+0.22$ au depending on the charge partitioning scheme used in the calculations.⁴³

The overall CD-NOCV analysis of the $\text{Al}-\text{Au}$ bond in **1** indicated only slight polarisation of $0.05e^-$ towards the Au centre.³⁵ This was partitioned into two charge fluxes, one from Au towards Al and the other from Al towards Au, that approximately counterbalanced each other (Fig. 2). Based on these observations and others, the authors assigned the $\text{Al}-\text{Au}$ bond in **1** as involving formally $\text{Al}(\text{II})$ and $\text{Au}(0)$ centres engaged in a bond with a high degree of covalency. Based on the bidirectional charge fluxes apparent by CD-NOCV, the authors also chose to describe the nature of **1** as “diradical-like” despite the calculated $\text{Al}-\text{Au}$ bond energy of 83 kcal mol^{-1} precluding the formation of any radicals at standard conditions.^{35,44} Indeed, complete active space self-consistent field (CASSCF) calculations (known to be more accurate than DFT) later reported




Fig. 1 Selected $\text{Al}-\text{M}$ bonds known to insert CO_2 to form metallacarboxylates with C -bound M centres.^{29,36–39} dipp = 2,6-di-isopropylphenyl.




Fig. 2 CD-NOCV curves for the interaction between $[\text{Al}(\text{NON})]^\cdot$ and $[\text{AuP}^t\text{Bu}_3]^\cdot$ fragments, with isodensity surfaces ($1 \text{ me}^- a_0^{-3}$) shown for the two main deformation densities contributing to $\text{Al}-\text{Au}$ bonding (charge flux: red $\rightarrow$ blue). Image reproduced with permission from Sorbelli *et al.*³⁵

by Guo *et al.* indicated that the contributions of open-shell configurations are small and that the active electrons remain paired throughout the CO₂ activation pathway.⁴⁵ Thus, description of the chemistry of **1** as “diradical-like” is questionable, in our view.

In the 2021 study by Sorbelli *et al.*, the reaction pathway for **1** and CO₂ converting to **2** was calculated.³⁵ The calculated mechanism is, effectively, a [2 + 2] cycloaddition involving the Al–Au σ -bonding pair and one of the CO₂ π -bonding pairs, followed by rearrangement from κ^1 - to κ^2 -metallacarboxylate binding to aluminium. Interestingly, the calculated [2 + 2] transition state structure features the carbon centre of CO₂ interacting with *both* Au and Al (M–C Mayer bond orders of 0.26 and 0.16, respectively), consistent with the Al–Au bond (and not a formal auride centre) acting as the nucleophilic electron pair. An analogous transition state geometry was calculated in 2022 by Guo *et al.* (Fig. 3), who also found that the Al–C interaction is even more significant at the transition state for the reaction of an Al–Cu analogue.⁴⁵ ETS-NOCV calculations by Sorbelli *et al.* indicated that the major interaction

during CO₂ activation by **1** involves the transfer of electron density ($0.33e^-$) from *both* Al and Au (specifically, from the Al–Au σ -bond region) to CO₂ (specifically, its LUMO).³⁵ Intrinsic bond orbital (IBO) calculations, which track changes in NBOs as a function of intrinsic reaction coordinate,⁴⁶ were performed by Guo *et al.*⁴⁵ The results (Fig. 4) agree with the results of Sorbelli *et al.* conceptually and are readily interpreted as involving flow of two reactive electron pairs: transformation of the Al–Au σ -bonding orbital into an Al–C σ -bonding orbital, and transformation of the CO₂ π -bonding orbital into an Al–O σ -bonding orbital. Collectively, these computational models converge on the concept of the Al–Au σ -bond being the nucleophilic electron pair that adds to the CO₂ carbon centre *via* a [2 + 2]-like transition state, with the C-bound rather than O-bound aurocarboxylate forming due to a secondary Lewis acid/base interaction between the Al site and the CO₂ oxygen centre rather than being indicative of nucleophilic character at any individual atom within **1**.




Fig. 3 Calculated transition state for CO₂ activation by **1**.^{35,44,45} Image reproduced with permission from Guo *et al.*⁴⁵

Cooperative CO₂ activation at a weak Al–Fe bond: frustrated radical pair behaviour

For about a decade (2012–2022), our group had studied (NHC) Cu–Fp complexes (NHC = *N*-heterocyclic carbene, Fp = FeCp(CO)₂) with reactive Cu–Fe bonds.^{5,47,48} All physical and computational data indicated that such complexes contained polar Cu ^{δ^+} –Fe ^{δ^-} bonds with Wiberg bond orders⁴⁹ of <0.5 and with





Fig. 4 IBO analysis of the reaction between **1** and CO₂. Image reproduced with permission from Guo *et al.*⁴⁵



Scheme 2 Synthesis and reactivity of a weak, covalent Al–Fe bond reported by Sinhababu *et al.*⁵⁹ dipp = 2,6-di-iso-propylphenyl.

an accumulation of negative charge at the Fe centres.^{50–52} (Note that the proposed $\text{Cu}^{\delta+}\text{--Fe}^{\delta-}$ polarisation opposes the relative Pauling electronegativity values: Cu, 1.90; Fe, 1.83.) As such, depending on reaction conditions, we typically proposed either that (NHC)Cu–Fp complexes react as Fe(0) nucleophiles (akin to the auride proposal of Hicks *et al.* discussed above) or that they dissociate heterolytically to reveal $[(\text{NHC})\text{Cu}]^+[\text{Fp}]^-$ ion pairs with frustrated Lewis pair (FLP)^{53,54} behaviour.^{47,55–58}

In 2022, our group synthesised $\text{LAl}(\text{Me})\text{Fp}$ (**3**) by reacting the metallonucleophile, KFp , with the Al(III) β -diketiminate complex, $\text{LAl}(\text{Me})\text{I}$ (Scheme 2, $\text{L} = \text{HC}[\text{MeCNdipp}]_2^-$, dipp = 2,6-di-iso-propylphenyl).⁵⁹ The resulting Al–Fe bond was found to be among the shortest on record (2.48 Å). Computational modelling revealed that the Al–Fe bond is significantly more covalent than the previously studied Cu–Fe systems, with a Wiberg bond order of >0.8 based on NBO calculations and with relatively less accumulation of negative charge at Fe according to theoretical QTAIM analysis. Moreover, whereas the heterolytic bond dissociation energy (BDE) of **3** to $[\text{LAlMe}]^+[\text{Fp}]^-$ was calculated to be quite high (59 kcal mol^{−1}), homolytic dissociation of **3** to the $[\text{LAlMe}]/[\text{Fp}]^\bullet$ radical pair was calculated to be energetically accessible (25 kcal mol^{−1}) at standard conditions. Thus, while the polar Cu–Fe bonds previously studied by our group were viewed as masked FLPs, the covalent Al–Fe bond in **3** can be regarded as a masked frustrated radical pair (FRP).^{60–62} Due to the covalent nature of the Al–Fe bond, oxidation state assignments have not been proposed in the literature for **3** and, arguably, are irrelevant to understanding chemical reactivity.

Exposure of **3** to CO_2 gas provided metallacarboxylate species **4** (Scheme 2).⁵⁹ The mechanism for this transformation proposed by Sinhababu *et al.* is shown in Scheme 2. A reasonable pathway was calculated by DFT and, in accord with recent findings by Mears *et al.* on an Al–Cu system,²¹ required the use of dispersion corrections to the DFT functional along with the inclusion of all ligand substituents (as opposed to use of simplified models). In the proposed mechanism, spontaneous homolytic Al–Fe dissociation from **3** releases $[\text{Fp}]^\bullet$ and forms $[\text{LAlMe}]^\bullet$ intermediate **A**. While the $[\text{LAlMe}]^\bullet$ formula

would indicate an Al(II) formal oxidation state, DFT calculation of spin density indicated that **A** is better assigned as having an Al(III) centre bound to the $1e^-$ reduced form of **L**. As such, the redox non-innocence of **L** must contribute significantly to the low homolytic BDE of **3**. Coordination of CO_2 to the Lewis acidic Al(III) site in **A** produces $[\text{LAl}(\text{Me})(\text{O}=\text{C}=\text{O})]^\bullet$ intermediate **B**. Finally, net $2e^-$ reduction of CO_2 involves coupled delivery of $1e^-$ from the non-innocent **L** ligand and $1e^-$ from $[\text{Fp}]^\bullet$ to generate κ^1 -metallacarboxylate **C**, which rearranges to κ^2 -isomer **4**.

In support of this mechanistic proposal, a synthetic model of **B**, namely $[\text{LAl}(\text{Me})(\text{O}=\text{CPh}_2)]^\bullet$, was isolated experimentally by replacing CO_2 with benzophenone and characterised spectroscopically.⁵⁹ An alternative mechanism involving heterolytic dissociation of **3** to the $[\text{LAlMe}]^+[\text{Fp}]^-$ ion pair followed by FLP-like CO_2 activation was ruled out based on the high heterolytic BDE calculated by DFT (see above) as well as FRP-like regioselectivity during ring opening of propylene oxide by **3**.⁶³ A concerted, $[2 + 2]$ -like mechanism (such as that discussed in the previous section for **1** proposed by Sorbelli *et al.* and Guo *et al.*) involving direct engagement of CO_2 with the covalent Al–Fe bond of **3** was ruled out based on the large and positive entropy of activation for CO_2 insertion measured experimentally, as well as the fact that the Al–Fe homolytic BDE was computed to be lower than the $[2 + 2]$ activation energy by ~ 15 kcal mol^{−1}.

Conclusions

Since heterobinuclear metal–metal bonds between p-block and d-block metals are historically less studied than metal–metal bonds between two d-block metals, it is unsurprising that frontier Al–M complexes like **1** and **3** would challenge traditional bonding paradigms, require careful iterations of interpretation and re-interpretation, and give rise to unconventional reaction pathways. As briefly reviewed here, complex **1** was originally assigned as having a polar $\text{Al}^{\delta+}\text{--Au}^{\delta-}$ bond featuring a nucleophilic gold centre assigned as formally Au(–I).

Subsequent re-evaluations led to reassignment as a covalently bonded, nucleophilic [Al–Au] unit best formulated as involving formal Al(II) and Au(0) centres. Lessons from the analysis of **1** and its reaction pathways informed studies of Al–Fe complex **3**, which in turn provided additional lessons for the community to consider.

When analysing the nature of Al–M bonds, certain trends may be emerging based on case studies presented here:

- Al–M bonds tend to be more covalent than analogous M–M bonds involving only d-block metals. This may be due to the stronger valence orbital overlap imparted by greater s/p character at aluminium relative to predominantly d character of a transition metal substitute.

- These covalent Al–M bonds are best analysed theoretically relative to their neutral (radical) fragments, even if they are synthesised from ionic precursors experimentally. Often, this is evident by comparing homolytic vs. heterolytic BDEs computationally.

- Atomic electronegativity values can be misleading since they do not necessarily reflect the properties of an atom in a molecular setting.

- Partial atomic charge calculations are method-dependent,⁴³ making their absolute values sometimes misleading. It is more appropriate to focus on trends in partial atomic charges across a series of interrelated compounds.

- Al–M bonds can involve anomalously high levels of dispersion forces.²¹ Partly for this reason, it can be useful to base theoretical analyses on full molecules rather than truncated model systems.

Additionally, some lessons emerge from these case studies regarding the analysis of reaction pathways (e.g. CO₂ activation):

- For CO₂ activation by Al–M bonds, it is not necessary for there to be an accumulation of negative charge at M. Additionally, the formation of a C-bound metallacarboxylate does not necessarily imply that the M centre is, by itself, a nucleophilic site.

- For covalently bonded [Al–M] units, the assignment of oxidation states can be irrelevant or even misleading for interpreting reactivity behaviour. A parallel can be drawn to the Enemark–Feltham system for analysing covalent [M–NO]ⁿ units.⁶⁴

- When making proposals about reaction mechanisms, it is more reliable to analyse *changes* in computed parameters over an entire reaction coordinate rather than focusing on the reactant Al–M bond itself.

- Where possible, one should measure experimental observables (e.g., activation parameters) reporting on the nature of a transition state to validate theoretical conclusions and help rule out alternative mechanisms.

Given the vibrant activity in the Al–M space, we expect new discoveries in small molecule activation and catalysis to continue emerging from studies of these fascinating metal–metal bonds. We hope that the brief reflection presented here will aid this research community as it continues to progress.

Author contributions

S. M. S. conceptualised the article and wrote the first draft. N. P. M. edited the manuscript and acquired funding.

Data availability

No primary research results have been included and no new data were generated as part of this review.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Grant DE-SC0021055.

References

- 1 J. Campos, *Nat. Rev. Chem.*, 2020, **4**, 696–702.
- 2 I. G. Powers and C. Uyeda, *ACS Catal.*, 2017, **7**, 936–958.
- 3 J. A. Chipman and J. F. Berry, *Chem. Rev.*, 2020, **120**, 2409–2447.
- 4 B. G. Cooper, J. W. Napoline and C. M. Thomas, *Catal. Rev.*, 2012, **54**, 1–40.
- 5 H.-C. Yu and N. P. Mankad, *Synthesis*, 2020, 1409–1422.
- 6 S. Sinhababu, Y. Lakliang and N. P. Mankad, *Dalton Trans.*, 2022, **51**, 6129–6147.
- 7 R. C. Cammarota, L. J. Clouston and C. C. Lu, *Coord. Chem. Rev.*, 2017, **334**, 100–111.
- 8 R. M. Charles and T. P. Brewster, *Coord. Chem. Rev.*, 2021, **433**, 213765.
- 9 R. P. Singh, S. Sinhababu and N. P. Mankad, *ACS Catal.*, 2023, **13**, 12519–12542.
- 10 N. Hara, K. Semba and Y. Nakao, *ACS Catal.*, 2022, **12**, 1626–1638.
- 11 Y.-X. Luan and M. Ye, *Chem. Commun.*, 2022, **58**, 12260–12273.
- 12 Y. Nakao, *Chem. Rev.*, 2021, **121**, 327–344.
- 13 R. Y. Kong and M. R. Crimmin, *Dalton Trans.*, 2021, **50**, 7810–7817.
- 14 N. Gorgas, A. J. P. White and M. R. Crimmin, *J. Am. Chem. Soc.*, 2022, **144**, 8770–8777.
- 15 N. Gorgas, A. J. P. White and M. R. Crimmin, *Chem. Commun.*, 2022, **58**, 10849–10852.
- 16 B. Stadler, N. Gorgas, A. J. P. White and M. R. Crimmin, *Angew. Chem., Int. Ed.*, 2023, **62**, e202219212.
- 17 R. C. Cammarota and C. C. Lu, *J. Am. Chem. Soc.*, 2015, **137**, 12486–12489.

- 18 A. B. Thompson, D. R. Pahls, V. Bernales, L. C. Gallington, C. D. Malonzo, T. Webber, S. J. Tereniak, T. C. Wang, S. P. Desai, Z. Li, I. S. Kim, L. Gagliardi, R. L. Penn, K. W. Chapman, A. Stein, O. K. Farha, J. T. Hupp, A. B. F. Martinson and C. C. Lu, *Chem. Mater.*, 2016, **28**, 6753–6762.
- 19 B. J. Graziano, M. V. Vollmer and C. C. Lu, *Angew. Chem.*, 2021, **133**, 15214–15221.
- 20 M. V. Vollmer, J. Ye, J. C. Linehan, B. J. Graziano, A. Preston, E. S. Wiedner and C. C. Lu, *ACS Catal.*, 2020, **10**, 2459–2470.
- 21 K. L. Mears, C. R. Stennett, E. K. Taskinen, C. E. Knapp, C. J. Carmalt, H. M. Tuononen and P. P. Power, *J. Am. Chem. Soc.*, 2020, **142**, 19874–19878.
- 22 J. Hansen and M. Sato, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 16109–16114.
- 23 A. M. Appel, J. E. Bercaw, A. B. Bocarsly, H. Dobbek, D. L. DuBois, M. Dupuis, J. G. Ferry, E. Fujita, R. Hille, P. J. A. Kenis, C. A. Kerfeld, R. H. Morris, C. H. F. Peden, A. R. Portis, S. W. Ragsdale, T. B. Rauchfuss, J. N. H. Reek, L. C. Seefeldt, R. K. Thauer, G. L. Waldrop, A. M. Appel, J. E. Bercaw, A. B. Bocarsly, H. Dobbek, D. L. DuBois, M. Dupuis, J. G. Ferry, E. Fujita, R. Hille, P. J. A. Kenis, C. A. Kerfeld, R. H. Morris, C. H. F. Peden, A. R. Portis, S. W. Ragsdale, T. B. Rauchfuss, J. N. H. Reek, L. C. Seefeldt, R. K. Thauer and G. L. Waldrop, *Chem. Rev.*, 2013, **113**, 6621–6658.
- 24 M. D. Burkart, N. Hazari, C. L. Tway and E. L. Zeitler, *ACS Catal.*, 2019, **9**, 7937–7956.
- 25 A. Paparo and J. Okuda, *Coord. Chem. Rev.*, 2017, **334**, 136–149.
- 26 D. H. Gibson, *Chem. Rev.*, 1996, **96**, 2063–2096.
- 27 G. Fachinetti, C. Floriani and P. F. Zanazzi, *J. Am. Chem. Soc.*, 1978, **100**, 7405–7407.
- 28 S. Gambarotta, F. Arena, C. Floriani and P. F. Zanazzi, *J. Am. Chem. Soc.*, 1982, **104**, 5082–5092.
- 29 J. Hicks, A. Mansikkamäki, P. Vasko, J. M. Goicoechea and S. Aldridge, *Nat. Chem.*, 2019, **11**, 237–241.
- 30 V. R. Landaeta, T. M. Horsley Downie and R. Wolf, *Chem. Rev.*, 2024, **124**, 1323–1463.
- 31 R. F. W. Bader, *Acc. Chem. Res.*, 1985, **18**, 9–15.
- 32 R. F. W. Bader, *Atoms in molecules: a quantum theory*, Clarendon Press, Oxford University Press, Oxford, England, New York, 1994.
- 33 F. Weinhold, *J. Comput. Chem.*, 2012, **33**, 2363–2379.
- 34 M. Mitoraj and A. Michalak, *J. Mol. Model.*, 2007, **13**, 347–355.
- 35 D. Sorbelli, L. Belpassi and P. Belanzoni, *J. Am. Chem. Soc.*, 2021, **143**, 14433–14437.
- 36 C. McManus, J. Hicks, X. Cui, L. Zhao, G. Frenking, J. M. Goicoechea and S. Aldridge, *Chem. Sci.*, 2021, **12**, 13458–13468.
- 37 H. Liu, R. J. Schwamm, M. S. Hill, M. F. Mahon, C. L. McMullin and N. A. Rajabi, *Angew. Chem., Int. Ed.*, 2021, **60**, 14390–14393.
- 38 H.-Y. Liu, S. E. Neale, M. S. Hill, M. F. Mahon and C. L. McMullin, *Dalton Trans.*, 2022, **51**, 3913–3924.
- 39 M. M. D. Roy, J. Hicks, P. Vasko, A. Heilmann, A.-M. Baston, J. M. Goicoechea and S. Aldridge, *Angew. Chem., Int. Ed.*, 2021, **60**, 22301–22306.
- 40 M. P. Mitoraj, A. Michalak and T. Ziegler, *J. Chem. Theory Comput.*, 2009, **5**, 962–975.
- 41 G. Bistoni, L. Belpassi and F. Tarantelli, *J. Chem. Theory Comput.*, 2016, **12**, 1236–1244.
- 42 G. Bistoni, S. Rampino, F. Tarantelli and L. Belpassi, *J. Chem. Phys.*, 2015, **142**, 084112.
- 43 M. Cho, N. Sylvetsky, S. Eshafi, G. Santra, I. Efremenko and J. M. L. Martin, *ChemPhysChem*, 2020, **21**, 688–696.
- 44 D. Sorbelli, L. Belpassi and P. Belanzoni, *Inorg. Chem.*, 2022, **61**, 1704–1716.
- 45 X. Guo, T. Yang, Y. Zhang, F. K. Sheong and Z. Lin, *Inorg. Chem.*, 2022, **61**, 10255–10262.
- 46 G. Knizia, *J. Chem. Theory Comput.*, 2013, **9**, 4834–4843.
- 47 N. P. Mankad, *Chem. Commun.*, 2018, **54**, 1291–1302.
- 48 N. J. Leon, H.-C. Yu, T. J. Mazzacano and N. P. Mankad, *Synlett*, 2020, 125–132.
- 49 K. B. Wiberg, *Tetrahedron*, 1968, **24**, 1083–1096.
- 50 M. K. Karunananda, F. X. Vázquez, E. E. Alp, W. Bi, S. Chattopadhyay, T. Shibata and N. P. Mankad, *Dalton Trans.*, 2014, **43**, 13661–13671.
- 51 S. Banerjee, M. K. Karunananda, S. Bagherzadeh, U. Jayarathne, S. R. Parmelee, G. W. Waldhart and N. P. Mankad, *Inorg. Chem.*, 2014, **53**, 11307–11315.
- 52 U. Jayarathne, T. J. Mazzacano, S. Bagherzadeh and N. P. Mankad, *Organometallics*, 2013, **32**, 3986–3992.
- 53 D. W. Stephan, *Science*, 2016, **354**, aaf7229.
- 54 A. R. Jupp and D. W. Stephan, *Trends Chem.*, 2019, **1**, 35–48.
- 55 M. K. Karunananda, S. R. Parmelee, G. W. Waldhart and N. P. Mankad, *Organometallics*, 2015, **34**, 3857–3864.
- 56 S. R. Parmelee, T. J. Mazzacano, Y. Zhu, N. P. Mankad and J. A. Keith, *ACS Catal.*, 2015, **5**, 3689–3699.
- 57 M. K. Karunananda and N. P. Mankad, *Organometallics*, 2017, **36**, 220–227.
- 58 D. R. Pye, L.-J. Cheng and N. P. Mankad, *Chem. Sci.*, 2017, **8**, 4750–4755.
- 59 S. Sinhababu, M. R. Radzhabov, J. Telser and N. P. Mankad, *J. Am. Chem. Soc.*, 2022, **144**, 3210–3221.
- 60 M. Ju, Z. Lu, L. F. T. Novaes, J. I. Martinez Alvarado and S. Lin, *J. Am. Chem. Soc.*, 2023, **145**, 19478–19489.
- 61 L. Liu, L. L. Cao, Y. Shao, G. Ménard and D. W. Stephan, *Chem*, 2017, **3**, 259–267.
- 62 A. Dasgupta, E. Richards and R. L. Melen, *Angew. Chem., Int. Ed.*, 2021, **60**, 53–65.
- 63 S. Sinhababu, R. P. Singh, M. R. Radzhabov, J. Kumawat, D. H. Ess and N. P. Mankad, *Nat. Commun.*, 2024, **15**, 1315.
- 64 J. H. Enemark and R. D. Feltham, *Coord. Chem. Rev.*, 1974, **13**, 339–406.