

PAPER

View Article Online
View Journal | View IssueCite this: *Dalton Trans.*, 2023, **52**,
1820Received 22nd December 2022,
Accepted 13th January 2023

DOI: 10.1039/d2dt04095f

rsc.li/dalton

Synthesis and lewis acidity of fluorinated triaryl borates†

Mashaal M. Alharbi,^{†a,b} Yara van Ingen,^{†a} Alberto Roldan,^a Tanja Kaehler^{*a}
and Rebecca L. Melen^{†a*}

A series of fluorinated triaryl borates $B(OAr^F)_3$ ($Ar^F = 2\text{-FC}_6\text{H}_4$, $3\text{-FC}_6\text{H}_4$, $4\text{-FC}_6\text{H}_4$, $2,4\text{-F}_2\text{C}_6\text{H}_3$, $3,5\text{-F}_2\text{C}_6\text{H}_3$, $2,3,4\text{-F}_3\text{C}_6\text{H}_2$, $2,4,6\text{-F}_3\text{C}_6\text{H}_2$, $3,4,5\text{-F}_3\text{C}_6\text{H}_2$) have been prepared and isolated from the reactions of the mono-, di-, or tri-fluorophenol with BCl_3 . The Lewis acidity of these borates has been determined by NMR spectroscopic and theoretical methods and compared to their well-established borane counterpart.

Introduction


Since 1923,¹ when Gilbert Lewis first suggested the concept of Lewis acids and bases, Lewis acids have been employed in all areas of chemistry in both the laboratory and on industrial scale. They are commonly employed to activate substrates both stoichiometrically and catalytically through adduct formation, lowering the LUMO of the reactive species. From a bond activation perspective, there is a strong correlation between the strength of a Lewis acid and substrate activation. However, this is not always observed when looking at an entire catalytic cycle.² Nevertheless, there has been a drive to develop strong Lewis acids for applications in catalysis. One particular area where strong Lewis acids have been developed is for applications in frustrated Lewis pair (FLP) chemistry.^{3–7} The strength of a Lewis acid can be quantified by several means, including NMR spectroscopy and computational methods.⁸ The Gutmann–Beckett^{9,10} and Childs¹¹ methods rely on adding an internal probe Et_3PO or crotonaldehyde (CA) and measuring the change in chemical shift ($\Delta\delta$) in the ^{31}P (Et_3PO) or ^1H (H3 atom of CA) NMR spectra respectively. Computational methods to quantify the Lewis acidity often investigate the fluoride ion affinity (FIA) or the hydride ion affinity (HIA).^{8,12} The latter being better probe for softer Lewis acids. Most often, the strength of a Lewis acid is defined by its

FIA value, and a molecule with an FIA stronger than SbF_5 (501 kJ mol^{-1}) is considered a Lewis superacid.⁸

First synthesised in the 1960s,¹³ $B(\text{C}_6\text{F}_5)_3$ is one of the most commonly used Lewis acids in FLP chemistry.^{3–7,14,15} However, with an FIA of 452 kJ mol^{-1} , $B(\text{C}_6\text{F}_5)_3$ falls short of being classified as a Lewis superacid.¹⁶ The heavier group 13 analogue $\text{Al}(\text{C}_6\text{F}_5)_3$ with an FIA of 530 kJ mol^{-1} , on the other hand, is a Lewis superacid.¹⁷ Much research has focused on making derivatives of $\text{E}(\text{C}_6\text{F}_5)_3$ ($\text{E} = \text{B}, \text{Al}$) with different fluorine or chlorine substitution patterns on the aryl rings (Ar^F or Ar^{Cl}).¹⁸ Adding an electronegative oxygen or nitrogen spacer before the

Fluorinated B/Al Lewis Acids		FIA (kJ mol^{-1})	Et_3PO $\Delta\delta$ (ppm)
F substitution	BPh_3	320 ²⁴	20.6 ³⁶
	$\text{B}(2,3,4\text{-F}_3\text{C}_6\text{H}_2)_3$	404 ²⁴	–
	$\text{B}(3,4,5\text{-F}_3\text{C}_6\text{H}_2)_3$	427 ²⁴	27.1 ²⁴
	$\text{B}(\text{C}_6\text{F}_5)_3$	452 ¹⁷	25.5 ¹⁷
	$(\text{C}_6\text{F}_5)_2\text{BOC}_6\text{F}_5$	–	33.6 ²²
O/N spacer	$\text{C}_6\text{F}_5\text{B}(\text{OC}_6\text{F}_5)_2$	–	34.1 ²²
	$\text{B}(\text{OC}_6\text{F}_5)_3$	419 ¹²	34.5 ²²
	$\text{Al}(\text{C}_6\text{F}_5)_3$	530 ¹⁷	–
	$\text{Al}(\text{OC}(\text{CF}_3)_3)_3$	547 ²⁰	–
	$\text{Al}(\text{O}(\text{C}(\text{C}_6\text{F}_5)_3)_3$	555 ²⁰	–
	$\text{Al}(\text{N}(\text{C}_6\text{F}_5)_2)_3$	555 ¹⁹	–

This work:



Scheme 1 Group 13 Lewis acids and their Lewis acidities from Fluoride Ion Affinity and Gutmann–Beckett (Et_3PO) metrics.

^aCardiff Catalysis Institute, School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT Cymru/Wales, UK. E-mail: KaehlerT@cardiff.ac.uk, MelenR@cardiff.ac.uk

^bDepartment of Chemistry, King Faisal University, College of Science, P.O. Box 400, Al-Ahsa 31982, Saudi Arabia

†Electronic supplementary information (ESI) available: Experimental procedures, NMR spectra, DFT data, and X-ray data. CCDC 2208652 (1) 2208553 (7) and 2208552 (8). For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d2dt04095f>

‡Both authors contributed equally.




halogenated aryl ring has been shown to increase the Lewis acidity (Scheme 1). For example, Sundermeyer reported an $\text{Al}(\text{N}(\text{C}_6\text{F}_5)_2)_3$ FIA Lewis acidity of 555 kJ mol^{-1} .¹⁹ Other notable aluminium Lewis superacids include Krossing's $\text{Al}(\text{OC}(\text{CF}_3)_3)_3$ (FIA = 547 kJ mol^{-1})¹⁶ and Beckmann's $\text{Al}(\text{OC}(\text{C}_6\text{F}_5)_3)_3$ (FIA = 555 kJ mol^{-1}).²⁰ Relating to this study, $\text{B}(\text{OC}_6\text{F}_5)_3$ was first reported in 1992 by Naumann.²¹ Britovsek later prepared the intermediate borinic $[(\text{C}_6\text{F}_5)_2\text{BOC}_6\text{F}_5]$ and boronic $[\text{C}_6\text{F}_5\text{B}(\text{OC}_6\text{F}_5)_2]$ acids and analysed their Lewis acidity from Gutmann–Beckett method ($\Delta\delta \text{ Et}_3\text{PO} = 33.6, 34.1 \text{ ppm}$, respectively).²² The increased Lewis acidity of the boron centre when including an O-atom spacer is attributed to the reduced $\text{p}\pi\text{--p}\pi$ bonding between the oxygen and the boron centre due to the high electron-withdrawing nature of the C_6F_5 groups. Surprisingly, the FIA for $\text{B}(\text{OC}_6\text{F}_5)_3$ is 419 kJ mol^{-1} , significantly lower than its borane counterpart $\text{B}(\text{C}_6\text{F}_5)_3$.¹² In contrast, the reported Gutmann–Beckett value of $\Delta\delta = 34.5 \text{ ppm}$ (ref. 22) for $\text{B}(\text{OC}_6\text{F}_5)_3$ showed a higher Lewis acidity when compared to $\Delta\delta = 25.5 \text{ ppm}$ for $\text{B}(\text{C}_6\text{F}_5)_3$.²¹

This discrepancy between computational and experimental methods for the Lewis acidity when including an O/N-atom spacer has not previously been investigated to the best of our knowledge. Previous studies in our laboratory have focused on synthesising $\text{E}(\text{Ar}^{\text{F}})_3$ compounds where $\text{Ar}^{\text{F}} \neq \text{C}_6\text{F}_5$.^{23,24} These derivatives have sometimes shown increased catalytic activity and/or stability, for example, for $\text{B}(3,4,5\text{-F}_3\text{C}_6\text{H}_2)_3$ (Scheme 1).^{23–27} In the present study, we were interested in synthesising a series of $\text{B}(\text{OAr}^{\text{F}})_3$ derivatives of varying Lewis acidities.

The fluorinated triaryl borates **1–8** were synthesised by the general procedure from the 3:1 stoichiometric reaction of mono-, di-, or tri-fluorophenol with BCl_3 . The fluorinated triaryl borate products were then purified by crystallisation from a concentrated solution in pentane at -35°C and washed with either cold pentane or CH_2Cl_2 to give the products in 74–91% yields (Scheme 2). All the compounds showed similar ^{11}B NMR chemical shifts for the boron environment, with a broad peak appearing between $\delta = 15.7\text{--}17.7 \text{ ppm}$ in the ^{11}B NMR spectrum (Table 1), in good agreement with previously reported $\text{B}(\text{OPh})_3$ ²⁸ ($\delta = 16.5 \text{ ppm}$) and $\text{B}(\text{OC}_6\text{F}_5)_3$ ²⁹ ($\delta = 15 \text{ ppm}$).

With a series of fluorinated borates in hand, we next investigated their Lewis acidity and compared them to literature reported $\text{B}(\text{OC}_6\text{F}_5)_3$, $\text{B}(\text{OPh})_3$, and $\text{B}(\text{C}_6\text{F}_5)_3$. Firstly, we analysed the Lewis acidity by use of the Gutmann–Beckett method. Et_3PO (0.6 equiv.) was added to an NMR tube containing PPh_3 as internal standard, and the fluorinated triaryl borates **1–8** (1.0 equiv.) in CDCl_3 to generate $\text{Et}_3\text{P}=\text{O} \rightarrow \text{B}(\text{OAr}^{\text{F}})_3$. The ^{31}P NMR chemical shift ($\delta \text{ ppm}$) was measured as well as the change in ^{31}P NMR chemical shift between the free phosphine oxide ($\delta = 52.5 \text{ ppm}$) and the adduct ($\Delta\delta$) (Table 1). The Lewis acidity was found to decrease in the order $6 = 7 > 8 > 5 > 1 > 4 > 2 > 3$, from which it can be deduced that reducing the number of electron withdrawing F-atoms decreases the Lewis acidity, however there appeared to be little correlation for the



Scheme 2 Synthesis of fluorinated triaryl borates and commercially available $\text{B}(\text{OPh})_3$.

substitution pattern. When comparing the synthesised borates to previously reported borates and boranes, the following trend is observed; $\text{B}(\text{OC}_6\text{F}_5)_3 > \mathbf{8}, \mathbf{7}, \mathbf{6}, \mathbf{5} > \text{B}(\text{C}_6\text{F}_5)_3 \gg \text{B}(\text{OPh})_3$. These results align with Britovsek's findings, *i.e.*, introducing an O-atom spacer increases Lewis acidity and verifies the importance of tuning acidity through fluorine substitutions.²²

The Lewis acidity of compounds **1–8** was also experimentally determined *via* the Childs method using CA as the Lewis base. However, only slight differences in the chemical shift of the H3 atom of CA in the ^1H NMR spectra were observed, $\Delta\delta = 0.02\text{--}0.03 \text{ ppm}$, when comparing fluorination patterns of the borates (Table 1). A comparison of the change in chemical shift ($\Delta\delta$) of CA for the perfluorinated borate $\text{B}(\text{OC}_6\text{F}_5)_3$, at 0.40 ppm, to the perfluorinated borane $\text{B}(\text{C}_6\text{F}_5)_3$, at 1.05 ppm, suggests that the borane is a stronger Lewis acid than the borate from the Childs method, a difference with the Gutmann–Beckett method. An expression of the Gutmann–Beckett and Childs metrics as percentages of $\text{B}(\text{C}_6\text{F}_5)_3$ are presented in Table 2. The hard soft acid base theory (HSAB) introduced by Pearson invokes the charge density and polarizability of an acid or base, and can be used to make qualitative predictions of acids and bases and how they may prefer to react.^{30–32} Et_3PO is considered hard due to its highly polarised $\text{P}=\text{O}$ bond, whereas CA is softer, hence the greater $\Delta\delta$ upon adduct



Table 1 Lewis acidity measurements for fluorinated triaryl borates **1–8** and previously reported B(OC₆F₅)₃, B(OPh)₃, and B(C₆F₅)₃

Compound	¹¹ B ^a (ppm)	Et ₃ PO ³¹ P ^a (ppm)	Et ₃ PO Δδ ^a (ppm)	CA ¹ H ^a (ppm)	CA Δδ ^a (ppm)	FIA ^b (kJ mol ^{−1})	HIA ^b (kJ mol ^{−1})	GEI ^c (eV)
1	16.5	76.3	23.8	6.87	0.02	351	339	0.88
2	15.9	74.5	22.0	— ^d	— ^d	368	340	1.19
3	16.3	66.6	14.1	6.87	0.02	359	329	1.09
4	16.7	76.2	23.7	— ^d	— ^d	376	360	1.24
5	15.6	78.9	26.4	6.87	0.02	404	377	1.31
6	16.5	79.6	27.1	6.88	0.03	402	374	1.30
7	17.7	79.6	27.1	6.88	0.03	372	347	1.23
8	15.7	79.1	26.6	6.88	0.03	433	414	1.45
B(OC ₆ F ₅) ₃	15 ²⁹	80.9 ²²	34.5 ²²	7.25 ³³	0.40 ²²	431	411	1.81
B(OPh) ₃	16.5 ²⁸	69.4 ²²	23.0 ²²	6.88 ²²	0.03 ²²	350	323	0.81
B(C ₆ F ₅) ₃	59 ²¹	78.0 ³³	25.5 ³³	7.90 ³³	1.05 ³³	454	497	3.65

^a In CDCl₃. ^b PBEh-3c//M06-2X/def2-QZVPP level. ^c B3LYP/def2-TZVP level. ^d Not determined.

Table 2 Comparison of Lewis acidity metrics of B(OC₆F₅)₃, **7** and **8** expressed as percentages relative to B(C₆F₅)₃

Compound	Et ₃ PO Δδ	CA Δδ	FIA	HIA	GEI
B(C ₆ F ₅) ₃	100%	100%	100%	100%	100%
B(OC ₆ F ₅) ₃	135%	38%	94%	82%	49%
7	78%	2%	81%	76%	33%
8	104%	2%	95%	83%	39%

formation of the borate with Et₃PO than CA would suggest that the borate is a harder Lewis acid than the borane. A key step in both these experimental Lewis acidity metrics is the pyramidalisation of the boron centre from trigonal planar to tetrahedral upon adduct formation with a Lewis base. The introduction of an O-spacer likely affects the ease of pyramidalization from two perspectives: the steric changes upon a larger distance between the B atom to the phenyl ring in borates when compared to boranes, as well as the attractive dispersion effects between the Lewis base and the O and fluorinated phenyl groups in the ligand sphere of the B centre. The smaller fluoride and hydride bases were considered to further probe these effects through the use of fluoride and hydride ion affinity (FIA/HIA) metrics.

Fluoride and hydride ion affinity (FIA/HIA) are computationally derived metrics of Lewis acidity, defined as the enthalpy of the reaction (or pairing) of a Lewis acid with a Lewis base (F[−] or H[−], respectively). Since the FIA of a Lewis acid considers the *hard* Lewis base F[−], this represents *hard* Lewis acidity, whilst the *soft* H[−] base, and consequently HIA of a Lewis acid, reflects *soft* Lewis acidity properties.

Both fluoride and hydride anions are small and lack directionality as a base upon coordination to an acid, a key difference in the pyramidalisation step to the bases considered in the Gutmann–Beckett and Childs metrics. Hence, the FIA/HIA values consider Lewis acidity with limited to no influence of effects of steric interactions or π -interactions, as highlighted by Greb in previous work on ion affinity metrics.^{12,34}

To determine the FIAs/HIAs, single point calculations at the M06-2X+D3(0)/def2-QZVPP level of theory^{35–37} were performed on the compounds optimised geometries at PBEh-3c,³⁸ previously VSEPR-optimised. Isodesmic affinities were obtained through anchoring the enthalpies to reference systems (see

ESI, section S6.2 for details†).^{12,39} A general trend of increasing FIA and HIA is observed with an increasing number of F-substituents, giving **1**, **3**, **2** < **4**, **5** < **6** < B(OC₆F₅)₃ < **8**. When the same number of F-atoms are present,⁴⁰ *meta*-substitution of fluorine induces higher FIA and HIAs than *para*- than *ortho*-substitution giving a trend of **1**, **3** < **2**, **4** < **5**, and **7** < **6** < **8**. Interestingly, **7** with a 2,4,6-F substitution has a lower Lewis acidity than compounds **4** and **5** which only have two F-atoms on the Ar^F rings. Likewise, the 3,4,5-F substitution in **8** gives higher FIA and HIA values than B(OC₆F₅)₃, at 433 and 431 kJ mol^{−1} (FIA) compared to 414 and 411 kJ mol^{−1} (HIA), respectively. In both cases, the presence of two *ortho*-F atoms appears to diminish the Lewis acidity.

Surprisingly, and in contrast to the Gutmann–Beckett and Childs methods, the FIA and HIA of borate B(OC₆F₅)₃ are significantly lower than that of borane B(C₆F₅)₃ (Table 2). As mentioned, the pyramidalisation of the boron centre is required for acid–base adduct formation, one that is unaffected by steric and dispersion effects of the Lewis base in the FIA and HIA metrics. The slightly lower Lewis acidity in FIA and HIA than for Gutmann–Beckett and Childs metrics for the borates, hence, suggest that the Lewis acid strength in the experimental methods is governed by either steric or dispersion effects or a combination of the two, as a result of the introduction of the O-spacer and its resultant ligand sphere interaction with the Lewis base. The two most widely accepted and applied metrics, FIA and Gutmann–Beckett are compared in Fig. 1, showing opposing strengths of Lewis acidity for the borates than the boranes.

A comparison between the hard FIA and soft HIA, by representing these strengths relative to B(C₆F₅)₃, show again that the borate is a harder Lewis acid than the borane (Table 2) the FIA of B(OC₆F₅)₃ is 94%, and the HIA is 82%. Furthermore, the inverse of HIA is hydride donor ability,⁴¹ and the involvement of electrophilic boron centres in stoichiometric and catalytic transformation of basic substrates often relies on a balance between Lewis acidity and hydride donor ability, hence the softer HIA is relevant in context other than Lewis acidity strength.

Another computational comparison of Lewis acidity is given by the global electrophilicity index (GEI).^{42,43} This index considers intrinsic Lewis acidity, independent from the coordi-






Fig. 1 Comparison of FIA and Gutmann–Beckett values for boranes and borates.





Fig. 2 Comparison of GEI, HIA, and FIA values for boranes and borates.

nation of a Lewis base, as a measure of the Lewis acid's ability to accept electrons by considering the HOMO and LUMO energy levels in the form of chemical hardness⁴² and Mulliken electronegativity^{44–46} (see ESI, section S6.3†). In this work, the Lewis acids were re-optimised at the B3LYP/def2-TZVP theory level,^{47,48} and the E_{HOMO} and E_{LUMO} were taken as the energies of corresponding Kohn–Sham orbitals.


In $\text{B}(\text{C}_6\text{F}_5)_3$, the LUMO is formed of a 2p-orbital that is orthogonal to the trigonal plane of the borane (Fig. 3A); in comparison, $\text{B}(\text{OC}_6\text{F}_5)_3$ does not exhibit a localised 2p-orbital as the LUMO (Fig. 3B). Instead, a delocalised orbital situated across the B–O–C atoms is observed, seemingly a distortion from the vacant 2p-orbital of the B atom as a likely result of a degree of $p\pi$ – $\pi\pi$ donation from O to B (Fig. 3B). The LUMO+1 of $\text{B}(\text{OC}_6\text{F}_5)_3$ does display a p-orbital perpendicular to the trigonal plane (Fig. 3C), yet most of the LUMO and LUMO+1 is localised on the ligand-sphere. Analysis of the unoccupied orbitals for all optimised borates showed that the most localised p-orbital on the B atom is in the LUMO+1, hence GEI values were calculated with the LUMO+1 for the borates (see ESI, Table S5† for a comparison), as has been done previously by Chitnis and co-workers.⁴⁹ The GEI values for the varying substitution patterns follow the same general trend as for FIA and HIA (Fig. 2). A lower GEI value (Table 2) for $\text{B}(\text{OC}_6\text{F}_5)_3$ in comparison to $\text{B}(\text{C}_6\text{F}_5)_3$ is consistent with the computed LUMO+1 of $\text{B}(\text{OC}_6\text{F}_5)_3$ lying at a higher energy than the LUMO of $\text{B}(\text{C}_6\text{F}_5)_3$ (see ESI, Table S5†).

An intrinsic Lewis acidity scale like GEI has benefits as it is readily available at low computational cost. However, whilst giving relevant insight into the unoccupied orbitals of the Lewis acid that participate in reactivity with a Lewis base, it in fact does not reflect the reactivity, and hence must be treated with some caution. Furthermore, computational metrics such as presented FIA and HIA were shown to give some insight into hard and soft characteristics of Lewis acids when com-

A. LUMO of $\text{B}(\text{C}_6\text{F}_5)_3$



B. LUMO of $\text{B}(\text{OC}_6\text{F}_5)_3$



C. LUMO+1 of $\text{B}(\text{OC}_6\text{F}_5)_3$




Fig. 3 Visual representation (side-view) of the: LUMO of $\text{B}(\text{C}_6\text{F}_5)_3$, top; the LUMO of $\text{B}(\text{OC}_6\text{F}_5)_3$, middle; and the LUMO+1 of $\text{B}(\text{OC}_6\text{F}_5)_3$ bottom, calculated at B3LYP/def2-TZVP level of theory (top-view can be found in the ESI, section S6.4†).



pared to each other. These affinity metrics can be used to describe the strength of a Lewis acid from a theoretical perspective but the Lewis base lacks directionality.

Compounds **1**, **7**, and **8** could be recrystallised from CH_2Cl_2 and their structures determined by single crystal X-ray diffraction (Fig. 4). The three-component positional disorder could be resolved for **1**, of which the major component is displayed in Fig. 1. As a result, the structural parameters of **1** will not be discussed. For compounds **7** and **8**, the boron centre occupies a trigonal planar geometry, with the boron atom lying merely 0.019 (**7**) and 0.012 Å (**8**) out of the O3-plane. This slight deviation implies higher planarity than exhibited in $\text{B}(\text{OC}_6\text{F}_5)_3$ (0.03 Å)²² and is comparable to that of $\text{B}(\text{OC}_6\text{H}_5)_3$ (0.014 Å),⁵⁰ as would be expected for lower F-substitution.

Varying Lewis acidity is expected from a $\text{p}\pi\text{-p}\pi$ donation from oxygen to boron reflected by the coplanarity between B and O, with a higher contribution reducing Lewis acidity. We assessed this by comparing the torsion angle between O1–O2–O3 and B–O–C planes of **7** and **8**, and that of $\text{B}(\text{OC}_6\text{F}_5)_3$ and $\text{B}(\text{OC}_6\text{H}_5)_3$. Torsion angles in **8** are smaller (B–O1 = 4°, B–O2 = 5°, B–O3 = 4°) than in $\text{B}(\text{OC}_6\text{F}_5)_3$, for which Britovsek observed one larger torsion angle (B–O1 = 5°, B–O2 = 21°, B–O3 = 7°),²² consistent with an extent of $\text{p}\pi\text{-p}\pi$ donation in **8**, and thus higher Lewis acidity than $\text{B}(\text{OC}_6\text{F}_5)_3$. For **7**, the torsion angles are larger than in **8** (B–O1 = 11°, B–O2 = 9°, B–O3 = 6°) and

approach those observed in $\text{B}(\text{OC}_6\text{H}_5)_3$ (B–O1 = 14°, B–O2 = 14°, B–O3 = 15°).⁵⁰ The differences are consistent with a lower Lewis acidity of **7** and $\text{B}(\text{OC}_6\text{H}_5)_3$ compared to **8** and $\text{B}(\text{OC}_6\text{F}_5)_3$. However, it must be noted that a key feature that determines structural parameters is the crystal packing within the unit cell, which is in part determined by $\pi\text{-}\pi$ and F– π interactions. As such, the torsion angles and inclination angles must be treated with some level of caution. The inclination angles of the phenyl rings compared to the BO_3 plane are larger and more uniform in **7** (69°, 64°, and 64°) than in **8** (21°, 11°, and 55°). The larger inclination angle for **7** is likely a result of steric effects of the *ortho*-F atoms.

Conclusions

In conclusion, we have demonstrated a simple synthesis of a series of fluorinated triaryl borates ($\text{B}(\text{OAr}^{\text{F}})_3$), for which Lewis acidity was determined by NMR (Gutmann–Beckett and Childs) methods as well as by computational means (HIA, FIA, and GEI). The Lewis acids were found to be stronger Lewis acids than the archetypical fluorinated boranes from Gutmann–Beckett, yet weaker when comparing their frontier orbitals by computational methods. Evidence for $\text{p}\pi\text{-p}\pi$ donation from oxygen to boron was seen in the borate-delocalised LUMO and trends in structural analysis of the torsion angles for **7** and **8**. The application of these compounds in Lewis acid catalysis and frustrated Lewis pair chemistry is the focus of our future studies.

Author contributions

MA performed the experimental work. YVI performed the calculations and the crystal structure analysis. YVI, TK, and RLM wrote the manuscript. RLM directed the project, and AR directed the computer simulations. MA, YVI, and TK wrote the ESI.† All authors proofread and edited the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

MA acknowledges financial support from the Saudi Ministry of Education and the Deanship of Scientific Research at King Faisal University, Saudi Arabia (Grant No. 1591). TK and RLM would like to thank the EPSRC (EP/R026912/1). YVI is grateful to Mr Vladimir Y. Vladimirov for guidance with calculations. YVI and AR acknowledge Supercomputing Wales for access to the Hawk HPC facility, part-funded by the European Regional Development Fund *via* the Welsh Government. Information about the data that underpins the results presented in this article can be found in the Cardiff University data catalogue at <http://doi.org/10.17035/d.2023.0237563437>.




Fig. 4 Crystal structures of **1** (top), **7** (middle), and **8** (bottom). Thermal ellipsoids shown at 50%. H atoms omitted for clarity. Carbon: black; oxygen: red; fluorine: light green; boron: pink.



References

- G. N. Lewis, *Valence and the Structure of Atoms and Molecules*, The Chemical Catalog Company, New York, 1923.
- S. Woodward, *Tetrahedron*, 2002, **58**, 1017–1050.
- D. W. Stephan, *J. Am. Chem. Soc.*, 2015, **137**, 10018–10032.
- D. W. Stephan, *Science*, 2016, **354**, 1248.
- J. Lam, K. M. Szkop, E. Mosafari and D. W. Stephan, *Chem. Soc. Rev.*, 2019, **48**, 3592–3612.
- D. W. Stephan, *Chem*, 2020, **6**, 1520–1526.
- D. W. Stephan, *J. Am. Chem. Soc.*, 2021, **143**, 20002–20014.
- L. Greb, *Chem. – Eur. J.*, 2018, **24**, 17881–17896.
- U. Mayer, V. Gutmann and W. Gerger, *Chem. Mon.*, 1975, **106**, 1235–1257.
- M. A. Beckett, G. C. Strickland, J. R. Holland and K. Sukumar Varma, *Polymer*, 1996, **37**, 4629–4631.
- R. F. Childs, D. L. Mulholland and A. Nixon, *Can. J. Chem.*, 1982, **60**, 809–812.
- P. Erdmann, J. Leitner, J. Schwarz and L. Greb, *ChemPhysChem*, 2020, **21**, 987–994.
- A. G. Massey and A. J. Park, *J. Organomet. Chem.*, 1964, **2**, 245–250.
- G. Erker, *Dalton Trans.*, 2005, 1883–1890.
- J. R. Lawson and R. L. Melen, *Inorg. Chem.*, 2017, **56**, 8627–8643.
- H. Böhrer, N. Trapp, D. Himmel, M. Schleep and I. Krossing, *Dalton Trans.*, 2015, **44**, 7489–7499.
- L. O. Müller, D. Himmel, J. Stauffer, G. Steinfeld, J. Slattery, G. Santiso-Quinones, V. Brecht and I. Krossing, *Angew. Chem., Int. Ed.*, 2008, **47**, 7659–7663.
- J. L. Carden, A. Dasgupta and R. L. Melen, *Chem. Soc. Rev.*, 2020, **49**, 1706–1725.
- J. F. Kögel, D. A. Sorokin, A. Khvorost, M. Scott, K. Harms, D. Himmel, I. Krossing and J. Sundermeyer, *Chem. Sci.*, 2018, **9**, 245–253.
- J. F. Kögel, A. Y. Timoshkin, A. Schröder, E. Lork and J. Beckmann, *Chem. Sci.*, 2018, **9**, 8178–8183.
- D. Naumann, H. Butler and R. Gnann, *Z. Anorg. Allg. Chem.*, 1992, **618**, 74–76.
- G. J. P. Britovsek, J. Ugoletti and A. J. P. White, *Organometallics*, 2005, **24**, 1685–1691.
- T. Kaehler and R. L. Melen, *Cell Rep. Phys. Sci.*, 2021, **2**, 100595.
- D. M. C. Ould, J. L. Carden, R. Page and R. L. Melen, *Inorg. Chem.*, 2020, **59**, 14891–14898.
- Q. Yin, Y. Soltani, R. L. Melen and M. Oestreich, *Organometallics*, 2017, **36**, 2381–2384.
- J. R. Lawson, L. C. Wilkins and R. L. Melen, *Chem. – Eur. J.*, 2017, **23**, 10997–11000.
- J. L. Carden, L. J. Gierlichs, D. F. Wass, D. L. Browne and R. L. Melen, *Chem. Commun.*, 2019, **55**, 318–321.
- B. Wrackmeyer, *Annual Reports on NMR Spectroscopy*, Elsevier, London, 1988.
- A. Sundararaman and F. Jäkle, *J. Organomet. Chem.*, 2003, **681**, 134–142.
- R. G. Pearson, *J. Am. Chem. Soc.*, 1963, **85**, 3533–3539.
- R. G. Pearson, *J. Chem. Educ.*, 1968, **45**, 581.
- R. G. Pearson, *J. Chem. Educ.*, 1968, **45**, 643.
- I. B. Sivaev and V. I. Bregadze, *Coord. Chem. Rev.*, 2014, **270–271**, 75–88.
- P. Erdmann and L. Greb, *ChemPhysChem*, 2021, **22**, 935–943.
- F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215–241.
- F. Weigend, F. Furche and R. Ahlrichs, *J. Chem. Phys.*, 2003, **119**, 12753–12762.
- F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2012, **2**, 73–78.
- P. Erdmann and L. Greb, *ChemPhysChem*, 2021, **22**, 935–943.
- C. Lewis, B. Fluoroarylboranes, M. M. Morgan, A. J. V. Marwitz, W. E. Piers and M. Parvez, *Organometallics*, 2013, **32**, 317.
- Z. M. Heiden and A. P. Lathem, *Organometallics*, 2015, **34**, 1818–1827.
- R. G. Parr, L. v. Szentpály and S. Liu, *J. Am. Chem. Soc.*, 1999, **121**, 1922–1924.
- A. R. Jupp, T. C. Johnstone and D. W. Stephan, *Dalton Trans.*, 2018, **47**, 7029–7035.
- R. G. Parr and R. G. Pearson, *J. Am. Chem. Soc.*, 1983, **105**, 7512–7516.
- R. G. Parr, R. A. Donnelly, M. Levy and W. E. Palke, *J. Chem. Phys.*, 1978, **68**, 3801–3807.
- R. S. Mulliken, *J. Chem. Phys.*, 1934, **2**, 782–793.
- P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623–11627.
- A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648–5652.
- K. M. Marczenko, S. Jee and S. S. Chitnis, *Organometallics*, 2020, **39**, 4287–4296.
- I. M. Malkowsky, R. Frohlich, U. Griesbach, H. Pütter and S. R. Waldvogel, *Eur. J. Inorg. Chem.*, 2006, 1690–1697.

