Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2017, 8, 6679

Porphyrinoid rotaxanes: building a mechanical picket fence†

T. H. Ngo, **D**ab* J. Labuta, **D**bc* G. N. Lim, *d* W. A. Webre, *d* F. D'Souza, **D**d* P. A. Karr, *e*
J. E. M. Lewis, **D**f* J. P. Hill, **D**b** K. Ariga **D**b** and S. M. Goldup **D**f**

Building on recent progress in the synthesis of functional porphyrins for a range of applications using the Cu-mediated azide—alkyne cycloaddition (CuAAC) reaction, we describe the active template CuAAC synthesis of interlocked triazole functionalised porphyrinoids in excellent yield. By synthesising interlocked analogues of previously studied porphyrin—corrole conjugates, we demonstrate that this approach gives access to rotaxanes in which the detailed electronic properties of the axle component are unchanged but whose steric properties are transformed by the mechanical "picket fence" provided by the threaded rings. Our results suggest that interlocked functionalised porphyrins, readily available using the AT-CuAAC approach, are sterically hindered scaffolds for the development of new catalysts and materials.

Received 19th July 2017 Accepted 3rd August 2017

DOI: 10.1039/c7sc03165c

rsc.li/chemical-science

Introduction

The copper-mediated alkyne–azide cycloaddition (CuAAC)¹ reaction has been widely applied in the synthesis of functionalised porphyrinoid macrocycles,² including multi-porphyrinoid arrays,³ for applications in biology,⁴ energy transfer,⁵ catalysis,⁶ self-assembly,⁻ and sensing.⁵ Indeed, the broad functional group tolerance, mild conditions and readily available starting materials of this archetypal "click"⁵ reaction make it an ideal tool for the synthesis of complex non-natural products.

The CuAAC reaction has also been widely applied in the synthesis of interlocked molecules, 10 including examples of rotaxanes and catenanes containing porphyrin sub-units. 11,12 To achieve this, triazole formation is often the final step that captures the interlocked structure by introducing a stopper unit in the case of rotaxanes or closing a macrocycle in the case of catenanes. A general feature of such "passive template" syntheses, 13 in which non-covalent interactions pre-organise the

covalent subcomponents in a threaded architecture prior to the CuAAC reaction, is that additional functionality must be included in the covalent structure of both components to provide the required pre-organisation. These functional groups remain in the interlocked product and, although the intercomponent interactions they often engender can be exploited in the design of molecular machines, ¹³ this approach imposes structural limitations on the products available for study.

The active template (AT) approach to interlocked molecules,14-16 removes the need for such templating units in the sub-components of the interlocked molecule. The potential of this methodology for the synthesis of porphyrin-containing architectures was demonstrated by Anderson and co-workers in the synthesis of diyne-linked porphyrin rotaxanes and catenanes using an AT-Glaser¹⁷ methodology.¹⁸ Furthermore, by reducing the size of the macrocycle employed, we have demonstrated that Leigh's active template modification of the CuAAC reaction (AT-CuAAC),15 in which a copper centre bound in the cavity of a bipyridine macrocycle mediates the formation of the triazole, is a general approach to functionalised and functional rotaxanes in excellent yield.19 Thus, although it has yet to be applied in this context,20 the AT-CuAAC reaction appears particularly appropriate for the synthesis of triazole functionalised porphyrin rotaxanes without altering their otherwise desirable properties.

Here we demonstrate the utility of the AT-CuAAC reaction in the synthesis of mechanically interlocked analogues of previously studied triazole-linked porphyrin–corrole conjugates,²¹ and that, because the covalent structure of the chromophores is not altered, photo-induced electron transfer between the tetrapyrrole chromophores is unaffected by mechanical bond formation.²² Conversely, the threaded macrocycles significantly

Street, Wayne, Nebraska, 68787, USA

[&]quot;International Center for Young Scientists (ICYS), WPI Center for Materials Nanoarchitectonics (WPI-MANA), National Institute for Materials Science, Namiki 1-1. Tsukuba. Ibaraki 305-0044. Iapan. E-mail: NGO.Huvnhthien@nims.go.ip

^bWPI Center for Materials Nanoarchitectonics, National Institute for Materials Science, Namiki 1-1, Tsukuba, Ibaraki 305-0044, Japan

^{&#}x27;International Center for Young Scientists (ICYS-SENGEN), National Institute for Materials Science, Sengen 1-2-1, Tsukuba, Ibaraki 305-0047, Japan

Materials Science, Sengen 1-2-1, Tsukuba, Ibaraki 305-0047, Japan

^aDepartment of Chemistry, University of North Texas, 1155 Union Circle, 305070,

Denton, TX 76203, USA. E-mail: Francis.Dsouza@unt.edu

*Department of Physical Sciences and Mathematics, Wayne State College, 111 Main

^fDepartment of Chemistry, University of Southampton, University Road, Highfield, Southampton, SO17 1BJ, UK. E-mail: s.goldup@soton.ac.uk

 $[\]dagger$ Electronic supplementary information (ESI) available: Full synthetic procedures and characterisation of all novel compounds. See DOI: 10.1039/c7sc03165c

modify the steric properties of the system, creating a "mechanical picket fence" motif that suppresses aggregation and ligand driven self-assembly.

Results and discussion

Synthesis and characterisation $4 \subseteq 3$, $5 \subseteq 3_2$ and $6 \subseteq 3_4$

Porphyrin-corrole diad 4⊂3 was synthesised in excellent yield (98% after size exclusion chromatography) by reaction of azide 1, alkyne 2 and macrocycle 3 in the presence of a Cu^I salt (Scheme 1). The extremely high efficiency of the AT-CuAAC reaction allowed this approach to be extended to triad [3]rotaxane $5 \subset 3_2$ and pentad [5]rotaxane $6 \subset 3_4$ in 96% and 70% yield, respectively after size exclusion chromatography, (98% and 91% yield per mechanical bond forming step).

The mass spectrum (MS) of [2]rotaxane dyad 4⊂3 shows a molecular ion at m/z = 1123.5 consistent with $[M + H]^{2+}$. Comparison of the ¹H NMR spectra (Fig. 1) of [2]rotaxane $4 \subseteq 3$ with non-interlocked thread 4 and macrocycle 3 further confirmed the formation of the mechanical bond; although many of the resonances associated with the axle remain unaffected by mechanical bond formation (Hn, Ho, Hp, Hq, and protons associated with Ar¹ and Ar²), which is in keeping with their location away from the threaded region of the axle, triazole proton H_k is shifted considerably to lower field ($\Delta \delta \sim 1.8$ ppm). This is consistent with previous observations of C-H·N hydrogen bonding between the polarised triazole-Hk and the Lewis basic pyridine nitrogen donors¹⁹ and suggests that the macrocycle is largely localised over the triazole unit. Conversely, benzylic protons Hi appear at higher field in the interlocked structure ($\Delta\delta\sim 1.2$ ppm) due to the close proximity of the induced magnetic field of the electron rich aromatic units of the macrocycle.

Resonances assigned to the macrocycle component also exhibit the expected changes on mechanical bond formation





Fig. 1 Partial ¹H NMR (CDCl₃, 300 MHz, 298 K) of (a) dyad axle 4, (b) [2]rotaxane $4 \subset 3$, (c) macrocycle 3, (d) [3]rotaxane $5 \subset 3_2$ and (e) triad axle 5. Peak assignments as shown in Scheme 1. Residual solvent signals are indicated in light grey.

including the dispersion of bipyridine protons HA, HB, HC, shielding of protons H_F, and H_G of the flanking aromatic units and the splitting of H_D and H_E into diastereotopic pairs due to the non-centrosymmetric axle desymmetrising the faces of the macrocycle on mechanical bond formation. Analysis of triad [3]rotaxane $5 \subset 3_2$ and pentad [5]rotaxane $6 \subset 3_4$ by MS also confirmed the presence of the corresponding molecular ions $(m/z = 1614.1 \text{ [M + H]}^{2+} \text{ and } 1870.7 \text{ [M + 3H]}^{3+}, \text{ respectively)}.$



Scheme 1 Synthesis of dyad $4 \subset 3$ and structures of triad [3]rotaxane $5 \subset 3_2$ and pentad [5]rotaxane $6 \subset 3_4$.

Their ¹H NMR spectra (Fig. 1d and 3a) compared with the non-interlocked components display broadly similar changes to that of $4 \subset 3$.

Electronic properties of interlocked corrole-porphyrin conjugates $4 \subseteq 3$, $5 \subseteq 3_2$ and $6 \subseteq 3_4$

Edge Article

Pleasingly, the electronic properties of $4 \subset 3$, $5 \subset 3_2$ and $6 \subset 3_4$ revealed no significant differences compared with the non-interlocked axles. The interlocked and non-interlocked compounds all display electronic absorption bands at ~ 428 , 557 and 598 nm associated with the Zn^{II} -porphyrin unit, and a band at 413 nm accompanied by broad features between 500 and 660 nm assigned to the Cu^{III} -corrole units (Fig. S17†). Furthermore, in all cases the emission associated with the excited singlet state of a $^1Zn^{II}$ -porphyrin* core was efficiently quenched in the interlocked corrole–porphyrin conjugates.

Femtosecond transient absorption spectroscopy of $4 \subseteq 3$, $5 \subseteq 3_2$ and $6 \subseteq 3_4$ confirmed that, as in the case of the corresponding non-interlocked axles,²¹ the quenching of $\mathrm{Zn^{II}}$ -porphyrin luminescence is due to efficient and rapid electron transfer from the $^1\mathrm{Zn^{II}}$ -porphyrin* excited state to the $\mathrm{Cu^{III}}$ -corrole moieties; transient peaks were observed corresponding to the reduced $\mathrm{Cu^{II}}$ -corrole moiety, and a broad peak appeared corresponding to the $\mathrm{Zn^{II}}$ -porphyrin* radical cation.²³ Using the transient signal of the $\mathrm{Cu^{II}}$ -corrole moiety, the rate constant for the charge separation process was evaluated to be $k_{\mathrm{CS}} \sim 10^{11} \ \mathrm{s^{-1}}$ for $4 \subseteq 3$, $5 \subseteq 3_2$ and $6 \subseteq 3_4$ with subsequent charge recombination rates of $k_{\mathrm{CR}} = 2.1 \times 10^{10} \ \mathrm{s^{-1}}$, $3.4 \times 10^9 \ \mathrm{s^{-1}}$, and $3.9 \times 10^9 \ \mathrm{s^{-1}}$ respectively (c.f. $k_{\mathrm{CS}} = 1.1 \times 10^{11} \ \mathrm{s^{-1}}$ and $k_{\mathrm{CR}} = 5.0 \times 10^{10} \ \mathrm{s^{-1}}$ for 4).²¹

These results clearly demonstrate that, as proposed, threading of the macrocycles around the arms of the porphyrin core does not significantly affect the electronic properties of the system.

Effect of threading on the steric properties of pentad $6 \subseteq 3_4$

Although the electronic properties of the interlocked products are unchanged compared with the axle moiety, rotaxanes $4 \subset 3$, $5 \subset 3_2$ and $6 \subset 3_4$ clearly have very different steric properties; encircling the triazole moieties with macrocycle 3 significantly increases the steric demand of the linker units. This difference is particularly striking in the case of pentad $6 \subset 3_4$ which lacks sterically bulky aryl groups on the central porphyrin unit; the space filling model of $6 \subset 3_4$ (Fig. 2a) shows that the Zn^{II}-




Fig. 2 (a) Truncated model (corrole moieties, two arms and axle protons removed for clarity) showing the steric influence of the threaded macrocycles; (b) variation of % $V_{\rm bur}$ of a sphere of radius r centered on Zn with respect to r.

porphyrin unit is significantly encumbered by the threaded macrocycles. This steric hindrance was quantified by determining the % buried volume (% $V_{\rm bur}$) of spheres centered on the central $\rm Zn^{II}$ ion (Fig. 2b).²⁴ At low sphere radii (r < 3 Å), the values of % $V_{\rm bur}$ for 6 and 6 \subset 34 are identical suggesting that the $\rm Zn^{II}$ center is still accessible to small molecules, as required for catalysis or ligand binding. However, as r increases, the values diverge as the threaded macrocycles lead to a higher excluded volume. The comparison between the variation in % $V_{\rm bur}$ of 6 \subset 34 and 6 suggests that although the interlocked macrocycles do not affect the accessible volume immediately around the $\rm Zn^{II}$ center, they provide a steric wall at higher radii similar to covalent picket fence porphyrinoids that have been developed for catalytic and photochemical applications.^{25,26}

An obvious consequence of the difference in steric demand of $6 \subset 3_4$ compared with 6 can be found in their 1H NMR spectra; non-interlocked axle 6 displays resonances that are broadened considerably compared with that of [5]rotaxane $6 \subset 3_4$ under the same experimental conditions (Fig. S31 and S32†). This difference is exacerbated as the concentration of the sample is increased; the signals of rotaxane $6 \subset 3_4$ remain sharp while those of non-interlocked axle 6 broaden and shift. The effect of concentration on the 1H NMR spectrum of non-interlocked pentad 6 is consistent with aggregation of the Zn^{II} -porphyrin unit through π -stacking interactions, as has been widely reported previously. Conversely, in the case of [5]rotaxane $6 \subset 3_4$, the macrocycles encircling the four arms of the porphyrin unsurprisingly appear to prevent the close approach of the Zn^{II} -porphyrin cores.

To further probe the steric effect of the threaded macrocycles we turned our attention to the well-established ability of the ditopic ligand DABCO (L) to direct the formation of [(ZnIIporphyrin)₂L] dimers.^{29,30} The ¹H NMR spectra (Fig. S36†) of noninterlocked pentad 6 displayed behaviour consistent with that previously reported as the quantity of L was varied: (i) at L:6 ratios up to 0.5:1 a signal was observed at -4.92 ppm in the 1 H NMR spectrum, along with a new signal corresponding to H_{n'} which is consistent with the formation of a $[6_2L]$ dimer in slow exchange on the NMR timescale; (ii) once the ratio of L:6 exceeded 0.5:1 the signal at -4.92 ppm disappeared and $H_{n'}$ moved to progressively lower field as further ligand was added, stabilising once L:6 = 1:1, consistent with fast ligand exchange once excess L is present and the formation of monomeric complex [6L] in competition with [62L]. This was further confirmed by cooling the equimolar solution of 6 and L; at 273 K a broad signal was observed at -2.96 ppm alongside the reappearance of the signal at -4.92 ppm, consistent with the monomeric species [6L] in equilibrium with [6₂L] at low temperature. Thus, the speciation of 6 (Fig. 4a) varies as expected with the ratio L: 6. Non-linear regression analysis (see ESI for details†) allowed the association constants for the stepwise association of 6 to ditopic guest DABCO to be determined as $K_1 \ge 5 \times 10^6 \text{ M}^{-1}$ and $K_2 \ge 4 \times 10^7 \text{ M}^{-1}$, albeit with relatively large associated errors of 30% and 27% respectively.31,32

The behavior of [5]rotaxane $6 \subset 3_4$ is significantly different (Fig. S34†). At 298 K progressive addition of L did not lead to the appearance of a signal around -5 ppm corresponding to

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 04 August 2017. Downloaded on 30.10.2025 15:49:30.

Chemical Science




Fig. 3 Partial ^1H NMR (CDCl3, 300 MHz) of (a) pentad [5]rotaxane $6{\subset}3_4$ (298 K); (b) $6{\subset}3_4$ + DABCO (0.4 equiv., 298 K); (c) $6{\subset}3_4$ + DABCO (0.4 equiv., 223 K); (d) $6{\subset}3_4$ + DABCO (1 equiv., 223 K). Peak assignments as shown in Scheme 1. Primed ("") and doubly primed labels refer to signals attributed to [(6 ${\subset}3_4$)L] and [(6 ${\subset}3_4$)₂L] respectively. Cartoon representations have been included to aid clarity but are not intended to be representative of the structures of the complexes formed.²8

 $[(6 \subset 3_4)_2 L]$ but instead the resonances corresponding to triazole proton H_k and porphyrin β -protons H_n underwent monotonic changes that continued until 1 equivalent of L had been added. Alongside these changes, a new broad signal appeared at -2.94 ppm (Fig. 3b) and increased in intensity until 1 equivalent L had been added at which point it disappeared. These changes are consistent with the formation of $[(6 \subset 3_4)L]$ that undergoes slow exchange with free porphyrin $6 \subset 3_4$ on the ¹H NMR timescale, progressing to fast exchange once excess L is present. Consistent with this, cooling an equimolar mixture of $6 \subset 3_4$ and L to 223 K (Fig. 3d) to reduce the rate of ligand exchange resulted in a spectrum consistent with $[(6 \subset 3_4)L]$. Thus, the speciation diagram of $6 \subset 3_4$ with respect to equivalents of L (Fig. 4b) is significantly different to that of 6. Nonlinear regression analysis (see ESI for details†) allowed us to determine K_1 to be $\geq 2 \times 10^5 \text{ M}^{-1}$ at 298 K.

To further examine this unusual observation, we performed a variable temperature 1H NMR study of $6\!\subset\! 3_4$ in the presence of 0.4 equiv. L (Fig. S38†). As expected, reducing the temperature to 273 K led a sharpening of the signal at -2.94 ppm, consistent with $[(6\!\subset\! 3_4)L]$. However, surprisingly, a small signal was also observed at -4.62 ppm. Reducing the temperature further to 248 K led to a number of significant changes consistent with the formation of $[(6\!\subset\! 3_4)_2L]$ alongside $[(6\!\subset\! 3_4)L]$ and unbound $6\!\subset\! 3_4$, including the appearance of resonances at -4.62 and 8.6 ppm attributed respectively to L and the porphyrin β protons H_n of the dimeric complex. Reducing the temperature further to 223 K (Fig. 3c) led to an increase in intensity of signals




Fig. 4 Speciation diagrams for (a) 6 with respect to equiv. L; (b) $6 \subset 3_4$ with respect to equiv. L; (c) L in a $6 \subset 3_4 + 0.4$ L mixture with respect to T; (d) L in a $6 + 6 \subset 3_4 + L$ (1:1:1) mixture with respect to T.

corresponding to $[(6 \subset 3_4)_2 L]$ at the expense of $[(6 \subset 3_4)L]$. Thus, as the temperature is decreased, the mixture of $[(6 \subset 3_4)L]$ and $6 \subset 3_4$ present at 298 K is converted to a mixture of $[(6 \subset 3_4)_2 L]$, $[(6 \subset 3_4)_2 L]$ and $6 \subset 3_4$ until an approximately equimolar mixture of $[(6 \subset 3_4)_2 L]$ and $[(6 \subset 3_4)_2 L]$ is produced at 223 K (Fig. 4c).

That the $[(\mathbf{6} \subset \mathbf{3}_4)_2 \mathbf{L}]$ complex forms at all is remarkable given the sterically hindered environment provided by the macrocycles. The effect of temperature suggests that the balance between negative steric interactions and the positive binding interaction between the \mathbf{Z}^{Π} center and the N donors of the DABCO ligand are finely balanced. Ultimately, the entropic cost of forming the ternary complex, along with the restrictions to conformational freedom associated with forming such a crowded structure, appear finely balanced against the enthalpic benefit of maximising Zn–N interactions, leading to a strongly temperature dependent self-assembly process.

Finally, we examined the speciation of mixtures of \mathbf{L} , $\mathbf{6}$ and $\mathbf{6} \subset \mathbf{3_4}$. In contrast to previous reports in which mixtures of different $\mathrm{Zn^{II}}$ -porphyrins in the presence of \mathbf{L} led to statistical mixtures of dimeric complexes, at low equivalents of \mathbf{L} there is a high selectivity for formation of $[\mathbf{6_2L}]$ (Fig. S38†). Furthermore, this selectivity is maintained in a 1:1:1 mixture of \mathbf{L} , $\mathbf{6}$ and $\mathbf{6} \subset \mathbf{3_4}$ as the temperature is varied; $\mathbf{6}$ is selectively consumed in formation of $[\mathbf{6_2L}]$ in keeping with the higher stability constant for dimerisation of the non-interlocked axle and we did not observe any evidence of hetero-complex formation. Thus, the mechanical picket fence provided by the macrocycles leads to self-sorting in a mixture of $\mathbf{Zn^{II}}$ -porphyrin hosts.

Conclusions

In conclusion, we have demonstrated that the ease and utility of the CuAAC reaction, which has led to its widespread use in the **Edge Article**

design of functional porphyrinoids for various applications,2 is maintained when the active template modification of this reaction is used to produce interlocked analogues. As the covalent structure of the axle is unaffected by mechanical bond formation, the electronic properties of the porphyrin-corrole dyad, triad and pentad reported here are not affected by threading through bipyridine macrocycles, suggesting that the macrocycle provides an alternative, electronically neutral site for structural diversification. Studies comparing the selfassembly behaviour of pentad [5]rotaxane and the corresponding non-interlocked axle component demonstrate that the mechanical bond provides a sterically hindered environment that can modulate intermolecular interactions including π stacking-driven aggregation and ligand-driven dimerisation. This ability to engineer the steric environment around triazolefunctionalised porphyrinoids, an important variable in determining their utility,25 without modifying their covalent structure, suggests that such readily available rotaxanes may play a role in the development of novel types of "picket fence" systems. In the longer term, by combining the steric properties demonstrated here with the well-developed chemistry of rotaxane molecular shuttles,13 it should be possible to extend these results to produce stimuli responsive systems in which the steric environment around the porphyrin core can be modu-

Acknowledgements

lated to produce "smart" materials and catalysts.

THN and JL thank the International Center for Young Scientists (ICYS) for support. This work was partly supported by the World Premier International Research Center Initiative (WPI Initiative), MEXT and JST CREST, Japan. THN thanks the Alexander von Humboldt Foundation and Institut für Chemie und Biochemie, Freie Universität Berlin for the continued support. SMG thanks the University of Southampton, EPSRC (EP/L016621/1) and Royal Society for financial support. JEML is an EU Marie Skłodowska-Curie Fellow, receiving financial support from the European Union's Horizon 2020 research and innovation programme under Marie Sklodowska-Curie grant agreement No 660731. SMG is a Royal Society Research Fellow. SMG acknowledges funding from the European Research Council (Consolidator Grant Agreement no. 724987). The authors are also grateful to the US-National Science Foundation (Grant No. 1401188 to FD) for support of this work.

Notes and references

- (a) C. W. Tornøe, C. Christensen and M. Meldal, *J. Org. Chem.*, 2002, 67, 3057; (b) V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, 41, 2596.
- For a recent review on the synthesis and applications of triazole-functionalised porphyrins see: K. Ladomenou,
 V. Nikolaou, G. Charalambidis and A. G. Coutsolelos, Coord. Chem. Rev., 2016, 306, 1.
- 3 (a) D.-M. Shen, C. Liu and Q.-Y. Chen, *Eur. J. Org. Chem.*, 2007, 1419; (b) M. Severac, L. L. Pleux, A. Scarpaci, E. Blart

- and F. Odobel, *Tetrahedron Lett.*, 2007, **48**, 6518; (c) S. Punidha, J. Sinha, A. Kumar and M. Ravikanth, *J. Org. Chem.*, 2008, **73**, 323; (d) V. S. Shetti and M. Ravikanth, *Eur. J. Org. Chem.*, 2010, 494.
- 4 (a) A. Nierth and M. A. Marletta, *Angew. Chem., Int. Ed.*, 2014, 53, 2611; (b) G. T. Mukosera, T. P. Adams, R. F. Rothbarth, H. Langat, S. Akanda, R. G. Barkley, R. D. Dolewski, J. V. Ruppel and N. L. Snyder, *Tetrahedron Lett.*, 2015, 56, 73.
- 5 (a) M. A. Fazio, O. P. Lee and D. I. Schuster, Org. Lett., 2008, 10, 4979; (b) C. Maeda, P. Kim, S. Cho, J. K. Park, J. M. Lim, D. Kim, J. Vura-Weis, M. R. Wasielewski, H. Shinokubo and A. Osuka, Chem.-Eur. J., 2010, 16, 5052; (c) L. C. Le Pleux, Y. Pellegrin, E. Blart, F. Odobel and A. Harriman, J. Phys. Chem. A, 2011, 115, 5069; (d) S. M. Aly, H. Guernon, B. Guérin and P. D. Harvey, J. Porphyrins Phthalocyanines, 2011, **15**, 871; (e) A. Eggenspiller, A. Takai, M. E. El-Khouly, K. Ohkubo, C. P. Gros, C. Bernhard, C. Goze, F. Denat, J. M. Barbe and S. Fukuzumi, J. Phys. Chem. A, 2012, 116, 3889; (f) A. Krawicz, J. Palazzo, G.-C. Wang and P. H. Dinolfo, RSC Adv., 2012, 2, 7513; (g) D. K. Singh and M. Nath, RSC Adv., 2015, 5, 68209; (h) V. Nikolaou, K. Karikis, Y. Farré, G. Charalambidis, F. Odobel and G. Coutsolelos, Dalton Trans., 2015, 13473; (i) A. Bukreev, K. Mikhailov, I. Shelaev, F. Gostev, Y. Polevaya, V. Tyurin, I. Beletskaya, S. Umansky and V. Nadtochenko, J. Phys. Chem. A, 2016, 120, 1961; (j) W.-D. Quan, A. Pitto-Barry, L. A. Baker, E. Stulz, R. Napier, R. K. O'Reilly and V. G. Stavros, Chem. Commun., 2016, 52, 1938.
- 6 (a) A. R. McDonald, N. Franssen, G. P. M. van Klink and G. van Koten, J. Organomet. Chem., 2009, 694, 2153; (b) J. Nakazawa, B. J. Smith and T. D. P. Stack, J. Am. Chem. Soc., 2012, 134, 2750; (c) T. Chinnusamy, V. Rodionov, F. E. Kühn and O. Reiser, Adv. Synth. Catal., 2012, 354, 1827.
 7 (a) C. Maeda, P. Kim, S. Cho, J. K. Park, J. M. Lim, D. Kim,
- J. Vura-Weis, M. R. Wasielewski, H. Shinokubo and A. Osuka, *Chem.-Eur. J.*, 2010, **16**, 5052; (*b*) D. A. Roberts, T. W. Schmidt, M. J. Crossley and S. Perrier, *Chem.-Eur. J.*, 2013, **19**, 12759; (*c*) D. A. Roberts, M. J. Crossley and S. Perrier, *Polym. Chem.*, 2014, **5**, 4016; (*d*) W.-D. Quan, A. Pitto-Barry, L. A. Baker, E. Stulz, R. Napier, R. K. O'Reilly and V. G. Stavros, *Chem. Commun.*, 2016, **52**, 1938.
- 8 (a) C. H. Lee, S. Lee, H. Yoon and W. D. Jang, Chem.-Eur. J.,
 2011, 17, 13898; (b) L. C. Gilday, N. G. White and P. D. Beer,
 Dalton Trans., 2012, 41, 7092; (c) D. Yim, J. Sung, S. Kim,
 J. Oh, H. Yoon, Y. M. Sung, D. Kim and W.-D. Jang, J. Am.
 Chem. Soc., 2017, 139, 993.
- 9 H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2001, **40**, 2004.
- 10 K. D. Hänni and D. A. Leigh, Chem. Soc. Rev., 2010, 39, 1240.
- 11 (a) D. Tuncel, N. Cindir and Ü. Koldemir, J. Inclusion Phenom. Macrocyclic Chem., 2006, 55, 373; (b) J. A. Faiz, V. Heitz and J.-P. Sauvage, Chem. Soc. Rev., 2009, 38, 422; (c) J.-S. Marois, K. Cantin, A. Desmarais and J.-F. Morin, Org. Lett., 2008, 10, 33; (d) J. D. Megiatto, D. I. Schuster, G. de Miguel, S. Wolfrum and D. M. Guldi, Chem. Mater., 2012, 24, 2472; (e) H. Zhang, Q. Liu, J. Li and D. H. Qu, Org. Lett., 2013, 15, 338.

12 For examples of porphyrin pseudo-rotaxanes see: (a) K. Kano, R. Nishiyabu, T. Asada and Y. Kuroda, *J. Am. Chem. Soc.*, 2002, 124, 9937; (b) K. Kano, R. Nishiyabu, T. Yamazaki and I. Yamazaki, *J. Am. Chem. Soc.*, 2003, 125, 10625; (c) R. Nishiyabu and K. Kano, *Eur. J. Org. Chem.*, 2004, 4985.

Chemical Science

- 13 For recent reviews on the synthesis of interlocked molecules and molecular machines see: (a) M. S. Vickers and P. D. Beer, Chem. Soc. Rev., 2007, 36, 211; (b) J. F. Stoddart, Chem. Soc. Rev., 2009, 38, 1802; (c) E. A. Neal and S. M. Goldup, Chem. Commun., 2014, 50, 5128; (d) S. Erbas-Cakmak, D. A. Leigh, C. T. McTernan and A. L. Nussbaumer, Chem. Rev., 2015, 115, 10081; (e) M. Xue, Y. Yang, X. Chi, X. Yan and F. Huang, Chem. Rev., 2015, 115, 7398; (f) C. J. Bruns and J. F. Stoddart, The Nature of the Mechanical Bond: From Molecules to Machines, Wiley, 2016; (g) J. E. M. Lewis, P. D. Beer, S. J. Loeb and S. M. Goldup, Chem. Soc. Rev., 2017, 46, 2577.
- 14 (a) J. D. Crowley, S. M. Goldup, A.-L. Lee, D. A. Leigh and R. T. McBurney, *Chem. Soc. Rev.*, 2009, 38, 1530; (b) M. Denis and S. M. Goldup, *Nat. Rev. Chem.*, 2017, DOI: 10.1038/s41570-017-0061.
- 15 (a) V. Aucagne, K. D. Hänni, D. A. Leigh, P. J. Lusby and D. B. Walker, *J. Am. Chem. Soc.*, 2006, 128, 2186; (b) V. Aucagne, J. Berna, J. D. Crowley, S. M. Goldup, K. D. Hänni, D. A. Leigh, P. J. Lusby, V. E. Ronaldson, A. M. Z. Slawin, A. Viterisi and D. B. Walker, *J. Am. Chem. Soc.*, 2007, 129, 11950.
- 16 For selected recent examples of the application of the AT-CuAAC reaction see ref. 19 and: (a) B. Lewandowski, G. De Bo, J. W. Ward, M. Papmeyer, S. Kuschel, M. J. Aldegunde, P. M. E. Gramlich, D. Heckmann, S. M. Goldup, D. M. D'Souza, A. E. Fenrandes and D. A. Leigh, Science, 2013, 339, 189; (b) G. De Bo, S. Kuschel, D. A. Leigh, B. Lewandowski, M. Papmeyer and J. W. Ward, J. Am. Chem. Soc., 2014, 136, 5811; (c) A. Noor, S. C. Moratti and J. D. Crowley, Chem. Sci., 2014, 5, 4283; (d) A. Noor, W. K. C. Lo, S. C. Moratti and J. D. Crowley, Chem. Commun., 2014, 50, 7044; (e) M. J. Langton, Y. Xiong and P. D. Beer, Chem.-Eur. J., 2015, 21, 18910; (f) J. Y. C. Lim, I. Marques, A. L. Thompson, K. E. Christensen, V. Félix and P. D. Beer, J. Am. Chem. Soc., 2017, 139, 3122.
- 17 (a) S. Saito, E. Takahashi and K. Nakazono, Org. Lett., 2006, 8, 5133; (b) Y. Sato, R. Yamasaki and S. Saito, Angew. Chem., Int. Ed., 2009, 48, 504; L. D. Movsisyan, D. V. Kondratuk, M. Franz, A. L. Thompson, R. R. Tykwinski and H. L. Anderson, Org. Lett., 2012, 14, 3424. (c) Y. Yamashita, Y. Mutoh, R. Yamasaki, T. Kasama and S. Saito, Chem.—Eur. J., 2014, 21, 2139; (d) R. Hayashi, K. Wakatsuki, R. Yamasaki, Y. Mutoh, T. Kasama and S. Saito, Chem. Commun., 2014, 50, 204; (e) L. D. Movsisyan, M. D. Peeks, G. M. Greetham, M. Towrie, A. L. Thompson, A. W. Parker and H. L. Anderson, J. Am. Chem. Soc., 2014, 136, 17996; (f) R. Hayashi, Y. Mutoh, T. Kasama and S. Saito, J. Org. Chem., 2015, 80, 7536–7546; (g) M. Franz, J. A. Januszewski, D. Wendinger, C. Neiss, L. D. Movsisyan, F. Hampel,

- H. L. Anderson, A. Görling and R. R. Tykwinski, *Angew. Chem., Int. Ed.*, 2015, **54**, 6645; (*h*) L. D. Movsisyan, M. Franz, F. Hampel, A. L. Thompson, R. R. Tykwinski and H. L. Anderson, *J. Am. Chem. Soc.*, 2016, **138**, 1366.
- 18 (a) M. J. Langton, J. D. Matichak, A. L. Thompson and H. L. Anderson, *Chem. Sci.*, 2011, 2, 1897; (b) D. R. Kohn, L. D. Movsisyan, A. L. Thompson and H. L. Anderson, *Org. Lett.*, 2017, 19, 348.
- (a) H. Lahlali, K. Jobe, M. Watkinson and S. M. Goldup, Angew. Chem., Int. Ed., 2011, 50, 4151; (b) J. Winn, A. Pinczewska and S. M. Goldup, J. Am. Chem. Soc., 2013, 135, 13318; (c) R. J. Bordoli and S. M. Goldup, J. Am. Chem. Soc., 2014, 136, 4817; (d) E. A. Neal and S. M. Goldup, Chem. Sci., 2015, 6, 2398; (e) M. Galli, J. E. M. Lewis and S. M. Goldup, Angew. Chem., Int. Ed., 2015, 54, 13545; (f) J. E. M. Lewis, R. J. Bordoli, M. Denis, C. J. Fletcher, M. Galli, E. A. Neal, E. M. Rochette and S. M. Goldup, Chem. Sci., 2016, 7, 3154; (g) E. A. Neal and S. M. Goldup, Angew. Chem., Int. Ed., 2016, 55, 12488; (h) J. E. M. Lewis, J. Winn, L. Cera and S. M. Goldup, J. Am. Chem. Soc., 2016, 138, 16329; (i) J. E. M. Lewis, J. Winn and S. M. Goldup, Molecules, 2017, 22, 89.
- 20 During the preparation of this manuscript an example of an AT-CuAAC reaction leading to a threaded strapped porphyrin rotaxane was disclosed: Y. Miyazaki, C. Kahlfuss, A. Ogawa, T. Matsumoto, J. A. Wytko, K. Oohora, T. Hayashi and J. Weiss, *Chem.-Eur. J.*, 2017, 4, DOI: 10.1002/chem.201702553.
- 21 T. H. Ngo, D. Zieba, W. A. Webre, G. N. Lim, P. A. Karr, S. Kord, S. Jin, K. Ariga, M. Galli, S. Goldup, J. P. Hill and F. D'Souza, *Chem.-Eur. J.*, 2016, **22**, 1301.
- 22 Conceptually this is similar to the stabilisation of dye molecules, conjugated polymers and other reactive species in threaded architectures. For selected examples see ref. 17b, e, g, h, 19b and: (a) J. E. H. Buston, J. R. Young and H. L. Anderson, Chem. Commun., 2000, 905; (b) T. Oku, Y. Furusho and T. Takata, Org. Lett., 2003, 5, 12384; (c) E. J. F. Klotz, T. D. W. Claridge and H. L. Anderson, J. Am. Chem. Soc., 2006, 128, 15374; (d) M. J. Frampton, G. Sforazzini, S. Brovelli, G. Latini, E. Townsend, C. C. Williams, A. Charas, L. Zalewski, N. S. Kaka, M. Sirish, L. J. Parrott, J. S. Wilson, F. Cacialli and H. L. Anderson, Adv. Funct. Mater., 2008, 18, 3367; (e) C. M. S. Yau, S. I. Pascu, S. A. Odom, J. E. Warren, E. J. F. Klotz, M. J. Frampton, C. Williams, V. Coropceanu, M. K. Kuimova, D. Phillips, S. Barlow, J.-L. Brédas, S. R. Marder, V. Millar and H. L. Anderson, Chem. Commun., 2008, 2897; (f) F. E. Oddy, S. Brovelli, M. T. Stone, E. J. F. Klotz, F. Cacialli and H. L. Anderson, J. Mater. Chem., 2009, 19, 2846; (g) J. Terao, S. Tsuda, Y. Tanaka, K. Okoshi, T. Fujihara, Y. Tsuji and N. Kambe, J. Am. Chem. Soc., 2009, 131, 16004; (h) C. B. Caputo, K. Zhu, V. N. Vukotic, S. J. Loeb and D. W. Stephan, Angew. Chem., Int. Ed., 2013, 52, 960; (i) J. Terao, Y. Konoshima, A. Matono, H. Masai, T. Fujihara and Y. Tsuji, Beilstein J. Org. Chem., 2014, 10, 2800.

Edge Article

23 In the case of $5 \subset 3_2$ and $6 \subset 3_4$ a transient signal at 690 nm corresponding to ${}^3\text{Cu}^{\text{III}}$ -corrole was also observed due to direct excitation of the corrole unit. See ESI† for full details.

- 24 Percent buried volume (% $V_{\rm bur}$) is defined as proportion of the volume of a sphere occupied by ligand atoms. It is typically used in the context of catalysis to determine the steric demand of phosphine and N-heterocyclic carbene ligands (A. Poater, B. Cosenza, A. Correa, S. Giudice, F. Ragone, V. Scarano and L. Cavallo, *Eur. J. Inorg. Chem.*, 2009, 1759). Here we use it to compare the steric environments of the interlocked and non-interlocked pentad.
- 25 For selected examples of the applications of picket fence porphyrins see: (a) J. P. Collman, R. R. Gagne, C. Reed, T. R. Halbert, G. Lang and W. T. Robinson, J. Am. Chem. Soc., 1975, 97, 1427; (b) J. Li, B. C. Noll, A. G. Oliver, C. E. Schulz and W. R. Scheidt, J. Am. Chem. Soc., 2013, 135, 15627; (c) L. C. Gilday, N. G. White and P. D. Beer, Dalton Trans., 2013, 42, 15766; (d) B. He, C. E. Schulz and J. Li, Dalton Trans., 2015, 44, 13651; (e) D. Intrieri, D. Carminati and E. Gallo, Dalton Trans., 2016, 45, 15746.
- 26 It should be noted that aggregation due to π - π stacking is observed by 1H NMR at significantly higher concentration (1 to 5 mM) than that at which the absorption and emission spectra are collected (2.5 μ M). Thus, the observed suppression of aggregation and the unchanged electronic properties of $6 \subset 3_4$ compared with 6 are consistent with one another.
- 27 For selected examples of aggregation in porphyrinoid systems see: (a) R. J. Abraham, B. Evans and K. M. Smith, Tetrahedron, 1978, 34, 1213; (b) U. Siggel, U. Bindig, C. Endisch, T. Komatsu, E. Tsuchida, J. Voigt and J.-H. Fuhrhop, Ber. Bunsenges. Phys. Chem., 1996, 2075, 2070; (c) S. Will, J. Lex, E. Vogel, H. Schmickler, J.-P. Gisselbrecht, C. Haubtmann, M. Bernard and M. Gorss, Angew. Chem., Int. Ed., 1997, 36, 357; (d) N. C. Maiti, S. Mazumdar and N. Periasamy, J. Phys. Chem. 1998, 102, 1528; (e) R. van Hameren, J. A. A. W. Elemans, D. Wyrostek, M. Tasior, D. T. Gryko, A. E. Rowan and R. J. M. Nolte, J. Mater. Chem., 2009, 19, 66; (f) A. Jana, H. B. Gobeze, M. Ishida, T. Mori, K. Ariga, J. P. Hill and F. D'Souza, *Dalton Trans.*, 2015, 44, 359; (g) S. Cantekin, A. J. Markvoort, J. A. A. W. Elemans, A. E. Rowan and R. J. M. Nolte, J. Am. Chem. Soc., 2015, 137, 3915.
- 28 In particular, it is reasonable to expect that the corrole arms and macrocycles of the porphyrin sub-units will twist relative to and bend away from one another in dimeric complex

- $6 \subset 3_4$. Simple molecular modelling suggests this is indeed the case (see ESI†).
- 29 For selected examples of DABCO-mediated assembly of ZnIIporphyrin dimers see: (a) C. A. Hunter, M. N. Meah and J. K. M. Sanders, J. Am. Chem. Soc., 1990, 112, 5773; (b) H. L. Anderson, Inorg. Chem., 1994, 33, 972; (c) C. C. Mak, N. Bampos and J. K. M. Sanders, Angew. Chem., Int. Ed., 1998, 37, 3020; (d) P. N. Taylor and H. L. Anderson, J. Am. Chem. Soc., 1999, 121, 11538; (e) C. C. Mak, N. Bampos, S. L. Darling, M. Montalti, L. Prodi and J. K. M. Sanders, J. Org. Chem., 2001, 66, 4476; (f) C. A. Hunter and R. Tregonning, Tetrahedron, 2002, 58, 691; (g) L. Baldini, P. Ballester, A. Casnati, R. M. Gomila, C. A. Hunter, F. Sansone and R. Ungaro, J. Am. Chem. Soc., 2003, 125, 14181; (h) M. C. Lensen, S. J. T. van Dingenen, J. A. A. W. Elemans, H. P. Dijkstra, G. P. M. van Klink, G. van Koten, J. W. Gerritsen, S. Speller, R. J. M. Nolte and A. E. Rowan, Chem. Commun., 2004, 762; (i) P. G. Plieger, A. K. Burrell, S. B. Hall and D. L. Officer, J. Inclusion Phenom., 2005, 53, 143; (j) P. Ballester, A. I. Oliva, A. Costa, P. M. Deyà, A. Frontera, R. M. Gomila and C. A. Hunter, J. Am. Chem. Soc., 2006, 128, 5560; (k) T. Kishida, N. Fujita, O. Hirata and S. Shinkai, Org. Biomol. Chem., 2006, 4, 1902; (1) L. Flamigni, B. Ventura, A. I. Oliva and P. Ballester, Chem.-Eur. J., 2008, 14, 4214.
- 30 Given that non-interlocked axle 6 exhibits the changes by ¹H NMR and simple binding equilibrium expected for a Zn^{II}-porphyrin in the presence of DABCO, any interaction between the Cu^{III}-corrole moiety and DABCO is assumed to be negligible. This is in keeping with the lack, to our knowledge, of any previous report of a DABCO-Cu^{III}-corrole complex.
- 31 The errors in the determination of K_1 and K_2 are relatively large due to the limitations of 1H NMR analysis. Unfortunately, it was not possible to use UV-vis analysis, which has been shown to be more precise in such measurements, to determine the association constants as the strong absorbance of the corrole units mask the expected changes in the porphyrin Soret band. $^{29a-d}$
- 32 With the caveat of the large errors associated with K_1 and K_2 , these values suggest a significant degree of positive cooperativity ($\alpha = 4K_2/K_1 \approx 15$), which is unusual in the DABCO mediated self-assembly of $\mathrm{Zn^{II}}$ -porphyrin units. $^{29a-d}$ The origin of this effect is uncertain and work is ongoing to establish the cause, but, if it is correct, may be due to interactions between the Cu corrole units that stabilise the dimeric complex.