

Showcasing research from Professors Tonner-Zech, Heine, and Asmis, members of the Research Training Group on Hydrogen Isotopes centred at Leipzig University, Saxony, Germany.

Direct evidence for ligand-enhanced activity of Cu(i) sites

In a combined experimental and computational effort, we study the strength and isotopologue selectivity of hydrogen adsorption on the undercoordinated copper (i) site.

Exploiting gas-phase complexes as model systems, we show that the high isotopologue selectivity in dihydrogen binding results from a large difference in the adsorption zero-point energies. We discuss the role of the environment and the coordination geometry of the adsorption site in achieving a high selectivity and the ramifications for identifying and designing future materials for adsorptive dihydrogen isotopologue separation.

### As featured in:



See Jiaye Jin, Ralf Tonner-Zech, Thomas Heine, Knut R. Asmis *et al.*, *Chem. Sci.*, 2024, **15**, 14635.

Cite this: *Chem. Sci.*, 2024, **15**, 14635

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 10th July 2024  
Accepted 9th August 2024

DOI: 10.1039/d4sc04582c  
[rsc.li/chemical-science](http://rsc.li/chemical-science)

## Introduction

Porous materials containing under-coordinated Cu(i) centres with a strong tendency to adsorb dihydrogen are promising candidates for efficient H<sub>2</sub>/D<sub>2</sub> separation.<sup>1–5</sup> A benchmark material with such centres is the metal-organic framework (MOF) Cu(i)-MFU-4*l*, containing trigonal pyramidal Cu<sup>+</sup> sites linked by BTDD (bis-(1*H*-1,2,3-triazolo-[4,5-*b*],[4',5'-*j*])dibenzof[1,4]-dioxin).<sup>2,4,5</sup> Its high D<sub>2</sub>/H<sub>2</sub> adsorption selectivity of 11 at the relatively high temperature of 100 K is due to the combination of high enthalpies of adsorption for H<sub>2</sub> (31 kJ mol<sup>–1</sup>) and D<sub>2</sub> (34 kJ mol<sup>–1</sup>) and the relatively high difference of these values between the two isotopologues. The latter is governed by the zero-point energy (ZPE) difference, and leads to a strong preference towards adsorption of the higher mass isotopologue.<sup>6–8</sup>

The rational design of high-performance separation materials requires a detailed understanding of the binding of

dihydrogen isotopologues to metal centres, particularly Cu(i), which has shown a strong dihydrogen affinity in many cases.<sup>2–4</sup> However, spectroscopic studies of dihydrogen coordinated to Cu(i) centres in bulk materials typically suffer from structural heterogeneity, complicating an in-depth understanding. This can be avoided by performing experiments on well-defined and isolated gas-phase complexes using vibrational action spectroscopy in combination with electronic structure calculations.<sup>9</sup> This approach allows characterization of the geometric structure of relevant M–H<sub>2</sub> motifs,<sup>10</sup> and also provides a deeper understanding of their binding nature as well as the isotopologue selectivity.

The adsorption thermodynamics of H<sub>2</sub> isotopologues at binding sites in porous materials are governed by the ZPE effects associated with the conversion of the three translational and two rotational degrees of freedom of free H<sub>2</sub> into five molecule-adsorption-site vibrations.<sup>11</sup> These low-frequency vibrational modes usually display strongly anharmonic behaviour and pronounced nuclear quantum effects,<sup>12</sup> raising a research interest in model complexes that show the influence of these vibrational modes on dihydrogen isotopologue separation.

A recent spectroscopic and theoretical study of the Cu<sup>+</sup>(H<sub>2</sub>)<sub>4</sub> complex and its isotopologues revealed a substantial red-shift of the dihydrogen stretching frequency ( $\nu_{\text{HH}} = 3729 \text{ cm}^{-1}$  and  $\nu_{\text{DD}} = 2678 \text{ cm}^{-1}$ ) upon complexation ( $\nu_{\text{HH}} = 4162 \text{ cm}^{-1}$  and  $\nu_{\text{DD}} = 2994 \text{ cm}^{-1}$ ).<sup>13</sup> The exceptionally high sequential bond dissociation energy (BDE) for Cu<sup>+</sup>(H<sub>2</sub>)<sub>*n*</sub> → Cu<sup>+</sup>(H<sub>2</sub>)<sub>*n*–1</sub> + H<sub>2</sub> with *n* = 1 of 64 kJ mol<sup>–1</sup> reduces to only 21 kJ mol<sup>–1</sup> with *n* = 4.<sup>13,14</sup> This

<sup>a</sup>Wilhelm-Ostwald-Institut für Physikalische und Theoretische Chemie, Universität Leipzig, Linnéstr. 2, 04103 Leipzig, Germany. E-mail: [jiaye.jin@uni-leipzig.de](mailto:jiaye.jin@uni-leipzig.de); [ralf.tonner@uni-leipzig.de](mailto:ralf.tonner@uni-leipzig.de); [knut.asmis@uni-leipzig.de](mailto:knut.asmis@uni-leipzig.de)

<sup>b</sup>Institute of Resource Ecology, Research Site Leipzig, Helmholtz-Zentrum Dresden-Rossendorf, Permoserstr. 15, 04318 Leipzig, Germany

<sup>c</sup>Faculty of Chemistry and Food Chemistry, School of Science, TU Dresden, 01062 Dresden, Germany. E-mail: [thomas.heine@tu-dresden.de](mailto:thomas.heine@tu-dresden.de)

<sup>d</sup>Department of Chemistry and ibs for Nanomedicine, Yonsei University, Seodaemun-gu, Seoul 120-749, Republic of Korea

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc04582c>

‡ These authors contributed equally.

study motivated the investigation of H<sub>2</sub>-affine Cu(i) complexes with oxygen-donor ligands that can significantly increase the Cu<sup>+</sup>-H<sub>2</sub> BDE.<sup>8,14,15</sup> The aqua complex Cu<sup>+</sup>(H<sub>2</sub>O), arguably the simplest such system, represents a useful model for an under-coordinated adsorption site in a zeolite or a MOF. The experimental BDE of the Cu<sup>+</sup>-H<sub>2</sub> bond in Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>) is 82 kJ mol<sup>-1</sup>,<sup>14</sup> which is 18 kJ mol<sup>-1</sup> larger than that in Cu<sup>+</sup>(H<sub>2</sub>). Such high BDEs are desirable for adsorptive separation of dihydrogen isotopologues, both because they allow higher operating temperatures and also because they typically result in larger ZPE differences, which lead to a higher selectivity.

In this paper, we disentangle why a single H<sub>2</sub>O ligand already markedly strengthens the Cu(i)-H<sub>2</sub> interaction. We first investigate the interaction of dihydrogen and its isotopologues with Cu<sup>+</sup>(H<sub>2</sub>O) by ion-trap mass spectrometry at variable temperature. We then combine cryogenic ion-trap vibrational action spectroscopy<sup>16,17</sup> of Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>)<sub>n</sub> isotopologues with quantum-chemical calculations to characterize their structures and vibrational properties. Finally, we make use of bond analysis methods to rationalize the strong chemical bonding of H<sub>2</sub> to the Cu(i) centre as well as the large ZPE difference of the dihydrogen isotopologues in order to identify the relevant factors that govern the isotopologue selectivity of dihydrogen adsorption.

## Results and analysis

### Ion-trap mass spectrometry

We begin our investigation by characterizing the adsorption yield of H<sub>2</sub> and D<sub>2</sub> on mass-selected Cu<sup>+</sup>(H<sub>2</sub>O) cations confined (on average) for 50 ms in an ion-trap reactor under multiple-collision conditions at ion-trap temperatures ranging from 295 K to 15 K. Under these conditions, Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>/D<sub>2</sub>)<sub>n</sub> complexes with  $n \leq 4 can be formed *via* three-body collisions. Their relative abundances are detected using time-of-flight (TOF) mass spectrometry. Integration over the corresponding normalized TOF-MS peaks yields the relative ion yields (see Fig. S1† for all TOF mass spectra and relative ion yields).$




Fig. 1 Total relative ion yield of Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>)<sub>n>0</sub> and Cu<sup>+</sup>(H<sub>2</sub>O)(D<sub>2</sub>)<sub>n>0</sub> for ion trap temperatures ranging from 15 K to 200 K, obtained from the TOF mass spectra shown in Fig. S1.† The TOF mass spectra were recorded after storing Cu<sup>+</sup>(H<sub>2</sub>O) cations on average 50 ms in a temperature-controlled ion trap filled with roughly 1 mbar of H<sub>2</sub> or D<sub>2</sub>.

Summation over all ion yields with  $n > 0$  represents the total relative yields of (H<sub>2</sub>/D<sub>2</sub>)-containing complexes, which is plotted as a function of ion-trap temperature between 15 K and 200 K in Fig. 1. The formation of dihydrogen complexes is already observed at room temperature, confirming the relatively strong interaction between the Cu(i) cation and dihydrogen.<sup>14</sup> Ion yields larger than ~4% are observed for temperatures of 200 K and below. Dihydrogen adsorption increases with decreasing temperature. Moreover, at sufficiently low temperatures (<80 K) the adsorption of additional dihydrogen molecules is possible leading to the formation of  $n > 2$  complexes (see Fig. S1†). The ion yields in Fig. 1 show that D<sub>2</sub>-containing complexes are more likely formed than the H<sub>2</sub>-containing complexes over the complete temperature range, revealing a substantial isotope effect in dihydrogen adsorption at Cu<sup>+</sup>(H<sub>2</sub>O). This isotopologue selectivity increases with decreasing temperature, reaching a maximum at 120 K. At lower temperatures the contribution of  $n > 2$  complexes and eventually saturation of the dihydrogen binding sites needs to be considered.

### Infrared photodissociation spectra

In order to characterize the nature of H<sub>2</sub>/D<sub>2</sub> adsorption on Cu<sup>+</sup>(H<sub>2</sub>O) in a more detailed way, we measured vibrational action spectra of the complexes using infrared




Fig. 2 Vibrational spectra of Cu<sup>+</sup>(D<sub>2</sub>O)(H<sub>2</sub>)<sub>2</sub> and Cu<sup>+</sup>(H<sub>2</sub>O)(D<sub>2</sub>)<sub>2</sub>. (A) and (B): Experimental IRPD spectra. (C) and (D): Predicted infrared spectra considering anharmonic contributions (VPT2/MP2/def2-TZVPP) as well as a 10 cm<sup>-1</sup> wide Gaussian lineshape function. Bands related to the excitation of the dihydrogen stretching fundamentals  $\nu_{HH}$  (a<sub>1</sub>) and  $\nu_{DD}$  (b<sub>1</sub>), are color-coded in blue and the corresponding combination bands in orange (see Table 1 for the band assignments).



photodissociation (IRPD) spectroscopy. For a comparison of the experimental IPRD spectra to computed IR spectra it proves helpful to measure the IPRD spectra in the linear absorption regime in the range of interest ( $2500\text{--}4500\text{ cm}^{-1}$ ), which covers the excitations of the O–H/O–D and H–H/D–D fundamental transitions. To avoid spectral overlap of the ligand-specific excitations, *e.g.*, O–H/O–D stretching transitions, we focus on the  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)$  complexes here.

The BDE for  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2) \rightarrow \text{Cu}^+(\text{D}_2\text{O}) + \text{H}_2$  is too high to allow for efficient single-photon dissociation. Therefore the IPRD spectra of the  $n = 2$  complexes,  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$ , were measured and these are compared to the corresponding calculated IR spectra in Fig. 2 (see Fig. S2† for the IPRD spectra of  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  and  $\text{Cu}^+(\text{D}_2\text{O})(\text{D}_2)_2$ ). The most prominent IPRD bands (see Table 1 for band positions) are assigned to the partly-resolved rovibrational transitions of the symmetric ( $\nu_{\text{OH/D}}^s$ ) and the antisymmetric O–H/O–D stretching fundamentals ( $\nu_{\text{OH/D}}^{\text{as}}$ ). The corresponding bands in the IPRD spectrum of  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$  are observed at  $3628\text{ cm}^{-1}$  ( $\nu_{\text{OH}}^s$ ) and  $3700\text{ cm}^{-1}$  ( $\nu_{\text{OH}}^{\text{as}}$ ), which is  $4\text{ cm}^{-1}$  higher than the previously reported values for  $\text{Cu}^+(\text{H}_2\text{O})\text{Ar}_2$ .<sup>18</sup> Likewise,  $\nu_{\text{OD}}^s$  and  $\nu_{\text{OD}}^{\text{as}}$  for  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$  are at  $2651\text{ cm}^{-1}$  and  $2751\text{ cm}^{-1}$ , only  $2\text{ cm}^{-1}$  different from the bands for  $\text{Cu}^+(\text{D}_2\text{O})\text{Ar}_2$ . This minor effect of dihydrogen adsorption on the O–H/O–D stretching frequencies in  $\text{Cu}^+(\text{H}_2\text{O}/\text{D}_2\text{O})$  is in line with  $\text{H}_2/\text{D}_2$  binding directly to the  $\text{Cu}^+$  cation.

The bands related to the excitation of H–H stretching mode ( $\nu_{\text{HH}}$ ) are best observed in the IPRD spectrum of  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$ , shown in Fig. 2A, and labelled  $a_{1\text{--}4}$ . The  $\nu_{\text{HH}}$  fundamental ( $a_1$ ) appears at  $3568\text{ cm}^{-1}$ , substantially lower than the  $\nu_{\text{HH}}$  fundamentals at  $3729\text{ cm}^{-1}$  in the previously reported IPRD spectrum of  $\text{Cu}^+(\text{H}_2)_4$ , indicating a stronger  $\text{Cu}^+$ – $\text{H}_2$  interaction in  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$  than in  $\text{Cu}^+(\text{H}_2)_4$ .<sup>13</sup> Features with excitation energies above  $a_1$  are observed at  $3693\text{ cm}^{-1}$  ( $a_2$ ),  $3733\text{ cm}^{-1}$  ( $a_3$ ) and  $4350\text{ cm}^{-1}$  ( $a_4$ ). These are tentatively assigned to combination bands (orange-coloured bands in Fig. 2B). Corresponding features are also observed in the IPRD spectrum of  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$  (Fig. 2B). They are assigned to the  $\nu_{\text{DD}}$  fundamental ( $b_1$ ,  $2567\text{ cm}^{-1}$ ) as well as the combination bands at  $2675\text{ cm}^{-1}$  ( $b_2$ ),  $2783\text{ cm}^{-1}$  ( $b_3$ ),  $3125\text{ cm}^{-1}$  ( $b_4$ ) and  $3414\text{ cm}^{-1}$  ( $b_5$ ).

## Computational results

**Structure and energetics.** MP2/def2-TZVPP minimum-energy structures and BDEs of  $\text{Cu}^+(\text{H}_2\text{O})_{0,1}(\text{H}_2/\text{D}_2)_{0-2}$  are shown in Fig. 3. The global minimum-energy structures of  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_{1,2}$  are of  $\text{C}_{2v}$  symmetry. Both hydrogen molecules bind directly to the  $\text{Cu}^+$  ion. A single  $\text{H}_2$  ligand binds to  $\text{Cu}^+(\text{H}_2\text{O})$  remarkably strongly with a BDE ( $D_0$ ) of  $76.8\text{ kJ mol}^{-1}$ , which is considerably higher than the  $55.5\text{ kJ mol}^{-1}$  found for one  $\text{H}_2$  bound to the bare  $\text{Cu}^+$  ion. This underscores the role of the *trans*-positioned water molecule in increasing the bond

**Table 1** Band positions (in  $\text{cm}^{-1}$ ) observed in the IPRD spectra of  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$  shown in Fig. 1, VPT2/MP2/def2-TZVPP vibrational frequencies and intensities (in  $\text{km mol}^{-1}$  in parenthesis) and band assignments

Label	IRPD	Calculated frequency (intensity)	Assignment <sup>a</sup>
<b><math>\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2</math></b>			
	2651 <sup>b</sup>	2650 (112)	$\nu_{\text{OD}}^s$
	2751 <sup>b</sup>	2754 (165)	$\nu_{\text{OD}}^a$
	2840	2857 (7)	$\nu_{\text{OD}}^{\text{as}} + \delta_{\text{OD}}^{\text{OOP}}$
	3177	3190 (2)	$\nu_{\text{OD}}^{\text{as}} + \delta_{\text{OD}}^{\text{IP}}$
$a_1$	3568	3564 (0.7), 3563 (0.2)	$\nu_{\text{HH}}$
$a_2$	3693	3698 (0.1)	$\nu_{\text{HH}} + \tau_{\text{OD}}^{\text{OOP}}$
$a_3$	3733	3733 (17), 3777 (18)	$\nu_{\text{HH}} + \delta_{\text{CuH}_2}^{\text{IP}}$
	3944	3945 (2), 4005 (0.3)	$\nu_{\text{OD}}^{\text{as}} + \delta_{\text{OD}}^{\text{DOD}}$
$a_4$	4350	4287 (4), 4362 (5), 4316 (2), 4338 (0.7)	$\nu_{\text{HH}} + \nu_{\text{CuH}_2}$
<b><math>\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2</math></b>			
	2567	2566 (0.4), 2568 (0.1)	$\nu_{\text{DD}}$
$b_1$	2675	2692 (0.2), 2695 (0.1)	$\nu_{\text{DD}} + \delta_{\text{CuD}_2}^{\text{IP}}$
$b_2$	2783	2800 (0.2), 2810 (0.1)	$\nu_{\text{DD}} + \delta_{\text{CuD}_2}^{\text{IP}}, \nu_{\text{DD}} + \delta_{\text{D}_2\text{CuD}_2}$
$b_4$	3125	3151 (0.3), 3151 (2), 3158 (1), 3165 (1)	$\nu_{\text{DD}} + \nu_{\text{CuD}_2}$
	3244	3222 (3)	$2 \times \delta_{\text{OD}}$
$b_5$	3414	3441 (0.5), 3442 (0.3), 3450 (0.1)	$\nu_{\text{DD}} + \nu_{\text{DCuD}}$
	3628 <sup>b</sup>	3622 (198)	$\nu_{\text{OH}}^s$
	3700 <sup>b</sup>	3703 (120)	$\nu_{\text{OH}}^a$
	3819	3826 (18)	$\nu_{\text{OH}}^a + \tau_{\text{OH}}^{\text{OOP}}$
	3889, 3900	3860 (4)	$\nu_{\text{OH}}^s + \delta_{\text{CuD}_2}^{\text{OOP}}$
	4207	4170 (0.1)	$\nu_{\text{OH}}^s + \nu_{\text{CuD}_2}^{\text{as}}$
	4283	4250 (3), 4286 (1)	

<sup>a</sup> Labeling of vibrational modes:  $\nu$  (stretching),  $\nu^s$  (symmetric stretching),  $\nu^{\text{as}}$  (antisymmetric stretching),  $\delta$  (bending),  $\delta^{\text{IP}}$  (in-plane bending),  $\delta^{\text{OOP}}$  (out-of-plane bending),  $\tau^{\text{OOP}}$  (out-of-plane hindered rotation, or libration). Only modes with IR intensities larger than  $0.1\text{ km mol}^{-1}$  are listed.

<sup>b</sup> Values are obtained from the fitting of the rovibrational profiles (see Fig. S2).



strength between  $\text{Cu}^+$  and  $\text{H}_2$ . However, only 17.9 kJ mol<sup>-1</sup> is gained when binding a second  $\text{H}_2$ , which is in line with a higher  $\text{Cu}^+ \text{--} \text{H}_2$  distance of 1.64 Å for the two  $\text{H}_2$  molecules in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  (compared to 1.58 Å in the complex with one  $\text{H}_2$ ). We will revisit the effect of  $\text{H}_2$  adsorption on the ion-ligand bond lengths in the bonding analysis section. The addition of a third  $\text{H}_2$  molecule yields only 2.37 kJ mol<sup>-1</sup>.

When compared to the bond length in the free hydrogen molecule, the H-H distance in the  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  complexes is elongated by 5.3 pm and 4.7 pm, respectively (see Table S6†). This activation of the H-H (D-D) bond is substantially larger than the 2.3–3.1 pm reported for  $\text{Cu}(\text{H}_2)_4^+$  and is in line with the much stronger redshift of the H-H (D-D) stretching fundamental observed for the present system.

**Vibrational spectra and band assignments.** In order to assign the bands observed in the IRPD spectra, we calculated IR spectra including anharmonic contributions using vibrational perturbation theory (see computational section for details of the VPT2/MP2/def2-TZVPP calculations). The IR spectra for  $\text{Cu}^+(\text{D}_2\text{O})(\text{H}_2)_2$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$  (shown in Fig. 2C and D) are determined from VPT2 frequencies and intensities (listed in Table 1) using a 10 cm<sup>-1</sup> wide Gaussian lineshape function.

Only the VPT2 IR bands involving excitation of  $\text{H}_2$  or  $\text{D}_2$  stretching modes, either as a fundamental (blue bands) or as part of a combination transition (orange bands), are color-coded. Based on the agreement between the experimental and computed vibrational spectra shown in Fig. 2, we can unambiguously assign all the observed bands (see Table 2). The excitation of the H-H (D-D) stretching fundamental  $a_1$  ( $b_1$ ) is observed at  $\nu_{\text{HH}} = 3568 \text{ cm}^{-1}$  ( $\nu_{\text{DD}} = 2567 \text{ cm}^{-1}$ ), resulting in a frequency ratio  $\nu_{\text{HH}}/\nu_{\text{DD}}$  of 1.390. This ratio is close to the value of 1.388 obtained from the VPT2 IR spectra, confirming the high quality of the MP2/def2-TZVPP potential along the vibrational

**Table 2** EDA-NOCV result for  $\text{Cu}^+$ – $\text{H}_2$  bond of  $\text{Cu}^+(\text{H}_2)_2(\text{H}_2\text{O})$ ,  $\text{Cu}^+(\text{H}_2)(\text{H}_2\text{O})$  and  $\text{Cu}^+(\text{H}_2)_2$ <sup>a</sup>

	$\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2(\text{H}_2)$	$\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$	$\text{Cu}^+(\text{H}_2)_2$
$\Delta E_{\text{int}}$	-71	-112	-95
$\Delta E_{\text{int}}(\text{disp})^b$	-3(4%)	-2(2%)	-2(2%)
$\Delta E_{\text{int}}(\text{elec})^b$	-68(96%)	-110(98%)	-93(98%)
$\Delta E_{\text{Pauli}}$	193	188	189
$\Delta E_{\text{elstat}}^c$	-151(58%)	-172(58%)	-155(55%)
$\Delta E_{\text{orb}}^c$	-110(42%)	-126(42%)	-127(45%)
$\Delta E_1(\sigma(\text{H}_2) \rightarrow \text{Cu}(\text{d}))^d$	-58(53%)	-79(63%)	-69(54%)
$\Delta E_2(\text{Cu}(\text{d}) \rightarrow \sigma^*(\text{H}_2))$	-24(22%)	-19(15%)	-38(30%)
$\Delta E_3(\text{Cu}(\text{d}) \rightarrow \sigma^*(\text{H}_2))$	-26(24%)	-25(20%)	-18(14%)
$\Delta E_{\text{prep}}$	44	5	4
$\Delta E_{\text{prep}}(\text{H}_2)$	4	5	4
$\Delta E_{\text{prep}}(\text{Cu}^+\text{L}_{0-3})$	40	0	0
$\Delta E_{\text{bond}}$	-27	-107	-91
$r(\text{Cu} \text{--} \text{H}_2)^e$	1.72	1.64	1.70


<sup>a</sup> All energies in kJ mol<sup>-1</sup>, bond distances in Å computed with B3LYP-D4/TZ2P. Fragments are closed-shell species  $\text{Cu}^+\text{L}_{0-3}$  and  $\text{H}_2$  ( $\text{L} = \text{H}_2\text{O}$ ,  $\text{H}_2$ ). <sup>b</sup> Percentage values give the relative contributions of dispersion and electronic effects to  $\Delta E_{\text{int}}$ . <sup>c</sup> Percentage values give the relative contributions to the attractive pEDA terms  $\Delta E_{\text{elstat}}$  and  $\Delta E_{\text{orb}}$ .

<sup>d</sup> Percentage values give the relative contributions of the NOCV to  $\Delta E_{\text{orb}}$ . <sup>e</sup> Distance between Cu and the centre of the  $\text{H}_2$  fragment.

mode. The combination bands  $a_2$  to  $a_4$ , observed at 3693 cm<sup>-1</sup>, 3733 cm<sup>-1</sup>, and 4350 cm<sup>-1</sup>, respectively, are assigned to combined excitations of the H-H stretching modes ( $\nu_{\text{HH}}$ ) with either of the following low-frequency vibrational modes: out-of-plane OD hindered rotation ( $\tau_{\text{OD}}^{\text{OOP}}$ ), in-plane  $\text{Cu}^+$ – $\text{H}_2$  bending ( $\delta_{\text{CuH}_2}^{\text{IP}}$ ), and  $\text{Cu}^+$ – $\text{H}_2$  stretching modes ( $\nu_{\text{CuH}_2}$ ). For the latter two modes, the present assignment is similar to the one made previously for the IRPD spectrum of  $\text{Cu}^+(\text{H}_2)_4$ .<sup>13</sup> Analogously, the combination bands  $b_2$  to  $b_5$ , observed at 2675 cm<sup>-1</sup>, 2783 cm<sup>-1</sup>, 3125 cm<sup>-1</sup> and 3414 cm<sup>-1</sup>, respectively, are assigned to combination excitations of one of the two D-D stretching modes ( $\nu_{\text{DD}}$ ) with  $\text{Cu}^+$ – $\text{D}_2$  bending ( $\delta_{\text{CuD}_2}^{\text{IP}}$  and  $\delta_{\text{CuD}_2}^{\text{OOP}}$ ) or the  $\text{Cu}^+$ – $\text{D}_2$  stretching excitations ( $\nu_{\text{DCuD}_2}^s$  and  $\nu_{\text{DCuD}_2}^a$ ).

The remarkably good agreement between the IRPD and VPT2 spectra in Fig. 2, which is discussed in detail in the ESI,† confirms that the chosen computational method provides an adequate description of also the low-frequency  $\text{M}^+$ – $\text{H}_2$ / $\text{D}_2$  vibrational modes, which play an important role in modelling the isotopologue selectivity. However, this appears to be in part due to a favourable error compensation between, *i.e.*, the neglect of substantial relativistic effects, basis set insufficiencies, and the overestimation of London dispersion interaction inherent in the MP2 method (see ESI†).

**Bonding analysis of  $\text{Cu}^+$ – $\text{H}_2$  and  $\text{Cu}^+$ – $\text{H}_2\text{O}$  bonds.** The nature of the  $\text{Cu}^+$ – $\text{H}_2$  and  $\text{Cu}^+$ – $\text{H}_2\text{O}$  interactions in the complexes investigated was analysed with an Energy Decomposition Analysis<sup>19–23</sup> together with the Natural Orbital for Chemical Valence approach (EDA-NOCV)<sup>20–24</sup> method. EDA allows the decomposition of the bonding energy  $\Delta E_{\text{bond}}$  between fragments, which correspond to negative dissociation energy, into chemically meaningful contributions. The bonding energy



**Fig. 3** (A) MP2/def2-TZVPP bond lengths (pm) in  $\text{Cu}^+(\text{H}_2)$ ,  $\text{Cu}^+(\text{H}_2\text{O})$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_1,2$  (colours: Cu orange, O red, H grey). (B) MP2/def2-TZVPP dissociation energies (kJ mol<sup>-1</sup>) with anharmonic (VPT2) ZPE correction. The red values (blue values in italic font) represent the dissociation energies of  $\text{H}_2$  ( $\text{D}_2$ ). Dissociation ZPE differences are highlighted.



$\Delta E_{\text{bond}}$  is represented as a sum of interaction energy  $\Delta E_{\text{int}}$  and preparation energy  $\Delta E_{\text{prep}}$  where the latter describes the deformation of the fragments during the bond formation. The interaction energy  $\Delta E_{\text{int}}$  can be further split into dispersion  $\Delta E_{\text{int}}(\text{disp})$  and electronic effects  $\Delta E_{\text{int}}(\text{elec})$ . The electronic effects can then be divided into quasi-electrostatic contribution  $\Delta E_{\text{elec}}$ , Pauli repulsion  $\Delta E_{\text{Pauli}}$  and orbital interaction  $\Delta E_{\text{orb}}$ . The NOCV extension helps to find the most important orbital interactions to the bond as further decomposition of  $\Delta E_{\text{orb}}$ .

The EDA-NOCV results for the complexes investigated in this work are shown in Table 2. They confirm a donor-acceptor-type interaction in the  $\text{Cu}^+ - \text{H}_2$  bond in all investigated complexes ( $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$ ,  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$ , and  $\text{Cu}^+(\text{H}_2)$ ). At first glance, the stronger bonding in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  compared to  $\text{Cu}^+(\text{H}_2)$  is due to an increase in the electrostatic attraction (difference  $\Delta \Delta E_{\text{elstat}} = -17 \text{ kJ mol}^{-1}$ ) while the orbital interaction ( $\Delta E_{\text{orb}}$ ) and Pauli repulsion ( $\Delta E_{\text{Pauli}}$ ) do not change significantly. However, a closer look reveals a more intriguing trend. The bond in  $\text{Cu}^+(\text{H}_2)$  is significantly longer compared to  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$ . If we analyse the main deformation densities (Fig. 4), we find that upon coordination of  $\text{H}_2\text{O}$ , the  $\sigma$ -donation from  $\sigma(\text{H}_2)$  into the d-orbitals at Cu ( $\Delta E_1$ ) increases by  $10 \text{ kJ mol}^{-1}$ . But the in-plane back-bonding ( $\Delta E_2$ ) decreases by a larger amount ( $-19 \text{ kJ mol}^{-1}$ ), while the out-of-plane back-bonding ( $\Delta E_3$ ) is slightly increased ( $+7 \text{ kJ mol}^{-1}$ ). This reduced in-plane back-bonding is due to the involvement of the Cu(d)-orbitals in the bonding to the  $\text{H}_2\text{O}$  ligand, which can be seen in the deformation densities  $\Delta \rho_1$  and  $\Delta \rho_2$  in Fig. 4. This leads to a lower Pauli repulsion and allows the  $\text{H}_2$  molecule to bind closer to the Cu centre. While orbital attraction and Pauli repulsion thus balance at the same values compared to  $\text{Cu}^+(\text{H}_2)$ , the electrostatic attraction is increased. The main reason is thus lowering of Pauli repulsion leading to a shorter Cu- $\text{H}_2$  distance in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$ .

The weakening of the  $\text{Cu}^+ - \text{H}_2$  bond in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  compared to  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  is attributed mainly to the large preparation energy resulting from the transition from a linear to




Fig. 4 Selected deformation densities ( $\Delta \rho_i$ ) from EDA-NOCVs for  $\text{Cu}^+ - \text{H}_2$  bond with energy contribution ( $\Delta E_i$ ) to  $\Delta E_{\text{orb}}$  in  $\text{kJ mol}^{-1}$  and eigenvalues ( $v_i$ ) Charge depletion (red) and charge accumulation (blue) with isosurface value 0.0025 for  $\text{Cu}(\text{H}_2)^+$ , 0.0020 for  $\text{Cu}(\text{H}_2)(\text{H}_2\text{O})^+$  and 0.0012 for  $\text{Cu}(\text{H}_2)_2(\text{H}_2\text{O})^+$ .

Table 3 Calculated CCSD(T)/aug-cc-pVTZ(-PP) BDEs ( $D_0$ ) including the desorption ZPE ( $\Delta_{\text{des}}\text{ZPE}$ ) calculated using VPT2 on the MP2/def2-TZVPP potential energy surface; anharmonic contribution (VPT2 level) for the dissociation of  $\text{H}_2$  isotopologues, *i.e.*  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2 \rightarrow \text{Cu}^+(\text{H}_2\text{O})(\text{H}_2) + \text{H}_2$ . All values reported in  $\text{kJ mol}^{-1}$

Species	$D_0$	$\Delta_{\text{des}}\text{ZPE}$	Anharmonic contribution
$\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$	16.9	-10.8	0.8
$\text{Cu}^+(\text{H}_2\text{O})(\text{D}_2)_2$	19.7	-8.0	0.4
$\text{Cu}^+(\text{H}_2\text{O})(\text{T}_2)_2$	21.0	-6.7	0.3
$\text{Cu}^+(\text{H}_2\text{O})(\text{HD})_2$	18.4	-9.3	0.6
$\text{Cu}^+(\text{H}_2\text{O})(\text{HT})_2$	19.2	-8.5	0.5
$\text{Cu}^+(\text{H}_2\text{O})(\text{DT})_2$	20.4	-7.3	0.3

a trigonal structure. Additionally, the larger bond lengths are due to the stronger Pauli repulsion between the higher number of ligands and contribute to this phenomenon because they weaken the attractive interactions compared to  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$ .

Note that the  $\text{Cu}^+ - \text{H}_2\text{O}$  bond is stronger in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  compared to  $\text{Cu}^+(\text{H}_2\text{O})$ , which is mainly due to reduced Pauli repulsion as an effect of the back-bonding contributions from  $\text{Cu}^+$  to  $\text{H}_2$  (see NOCV in Fig. S9 in ESI†). Adding another  $\text{H}_2$  molecule, *i.e.* the formation of  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$ , leads to a weaker  $\text{Cu}^+ - \text{H}_2\text{O}$  bond due to competition of the three ligands for the electrons at the central atom.

### Anharmonic contribution to the BDEs for different isotopologues


As shown in Table 3, going from  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  to  $\text{Cu}^+(\text{H}_2\text{O})(\text{T}_2)_2$ , the BDE of the  $\text{H}_2$  isotopologue increases due to the increasing mass of the isotopologues and the consequently lower ZPE. The difference in the adsorption ZPE between the  $\text{D}_2$  and the  $\text{H}_2$  complexes is  $2.8 \text{ kJ mol}^{-1}$  ( $1.0 \text{ kJ mol}^{-1}$  between  $\text{T}_2$  and  $\text{D}_2$ ), which is slightly larger than the one observed in  $\text{Cu}(\text{i})\text{-MFU-4l}$  ( $2.5 \text{ kJ mol}^{-1}$ ) by I. Weinrauch *et al.*<sup>4</sup> and about one-third greater than the ZPE change for the exchange reaction  $\text{Cu}^+(\text{D}_2)_3(\text{H}_2) + \text{D}_2 \rightarrow \text{Cu}^+(\text{D}_2)_4 + \text{H}_2$  ( $2.1 \text{ kJ mol}^{-1}$ ) observed in the previous study conducted by some of us.<sup>13</sup>

Our calculations also show that the anharmonic contribution to the  $\text{H}_2$  desorption ZPE for the  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2 \rightarrow \text{Cu}^+(\text{H}_2\text{O})(\text{H}_2) + \text{H}_2$  reaction is more than three times as big as the value obtained for free  $\text{H}_2$  ( $0.2 \text{ kJ mol}^{-1}$ ). Similar factors are observed for  $\text{D}_2$  (3.1) and  $\text{T}_2$  (3.1). This demonstrates the substantial influence of anharmonic contributions on the thermodynamic equilibrium of the isotopologue exchange reaction, reducing, *e.g.*, the energy difference between  $\text{H}_2$  and  $\text{D}_2$  adsorption by 12%. As expected, the anharmonic contributions to the vibrational frequencies and ZPEs are inversely proportional to the reduced mass of the dihydrogen isotopologue.

### $\text{H}_2$ isotopologue selectivity

As shown in Fig. 5, the predicted isotopologue selectivity of  $\text{H}_2$  adsorption is much higher at  $\text{Cu}^+(\text{H}_2\text{O})$  (B) than at  $\text{Cu}^+$  (A) and decreases again for a second  $\text{H}_2$  adsorbing at  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$ . We





**Fig. 5** Predicted selectivities for dihydrogen isotopologue adsorption at  $\text{Cu}^+$ ,  $\text{Cu}^+(\text{H}_2\text{O})$ , and  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  as a function of temperature. For the latter, two models are compared (see text). Literature data of  $\text{Cu}(\text{i})$ -MFU-4I are shown for comparison.<sup>4</sup>

study two scenarios for the resulting  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  complex: in the first one, we assume that the second  $\text{H}_2$  molecule adsorbs at the free space of a hypothetical  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)$  complex where  $\text{H}_2\text{O}$  and the first  $\text{H}_2$  are already positioned as in  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$ , thus mimicking a scenario as it could occur on an under-coordinated adsorption site in a porous solid (C). Here, only the six degrees of freedom of the considered  $\text{H}_2$  molecule are taken into account to calculate the isotopologue selectivity of dihydrogen adsorption, an approach that has been used in previous studies of MOFs.<sup>4</sup>

In the second one (D), all complexes are fully optimized and the degrees of freedom of all involved  $\text{H}_2$  moieties are considered, which can be compared directly with the IRPD experiments. As a consequence of the weaker binding energy per  $\text{H}_2$  molecule, the ZPE per  $\text{H}_2$  and finally the overall selectivity is reduced significantly.

We conclude that a model with a rigid ligand structure is insufficient when dealing with structurally flexible adsorption sites such as undercoordinated metal ions, as it will lead to spurious results due to ligand reorientation. Conversely, as

evidenced by prior studies,<sup>4,8</sup> rigid adsorption sites with structurally constraining ligands like in the hypothetical starting structure in Fig. 5C may be essential for achieving a high adsorption energy and isotopologue selectivity.

## Summary and conclusions

We elucidated the strong hydrogen isotopologue selectivity of undercoordinated  $\text{Cu}(\text{i})$  sites by examining the model complexes  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_n$  (with  $n = 0, 1, 2$ ) using experiment and theory. Both  $\text{Cu}^+(\text{H}_2\text{O})$  and  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2/\text{D}_2)$  show an obvious isotope effect in the adsorption of  $\text{H}_2$  and  $\text{D}_2$ . The observed vibrational spectra of  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_2$  and its isotopologues in the range from 2500–4500  $\text{cm}^{-1}$  match the calculated frequencies of the fundamentals and related combination bands of  $\text{H}_2/\text{D}_2$  when taking anharmonicity into account. This suggests that these systems are ideal model complexes for gas-phase studies of the chemistry at individual active sites as they occur in framework materials. The stronger  $\text{H}_2$  affinity of  $\text{Cu}(\text{i})$  coordinated by a single oxygen-donor ligand, in comparison to the bare cation, is explained by EDA and traced back to lowering of Pauli repulsion which allows the  $\text{H}_2$  molecule to bind closer to the  $\text{Cu}$  centre. Finally, we used the computational data to predict the dihydrogen isotopologue selectivity of the formation of  $\text{Cu}^+(\text{H}_2\text{O})(\text{H}_2)_n$  complexes, observing a significant selectivity decrease for  $n = 2$  vs.  $n = 1$ . The present study underlines that the structural environment of undercoordinated metal centres in framework materials play a crucial role in the local chemistry and demonstrate particular promise for rigid frameworks with highly exposed metal sites.

## Methods

### Experimental methods

The experiments were performed on the Leipzig 5 K ring-electrode ion-trap triple mass spectrometer described previously.<sup>16</sup>  $\text{Cu}^+(\text{H}_2\text{O})$  cations are transferred to the gas phase from a 5 mmol  $\text{CuSO}_4$  (Sigma Aldrich:  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  99% in  $\text{CH}_3\text{OH}$  99.9% and distilled water) solution using a nanospray ion source under open atmospheric conditions, while the  $\text{Cu}^+(\text{D}_2\text{O})$  cations are produced *via* H/D exchange in moist air conditions saturated with heavy water vapour (99.9%  $\text{D}_2\text{O}$ ). The beam of cations is skimmed, thermalized to room temperature in a Helium-filled radio-frequency ion guide, and then mass-selected in a quadrupole mass filter. Mass-selected  $\text{Cu}^+(\text{H}_2\text{O}/\text{D}_2\text{O})$  ions are continuously trapped in a radio-frequency ring-electrode ion trap, held at a temperature in-between 15 K and 295 K using a closed-cycle helium cryostat and filled with  $\approx 1$  mbar  $\text{H}_2(\text{D}_2)$ .  $\text{Cu}^+(\text{H}_2\text{O}/\text{D}_2\text{O})(\text{H}_2/\text{D}_2)_n$  adducts are formed by three-body collisions and thermalized to the ambient temperature by many collisions with other gas molecules.

All cations are extracted from the ion trap every 100 ms and weakly focused both temporally and spatially into the centre of the extraction region of an orthogonally mounted double-focusing reflectron time-of-flight (TOF) mass spectrometer. For the determination of the ion yields, the ions are extracted and accelerated without using the reflectron. Furthermore, TOF



mass spectra are recorded for TOF extraction delay times ranging from 100  $\mu$ s to 135  $\mu$ s and summed to account for incomplete temporal focusing of the extracted ions. The absolute ion yield  $I_j$  of the  $j$ 'th ion is obtained by integration of the corresponding TOF peak, which is assumed proportional to the number of detected cations of a particular mass-to-charge ratio. Multiple TOF mass spectra are measured and averaged to determine the relative ion yield  $Y_j(\%) = \frac{I_j}{\sum I_i} \times 100\%$ .

IRPD spectra are measured using the IR<sup>1</sup>MS<sup>2</sup> scheme.<sup>25</sup> To this end, the cation packet extracted from the ion trap is accelerated into the reflectron stage and refocused in the initial extraction region. Prior to reacceleration towards the MCP detector, ions with a particular mass-to-charge ratio are selectively irradiated with a properly timed, widely wavelength tunable (750–7000  $\text{cm}^{-1}$ ) IR laser pulse (bandwidth:  $\sim 3.5 \text{ cm}^{-1}$ ), supplied by an optical parametric oscillator/amplifier (LaserVision) laser system.<sup>26</sup> IRPD spectra are recorded by continuously scanning the laser wavelength that is monitored online using a HighFinesse WS6-600 wavelength meter. The scan speed is set to obtain an averaged TOF mass spectrum over 40–80 laser shots every 2  $\text{cm}^{-1}$ . Typically, four scans are measured and averaged to obtain each IRPD spectrum. The photodissociation cross section  $\sigma_{\text{IRPD}}$  determined as described previously.<sup>16,27</sup>

## Computational methods

Structure optimizations and single point energy calculations were performed using coupled cluster theory with single, double, and perturbative triple excitations, *i.e.* CCSD(T), utilizing CFOUR 2.1.<sup>28</sup> The heavily augmented correlation-consistent aug-cc-pVTZ basis set was used for all atoms except for copper, where the version with an effective core potential (ECP), *i.e.* aug-cc-pVTZ-PP, was used to take into consideration relativistic effects. The combination will be referred to as “aug-cc-pVTZ(-PP)” henceforth. Very tight SCF convergence criteria for the Hartree–Fock equations and the coupled cluster amplitudes equations were used ( $10^{-10}$  hartree, *i.e.* SCF\_CONV = 10 and CC\_CONV = 10). The geometry optimizations were performed using analytically evaluated gradients with good starting geometry in Z-matrix files and tighter thresholds. During the geometry optimizations, the RMS gradients were converged to  $10^{-9}$  hartree.

Vibrational analysis (calculation of vibrational frequencies and intensities as well as ZPEs) was performed using standard second-order vibrational perturbation theory (VPT2) without any resonance treatment on a potential energy surface obtained with second-order Møller–Plesset perturbation theory (MP2) in conjunction with the def2-TZVPP basis set using the Gaussian<sup>29</sup> program package. For the preceding geometry optimization, internal coordinates and very tight geometry convergence criteria were used. The frozen-core approximation was applied throughout by specifying five orbitals (DROPMO = 1 > 5), *i.e.* freezing the 1s, 2s, and the three 2p orbitals of the Cu atom. The full details of the computed structures for Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>)<sub>0,1,2</sub> complexes are given in the ESI.<sup>†</sup>

For calculation of dihydrogen isotopologue selectivities, harmonic vibrational frequencies obtained with the Gaussian

program package were used and two approaches are contrasted. One, which was used for all systems considered, follows a previous methodology,<sup>4,7</sup> which relies on the calculation of the partition functions for the individual contributions of each of the six normal modes of H<sub>2</sub> in the field of the frozen adsorption site structure. Here, we used the fragment structure of the fully optimized complex. The harmonic approximation was applied to analyse five of the six modes ( $\nu_{\text{HH}}$ ,  $\nu_{\text{CuH}_2}$ ,  $\delta_{\text{CuH}_2}^{\text{IP}}$ ,  $\delta_{\text{CuH}_2}^{\text{OOP}}$ ,  $\tau_{\text{CuH}_2}^{\text{OOP}}$ ) of the adsorbed hydrogen isotopologues, which are visualized in Fig. S6<sup>†</sup> (see ESI<sup>†</sup>). The sixth mode, *i.e.* the quasi-free rotation of H<sub>2</sub> about the Cu–H<sub>2</sub> axis, was treated using the model of the rigid rotor rotating in the plane. For Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>)<sub>2</sub>, this was contrasted with a second approach, which uses fully relaxed structures and considers the change of the vibrational properties of both H<sub>2</sub> upon binding of the second, *i.e.* considering twelve vibrational modes in Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>)<sub>2</sub>, six in Cu<sup>+</sup>(H<sub>2</sub>O)(H<sub>2</sub>) and the six in free H<sub>2</sub>.

The bonding analysis with the EDA-NOCV analysis, was performed using the Amsterdam Modeling Suite (AMS, version 2021.105),<sup>30</sup> where complexes and their fragments are optimized beforehand. All DFT calculations, were carried out using the B3LYP functional<sup>31</sup> with D4 dispersion correction of Grimme<sup>32</sup> and TZ2P basis set.<sup>33</sup> The selection of the xc-functional is based on the benchmark study presented in ESI.<sup>†</sup> Relativistic effects were considered using the zeroth-order regular approximation (ZORA).<sup>34</sup> Symmetry was not considered. The SCF convergence criterion was set to  $10^{-6} \text{ E}_h$ , indicated by the keyword ‘Very Good’ as a numerical quality indicator. Geometry optimization accounted for the energy criterion at  $3 \times 10^{-3} \text{ E}_h$  and the gradient criterion at  $10^{-3} \text{ E}_h \text{ \AA}^{-1}$ .

## Data availability

The data that support the findings of this study are openly available in zenodo at <https://doi.org/10.5281/zenodo.12554684>.

## Author contributions

EGD and SH contributed equally. SH and JJ performed the experiments. EGD performed the CCSD(T) and MP2 calculations, vibrational analysis and the selectivity calculations. FK carried out the bonding analysis. TH, KA and RTZ conceived the original idea of the project. All authors provided feedback and contributed to the final version of the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG) Project ID 443871192 – GRK 2721: “Hydrogen Isotopes, <sup>1,2,3</sup>H”. EGD and TH thank the Center for Information Services and High Performance Computing (ZIH) at TU Dresden for computational resources as well as Johannes

Hoja and A. Daniel Boese at University of Graz for fruitful scientific discussions and computational resources. JJ and KA thank André Fielicke (Fritz-Haber-Institut der Max-Planck-Gesellschaft) for helpful discussions of low-frequency  $M^+H_2$  modes. SH, JJ and KA thank Alexandra Giermann for the preparation of sample solutions.

## References

- (a) D. Sengupta, P. Melix, S. Bose, J. Duncan, X. Wang, M. R. Mian, K. O. Kirlikovali, F. Joodaki, T. Islamoglu, T. Yildirim, R. Q. Snurr and O. K. Farha, *J. Am. Chem. Soc.*, 2023, **145**, 20492; (b) B. R. Barnett, H. A. Evans, G. M. Su, H. Z. H. Jiang, R. Chakraborty, D. Banyeretse, T. J. Hartman, M. B. Martinez, B. A. Trump and J. D. Tarver, others, *J. Am. Chem. Soc.*, 2021, **143**, 14884.
- R. Xiong, L. Zhang, P. Li, W. Luo, T. Tang, B. Ao, G. Sang, C. Chen, X. Yan, J. Chen and M. Hirscher, *Chem. Eng. J.*, 2020, **391**, 123485.
- B. Ipek, R. A. Pollock, C. M. Brown, D. Uner and R. F. Lobo, *J. Phys. Chem. C*, 2018, **122**, 540.
- I. Weinrauch, I. Savchenko, D. Denysenko, S. M. Souliou, H.-H. Kim, M. Le Tacon, L. L. Daemen, Y. Cheng, A. Mavrandonakis, A. J. Ramirez-Cuesta, D. Volkmer, G. Schütz, M. Hirscher and T. Heine, *Nat. Commun.*, 2017, **8**, 14496.
- J. Teufel, H. Oh, M. Hirscher, M. Wahiduzzaman, L. Zhechkov, A. Kuc, T. Heine, D. Denysenko and D. Volkmer, *Adv. Mater.*, 2013, **25**, 635.
- H. Oh and M. Hirscher, *Eur. J. Inorg. Chem.*, 2016, **2016**, 4278.
- T. Wulf and T. Heine, *Int. J. Quantum Chem.*, 2018, **118**, e25545.
- T. Wulf and T. Heine, *J. Phys. Chem. C*, 2020, **124**, 9409.
- (a) K. R. Asmis, A. Fielicke, G. von Helden and G. Meijer, in *Atomic clusters*, ed. D. Woodruff, Elsevier, Amsterdam and Boston, 2007, pp. 327–375; (b) E. J. Bieske and O. Dopfer, *Chem. Rev.*, 2000, **100**, 3963; (c) J. M. Lisy, *Int. Rev. Phys. Chem.*, 1997, **16**, 267; (d) M. A. Duncan, *Int. Rev. Phys. Chem.*, 2003, **22**, 407; (e) A. Fielicke, *Chem. Soc. Rev.*, 2023, **52**, 3778.
- (a) V. Dryza and E. J. Bieske, *Int. Rev. Phys. Chem.*, 2013, **32**, 559; (b) V. Dryza, B. L. J. Poad and E. J. Bieske, *Phys. Chem. Chem. Phys.*, 2012, **14**, 14954.
- (a) G. Piccini and J. Sauer, *J. Chem. Theory Comput.*, 2014, **10**, 2479; (b) K. Sillar, A. Hofmann and J. Sauer, *J. Am. Chem. Soc.*, 2009, **131**, 4143.
- (a) F. Dahlmann, C. Lochmann, A. N. Marimuthu, M. Lara-Moreno, T. Stoecklin, P. Halvick, M. Raoult, O. Dulieu, R. Wild, S. Schlemmer and others, *J. Chem. Phys.*, 2021, 155; (b) F. Dahlmann, P. Jusko, M. Lara-Moreno, P. Halvick, A. N. Marimuthu, T. Michaelsen, R. Wild, K. Geistlinger, S. Schlemmer, T. Stoecklin, R. Wester and S. Brünken, *Mol. Phys.*, 2022, 120; (c) M. Jia, J. Vanbuel, P. Ferrari, E. M. Fernández, S. Gewinner, W. Schöllkopf, M. T. Nguyen, A. Fielicke and E. Janssens, *J. Phys. Chem. C*, 2018, **122**, 18247; (d) I. Swart, F. M. F. de Groot, B. M. Weckhuysen, P. Gruene, G. Meijer and A. Fielicke, *J. Phys. Chem. A*, 2008, **112**, 1139.
- J. Jin, T. Wulf, M. Jorewitz, T. Heine and K. R. Asmis, *Phys. Chem. Chem. Phys.*, 2023, **25**, 5262.
- P. R. Kemper, P. Weis, M. T. Bowers and P. Maître, *J. Am. Chem. Soc.*, 1998, **120**, 13494.
- (a) P. Maitre and C. W. Bauschlicher, *J. Phys. Chem.*, 1993, **97**, 11912; (b) R. H. Crabtree, *Chem. Rev.*, 2016, **116**, 8750; (c) G. J. Kubas, *Metal Dihydrogen and S-Bond Complexes: Structure, Theory, and Reactivity*, Springer Science & Business Media, 2001.
- N. Heine and K. R. Asmis, *Int. Rev. Phys. Chem.*, 2015, **34**, 1.
- K. R. Asmis, M. Brümmer, C. Kaposta, G. Santambrogio, G. von Helden, G. Meijer, K. Rademann and L. Wöste, *Phys. Chem. Chem. Phys.*, 2002, **4**, 1101.
- P. D. Carnegie, A. B. McCoy and M. A. Duncan, *J. Phys. Chem. A*, 2009, **113**, 4849.
- T. Ziegler and A. Rauk, *Theoret. Chim. Acta*, 1977, **46**, 1.
- K. Kitaura and K. Morokuma, *Int. J. Quantum Chem.*, 1976, **10**, 325.
- F. M. Bickelhaupt and E. J. Baerends, in *Reviews in Computational Chemistry*, ed. K. B. Lipkowitz and D. B. Boyd, Wiley, 2000, vol. 15, pp. 1–86.
- L. Zhao, M. von Hopffgarten, D. M. Andrade and G. Frenking, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2018, **8**, e1345.
- M. P. Mitoraj, A. Michalak and T. Ziegler, *J. Chem. Theory Comput.*, 2009, **5**, 962.
- T. Ziegler and A. Rauk, *Theoret. Chim. Acta*, 1977, **46**, 1.
- M. Mayer and K. R. Asmis, *J. Phys. Chem. A*, 2021, **125**, 2801.
- W. R. Bosenberg and D. R. Guyer, *J. Opt. Soc. Am. B*, 1993, **10**, 1716.
- N. Heine and K. R. Asmis, *Int. Rev. Phys. Chem.*, 2016, **35**, 507.
- J. F. Stanton, J. Gauss, L. Cheng, M. E. Harding, D. A. Matthews and P. G. Szalay, *CFOUR, Coupled-Cluster techniques for Computational Chemistry, a quantum chemical program package*, 2024.
- M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. W. Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16 (Revision C.01)*, Gaussian Inc., Wallingford CT, 2016.
- G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931.



31 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623.

32 E. Caldeweyher, S. Ehlert, A. Hansen, H. Neugebauer, S. Spicher, C. Bannwarth and S. Grimme, *J. Chem. Phys.*, 2019, **150**, 154122.

33 E. van Lenthe and E. J. Baerends, *J. Comput. Chem.*, 2003, **24**, 1142.

34 E. van Lenthe, A. Ehlers and E.-J. Baerends, *J. Chem. Phys.*, 1999, **110**, 8943.

